


MAIN

Looking on the bright side and seeing it vividly: interpretation bias and involuntary mental imagery are related to risk for bipolar disorder

Andrew D. Peckham^{1,*} , Kiana Modavi² and Sheri L. Johnson²

¹Department of Psychiatry, McLean Hospital and Harvard Medical School, 115 Mill Street, Mail Stop 113, Belmont, MA 02478, USA and ²Department of Psychology, University of California, Berkeley, 2121 Berkeley Way, Berkeley, CA 94720, USA

*Corresponding author. Email: adpeckham@mclean.harvard.edu

(Received 6 March 2019; revised 5 August 2019; accepted 15 August 2019; first published online 10 October 2019)

Abstract

Background: Involuntary mental imagery is elevated among people with bipolar disorder, and has been shown to shape biases in interpretation of ambiguous information. However, it is not clear whether biases in interpretation of ambiguous scenarios can be observed in those at risk for bipolar disorder, or whether involuntary imagery is related to such a bias.

Aims: In the present study, we extended a prominent model of bipolar cognition to an at-risk sample. We specifically tested whether positive interpretation bias and involuntary mental imagery are linked to a greater risk of bipolar disorder.

Method: Young adults ($N = 169$) completed measures of risk for bipolar disorder (the Hypomanic Personality Scale [HPS]), interpretation bias, and involuntary mental imagery.

Results: Higher scores on the HPS were significantly correlated with more positive interpretations of ambiguous scenarios ($\beta = 0.29$, $p < .01$) and more frequent involuntary mental imagery ($\beta = 0.22$, $p < .01$). There was no evidence of an interaction between interpretation bias and mental imagery in predicting HPS score, $\beta = .04$, $p = .62$.

Conclusions: Further research is warranted to determine if intrusive imagery or interpretation bias influence the development of bipolar disorder over time in those at risk.

Keywords: hypomania; interpretation biases; mental imagery; non-clinical

Introduction

According to cognitive models of psychopathology, emotional disorders are maintained by biases and distortions in processing information (Beck, 1976). One such bias is *interpretation bias*, in which ambiguous stimuli are interpreted as having an emotionally valenced quality – for example, people who are depressed may interpret neutral information as negative, while those with anxiety disorders may interpret neutral information as threatening (Hirsch *et al.*, 2016; Mathews and MacLeod, 2005). These findings have emerged in the context of many different types of ambiguous information, including facial expressions, written scenarios and interoceptive sensations (Hirsch *et al.*, 2016; Schoth and Lioffi, 2017). As research on this basic mechanism has flourished, evidence has subsequently emerged that directly modifying these biases via computerized tasks may be a viable intervention strategy for anxiety and depression (Holmes *et al.*, 2009; MacLeod and Mathews, 2012).

Compared with those diagnosed with anxiety or depressive disorders, much less is known about the degree to which bipolar disorder is characterized by interpretation bias. However, a prominent model of cognition in bipolar disorder posits that interpretation biases play a key

role in the development of mood symptoms. Holmes and colleagues argue that cognitive biases in interpretation are shaped and amplified by tendencies to experience more vivid and intrusive mental images (Holmes *et al.*, 2008). This theory includes the prediction that a combination of positive interpretation bias and vivid mental imagery of imagined positive events could amplify symptoms of mania. In the present study, we tested one component of this model among individuals at risk for bipolar disorder. We reasoned that if positive bias and vivid, positive imagery contribute to mania symptoms, similar cognitive factors involving aspects of positive bias and imagery might also confer higher vulnerability to bipolar disorder. As backdrop to this study, we first review two converging topics: interpretation bias in bipolar disorder and mental imagery in bipolar disorder.

Interpretation bias and bipolar disorder

Among individuals diagnosed with bipolar disorder, investigations of cognitive biases have yielded somewhat inconsistent results. At a broad level, some studies indicate that mania, the defining feature of bipolar disorder, is tied to positive cognitive biases (García-Blanco *et al.*, 2013; Murphy *et al.*, 1999), and there is evidence that people with bipolar disorder show overly positive appraisals of internal states (Dodd *et al.*, 2011; Jones *et al.*, 2006; Kelly *et al.*, 2011; Mansell, 2006). In contrast, some studies find no evidence of positive biases (e.g. Peckham *et al.*, 2016; Purcell *et al.*, 2018), while others find that some with bipolar disorder show negative biases and negative appraisals of internal states (e.g. Lyon *et al.*, 1999; Mansell *et al.*, 2007). However, very few studies have tested interpretation biases using stimuli such as ambiguous scenarios or words, methods that are commonly used to identify biases in mood and anxiety disorders (cf. Hirsch *et al.*, 2016). The few studies that have used such measures do not show evidence for positive bias during remission or depressive episodes (Di Simplicio *et al.*, 2016; Holmes *et al.*, 2011).

In comparison with diagnosed samples, more consistent evidence supports positive interpretation bias in people at risk for bipolar disorder. Evidence shows that positive cognitive biases are present in at-risk samples for various types of ambiguous stimuli, including ambiguous physical touch (Piff *et al.*, 2012), positive perceptions in dyadic couples' interactions (Dutra *et al.*, 2014), and positive appraisals of facial affect (Campellone *et al.*, 2018; Trevisani *et al.*, 2008). Similar to diagnosed samples, people at risk for bipolar disorder also report more positive interpretations of ambiguous internal states (Johnson and Jones, 2009; Jones *et al.*, 2006; Kelly *et al.*, 2012).

Although these findings indicate that people at risk for bipolar disorder may show a positive bias to some ambiguous stimuli, most of these studies have used tasks that measure interpretation using ambiguous faces or assessment of internal states. Interpretation bias paradigms that rely on facial stimuli may be problematic because some with bipolar disorder show difficulties with facial affect recognition of non-ambiguous stimuli (Kohler *et al.*, 2011). Some have argued that ambiguous scenarios are a more ecologically valid approach to assessing interpretations of ambiguity (Schoth and Liossi, 2017); accordingly, scenarios have been the more common approach to assessing interpretation bias in other disorders.

To our knowledge, ambiguous scenarios have not been used to probe interpretation bias in people at risk for bipolar disorder, and have only been used in one instance among those diagnosed with bipolar disorder (Di Simplicio *et al.*, 2016). To best assess the contribution of imagery and interpretation bias to risk for bipolar disorder, one goal of the present study was to assess interpretation bias among those at risk for bipolar disorder using a validated measure of interpretation of ambiguous scenarios.

Imagery and bipolar disorder

Empirically, diagnoses and risk of bipolar disorder are associated with elevated tendencies to experience mental images as vivid (Gregory *et al.*, 2010; Holmes *et al.*, 2011), intrusive (e.g. involuntary), and future-oriented (Deeprouse *et al.*, 2011; Di Simplicio *et al.*, 2016; Hales *et al.*, 2011; Holmes *et al.*, 2011). Among those diagnosed with bipolar disorder, involuntary imagery in particular is linked to mood instability (Holmes *et al.*, 2011). Basic findings outside of bipolar disorder reveal that imagery has been shown to increase tendencies toward interpretation biases. That is, participants prompted to form images of positive stimuli showed more positive interpretations of ambiguous words on a transfer task (Pictet *et al.*, 2011), and participants trained to show negative interpretations of ambiguous stimuli report more negative mental imagery (Hertel *et al.*, 2003). Given the prominent role for involuntary imagery in bipolar disorder, and increasing evidence for a prominent role for involuntary imagery in psychopathology more broadly (Brewin *et al.*, 2010), understanding how interpretations biases and imagery jointly contribute to symptoms and mood episodes may have important clinical implications.

Little is known, however, about how individual differences in imagery guide cognitive interpretation biases within bipolar disorder. In one study, those with bipolar disorder rated ambiguous scenarios *less* positively than those without the disorder, but among those with bipolar disorder, those who reported more vivid images of a scenario showed more positive bias for that scenario (Holmes *et al.*, 2011). In addition, imagery has been linked to memory bias for positive, high-activation words among people at risk for bipolar disorder (Pyle and Mansell, 2010). However, no study to date has evaluated whether positive interpretation bias and involuntary imagery jointly influence greater risk for bipolar disorder. Extending a test of Holmes' and colleagues (2008) hypothesis to an at-risk sample is a critical step to understanding if positive interpretation bias and mental imagery can help to explain vulnerability for bipolar disorder; this was the primary goal for the present study.

Aims and hypotheses

In the present study, we tested the degree to which interpretation bias and involuntary mental imagery are separately or jointly related to higher scores on a self-report measure previously linked to risk for bipolar disorder. In a non-clinical sample, we administered a battery of self-report measures to assess risk for mania, interpretation of ambiguous scenarios, and involuntary imagery about future self-relevant events. Given the inherent difficulty in isolating the role of imagery in measures of ambiguous scenarios, we chose a measure of interpretation bias (described below) that explicitly asks participants to rate both their interpretation and the vividness of the scenarios, so that the effects of image vividness could be accounted for in analyses of interpretation bias. Given evidence for some overlap between measures of positive affect and hypomania risk (Kirkland *et al.*, 2015), we also included a separate measure of positive affect to better isolate effects of hypomania risk on positive bias. We hypothesized that risk for mania would be related to more positive interpretations of ambiguous situations and more frequent involuntary mental imagery, and to the interaction of involuntary imagery and positive interpretations of ambiguity, such that high levels of both would be particularly related to increased mania risk.

Method

Participants and design

Participants ($N = 206$) were undergraduate college students who completed written informed consent, and then online demographic items and the self-report measures described below.

Table 1. Descriptive statistics for primary study variables

Demographic and clinical characteristics	<i>n</i>	(%) ¹	Mean	(SD)
Gender (% female)	133	(80%)		
Race				
African-American/Black	3	(2%)		
Caucasian	49	(30%)		
Asian	90	(55%)		
Native American	1	(1%)		
More than one race	21	(12%)		
Ethnicity				
Not Hispanic or Latino/a	138	(84%)		
Hispanic or Latino/a	27	(16%)		
Age			21.09	(3.34)
HPS score			19.99	(5.45)
AST-Pleasantness			5.31	(0.95)
AST-Vividness			4.77	(0.89)
IDDD total symptoms			1.60	(2.18)
IFES total score			1.52	(0.57)
IFES number of negative future events			1.04	(0.91)
PANAS-PA			24.59	(8.67)
PANAS-NA			16.74	(7.73)

AST, Ambiguous Scenarios Test; IFES, Impact of Future Events Scale; PANAS, Positive and Negative Affect Scale; RPA, Responses to Positive Affect Scale.

¹Percentage of valid data not including missing cases.

Participants received partial course credit in psychology classes for taking part in this study. Measures were delivered through Qualtrics (Qualtrics, Provo, UT, USA). Some participants (17.96%; $n = 37$) either failed to respond correctly to a validity item ('Please respond 3 to this question') or failed to complete one or more of the primary study measures; these participants were excluded, leaving a final sample of 169 (78.70% female, mean age = 21.09 years, $SD = 3.34$, range: 18–50). Race and ethnicity are described in Table 1.

Measures

Ambiguous Scenarios Test-Depression (AST-D; Berna et al., 2011)

The AST-D is a self-report measure of interpretation bias. Participants read 24 ambiguous scenarios (e.g. 'You go to a place you visited as a child. Walking around makes you feel emotional'), were asked to form an image of that scenario happening to them, and then rate how pleasant and vivid the image is. Pleasantness ratings, made on a 9-point Likert scale from 'extremely unpleasant' to 'extremely pleasant', are considered the index of interpretation bias. Vividness is rated on a 7-point Likert scale from 'not vivid at all' to 'extremely vivid'. Although initially designed to capture potential depression-related biases (Berna *et al.*, 2011), the AST has shown good validity in unselected college samples such as the present study (e.g. Cooper and Wade, 2015). Internal consistency was high for both subscales across the 24 scenarios (pleasantness $\alpha = 0.85$, vividness $\alpha = 0.93$).

Hypomanic Personality Scale (HPS; Eckblad and Chapman, 1986)

The HPS is a measure of lifetime experiences of mild hypomanic symptoms and of personality traits designed to identify those 'predisposed to both hypomanic episodes and bipolar disorder' (Eckblad and Chapman, 1986). The measure includes 48 items to assess risk for mania (e.g. 'I frequently find that my thoughts are racing'); in the present study, these items were rated on a 0 (*strongly disagree*) to 3 (*strongly agree*) scale validated in previous work (e.g. Giovanelli *et al.*, 2013). Participants are instructed to include only experiences that occurred when they

were not under the influence of drugs. Across two longitudinal studies, higher HPS scores related concurrently and predictively to bipolar disorder and bipolar spectrum disorder diagnosis (Kwapil *et al.*, 2000; Walsh *et al.*, 2015). HPS scores are also linked to genetic polymorphisms related to bipolar disorder (Johnson *et al.*, 2014a). We found the scale to have excellent internal consistency in our sample ($\alpha = 0.91$). HPS scores were normally distributed; however, all but one participant in the study scored below the 'high-risk' range of 35 or higher used by previous studies (e.g. Johnson *et al.*, 2014a).

Impact of Future Events Scale (IFES; Deeproose and Holmes, 2010)

The IFES is a measure of the degree to which a person experiences intrusive (involuntary), future-oriented mental imagery (Deeproose and Holmes, 2010). Participants are asked to identify three future events that they have been thinking about over the past 7 days, and to indicate whether each future event is positive or negative. They then answer 24 questions about the extent to which they experienced future-oriented imagery of these events over the past week (e.g. 'Pictures about the future popped into my mind') on a 0 (*not at all*) to 4 (*extremely*) scale. Internal consistency for this sample was good ($\alpha = 0.88$). The IFES yields two scores: the valence of the three future events (i.e. how many of the three events were negative or positive), and the average level of involuntary future-oriented imagery from the 24-item scale. As in past research using the IFES, we focused on the average of the 24-item scale in our analysis, but controlled for the valence of negative and positive events. Higher IFES scores have been shown to relate to risk for bipolar disorder (Deeproose *et al.*, 2011) and to bipolar disorder diagnoses (Holmes *et al.*, 2011).

Inventory to Diagnose Depression (IDD)

Participants completed the IDD, a measure of current depression symptoms. Participants were asked to indicate the severity of symptoms of depression in the past week. For each of 22 symptoms, they select one of five possible responses describing the severity of the symptom (e.g. 'My energy level was normal' to 'I felt tired or exhausted almost all of the time'). For each symptom endorsed, they were asked to indicate whether the symptom persisted at least 2 weeks. IDD scores cover the DSM symptoms of major depressive disorder (range: 0 to 9) and are significantly associated with diagnosis of major depressive disorder (Zimmerman and Coryell, 1987). Internal consistency for this sample was excellent ($\alpha = 0.93$).

Positive and Negative Affect Schedule - 24 Item (PANAS-24)

As a measure of current mood, participants were asked to rate their current feelings using 12 positive (PA) and 12 negative affective (NA) state adjectives selected from the PANAS-X (12 negative and 12 positive; from Watson and Clark, 1999). Each item was rated on a Likert scale from 1 (*very slightly or not at all*) to 5 (*extremely*). Internal consistency for our sample was excellent (PA $\alpha = 0.93$, NA $\alpha = 0.93$).

Analysis plan

Analyses were completed using IBM SPSS version 24.0 and Jamovi version 0.9. Before conducting tests of hypotheses, variables were graphed and their distributions checked for normality. Bivariate relationships between study variables were examined using Pearson correlations. To consider the hypothesis that interpretation bias, mental imagery, and their interaction were related to mania risk (HPS), we used a hierarchical linear regression. Independent variables were centred prior to analyses. Potential multivariate outliers were identified via evaluation of Mahalanobis distances. Additional regression diagnostics [tolerance and variance inflation factor (VIF) values] were evaluated to assess for the potential of multicollinearity.

Table 2. Pearson correlations between primary study variable

	HPS	AST Pleasantness	AST Vividness	IFES total score	PANAS Positive Affect	PANAS Negative Affect	IFES Negative Events	IDD total symptoms
HPS	—	0.27 ***	0.05	0.23 **	0.15 *	0.01	-0.26 ***	0.09
AST Pleasantness		—	0.41 ***	-0.10	0.26 ***	-0.25 **	-0.32 ***	-0.23 **
AST Vividness			—	0.18 *	0.29 ***	0.06	-0.15	0.06
IFES total score				—	0.04	0.25 **	-0.01	0.40 ***
PANAS Positive Affect					—	0.29 ***	-0.15	-0.09
PANAS Negative Affect						—	-0.03	0.29 ***
IFES negative events							—	0.17 *
IDD total symptoms								—

* $p < .05$, ** $p < .01$, *** $p < .001$.

AST, Ambiguous Scenarios Test; IFES, Impact of Future Events Scale; PANAS, Positive and Negative Affect Scale; RPA, Responses to Positive Affect Scale.




Figure 1. Scatter plot of Interpretation Bias and Hypomanic Personality Scale (HPS). AST Pleasantness ratings on the y-axis refer to average ratings of the pleasantness of the 24 scenarios in the Ambiguous Scenarios Test. The shaded line shows the standard error of the mean.

Results

Descriptive statistics for the sample are shown in Table 1. Table 2 shows zero-order correlations between variables of interest. As would be expected, the two scales of the AST (Pleasantness and Vividness) were significantly correlated. HPS scores showed small, significant correlations with more positive interpretations of ambiguous scenarios (Fig. 1) and involuntary future-oriented mental imagery. Correlations of age and sex, and ANOVAs of race and ethnicity showed no significant relationship to HPS score ($p > .25$). Current depression symptoms were significantly correlated with more frequent involuntary imagery of future events.

Are interpretation bias, intrusive imagery, and their interaction related to risk for bipolar disorder?

Hierarchical multiple linear regression was used to test whether more positive interpretation of ambiguous scenarios (AST-pleasantness score), more frequent involuntary imagery of future events (IFES score), and their interaction would relate to higher scores on the HPS measure. An initial regression model revealed two potential multivariate outliers (Mahalanobis distances exceeding a threshold of $\chi^2(10) = 29.59, p < .001$); these two participants were removed from analysis.¹ AST-pleasantness score and involuntary imagery (IFES total score) were entered in block 1, as well as gender and age. The interaction term of Involuntary imagery \times Interpretation bias was entered in block 2. State NA and PA (PANAS), depression symptoms (IDD), vividness ratings (AST) and overall number of negative future events (IFES) were entered as control variables in block 3, to evaluate if any observed relationships between HPS score, interpretation, and involuntary imagery remained significant after accounting for these additional variables. The overall model was significant; $F(4,158) = 4.98, p < .001$, adjusted $R^2 = 0.09$. HPS scores were unrelated to gender or age. However, both involuntary mental imagery ($\beta = 0.25, p < .01$) and more positive interpretation bias ($\beta = 0.25, p < .01$) each related significantly to higher HPS scores in block 1 (R^2 change = .11). The interaction term entered in block 2 (AST-pleasantness \times IFES total score) did not significantly improve model fit: $\beta = .05; R^2$ change = .002, $F(1,157) = .44, p = .51$. In the final model [$F(10,152) = 4.08, p < .001$, adjusted $R^2 = 0.16$], HPS scores were related to fewer future negative events on the IFES ($\beta = -0.20, p = .02$) and to lower vividness ratings on the AST ($\beta = -0.24, p < .01$). (The effect for vividness ratings was in the opposing direction to that shown in bivariate correlations.) HPS scores were not significantly related to positive or negative affect (PANAS), nor to current depressive symptoms ($p > .07$). However, both positive interpretation bias ($\beta = .29, p < .01$) and involuntary imagery ($\beta = .22, p < .01$) remained significantly related to HPS score in the presence of these additional control variables. The interaction term (AST-pleasantness \times IFES total score) was not significant in this model; $\beta = .04, p = .62$. Inspection of regression diagnostics in this final model did not suggest the presence of multicollinearity (tolerance exceeded .58, with VIF less than 1.72 for all variables).

Discussion

Overly positive interpretation of ambiguous stimuli and involuntary mental imagery are proposed contributors to symptoms of mania (Holmes *et al.*, 2011), yet little research has considered whether these two factors jointly contribute to risk for bipolar disorder. In the present study, both a more positive interpretation of ambiguous scenarios and enhanced involuntary mental imagery were associated with higher scores on the HPS measure, which has been shown to be associated with and to predict diagnosis of bipolar disorder (Kwapil *et al.*, 2000; Walsh *et al.*, 2015). These two dimensions did not interact. Strengths of this study include use of well-validated measures of imagery and interpretation bias, a large sample size with sufficient variability in HPS scores, and careful consideration of potential confounds (depression symptoms and mood state).

Previous findings link risk for bipolar disorder with more positive responses to other types of ambiguous stimuli, such as faces, internal sensations, and touch (Campellone *et al.*, 2018; Dutra *et al.*, 2014; Jones *et al.*, 2006; Piff *et al.*, 2012; Trevisani *et al.*, 2008). The present study extends these findings by showing that scores on the Hypomanic Personality Scale, a well-validated scale associated with risk for bipolar disorder, were significantly correlated with more positive interpretations of ambiguous scenarios. Although the magnitude of this effect was modest, the

¹Removal of these two multivariate outliers did not influence the significance or direction of any results.

relationship between HPS scores and interpretation bias remained even when concurrently controlling for state affect, depression symptoms, and vividness of mental imagery. Across the present study and others in the field, consistent evidence is emerging that higher HPS scores are significantly associated with tendencies to perceive greater positivity in ambiguous situations. This pattern complements findings that subclinical depression and anxiety symptoms are tied to a more negative interpretation bias (e.g. Berna *et al.*, 2011; Hindash and Amir, 2012; Steinman and Teachman, 2010).

These findings are important to consider in the context of previous research on positivity-related traits in bipolar disorder. Consistent with the present findings, elevations in personality traits related to positive affect are also observed among unaffected first-degree relatives of those with bipolar disorder (Higier *et al.*, 2014), indicating that elevated trait-like positivity may be related to aspects of vulnerability for the disorder. However, caution is warranted in assessing the results of undiagnosed samples, particularly given the partial overlap between measures of mania risk (such as the HPS) with positivity-relevant measures of extraversion and positive affect (e.g. Kirkland *et al.*, 2015). It is also important to consider that the present study evaluated only two aspects of positivity. Other research demonstrates the importance of considering related constructs such as reactivity to positive stimuli (Gruber *et al.*, 2011a), impulsivity during positive mood (Muhtadie *et al.*, 2014), and regulation of positive affect (Feldman *et al.*, 2008). Finally, in the present study, higher scores on the HPS measure were modestly correlated with fewer negative future events listed on the IFES measure, potentially consistent with previous research indicating that higher scores on measures related to bipolar disorder are linked to optimism, predictions of success, and higher goal-setting for future rewarding events (Johnson and Carver, 2006; Schönfelder *et al.*, 2017). Future studies would do well to conjointly consider how different aspects of positivity relate to risk for bipolar disorder, and from the use of prospective designs to understand whether these traits contribute to onset of the disorder.

Attempts to understand the nature of positivity-related traits in bipolar disorder must also contend with significant heterogeneity in outcomes and presentation. This heterogeneity is evident in the literature on positivity-related traits. Low levels of positive affect predict lower quality of life in people with bipolar disorder (Johnson *et al.*, 2016), yet other aspects of positivity are linked to negative outcomes. For example, heightened rumination on positive affect is related to more frequent episodes of mania (Gruber *et al.*, 2011b). Another source of heterogeneity stems from the observations that many (but not all) individuals with bipolar disorder experience episodes of depression (Cuellar *et al.*, 2005). In the present study, depression symptoms showed the inverse relationship to interpretation bias compared with the HPS measure: that is, higher levels of depression were correlated with a more negative interpretation bias, consistent with previous research (Hindash and Amir, 2012). Bipolar disorder and unipolar depression share a number of similar risk factors (Johnson *et al.*, 2014b), and it is unknown how interpretation bias measures may help to specify risk profiles for depression *vs* mania. However, previous research indicates that negative interpretation bias predicts future depressive symptoms (Rude *et al.*, 2002), while positive bias may protect against depression (Kleim *et al.*, 2014). Finally, although the present study focused on positive bias, future studies would do well to consider findings that some individuals with bipolar disorder may also show biases towards negative stimuli or to endorse negative beliefs about internal states (e.g. Lyon *et al.*, 1999; Mansell *et al.*, 2007). Together, these findings support the need for multifaceted, longitudinal studies that consider positive-affect relevant traits alongside other well-established risk variables to help understand this significant variability in outcomes.

Separately, results of the present study showed that higher HPS scores are associated with more frequent experiences of involuntary future-oriented mental imagery, replicating previous research in this area (Deepröse *et al.*, 2011; McCarthy-Jones *et al.*, 2012). This relationship between

involuntary imagery and HPS scores remained significant even when controlling for interpretation bias, concurrent depression symptoms, and the vividness of images during the interpretation bias task. This finding is consistent with theoretical and empirical work emphasizing the importance of involuntary, prospective imagery in bipolar disorder (Holmes *et al.*, 2011). In contrast to our hypotheses, imagery and interpretation bias did not interact. Comparisons to Holmes and colleagues' (2008) model of imagery and bipolar disorder, however, are somewhat limited by the scope of the present study. First, the interpretation bias measure included in this study (the AST) inherently includes both an imagery rating and an interpretation rating, which may have limited our ability to detect any additional interaction of these dimensions. Also, this study evaluated one relatively specific aspect of imagery (i.e. future-oriented and involuntary), whereas the imagery-interpretation theory described by Holmes and colleagues describes a more detailed process that includes initial mood elevation, increased goal pursuit, and increased imagery in response to positive cues (Holmes *et al.*, 2008). Given that interpretation of ambiguous stimuli has been shown to be more positive after a mood induction for those at risk of bipolar disorder (Trevisani *et al.*, 2008), future studies could test a more detailed set of hypotheses drawn from the theory of Holmes and colleagues by testing the impact of mood inductions on the effects of being high in both positive interpretation bias and imagery.

Results of the present study also support previous findings in isolating the particular types of mental imagery associated with mania risk. In the multiple regression, higher vividness ratings during the interpretation task correlated with lower scores on the HPS, yet involuntary, future-oriented mental imagery significantly correlated with higher scores. The inverse relationship between vividness ratings and HPS scores that emerged in regression analyses was somewhat surprising, given some previous studies showing elevated vividness ratings among individuals diagnosed with bipolar disorder (Gregory *et al.*, 2010; Holmes *et al.*, 2011). While speculative, the divergent findings regarding HPS scores and vividness *vs* volition of imagery indicate that enhanced vividness may be a feature of imagery more relevant to diagnosed samples as opposed to the student sample used in the present study. In contrast, the positive relationship between HPS scores and involuntary imagery is consistent with other studies showing a unique link between involuntary future-oriented forms of mental imagery and mood instability in bipolar disorder (Holmes *et al.*, 2011). Moreover, this finding supports theory and research indicating that involuntary aspects of mental imagery in particular are relevant for psychological disorders (Brewin *et al.*, 2010; Holmes and Mathews, 2010). Indeed, frequency of involuntary mental imagery was also correlated with current symptoms of depression in the present study, consistent with prior research (e.g. Deeprose and Holmes, 2010). However, the relationship between HPS scores and intrusive imagery remained even when controlling for depression symptoms and negative affect.

Several limitations are important to consider. The web-based data collection in the present study is less controlled than laboratory assessments of interpretation bias, which could affect the degree to which participants fully concentrated on imagining the scenarios used in the task. Second, the absence of follow-up prevents us from testing whether positive interpretation bias in those with higher HPS scores may lead to the onset of bipolar disorder over time. Third, HPS scores reported in this sample were largely below the level of cut-off scores used to index high levels of mania risk in other studies, indicating that despite the variability in HPS scores captured in our sample, it did not contain individuals at the highest risk for bipolar disorder. This limitation in range raises the possibility that the magnitude of correlations of the HPS with other measures is restricted, limiting the strength of findings. The use of a college student sample may have limited our ability to recruit individuals at the highest risk for bipolar disorder and may thus constrain the generalizability of study findings, although given the high rates of psychological disorders in college students (cf. Auerbach *et al.*, 2016), the sample was probably appropriate for testing hypotheses of psychopathology.

Many previous studies have used selective recruitment strategies to target high-scoring individuals; this strategy could be used in future studies testing interpretation bias and imagery. Finally, current symptoms other than depression were not assessed among study participants. Anxiety symptoms are linked with more negative interpretation bias in many studies (Holmes *et al.*, 2008; Mathews and MacLeod, 2005), and are thought to be particularly relevant for bipolar disorder (Holmes *et al.*, 2008). The limited number of symptoms assessed also pre-empts any evaluation of trauma-relevant imagery; the present study focused only on imagery of future events, which does not allow for a test of how post-traumatic stress disorder or other trauma-related conditions may also influence involuntary images of past trauma. This may be particularly important for future studies assessing bipolar disorder, as trauma-related disorders are frequently co-morbid with bipolar disorder (Palmier-Claus *et al.*, 2016).

Conclusions

In summary, this study provides evidence that more positive interpretations of ambiguous scenarios and more frequent involuntary mental imagery are relatively independent correlates of greater scores on the HPS. Studies assessing the prospective relationship between interpretative style and development of mood symptoms will be an important future step in evaluating the meaning of enhanced positive interpretations of ambiguity. Such work is needed, given that interpretation bias is a viable treatment target.

Acknowledgements. We thank Caroline Kinsey and Aly DiRocco for their help in preparing the measures used in this study.

Financial support. Dr Peckham was supported by NIMH grant F32 MH 115530 during preparation of this manuscript.

Conflicts of interest. Dr Peckham, Ms Modavi and Dr Johnson have no conflicts of interest with respect to this publication.

Ethical statements. All authors have abided by the Ethical Principles of Psychologists and Code of Conduct as set out by the APA. This research was approved by the Committee for the Protection of Human Subjects at the University of California, Berkeley (protocol no. 2014-07-6515).

References

- Auerbach, R. P., Alonso, J., Axinn, W. G., Cuijpers, P., Ebert, D. D., Green, J. G., Hwang, I., Kessler, R. C., Liu, H., Mortier, P., Nock, M. K., Pinder-Amaker, S., Sampson, N. A., Aguilar-Gaxiola, S., Al-Hamzawi, A., Andrade, L. H., Benjet, C., Caldas-de-Almeida, J. M., Demyttenaere, K., Florescu, S., de Girolamo, G., Gureje, O., Haro, J. M., Karam, E. G., Kiejna, A., Kovess-Masfety, V., Lee, S., McGrath, J. J., O'Neill, S., Pennell, B.-E., Scott, K., ten Have, M., Torres, Y., Zaslavsky, A. M., Zarkov, Z., & Bruffaerts, R. (2016). Mental disorders among college students in the World Health Organization World Mental Health Surveys. *Psychological Medicine*, 46, 2955–2970. <https://doi.org/10.1017/S0033291716001665>
- Beck, A. T. (1976). *Cognitive Therapy and the Emotional Disorders*. Oxford, UK: International Universities Press.
- Berna, C., Lang, T. J., Goodwin, G. M., & Holmes, E. A. (2011). Developing a measure of interpretation bias for depressed mood: an ambiguous scenarios test. *Personality and Individual Differences*, 51, 349–354. <https://doi.org/10.1016/j.paid.2011.04.005>
- Brewin, C. R., Gregory, J. D., Lipton, M., & Burgess, N. (2010). Intrusive images in psychological disorders: characteristics, neural mechanisms, and treatment implications. *Psychological Review*, 117, 210–232. <https://doi.org/10.1037/a0018113>
- Campellone, T. R., Peckham, A. D., & Johnson, S. L. (2018). Parsing positivity in the bipolar spectrum: the effect of context on social decision-making. *Journal of Affective Disorders*, 235, 316–322. <https://doi.org/10.1016/j.jad.2018.02.020>
- Cooper, J. L., & Wade, T. D. (2015). The relationship between memory and interpretation biases, difficulties with emotion regulation, and disordered eating in young women. *Cognitive Therapy and Research*, 39, 853–862. <https://doi.org/10.1007/s10608-015-9709-1>
- Cuellar, A. K., Johnson, S. L., & Winters, R. (2005). Distinctions between bipolar and unipolar depression. *Clinical Psychology Review*, 25, 307–339. <https://doi.org/10.1016/j.cpr.2004.12.002>
- Deepröse, C., & Holmes, E. A. (2010). An exploration of prospective imagery: the Impact of Future Events Scale. *Behavioural and Cognitive Psychotherapy*, 38, 201–209. <https://doi.org/10.1017/S1352465809990671>

- Deepröse, C., Malik, A., & Holmes, E. A. (2011). Measuring intrusive prospective imagery using the Impact of Future Events Scale (IFES): psychometric properties and relation to risk for bipolar disorder. *International Journal of Cognitive Therapy*, 4, 187–196. <https://doi.org/10.1521/ijct.2011.4.2.187>
- Di Simplicio, M., Renner, F., Blackwell, S. E., Mitchell, H., Stratford, H. J., Watson, P., ... & Holmes, E. A. (2016). An investigation of mental imagery in bipolar disorder: exploring ‘the mind’s eye’. *Bipolar Disorders*, 18, 669–683. <https://doi.org/10.1111/bdi.12453>
- Dodd, A. L., Mansell, W., Morrison, A. P., & Tai, S. (2011). Extreme appraisals of internal states and bipolar symptoms: the Hypomanic Attitudes and Positive Predictions Inventory. *Psychological Assessment*, 23, 635–645. <https://doi.org/10.1037/a0022972>
- Dutra, S. J., West, T. V., Impett, E. A., Oveis, C., Kogan, A., Keltner, D., & Gruber, J. (2014). Rose-colored glasses gone too far? Mania symptoms predict biased emotion experience and perception in couples. *Motivation and Emotion*, 38, 157–165. <https://doi.org/10.1007/s11031-013-9363-4>
- Eckblad, M. & Chapman, L.J. (1986). Development and validation of a scale for hypomanic personality. *Journal of Abnormal Psychology*, 95, 214–222. <http://dx.doi.org/10.1037/0021-843X.95.3.214>
- Feldman, G. C., Joormann, J., & Johnson, S. L. (2008). Responses to positive affect: a self-report measure of rumination and dampening. *Cognitive Therapy and Research*, 32, 507–525. <https://doi.org/10.1007/s10608-006-9083-0.Responses>
- García-Blanco, A. C., Perea, M., & Livianos, L. (2013). Mood-congruent bias and attention shifts in the different episodes of bipolar disorder. *Cognition & Emotion*, 27, 1114–1121. <https://doi.org/10.1080/02699931.2013.764281>
- Giovannelli, A., Hoerger, M., Johnson, S. L., & Gruber, J. (2013). Impulsive responses to positive mood and reward are related to mania risk. *Cognition & Emotion*, 27, 1091–1104. <https://doi.org/10.1080/02699931.2013.772048>
- Gregory, J. D., Brewin, C. R., Mansell, W., & Donaldson, C. (2010). Intrusive memories and images in bipolar disorder. *Behaviour Research and Therapy*, 48, 698–703. <https://doi.org/10.1016/j.brat.2010.04.005>
- Gruber, J., Dutra, S., Eidelman, P., Johnson, S. L., & Harvey, A. G. (2011a). Emotional and physiological responses to normative and idiographic positive stimuli in bipolar disorder. *Journal of Affective Disorders*, 133, 437–442. <https://doi.org/10.1016/j.jad.2011.04.045>
- Gruber, J., Eidelman, P., Johnson, S. L., Smith, B., & Harvey, A. G. (2011b). Hooked on a feeling: rumination about positive and negative emotion in inter-episode bipolar disorder. *Journal of Abnormal Psychology*, 120, 956–961. <https://doi.org/10.1037/a0023667>
- Hales, S. A., Deepröse, C., Goodwin, G. M., & Holmes, E. A. (2011). Cognitions in bipolar affective disorder and unipolar depression: imagining suicide. *Bipolar Disorders*, 13, 651–661. <https://doi.org/10.1111/j.1399-5618.2011.00954.x>
- Hertel, P. T., Mathews, A., Peterson, S., & Kintner, K. (2003). Transfer of training emotionally biased interpretations. *Applied Cognitive Psychology*, 17, 775–784. <https://doi.org/10.1002/acp.905>
- Higier, R. G., Jimenez, A. M., Hultman, C. M., Borg, J., Roman, C., Kizling, I., ... & Cannon, T. D. (2014). Enhanced neurocognitive functioning and positive temperament in twins discordant for bipolar disorder. *American Journal of Psychiatry*, 171, 1191–1198. <https://doi.org/10.1176/appi.ajp.2014.13121683>
- Hindash, A. H. C., & Amir, N. (2012). Negative interpretation bias in individuals with depressive symptoms. *Cognitive Therapy and Research*, 36, 502–511. <https://doi.org/10.1007/s10608-011-9397-4>
- Hirsch, C. R., Meeten, F., Krahé, C., & Reeder, C. (2016). Resolving ambiguity in emotional disorders: the nature and role of interpretation biases. *Annual Review of Clinical Psychology*, 12, 281–305. <https://doi.org/10.1146/annurev-clinpsy-021815-093436>
- Holmes, E. A., Deepröse, C., Fairburn, C. G., Wallace-Hadrill, S. M. A., Bonsall, M. B., Geddes, J. R., & Goodwin, G. M. (2011). Mood stability versus mood instability in bipolar disorder: a possible role for emotional mental imagery. *Behaviour Research and Therapy*, 49, 707–713. <https://doi.org/10.1016/j.brat.2011.06.008>
- Holmes, E. A., Geddes, J. R., Colom, F., & Goodwin, G. M. (2008). Mental imagery as an emotional amplifier: application to bipolar disorder. *Behaviour Research and Therapy*, 46, 1251–1258. <https://doi.org/10.1016/j.brat.2008.09.005>
- Holmes, E. A., Lang, T. J., & Shah, D. M. (2009). Developing interpretation bias modification as a ‘cognitive vaccine’ for depressed mood: imagining positive events makes you feel better than thinking about them verbally. *Journal of Abnormal Psychology*, 118, 76–88. <https://doi.org/10.1037/a0012590>
- Holmes, E. A., & Mathews, A. (2010). Mental imagery in emotion and emotional disorders. *Clinical Psychology Review*, 30, 349–362. <https://doi.org/10.1016/j.cpr.2010.01.001>
- Johnson, S. L., & Carver, C. S. (2006). Extreme goal setting and vulnerability to mania among undiagnosed young adults. *Cognitive Therapy and Research*, 30, 377–395. <https://doi.org/10.1007/s10608-006-9044-7>
- Johnson, S. L., Carver, C. S., Joormann, J., & Cuccaro, M. (2014a). A genetic analysis of the validity of the Hypomanic Personality Scale. *Bipolar Disorders*, 17, 331–339. <https://doi.org/10.1111/bdi.12251>
- Johnson, S. L., Cuellar, A. K., & Peckham, A. D. (2014b). Risk factors for bipolar disorder. In I. H. Gotlib & C. L. Hammen (eds), *Handbook of Depression* (3rd edn, pp. 315–333). New York, New York, USA: Guilford Press.
- Johnson, S. L., & Jones, S. (2009). Cognitive correlates of mania risk: are responses to success, positive moods, and manic symptoms distinct or overlapping? *Journal of Clinical Psychology*, 65, 891–905. <https://doi.org/10.1002/jclp>

- Johnson, S. L., Tharp, J. A., Peckham, A. D., & McMaster, K. J. (2016). Emotion in bipolar I disorder: implications for functional and symptom outcomes. *Journal of Abnormal Psychology, 125*, 40–52. <https://doi.org/10.1037/abn0000116>
- Jones, S., Mansell, W., & Waller, L. (2006). Appraisal of hypomania-relevant experiences: development of a questionnaire to assess positive self-dispositional appraisals in bipolar and behavioural high risk samples. *Journal of Affective Disorders, 93*, 19–28. <https://doi.org/10.1016/j.jad.2006.01.017>
- Kelly, R. E., Mansell, W., Sadhmani, V., & Wood, A. M. (2012). Positive and negative appraisals of the consequences of activated states uniquely relate to symptoms of hypomania and depression. *Cognition & Emotion, 26*, 899–906. <https://doi.org/10.1080/02699931.2011.613918>
- Kelly, R. E., Mansell, W., Wood, A. M., Alatiq, Y., Dodd, A., & Searson, R. (2011). Extreme positive and negative appraisals of activated states interact to discriminate bipolar disorder from unipolar depression and non-clinical controls. *Journal of Affective Disorders, 134*, 438–443. <https://doi.org/10.1016/j.jad.2011.05.042>
- Kirkland, T., Gruber, J., & Cunningham, W. A. (2015). Comparing happiness and hypomania risk: a study of extraversion and neuroticism aspects. *PLoS ONE, 10*, 1–18. <https://doi.org/10.1371/journal.pone.0132438>
- Kleim, B., Thörn, H. A., & Ehler, U. (2014). Positive interpretation bias predicts well-being in medical interns. *Frontiers in Psychology, 5*, 640. <https://doi.org/10.3389/fpsyg.2014.00640>
- Kohler, C. G., Hoffman, L. J., Eastman, L. B., Healey, K., & Moberg, P. J. (2011). Facial emotion perception in depression and bipolar disorder: a quantitative review. *Psychiatry Research, 188*, 303–309. <https://doi.org/10.1016/j.psychres.2011.04.019>
- Kwapil, T. R., Miller, M. B., Zinser, M. C., Chapman, L. J., Chapman, J., & Eckblad, M. (2000). A longitudinal study of high scorers on the hypomanic personality scale. *Journal of Abnormal Psychology, 109*, 222–226. <https://doi.org/10.1037//0021-843X.109.2.222>
- Lyon, H. M., Startup, M., & Bentall, R. P. (1999). Social cognition and the manic defense: attributions, selective attention, and self-schema in bipolar affective disorder. *Journal of Abnormal Psychology, 108*, 273–282.
- MacLeod, C., & Mathews, A. (2012). Cognitive bias modification approaches to anxiety. *Annual Review of Clinical Psychology, 8*, 189–217. <https://doi.org/10.1146/annurev-clinpsy-032511-143052>
- Mansell, W. (2006). The Hypomanic Attitudes and Positive Predictions Inventory (HAPPI): a pilot study to select cognitions that are elevated in individuals with bipolar disorder compared to non-clinical controls. *Behavioural and Cognitive Psychotherapy, 34*, 467–476. <https://doi.org/10.1017/S1352465806003109>
- Mansell, W., Morrison, A. P., Reid, G., Lowens, I., & Tai, S. (2007). The interpretation of, and responses to, changes in internal states: an integrative cognitive model of mood swings and bipolar disorders. *Behavioural and Cognitive Psychotherapy, 35*, 515–539. <https://doi.org/10.1017/S1352465807003827>
- Mathews, A., & MacLeod, C. (2005). Cognitive vulnerability to emotional disorders. *Annual Review of Clinical Psychology, 1*, 167–195. <https://doi.org/10.1146/annurev.clinpsy.1.102803.143916>
- McCarthy-Jones, S., Knowles, R., & Rowse, G. (2012). More than words? Hypomanic personality traits, visual imagery and verbal thought in young adults. *Consciousness and Cognition, 21*, 1375–1381. <https://doi.org/10.1016/j.concog.2012.07.004>
- Muhtadie, L., Johnson, S. L., Carver, C. S., Gotlib, I. H., & Ketter, T. A. (2014). A profile approach to impulsivity in bipolar disorder: the key role of strong emotions. *Acta Psychiatrica Scandinavica, 129*, 100–108. <https://doi.org/10.1111/acps.12136>
- Murphy, F. C., Sahakian, B. J., Rubinsztein, J. S., Michael, A., Rogers, R. D., Robbins, T. W., & Paykel, E. S. (1999). Emotional bias and inhibitory control processes in mania and depression. *Psychological Medicine, 29*, 1307–1321.
- Palmier-Claus, J. E., Berry, K., Bucci, S., Mansell, W., & Varese, F. (2016). Relationship between childhood adversity and bipolar affective disorder: systematic review and meta-analysis. *British Journal of Psychiatry, 209*, 454–459. <https://doi.org/10.1192/bjp.bp.115.179655>
- Peckham, A. D., Johnson, S. L., & Tharp, J. A. (2016). Eye tracking of attention to emotion in bipolar I disorder: links to emotion regulation and anxiety comorbidity. *International Journal of Cognitive Therapy, 9*, 1–18.
- Pictet, A., Coughtrey, A. E., Mathews, A., & Holmes, E. A. (2011). Fishing for happiness: the effects of generating positive imagery on mood and behaviour. *Behaviour Research and Therapy, 49*, 885–891. <https://doi.org/10.1016/j.brat.2011.10.003>
- Piff, P. K., Purcell, A., Gruber, J., Hertenstein, M. J., & Keltner, D. (2012). Contact high: mania proneness and positive perception of emotional touches. *Cognition & Emotion, 26*, 1116–1123. <https://doi.org/10.1080/02699931.2011.633987>
- Purcell, J. R., Lohani, M., Musket, C., Hay, A. C., Isaacowitz, D. M., & Gruber, J. (2018). Lack of emotional gaze preferences using eye-tracking in remitted bipolar I disorder. *International Journal of Bipolar Disorders, 6*, 15. <https://doi.org/10.1186/s40345-018-0123-y>
- Pyle, A., & Mansell, W. (2010). Cognitive biases in hypomanic personality: preliminary findings indicating the relevance of self-versus-other encoding and high-versus-low levels of activation. *Behavioural and Cognitive Psychotherapy, 38*, 459–472. <https://doi.org/10.1017/S1352465810000299>
- Rude, S. S., Wenzlaff, R. M., Gibbs, B., Vane, J., & Whitney, T. (2002). Negative processing biases predict subsequent depressive symptoms. *Cognition & Emotion, 16*, 423–440. <https://doi.org/10.1080/02699930143000554>
- Schönfelder, S., Langer, J., Schneider, E. E., & Wessa, M. (2017). Mania risk is characterized by an aberrant optimistic update bias for positive life events. *Journal of Affective Disorders, 218*, 313–321. <https://doi.org/10.1016/j.jad.2017.04.073>

- Schoth, D. E., & Liossi, C.** (2017). A systematic review of experimental paradigms for exploring biased interpretation of ambiguous information with emotional and neutral associations. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.00171>
- Steinman, S. A., & Teachman, B. A.** (2010). Modifying interpretations among individuals high in anxiety sensitivity. *Journal of Anxiety Disorders*, 24, 71–78. <https://doi.org/doi:10.1016/j.janxdis.2009.08.008>
- Trevisani, D. P., Johnson, S. L., & Carver, C. S.** (2008). Positive mood induction and facial affect recognition among students at risk for mania. *Cognitive Therapy and Research*, 32, 639–650. <https://doi.org/10.1007/s10608-007-9140-3>
- Walsh, M. A., DeGeorge, D. P., Barrantes-Vidal, N., & Kwapil, T. R.** (2015). A 3-year longitudinal study of risk for bipolar spectrum psychopathology. *Journal of Abnormal Psychology*, 124, 486–497. <https://doi.org/10.1037/abn0000045>
- Watson, D., & Clark, L. A.** (1999). *The PANAS-X: Manual for the Positive and Negative Affect Schedule-Expanded Form*. Ames: The University of Iowa.
- Zimmerman, M., & Coryell, W.** (1987). The Inventory to Diagnose Depression (IDD): a self-report scale to diagnose major depressive disorder. *Journal of Consulting and Clinical Psychology*, 55, 55–59. <https://doi.org/10.1037/0022-006X.55.1.55>

Cite this article: Peckham AD, Modavi K, and Johnson SL (2020). Looking on the bright side and seeing it vividly: interpretation bias and involuntary mental imagery are related to risk for bipolar disorder. *Behavioural and Cognitive Psychotherapy* 48, 203–215. <https://doi.org/10.1017/S1352465819000559>