

Original Article

Cite this article: Men VY, Emery CR, Lam T-C, Yip PSF (2022). Suicidal/self-harm behaviors among cancer patients: a population-based competing risk analysis. *Psychological Medicine* **52**, 2342–2351. <https://doi.org/10.1017/S0033291720004250>

Received: 25 July 2020
Revised: 25 September 2020
Accepted: 22 October 2020
First published online: 23 November 2020


Key words:

palliative care; psycho-oncology; suicide prevention

Author for correspondence:

Paul Siu Fai Yip, E-mail: sfpyip@hku.hk

Suicidal/self-harm behaviors among cancer patients: a population-based competing risk analysis

Vera Yu Men¹ , Clifton Robert Emery¹, Tai-Chung Lam² and Paul Siu Fai Yip^{1,3}

¹Department of Social Work and Social Administration, The University of Hong Kong, Pokfulam, Hong Kong SAR;

²Department of Clinical Oncology, The University of Hong Kong, Pokfulam, Hong Kong SAR and ³Centre for Suicide Research and Prevention, The University of Hong Kong, Pokfulam, Hong Kong SAR

Abstract

Background. Cancer patients had elevated risk of suicidality. However, few researches studied the risk/protective factors of suicidal/self-harm behaviors considering the competing risk of death. The objective of this study is to systematically investigate the risk of suicidal/self-harm behaviors among Hong Kong cancer patients as well as the contributing factors.

Methods. Patients aged 10 or above who received their first cancer-related hospital admission (2002–2009) were identified and their inpatient medical records were retrieved. They were followed for 365 days for suicidal/self-harm behaviors or death. Cancer-related information and prior 2-year physical and psychiatric comorbidities were also identified. Competing risk models were performed to explore the cumulative incidence of suicidal/self-harm behavior within 1 year as well as its contributing factors. The analyses were also stratified by age and gender.

Results. In total, 152 061 cancer patients were included in the analyses. The cumulative incidence of suicidal/self-harm behaviors within 1 year was 717.48/100 000 person-years. Overall, cancer severity, a history of suicidal/self-harm behaviors, diabetes and hypertension were related to the risk of suicidal/self-harm behaviors. There was a U-shaped association between age and suicidal/self-harm behaviors with a turning point at 58. Previous psychiatric comorbidities were not related to the risk of suicidal/self-harm behaviors. The stratified analyses confirmed that the impact of contributing factors varied by age and gender.

Conclusions. Cancer patients were at risk of suicidal/self-harm behaviors, and the impacts of related factors varied by patients' characteristics. Effective suicide prevention for cancer patients should consider the influence of disease progress and the differences in age and gender.

Introduction

Cancer is a serious public health burden especially in developed countries and regions including Hong Kong. Improvements in cancer survival have been identified in the past 20 years (Arnold et al., 2019), but they do not necessarily equate with better quality of life. Previous research found that suicide risk among cancer patients was higher compared to that in the general population (Henson et al., 2019; Zaorsky et al., 2019). Cancer metastasis (Rahouma et al., 2018; Zhong et al., 2017), other commonly reported physical and psychiatric comorbidities among cancer patients such as hypertension, diabetes, heart diseases and depression (Roy, Vallepu, Barrios, & Hunter, 2018; Sarfati et al., 2016) were also documented as independent risk factors for suicide in previous research (Ahmedani et al., 2017; Hu & Lin, 2020; Ribeiro, Huang, Fox, & Franklin, 2018; Wang, An, Shi, & Zhang, 2017). Palliative care, which is provided to patients with life-threatening diseases to improve their last-stage quality of life (Hong Kong Hospital Authority, 2020), was known to alleviate the risk of suicide among patients with advanced cancer (Sullivan et al., 2018).

Currently, the suicide prevention strategies in the Hong Kong medical system mainly focus on mental health screening and depression management (Ma & Mak, 2012; Yeung, 2015). However, it may not be effective for Hong Kong cancer patients if there is no comprehensive understanding of their suicide risk and contributing factors. Inpatient services usually provide treatments such as chemotherapy, radiotherapy and symptom control to the cancer patients (Mix et al., 2017). Patients who require inpatient services may have different disease progress and experience compared to their counterparts using outpatient and accident and emergency (A&E) services (Hinz et al., 2018; Joo, Rha, Ahn, & Kang, 2011). Moreover, as cancer patients with inpatient admission usually have a longer length of stay in hospital, hospitals should be considered as important sites for suicide prevention.

To date, the majority of suicide research among cancer patients focuses on suicide deaths and suicidal ideation (Henson et al., 2019; Ravaioli et al., 2020) as opposed to suicidal behaviors. Previous research also found mixed results regarding suicide risk by gender (Bowden

et al., 2017; Komic et al., 2017; Smalyte et al., 2013). Regarding age, previous research identified age as a linear factor associated with suicidality among cancer patients (Anderson, Park, Rosenstein, & Nichols, 2018; Samawi et al., 2017). However, we hypothesized that there would be different risk factors among different age groups especially between the young and the elderly since the suicide pattern and disease progress between the two groups are different (Barbas et al., 2012; Chou et al., 2011). Hence, we conducted stratified analysis based on age (before and after 60 years old) and gender.

As far as we know, this is the first research to systematically investigate suicidal/self-harm behaviors and the contributing factors among Chinese cancer patients. In detail, the first aim is to estimate the incidence of suicidal/self-harm behaviors among cancer patients within the first year of first cancer inpatient admission. Another aim is to identify risk and protective factors for suicidal/self-harm behaviors among cancer patients, including cancer severity, physical and psychiatric comorbidities and other sociodemographic factors, and how the impact of these factors varies by gender and age groups.

Methods

Data source

This study used the inpatient electronic medical records from the Hong Kong Hospital Authority (HA) Clinical Data Analysis and Reporting System (CDARS) between 1 January 2000 and 31 December 2010. According to the HA, the public hospitals covered approximately 90% of the inpatient services in Hong Kong, and the inpatient medical records were captured by CDARS (Leong, *n.d.*). On each record, essential clinical information such as patients' age at diagnosis, gender, hospital, admission and discharge date, number of days of stay at the hospital, up to 15 diagnoses and death status were included. To protect patients' confidentiality, each patient was assigned with a unique patient ID for data retrieval. The International Classification of Diseases, Ninth Revision (ICD-9) was used for diagnoses in the system. The CDARS medical records have been used in various epidemiological and population studies (Chai et al., 2020; Wong et al., 2020; Yung et al., 2020).

Study population and records retrieval

The study population was composed of any patients diagnosed of primary malignant tumor with their first cancer-related inpatient admission between 1 January 2002 and 31 December 2009 and were 10 years or older at the admission. For case selection, first, all the inpatient records with any primary malignant tumor diagnosis code (ICD-9 code: 140–208, exclude 196–198) between 1 January 2000 and 31 December 2010 were extracted. Then, all the cancer records were grouped by patients' reference ID, and the earliest cancer record of each patient was kept. This study focused on the impact of first cancer admission. Patients whose first admission was between 2000 and 2001 may have cancer admissions before 2000, and those records could not be captured by our data. To address this left censoring issue, patients whose first admission was in and after 2002 were included in the study. Patients who were first admitted in 2010 were excluded from the study because they did not have a full year of follow-up. Patients who were younger than 10 years at the time of admission were excluded from the study because young children do not have




Fig. 1. Study population selection process.

a basic understanding of suicide (Normand & Mishara, 1992) and suicidal ideation and attempts are rarely reported before adolescents in most countries (Borges et al., 2012). For the remaining cancer patients, all of their inpatient medical records between 2000 and 2010 were retrieved and matched based on the reference ID (Fig. 1).

Outcomes

The main outcomes were being admitted for suicidal/self-harm behaviors or death within 1 year after the discharge of the first cancer inpatient admission. A broader diagnosis in ICD-9 was used in this study (E950–959: suicide and self-inflicted injury; E980–989: injury undetermined whether accidentally or purposely inflicted) as the number of strictly defined cases (E95X) in the sample was low ($N = 198$). Meanwhile, suicides are often underreported or misclassified as with undetermined intent or accidents in Asia because of stigma against suicide and cultural and religious influences (Chang, Sterne, Lu, & Gunnell, 2010; Chen, Wu, Yousuf, & Yip, 2012). The date of the suicidal/self-harm behaviors was recorded as the admission date of the record. The date of death was recorded as the discharge date of the record indicating death.

Study variables

The severity of the cancer was the exposure of interest in the study. Since there was no detailed information on the stage of cancer in the diagnosis code, the 5-year relative survival rate for each cancer site was used as a proxy of cancer severity in our study.

The 5-year relative survival rate for each cancer site by age group based on the 1996–2001 cancer cases in Hong Kong was identified in the World Health Organization database (Law & Mang, 2011). Each cancer patient was matched with the corresponding 5-year relative survival rate based on their first primary cancer site and the age of the admission. If a patient had multiple primary cancer diagnoses on the first inpatient admission record, the lowest survival rate was used. Other cancer-related variables of interest included whether the patients received a metastatic cancer diagnosis (ICD-9 code: 196–198) or encounter for palliative care (ICD-9 code: V66.7) during the first inpatient admission and the number of days at the hospital during the first admission. The information could reflect the severity of the patients' cancer diagnosis.

Other physical and psychiatric comorbidities 730 days prior to the first cancer inpatient admission were also identified, including previous suicidal/self-harm behaviors, cardiovascular conditions, hypertension, renal problems, coagulopathy, diabetes, lipid metabolic problems, chronic obstructive pulmonary disease (COPD), depression, psychosis, drug and alcohol abuse. The diagnose codes for the abovementioned comorbidities were categorized based on Quan's Enhanced ICD-9-CM coding algorithms for Charlson comorbidities (Quan *et al.*, 2005). Demographic information such as age at diagnosis and gender was also included.

Statistical analysis

The count with percentage or mean with standard deviation (s.d.) were reported for categorical and continuous variable of interest respectively. For the study population, the differences in those variables were compared between cancer patients with and without suicidal/self-harm behaviors within 1 year after the hospital admission. Independent *t* test and χ^2 test were applied for continuous and categorical variables respectively. The Fine and Gray multivariable competing risk model was used to predict the probability of having suicidal/self-harm behaviors within 365 days of the first cancer hospitalization, taking into account the competing risk of death. It is preferred over the Cox proportional hazard regression model because cancer is a disease with high mortality rate and death precludes the occurrence of the primary outcome (suicidal/self-harm behaviors). The competing risk model provides more accurate estimation in this circumstance (Austin & Fine, 2017). The follow-up time was calculated as the number of days between the discharge date of the first admission and the outcome events (suicidal/self-harm behaviors or death) within 365 days, or 365 days if the patient was event-free and censored. For cancer patients who received a diagnosis of suicidal/self-harm behaviors on the same record of the first cancer admission, this outcome event was not counted, and they were followed for future events within the study period.

First, the cumulative incidence function for the primary outcome was generated in the competing risk analysis. Then, the bivariate and multivariable competing risk models were run for all the variables of interest. A squared term of age was also included in the multivariable model to account for the non-linear association of age. Further subgroup analyses were performed to look at whether the risk of having suicidal/self-harm behaviors varied by gender or age group (younger than 60 years old; 60 years or older). The age of 60 was chosen as the cut-off point to maintain high statistical power for subgroup analyses as developing cancer is a rare event among young adults. In Hong Kong, the incidence of cancer became more prominent after turning 60

years old (Hong Kong Cancer Registry, 2017). The crude and adjusted cause-specific hazard ratios (HRs) and the corresponding 95% confidence intervals (95% CIs) for each model were reported. Due to the large number of hypothesis tests performed, a two-sided *p* value less than 0.01 (Bonferroni corrected for five multivariable competing risk models: overall, male, female, younger than 60, 60 and older) was considered statistically significant instead of the usual 0.05 level to reduce the possibility of type I error. SAS (version 9.4; SAS Institute, Inc., Cary, NC) was used for all statistical analyses.

Result

In total, 152 061 patients aged 10 years or older received their first cancer-related inpatient admission between 2002 and 2009, among which 44.75% were female. The mean and median age of the study population was 65.20 and 68 years, respectively, and the most prevalent type of cancer was lung cancer (18.84%). The study population contributed a total of 39 926 729 person-days (109 388.30 person-years), and the mean (s.d.) and the median of the follow-up time were 262.57 (145.24) and 365 days, respectively. During the 1-year follow-up period, 1091 (0.72%) patients had an admission of suicidal/self-harm behaviors, and 56 555 (37.22%) were dead. Among the dead cancer patients, 13 098 (out of the 152 061 patients in the study) died during their first cancer admission. The characteristics of the study population are summarized in Table 1.

The baseline characteristics of the cancer patients with and without suicidal/self-harm behaviors are compared and summarized in Table 1. Compared to the cancer patients who were suicidal/self-harm free, those with suicidal/self-harm behaviors were generally older ($p < 0.001$) and more likely to be male ($p = 0.012$). They were more likely to be diagnosed with cancer with lower relative survival rate and their first cancer hospitalization was longer ($p < 0.001$). They were more likely to have a history of other physical conditions ($p < 0.05$) and more likely to have previous suicidal/self-harm behaviors ($p < 0.001$). The two groups did not differ statistically in prior 2-year psychiatric conditions.

The cumulative incidence of suicidal/self-harm behaviors within 1 year was 717.48 per 100 000 person-years (95% CI 676.05–760.91). The cumulative incidence curve is shown in Fig. 2. The results from the crude and multivariable models are summarized in Table 2. In the crude competing risk model, the cancer severity was positively associated with the risk of suicidal/self-harm behaviors, and the relationship remained significant in the multivariable analysis. In the adjusted model, for every 1% decrease in the 5-year relative survival rate, the risk of suicidal/self-harm behaviors increased by 1.1% (HR = 0.989, [0.987–0.991]). Besides lower cancer survival, having a metastatic cancer diagnosis (HR = 1.295, [1.101–1.523]), an encounter for palliative care (HR = 1.466, [1.228–1.751]), and a longer stay during the first cancer hospital admission (HR = 1.002, [1.002–1.003]) were also strongly associated with increased risk of suicidal/self-harm behaviors.

The increase in age at diagnosis was associated with elevated risk of suicidal/self-harm behaviors in the crude model (HR = 1.016, [1.012–1.020]). However, the relationship was reversed in the adjusted model. The significance of squared term of age indicated that there was a non-linear relationship between age and the risk of suicidal/self-harm behaviors after adjusting for all other variables of interest. The turning point was estimated by the derivative of the function with respect to age and setting the first derivative

Table 1. The characteristics of the study population and the comparisons between cancer patients with and without suicidal/self-harm behaviors

	Overall N = 152 061 N (%) / mean (±s.d.)	Suicidal behaviors N = 1091 N (%) / mean (±s.d.)	Non suicidal behaviors N = 150 970 N (%) / mean (±s.d.)	<i>p</i> value
Cancer survival rate	51.54 (±27.92) ^a	46.04 (±26.19) ^a	51.58 (±27.92) ^a	<0.001
Age at diagnosis	65.20 (±15.20) ^a	66.78 (±15.48) ^a	63.32 (±14.78) ^a	<0.001
Gender (female)	68 042 (44.75)	447 (40.97)	67 595 (44.77)	0.012
Length of first admission (days)	9.90 (±20.90) ^a	15.30 (±74.17) ^a	9.85 (±20.01) ^a	<0.001
Metastatic cancer	29 100 (19.14)	212 (19.43)	28 888 (19.13)	0.062
Encounter for palliative care	26 980 (17.74)	198 (18.15)	26 872 (17.74)	0.124
Previous suicide attempts	874 (0.57)	38 (3.48)	836 (0.55)	<0.001
Cardiovascular conditions	6452 (4.24)	68 (6.23)	6384 (4.23)	0.001
Hypertension	9568 (6.29)	105 (9.62)	9463 (6.27)	<0.001
Renal problems	1566 (1.03)	21 (1.92)	1545 (1.02)	0.003
Coagulopathy	349 (0.23)	6 (0.55)	343 (0.23)	0.026
Diabetes	7102 (4.67)	181 (16.59)	6921 (4.58)	<0.001
Lipid metabolic problems	1525 (1.00)	17 (1.56)	1508 (1.00)	0.065
COPD	4826 (3.17)	52 (4.77)	4774 (3.16)	0.003
Depression	501 (0.33)	4 (0.37)	497 (0.33)	0.830
Psychosis	583 (0.38)	8 (0.73)	575 (0.38)	0.061
Drug abuse	77 (0.05)	4 (0.37)	73 (0.05)	<0.001
Alcohol abuse	788 (0.52)	9 (0.82)	779 (0.52)	0.157

Bold and italic indicate that the *p* value is significant (i.e. *p* < 0.01).

^aMean and s.d. are shown.

to zero. A positive second derivative (0.0009048) indicated a U-shaped curve with the inflection point being a minimum:

$$Y = \beta_2(\text{age}) + \beta_1(\text{age})^2$$

$$Y = -0.05212(\text{age}) + 0.0004524(\text{age})^2$$

$$\frac{dY}{d(\text{age})} = -0.05212 + 2 \times 0.0004524(\text{age})$$

$$0 = 2 \times 0.0004524(\text{age}) - 0.05212$$

$$\text{age}(\text{min}) = 57.60$$

Hence, the increase in age among cancer patients was associated with decreased risk of suicidal/self-harm behaviors starting from age 10 years until about 58 years. The relationship then turned positive, indicating that the age increase was associated with elevated risk of suicidal/self-harm behaviors at older age.

Regarding prior physical and psychiatric comorbidities, in the crude models, cancer patients with previous diagnoses of physical conditions had a higher risk of developing suicidal/self-harm behaviors. The associations attenuated after adjusting for other covariates except diabetes, which was still strongly associated

with increased risk of suicidal/self-harm behaviors (HR = 4.350, [3.597–5.261]). The influence of hypertension was reversed after adjusting for covariates, meaning that having a history of hypertension was associated with decreased risk of suicidal/self-harm behaviors (HR = 0.706, [0.556–0.895]). Previous psychiatric comorbidities did not associate with elevated risk of suicidal/self-harm behaviors among cancer patients in the crude models, except for drug abuse (HR = 8.546, [3.202–22.809]). However, its influence weakened to marginal significant ($0.01 < p < 0.05$) in the multivariable model.

The relationships between study variables and the risk of suicidal/self-harm behaviors varied between males and females when stratifying by gender. The adjusted HRs and 95% CIs are summarized in Table 3. For both genders, the risk of suicidal/self-harm behaviors was higher among patients with lower survival rate, longer stay during the first cancer admission, an encounter for palliative care and history of suicidal/self-harm behaviors or diabetes. However, the age at diagnosis and having a history of hypertension were strongly associated with the risk of suicidal/self-harm behaviors among female patients (age = 0.936 [0.907–0.966]; hypertension = 0.557 [0.380–0.817]) but not males. Receiving a metastatic cancer and a history of psychosis were associated with increased risk of suicidal/self-harm behaviors among female patients with a marginal significance ($0.01 < p < 0.05$).

The results for the age-stratified models are also reported in Table 3. The risk of suicidal/self-harm behaviors was higher among cancer patients with lower 5-year relative survival rate, longer stay during the first cancer admission and history of suicidal/self-harm behaviors or diabetes for both age groups.




Fig. 2. Cumulative incidence curve and 95% CI of suicidal/self-harm behaviors among cancer patients within 1 year after inpatient hospital admission in Hong Kong, 2002–2009.

Table 2. Crude and adjusted HRs and 95% CIs for the association between variables of interest and suicidal/self-harm behaviors among cancer patients

	Crude HR (95% CI)	Adjusted HR (95% CI)
Cancer survival rate	0.985 (0.983–0.987)^a	0.989 (0.987–0.991)^a
Age at diagnosis	1.016 (1.012–1.020)^a	0.949 (0.928–0.971)^a
Age squared	1.000 (1.000–1.000)^a	1.000 (1.000–1.001)^a
Gender (female as reference)	1.298 (1.150–1.464)^a	1.130 (0.997–1.281)
Length of first admission (days)	1.002 (1.002–1.003)^a	1.002 (1.002–1.003)^a
Metastatic cancer	1.622 (1.395–1.886)^a	1.295 (1.101–1.523)^b
Encounter for palliative care	2.397 (2.050–2.802)^a	1.466 (1.228–1.751)^a
Previous suicide attempts	7.886 (5.705–10.901)^a	3.954 (2.795–5.594)^a
Cardiovascular conditions	1.857 (1.453–2.374)^a	1.040 (0.800–1.352)
Hypertension	1.844 (1.508–2.255)^a	0.706 (0.556–0.895)^a
Renal problems	2.281 (1.481–3.513)^a	1.185 (0.759–1.851)
Coagulopathy	2.785 (1.249–6.213) ^c	1.760 (0.787–3.936)
Diabetes	4.746 (4.046–5.567)^a	4.350 (3.597–5.261)^a
Lipid metabolic problems	1.653 (1.024–2.669) ^c	0.801 (0.487–1.316)
COPD	2.066 (1.564–2.729)^a	1.269 (0.953–1.690)
Depression	1.337 (0.501–3.568)	0.706 (0.261–1.907)
Psychosis	2.234 (1.116–4.475) ^c	1.644 (0.816–3.314)
Drug abuse	8.546 (3.202–22.809)^a	2.867 (1.033–7.961) ^c
Alcohol abuse	2.017 (1.047–3.886) ^c	1.068 (0.549–2.075)

Bold and italic indicate that the *p* value is significant (i.e. *p* < 0.01).

^a*p* value less than 0.001.

^b*p* value between 0.01 and 0.05.

^c*p* value between 0.001 and 0.01.

Regarding the influence of age at diagnosis, increases in age were negatively associated with the risk of suicidal/self-harm behaviors among the younger patients (HR = 0.980, [0.969–0.992]). However, the association turned positive for older patients (HR = 1.012, [1.003–1.022]). Among older patients, an encounter for palliative care during the first admission (HR = 1.575, [1.294–1.917]) and a history of hypertension (HR = 0.682, [0.532–0.874]) were strongly associated with risk of suicidal/self-harm behaviors. However, the associations were not statistically significant among younger patients. Receiving a metastatic cancer diagnosis during the first admission was associated with increased risk of suicidal/self-harm behaviors among younger patients (HR = 1.699, [1.273–2.266]) but not among their older counterparts.

Discussion

This study employed competing risk models to explore the risk of developing suicidal/self-harm behaviors within 1 year after patients' first cancer inpatient hospital admission and investigate the contributing factors using the electronic medical records in Hong Kong public hospitals. Our results indicated that cancer severity, age at diagnosis, previous suicidal/self-harm behaviors and some physical comorbidities were significant predictors of suicidal/self-harm behaviors. Some risk and protective factors also varied by gender and different age groups.

In our sample, the cumulative incidence of suicidal/self-harm behaviors within first year of cancer admission was 717.48/100 000 person-years. Previous literature looking at suicide deaths among cancer patients reported the rate ranging from 31.4/100 000 to 274.7/100 000 person-years (Anguiano, Mayer, Piven, & Rosenstein, 2012; Kam *et al.*, 2015; Misono, Weiss, Fann, Redman, & Yueh, 2008; Siracuse, Gorgy, Ruskin, & Beebe, 2017). Another study investigating the self-harm behaviors among UK primary care patients reported the incidence being 123 and 179/100 000 person-years for male and female patients respectively (Carr *et al.*, 2016). The high incidence of suicidal/self-harm behaviors among cancer patients suggests that cancer patients may be vulnerable to suicidality because of their physical

Table 3. Adjusted HRs and 95% CIs for the association between variables of interest and suicidal/self-harm behaviors among cancer patients stratified by gender and age.

	Gender		Age	
	Male N = 84 019 HR (95% CI)	Female N = 68 042 HR (95% CI)	Younger than 60 N = 52 863 HR (95% CI)	60 and older N = 99 198 HR (95% CI)
Cancer survival rate	0.990 (0.986–0.993)^a	0.988 (0.984–0.992)^a	0.985 (0.980–0.990)^a	0.990 (0.987–0.993)^a
Age at diagnosis	0.965 (0.932–0.998) ^b	0.936 (0.907–0.966)^a	0.980 (0.969–0.992)^a	1.012 (1.003–1.022)^c
Age squared	1.000 (1.000–1.001) ^b	1.001 (1.000–1.001)^a	NA	NA
Gender (female as reference)	NA	NA	0.973 (0.763–1.240)	1.171 (1.008–1.359) ^b
Length of first admission (days)	1.002 (1.001–1.003)^a	1.004 (1.003–1.005)^a	1.004 (1.003–1.005)^a	1.002 (1.001–1.003)^a
Metastatic cancer	1.234 (0.999–1.525)	1.387 (1.074–1.791) ^b	1.699 (1.273–2.266)^a	1.129 (0.926–1.375)
Encounter for palliative care	1.435 (1.138–1.809)^c	1.499 (1.137–1.978)^c	1.304 (0.854–1.990)	1.575 (1.294–1.917)^a
Previous suicide attempts	3.825 (2.412–6.063)^a	4.133 (2.431–7.027)^a	8.123 (3.964–16.644)^a	3.329 (2.223–4.955)^a
Cardiovascular conditions	0.986 (0.699–1.392)	1.112 (0.740–1.672)	0.361 (0.050–2.623)	1.089 (0.834–1.421)
Hypertension	0.834 (0.616–1.128)	0.557 (0.380–0.817)^c	0.942 (0.409–2.173)	0.682 (0.532–0.874)^c
Renal problems	0.930 (0.504–1.718)	1.655 (0.862–3.180)	1.207 (0.292–4.995)	1.186 (0.741–1.896)
Coagulopathy	1.362 (0.437–4.243)	2.424 (0.771–7.614)	2.676 (0.657–10.891)	1.499 (0.560–4.013)
Diabetes	4.292 (3.367–5.472)^a	4.469 (3.297–6.058)^a	3.018 (1.713–5.318)^a	4.605 (3.758–5.643)^a
Lipid metabolic problems	0.977 (0.547–1.777)	0.555 (0.225–1.371)	0.739 (0.099–5.507)	0.809 (0.484–1.351)
COPD	1.196 (0.861–1.662)	1.542 (0.859–2.770)	2.786 (1.088–7.127)	1.184 (0.876–1.600)
Depression	0.406 (0.057–2.909)	0.863 (0.265–2.811)	2.383 (0.299–5.493) ^b	0.465 (0.115–1.880)
Psychosis	0.439 (0.062–3.128)	2.740 (1.268–5.923) ^b	1.549 (0.478–4.960)	1.744 (0.720–4.221)
Drug abuse	2.873 (0.870–9.486)	2.363 (0.307–18.190)	1.861 (0.491–7.060)	1.757 (0.240–12.849)
Alcohol abuse	1.039 (0.512–2.110)	1.493 (0.205–10.854)	1.576 (0.617–4.027)	0.729 (0.271–1.963)

NA, not applicable.

Bold and italic indicate that the *p* value is significant (i.e. *p* < 0.01).^a*p* value less than 0.001.^b*p* value between 0.01 and 0.05.^c*p* value between 0.001 and 0.01.

conditions. Special attention and care are required during hospital visits and follow-ups to allow early prevention.

Our results indicated that the cancer severity, such as low survival rate, receiving a metastatic cancer diagnosis and longer stay in hospital elevated the risk of suicidal/self-harm behaviors among patients. The findings are supported by previous literature, which found that the suicide risk was higher among cancer patients with severe disease progress (Urban et al., 2013; Vyssoki et al., 2015). The effect of receiving a metastatic diagnosis varied by age, as having a metastatic cancer diagnosis increased the risk of suicidal/self-harm behaviors among younger patients, whereas the effect was insignificant among older patients. Younger cancer patients may perceive a metastatic cancer diagnosis as a more serious negative life event, which may explain the variation between age groups. Length of admission was not considered in previous research. However, the length of first admission may proxy the seriousness of the situation since a longer stay might indicate the patients require more medical care. Moreover, increasing the length of stay exposes the patients to the uncomfortable medical environment for a longer period.

Palliative care aims at improving the quality of life of the patients during the last stage of their illnesses. In contrast to the previous finding (Sullivan et al., 2018), this study concluded

that encountering palliative care increased the risk of suicidal/self-harm behaviors among the patients overall. The discrepancy in findings may be explained by the difference in study population and outcomes, as previous research focused on suicide death among US lung cancer patients whereas we investigated suicidal/self-harm behaviors among all Hong Kong cancer patients. The positive relationship between palliative care and the risk of suicidal/self-harm behaviors in our study may occur because palliative care reduces pain (Kassianos, Ioannou, Koutsantoni, & Charalambous, 2018) but also alters the patients' self-assessment of long-term survival (Zimmermann et al., 2016). The stratified analysis provided further evidence that the impact of palliative care was stronger among older cancer patients. Patients who encountered palliative care were at the end stage of their life suffering from more severe physical pain compared to their counterparts who did not require the services (Riechelmann, Krzyzanowska, O'Carroll, & Zimmermann, 2007). Moreover, cancer patients receiving palliative care often reported experiencing spiritual pain (Mako, Galek, & Poppito, 2006), and they were more likely to suffer from depression and anxiety symptoms (Wilson et al., 2007). Therefore, they may be more likely to attempt suicidal/self-harm behaviors to end their suffering. On the other hand, palliative care may be interpreted differently in

various cultural settings. Palliative care is relatively new in the Chinese society. Dying at home is a traditional concept. A previous study in Taiwan documented that about two-thirds of the cancer patients preferred dying at home since it was a place with a sense of belonging and security, greater control and more autonomy (Tang, 2000). Compared to the younger generation, this concept may be more widely accepted in the older population. Hence, elderly patients who encountered palliative care may feel a loss of autonomy and insecurity which may increase their risk of suicidal/self-harm behaviors.

Our study hypothesized that the relationship between age and suicidal/self-harm behaviors may not be linear, and the study results suggested a U-shaped relationship with a turning point at around 58 years old. The stratified analysis further showed that there is a negative association between age and suicidal/self-harm behaviors for patients younger than 60 years old, but the association reversed over the age of 60 years. The higher risk among the younger patients may be because they are just at the beginning of their journey in life. They are more impulsive and vulnerable to risk taking compared to people at older age (Steinberg *et al.*, 2008). Being admitted for cancer may be more devastating compared to those who are middle-aged. Among the old cancer patients, their physical conditions have already deteriorated. The feeling of being a burden to the family, low quality of life and loss of autonomy may trigger a wish for hastened death (McPherson, Wilson, & Murray, 2007; Mystakidou, Parpa, Katsouda, Galanos, & Vlahos, 2004; Olden, Pessin, Lichtenthal, & Breitbart, 2012). Aged 60 years is the usual retirement age of many Hong Kong working adults and suicidal risk among older adults is also about two times than that of the general population in Hong Kong (Yip & Zheng, 2020). The presence of cancer among older adults will certainly increase the suicidal risk. The risk of suicidal/self-harm behaviors is relatively low among middle-aged patients, as they may have more resources for their disease treatment and autonomy for decision-making (Sheldon, Houser-Marko, & Kasser, 2006), and they may carry more responsibility for their families. This is crucial for suicide prevention since medical and public health professionals may currently focus more on the elderly but may overlook the risk among young people.

In our sample, the patients with previous history of suicidal/self-harm behaviors had a higher risk of repeating behaviors, regardless of patients' age and gender. Past literature is consistent with our findings in showing that previous self-harm behaviors and suicide attempts are strong predictors for future suicidal behaviors among cancer patients (Camidge *et al.*, 2007).

In the past research, Schneider and Shenassa (2008) found that cancer patients with one or more chronic diseases were more likely to develop suicidal ideation. Our findings confirmed that a history of diabetes was strongly associated with the elevated risk of suicidal/self-harm behaviors. Diabetes is a common physical comorbidity among cancer patients (Sarfati *et al.*, 2016), and it is an independent risk factor for suicidality (Sarkar & Balhara, 2014). Unexpectedly, having a history of hypertension was a protective factor of suicidal/self-harm behaviors, especially among female and older cancer patients. A possible explanation is that patients with hypertension are usually advised to manage their stress, perform relaxation techniques (Abgrall-Barbry & Consoli, 2006), and engage in more physical activities (Semlitsch *et al.*, 2013) for symptom control, all of which are beneficial to people's mental health (Chiesa & Serretti, 2009; Dinas, Koutedakis, & Flouris, 2011; Song, Xu, Zhang, Ma, & Zhao, 2013) and may potentially decrease the risk of suicidal/self-harm behaviors.

Many other studies have identified a positive relationship between depression and mental distress and suicidal ideation (Tang *et al.*, 2016; Zhong *et al.*, 2017) or suicide death (Aboumrad, Shiner, Riblet, Mills, & Watts, 2018; Klaassen *et al.*, 2018) among cancer patients. However, our results indicated that having a history of psychiatric comorbidities was not associated with the risk of suicidal/self-harm behaviors. The discrepancy in results may be explained by the difference in the outcome measures, as our study focuses on suicidal/self-harm behaviors while previous studies mainly looked at suicidal ideation and death. Another possible explanation is that our study used inpatient medical records to look for psychiatric comorbidities, so that only the severe mental conditions resulting in clinical diagnosis were captured. Meanwhile, the diagnosis of depression can be difficult among cancer patients as the symptoms are overlapping (Weinberger, Roth, & Nelson, 2009), and the Chinese have stigma against depression (Georg Hsu *et al.*, 2008) and express it differently compared to the West (Parker, Gladstone, & Chee, 2001), all of which may contribute to the underdiagnosis of depression in our sample. This may disproportionately affect cancer patients with suicidal/self-harm behaviors if depression is a significant risk factor, leading the effect size toward the null. Therefore, the influence of psychiatric comorbidities on suicidal/self-harm behaviors may be underestimated in our study. However, it is also possible that mental health status may not be a significant predictor of suicide among cancer patients as Cheung, Douwes, and Sundram (2017) found out that people who died from suicide with terminal cancer were less likely to have depression or have previous contact with mental health services compared with their cancer-free counterparts.

The current study has numerous strengths. To our knowledge, this is the first study systematically investigating suicidal/self-harm behaviors and related factors among Hong Kong cancer patients. Unlike previous studies which mainly focused on suicide deaths, the main outcomes of this study are suicidal/self-harm behaviors. Exploring the risk and protective factors for suicidal/self-harm behaviors instead of suicide deaths allows the implementation of earlier prevention. The stratified analysis and the use of competing risk models provide better understanding and more accurate estimation of the risk of suicidal/self-harm behaviors as well as the identification of the risk and protective factors. The large and representative sample ensures adequate statistical power and allows the study results to be generalized to other populations especially the Chinese and Asian populations.

Our study has several limitations. This study only used inpatient medical records to retrieve diagnosis of suicidal/self-harm behaviors and previous physical and psychiatric comorbidities, which means that cancer patients' private hospital visits, outpatient and A&E services utilization were not included. As a result, the cumulative incidence of suicidal/self-harm behaviors may be underestimated because the less severe cases are usually treated and discharged in the A&E department. This study also adopted a broader definition of suicidal/self-harm behaviors (E95X and E98X) due to a low number of strict defined cases (E95X) in our sample. In the inpatient medical records, there is no detailed information on patients' cancer disease progress, such as the stage of cancer and medicine prescription records. To address this issue, in our analysis, the age-specific 5-year relative survival rate for each cancer diagnosis was used to proxy cancer severity during patients' first admission. Additionally, whether receiving a metastatic cancer diagnosis, an encounter for palliative care and the length of stay were used to supplement information

on the disease progress. Future research may provide more comprehensive understanding of the suicidal/self-harm behaviors among cancer patients by linking inpatient, outpatient and A&E records together using a strict defined definition.

Suicide usually progresses from suicidal ideation to attempting suicidal/self-harm behaviors and to tragic death. Suicidal/self-harm behaviors have been proved to be strong risk factors for suicide death (Centers for Disease Control and Protection, 2019). Our study has identified numerous contributing factors of suicidal/self-harm behaviors among cancer patients, making early warning systems and interventions for suicide viable. Special attention should be paid to cancer patients with low survival as they have a higher risk of attempting suicidal/self-harm behaviors. Although a longer stay in hospital is found to be a risk factor for suicidal/self-harm behaviors in the study, it can be an opportunity for suicide prevention. Currently, suicide prevention among cancer patients focuses on mental health screening and depression management not only in Hong Kong but also in many developed nations (Butow et al., 2015; Walker & Sharpe, 2009). However, the current study suggests that psychiatric conditions are not significant risk factors. Therefore, the current suicide prevention strategies may overlook some needs of the cancer patients. Patients with previous suicide attempts must receive additional attention. The current study also provides evidence that both genders require equal attention. Numerous studies have focused on the issue of suicide among elderly cancer patients. This study confirms that the elderly is one high-risk population, but more importantly, the higher risk among the younger population cannot be overlooked in suicide prevention. Healthcare professionals should not omit the potential suicidal/self-harm risks of the younger cancer patients, and more information and resources should be provided to this group. Palliative care is a relatively new concept in Chinese society. It aims at improving the quality of life of the last-stage patients. However, palliative care may conflict with the traditional Chinese concept of dying at home. It is essential for healthcare professionals to understand Chinese cultural beliefs and values related to dying and have comprehensive communication with the patients and their families to provide appropriate end-of-life care to enhance their sense of control over the unknown process of dying.

Acknowledgements. We thank Hong Kong Hospital Authority for their supports in getting access to the data. We would also like to show our gratitude to Cheuk Yui Yeung, M.P.H. at the Department of Social Work and Social Administration, The University of Hong Kong for helpful discussion and comments that greatly improved the manuscript.

Financial support. This work was supported by the Li Ka Shing Foundation (AR180055) and Hong Kong General Research Fund (GRF) (17103620).

Conflict of interest. None.

References

- Abgrall-Barbry, G., & Consoli, S. M. (2006). L'approche psychologique dans le traitement de l'hypertension artérielle. *La Presse Médicale*, 35(6), 1088–1094. [https://doi.org/10.1016/S0755-4982\(06\)74753-4](https://doi.org/10.1016/S0755-4982(06)74753-4).
- Aboumrad, M., Shiner, B., Riblet, N., Mills, P. D., & Watts, B. V. (2018). Factors contributing to cancer-related suicide: A study of root-cause analysis reports. *Psycho-Oncology*, 27(9), 2237–2244. <https://doi.org/10.1002/pon.4815>.
- Ahmedani, B. K., Peterson, E. L., Hu, Y., Rossom, R. C., Lynch, F., Lu, C. Y., ... Simon, G. E. (2017). Major physical health conditions and risk of suicide. *American Journal of Preventive Medicine*, 53(3), 308–315. <https://doi.org/10.1016/j.amepre.2017.04.001>.
- Anderson, C., Park, E. M., Rosenstein, D. L., & Nichols, H. B. (2018). Suicide rates among patients with cancers of the digestive system. *Psycho-Oncology*, 27(9), 2274–2280. <https://doi.org/10.1002/pon.4827>.
- Anguiano, L., Mayer, D. K., Piven, M. L., & Rosenstein, D. (2012). A literature review of suicide in cancer patients. *Cancer Nursing*, 35(4), E14–E26. <https://doi.org/10.1097/NCC.0b013e31822fc76c>.
- Arnold, M., Rutherford, M. J., Bardot, A., Ferlay, J., Andersson, T. M. L., Myklebust, T. Å., ... Bray, F. (2019). Progress in cancer survival, mortality, and incidence in seven high-income countries 1995–2014 (ICBP SURVMARK-2): A population-based study. *The Lancet Oncology*, 20(11), 1493–1505. [https://doi.org/10.1016/S1470-2045\(19\)30456-5](https://doi.org/10.1016/S1470-2045(19)30456-5).
- Austin, P. C., & Fine, J. P. (2017). Practical recommendations for reporting Fine-Gray model analyses for competing risk data. *Statistics in Medicine*, 36(27), 4391–4400. <https://doi.org/10.1002/sim.7501>.
- Barbas, A. S., Turley, R. S., Ceppa, E. P., Reddy, S. K., Blazer, D. G., Clary, B. M., ... Lagoo, S. A. (2012). Comparison of outcomes and the use of multi-modality therapy in young and elderly people undergoing surgical resection of pancreatic cancer. *Journal of the American Geriatrics Society*, 60(2), 344–350. <https://doi.org/10.1111/j.1532-5415.2011.03785.x>.
- Borges, G., Chiu, W. T., Hwang, I., Panchal, B., Ono, Y., Sampson, N. A., ... Nock, M. K. (2012). Prevalence, onset, and transitions among suicidal behaviors. In M. K. Nock, G. Borges, & Y. Ono (Eds.), *Suicide: Global perspectives from the WHO world mental health surveys* (p. 66). Cambridge, United Kingdom: Cambridge University Press.
- Bowden, M. B., Walsh, N. J., Jones, A. J., Talukder, A. M., Lawson, A. G., & Kruse, E. J. (2017). Demographic and clinical factors associated with suicide in gastric cancer in the United States. *Journal of Gastrointestinal Oncology*, 8(5), 897–901. <https://doi.org/10.21037/jgo.2017.08.02>.
- Butow, P., Price, M. A., Shaw, J. M., Turner, J., Clayton, J. M., Grimison, P., ... Kirsten, L. (2015). Clinical pathway for the screening, assessment and management of anxiety and depression in adult cancer patients: Australian guidelines. *Psycho-Oncology*, 24(9), 987–1001. <https://doi.org/10.1002/pon.3920>.
- Camidge, D. R., Stockton, D. L., Frame, S., Wood, R., Bain, M., & Bateman, D. N. (2007). Hospital admissions and deaths relating to deliberate self-harm and accidents within 5 years of a cancer diagnosis: A national study in Scotland, UK. *British Journal of Cancer*, 96(5), 752–757. <https://doi.org/10.1038/sj.bjc.6603617>.
- Carr, M. J., Ashcroft, D. M., Kontopantelis, E., Awenat, Y., Cooper, J., Chew-Graham, C., ... Webb, R. T. (2016). The epidemiology of self-harm in a UK-wide primary care patient cohort, 2001–2013. *BMC Psychiatry*, 16(1), 53. <https://doi.org/10.1186/s12888-016-0753-5>.
- Centers for Disease Control and Protection. (2019). *Risk and Protective Factors for Suicide*. Retrieved from <https://www.cdc.gov/violenceprevention/suicide/riskprotectivefactors.html>.
- Chai, Y., Luo, H., Wong, G. H. Y., Tang, J. Y. M., Lam, T.-C., Wong, I. C. K., & Yip, P. S. F. (2020). Risk of self-harm after the diagnosis of psychiatric disorders in Hong Kong, 2000–10: A nested case-control study. *The Lancet Psychiatry*, 7(2), 135–147. [https://doi.org/10.1016/S2215-0366\(20\)30004-3](https://doi.org/10.1016/S2215-0366(20)30004-3).
- Chang, S.-S., Sterne, J. A. C., Lu, T.-H., & Gunnell, D. (2010). 'Hidden' suicides amongst deaths certified as undetermined intent, accident by pesticide poisoning and accident by suffocation in Taiwan. *Social Psychiatry and Psychiatric Epidemiology*, 45(2), 143–152. <https://doi.org/10.1007/s00127-009-0049-x>.
- Chen, Y. Y., Wu, K. C.-C., Yousuf, S., & Yip, P. S. F. (2012). Suicide in Asia: Opportunities and challenges. *Epidemiologic Reviews*, 34(1), 129–144. <https://doi.org/10.1093/epirev/mxr025>.
- Cheung, G., Douwes, G., & Sundram, F. (2017). Late-life suicide in terminal cancer: A rational act or underdiagnosed depression? *Journal of Pain and Symptom Management*, 54(6), 835–842. <https://doi.org/10.1016/j.jpainsymman.2017.05.004>.
- Chiesa, A., & Serretti, A. (2009). Mindfulness-based stress reduction for stress management in healthy people: A review and meta-analysis. *The Journal of Alternative and Complementary Medicine*, 15(5), 593–600. <https://doi.org/10.1089/acm.2008.0495>.
- Chou, C.-L., Chang, S.-C., Lin, T.-C., Chen, W.-S., Jiang, J.-K., Wang, H.-S., ... Lin, J.-K. (2011). Differences in clinicopathological characteristics of

- colorectal cancer between younger and elderly patients: An analysis of 322 patients from a single institution. *The American Journal of Surgery*, 202(5), 574–582. <https://doi.org/10.1016/j.amjsurg.2010.10.014>.
- Dinas, P. C., Koutedakis, Y., & Flouris, A. D. (2011). Effects of exercise and physical activity on depression. *Irish Journal of Medical Science*, 180(2), 319–325. <https://doi.org/10.1007/s11845-010-0633-9>.
- Georg Hsu, L. K., Wan, Y. M., Chang, H., Summergrad, P., Tsang, B. Y. P., & Chen, H. (2008). Stigma of depression is more severe in Chinese Americans than Caucasian Americans. *Psychiatry: Interpersonal and Biological Processes*, 71(3), 210–218. <https://doi.org/10.1521/psyc.2008.71.3.210>.
- Henson, K. E., Brock, R., Charnock, J., Wickramasinghe, B., Will, O., & Pitman, A. (2019). Risk of suicide after cancer diagnosis in England. *JAMA Psychiatry*, 76(1), 51–60. <https://doi.org/10.1001/jamapsychiatry.2018.3181>.
- Hinz, A., Weis, J., Faller, H., Brähler, E., Härter, M., Keller, M., ... Mehnert, A. (2018). Quality of life in cancer patients – a comparison of inpatient, outpatient, and rehabilitation settings. *Supportive Care in Cancer*, 26(10), 3533–3541. <https://doi.org/10.1007/s00520-018-4211-4>.
- Hong Kong Cancer Registry. (2017). Overview of 2009 Hong Kong Cancer Statistics. Retrieved from <https://www3.ha.org.hk/cancereg/pdf/overview/SummaryofCanStat2009.pdf>.
- Hong Kong Hospital Authority. (2020). Palliative Care Service. Retrieved from https://www.ha.org.hk/visitor/ha_visitor_text_index.asp?Content_ID=10096&Lang=ENG&Dimension=100.
- Hu, W.-S., & Lin, C.-L. (2020). A retrospective study of suicide attempts in patients with pulmonary hypertension. *Heart, Lung and Circulation*, 29(9), e231–e237. <https://doi.org/10.1016/j.hlc.2020.02.005>.
- Joo, E.-H., Rha, S.-Y., Ahn, J. B., & Kang, H.-Y. (2011). Economic and patient-reported outcomes of outpatient home-based versus inpatient hospital-based chemotherapy for patients with colorectal cancer. *Supportive Care in Cancer*, 19(7), 971–978. <https://doi.org/10.1007/s00520-010-0917-7>.
- Kam, D., Salib, A., Gorgy, G., Patel, T. D., Carniol, E. T., Eloy, J. A., ... Park, R. C. W. (2015). Incidence of suicide in patients with head and neck cancer. *JAMA Otolaryngology – Head and Neck Surgery*, 141(12), 1075–1081. <https://doi.org/10.1001/jamaoto.2015.2480>.
- Kassianos, A. P., Ioannou, M., Koutsantoni, M., & Charalambous, H. (2018). The impact of specialized palliative care on cancer patients' health-related quality of life: A systematic review and meta-analysis. *Supportive Care in Cancer*, 26(1), 61–79. <https://doi.org/10.1007/s00520-017-3895-1>.
- Klaassen, Z., Arora, K., Wilson, S. N., King, S. A., Madi, R., Neal, D. E., ... Terris, M. K. (2018). Decreasing suicide risk among patients with prostate cancer: Implications for depression, erectile dysfunction, and suicidal ideation screening. *Urologic Oncology: Seminars and Original Investigations*, 36(2), 60–66. <https://doi.org/10.1016/j.urolonc.2017.09.007>.
- Komic, A., Taludker, A., Walsh, N., Jones, A., Lawson, A., Bateson, B., & Kruse, E. J. (2017). Suicide risk in melanoma: What do we know? *The American Surgeon*, 83(11), e435–e437. <http://www.ncbi.nlm.nih.gov/pubmed/30454229>.
- Law, S. C., & Mang, O. W. (2011). Cancer survival in Hong Kong SAR, China, 1996–2001. In R. Sankaranarayanan, & R. Swaminathan (Eds.), *Cancer survival in Africa, Asia, the Caribbean and Central America* (p. 40). Lyon, France: International Agency for Research on Cancer, World Health Organization.
- Leong, J. C. Y. (n.d.). Hospital Authority of Hong Kong. Retrieved from https://www.ha.org.hk/haho/ho/hesd/attachments/oppo_HospitalAuthority-Overview.pdf.
- Ma, Y., & Mak, K. (2012). Management of depression in cancer patients. *The Hong Kong Practitioner*, 34(4), 154–159.
- Mako, C., Galek, K., & Poppito, S. R. (2006). Spiritual pain among patients with advanced cancer in palliative care. *Journal of Palliative Medicine*, 9(5), 1106–1113. <https://doi.org/10.1089/jpm.2006.9.1106>.
- McPherson, C. J., Wilson, K. G., & Murray, M. A. (2007). Feeling like a burden to others: A systematic review focusing on the end of life. *Palliative Medicine*, 21(2), 115–128. <https://doi.org/10.1177/0269216307076345>.
- Misono, S., Weiss, N. S., Fann, J. R., Redman, M., & Yueh, B. (2008). Incidence of suicide in persons with cancer. *Journal of Clinical Oncology*, 26(29), 4731–4738. <https://doi.org/10.1200/JCO.2007.13.8941>.
- Mix, J. M., Granger, C. V., LaMonte, M. J., Niewczyk, P., DiVita, M. A., Goldstein, R., ... Freudenheim, J. L. (2017). Characterization of cancer patients in inpatient rehabilitation facilities: A retrospective cohort study. *Archives of Physical Medicine and Rehabilitation*, 98(5), 971–980. <https://doi.org/10.1016/j.apmr.2016.12.023>.
- Mystakidou, K., Parpa, E., Katsouda, E., Galanos, A., & Vlahos, L. (2004). Influence of pain and quality of life on desire for hastened death in patients with advanced cancer. *International Journal of Palliative Nursing*, 10(10), 476–483. <https://doi.org/10.12968/ijpn.2004.10.10.16211>.
- Normand, C. L., & Mishara, B. L. (1992). The development of the concept of suicide in children. *OMEGA – Journal of Death and Dying*, 25(3), 183–203. <https://doi.org/10.2190/7TG9-NVG3-1RBJ-UDAH>.
- Olden, M., Pessin, H., Lichtenthal, W. G., & Breitbart, W. (2012). Suicide and desire for hastened death in the terminally ill. In H. M. Chochinov, & W. Breitbart (Eds.), *Handbook of psychiatry in palliative medicine* (p. 103). Oxford, United Kingdom: Oxford University Press.
- Parker, G., Gladstone, G., & Chee, K. T. (2001). Depression in the planet's largest ethnic group: The Chinese. *American Journal of Psychiatry*, 158(6), 857–864. <https://doi.org/10.1176/appi.ajp.158.6.857>.
- Quan, H., Sundararajan, V., Halfon, P., Fong, A., Burnand, B., Luthi, J.-C., ... Ghali, W. A. (2005). Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Medical Care*, 43(11), 1130–1139. <https://doi.org/10.1097/01.mlr.0000182534.19832.83>.
- Rahouma, M., Kamel, M., Abouarab, A., Eldessouki, I., Nasar, A., Harrison, S., ... Port, J. L. (2018). Lung cancer patients have the highest malignancy-associated suicide rate in USA: A population-based analysis. *Ecancermedicalscience*, 12, 1–12. <https://doi.org/10.3332/ecancer.2018.859>.
- Ravaioli, A., Crocetti, E., Mancini, S., Baldacchini, F., Giuliani, O., Vattiato, R., ... Falcini, F. (2020). Suicide death among cancer patients: New data from northern Italy, systematic review of the last 22 years and meta-analysis. *European Journal of Cancer*, 125, 104–113. <https://doi.org/10.1016/j.ejca.2019.08.019>.
- Ribeiro, J. D., Huang, X., Fox, K. R., & Franklin, J. C. (2018). Depression and hopelessness as risk factors for suicide ideation, attempts and death: Meta-analysis of longitudinal studies. *British Journal of Psychiatry*, 212(5), 279–286. <https://doi.org/10.1192/bjp.2018.27>.
- Riechelmann, R. P., Krzyzanowska, M. K., O'Carroll, A., & Zimmermann, C. (2007). Symptom and medication profiles among cancer patients attending a palliative care clinic. *Supportive Care in Cancer*, 15(12), 1407–1412. <https://doi.org/10.1007/s00520-007-0253-8>.
- Roy, S., Vallepu, S., Barrios, C., & Hunter, K. (2018). Comparison of comorbid conditions between cancer survivors and age-matched patients without cancer. *Journal of Clinical Medicine Research*, 10(12), 911–919. <https://doi.org/10.14740/jocmr3617w>.
- Samawi, H. H., Shaheen, A. A., Tang, P. A., Heng, D. Y. C., Cheung, W. Y., & Vickers, M. M. (2017). Risk and predictors of suicide in colorectal cancer patients: A surveillance, epidemiology, and end results analysis. *Current Oncology*, 24(6), e513–e517. <https://doi.org/10.3747/co.24.3713>.
- Sarfati, D., Gurney, J., Lim, B. T., Bagheri, N., Simpson, A., Koea, J., & Dennett, E. (2016). Identifying important comorbidity among cancer populations using administrative data: Prevalence and impact on survival. *Asia-Pacific Journal of Clinical Oncology*, 12(1), e47–e56. <https://doi.org/10.1111/ajco.12130>.
- Sarkar, S., & Balhara, Y. P. (2014). Diabetes mellitus and suicide. *Indian Journal of Endocrinology and Metabolism*, 18(4), 468. <https://doi.org/10.4103/2230-8210.137487>.
- Schneider, K. L., & Shenassa, E. (2008). Correlates of suicide ideation in a population-based sample of cancer patients. *Journal of Psychosocial Oncology*, 26(2), 49–62. https://doi.org/10.1300/J077v26n02_04.
- Semlitsch, T., Jeitler, K., Hemkens, L. G., Horvath, K., Nagele, E., Schuermann, C., ... Siebenhofer, A. (2013). Increasing physical activity for the treatment of hypertension: A systematic review and meta-analysis. *Sports Medicine*, 43(10), 1009–1023. <https://doi.org/10.1007/s40279-013-0065-6>.
- Sheldon, K. M., Houser-Marko, L., & Kasser, T. (2006). Does autonomy increase with age? Comparing the goal motivations of college students and their parents. *Journal of Research in Personality*, 40(2), 168–178. <https://doi.org/10.1016/j.jrp.2004.10.004>.
- Siracuse, B. L., Gorgy, G., Ruskin, J., & Beebe, K. S. (2017). What is the incidence of suicide in patients with bone and soft tissue cancer?: Suicide and sarcoma. *Clinical Orthopaedics and Related Research*, 475(5), 1439–1445. <https://doi.org/10.1007/s11999-016-5171-y>.

- Smailyte, G., Jasilionis, D., Kaceniene, A., Krilaviciute, A., Ambrozaitiene, D., & Stankuniene, V. (2013). Suicides among cancer patients in Lithuania: A population-based census-linked study. *Cancer Epidemiology*, 37(5), 714–718. <https://doi.org/10.1016/j.canep.2013.05.009>.
- Song, Q.-H., Xu, R.-M., Zhang, Q.-H., Ma, M., & Zhao, X.-P. (2013). Relaxation training during chemotherapy for breast cancer improves mental health and lessens adverse events. *International Journal of Clinical and Experimental Medicine*, 6(10), 979–984. <http://www.ncbi.nlm.nih.gov/pubmed/24260607>.
- Steinberg, L., Albert, D., Cauffman, E., Banich, M., Graham, S., & Woolard, J. (2008). Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: Evidence for a dual systems model. *Developmental Psychology*, 44(6), 1764–1778. <https://doi.org/10.1037/a0012955>.
- Sullivan, D. R., Forsberg, C. W., Golden, S. E., Ganzini, L., Dobscha, S. K., & Slatore, C. G. (2018). Incidence of suicide and association with palliative care among patients with advanced lung cancer. *Annals of the American Thoracic Society*, 15(11), 1357–1359. <https://doi.org/10.1513/AnnalsATS.201805-299RL>.
- Tang, S. T. (2000). Meanings of dying at home for Chinese patients in Taiwan with terminal cancer. *Cancer Nursing*, 23(5), 367–370. <https://doi.org/10.1097/00002820-200010000-00007>.
- Tang, G. X., Yan, P. P., Yan, C. L., Fu, B., Zhu, S. J., Zhou, L. Q., ... Lei, J. (2016). Determinants of suicidal ideation in gynecological cancer patients. *Psycho-Oncology*, 25(1), 97–103. <https://doi.org/10.1002/pon.3880>.
- Urban, D., Rao, A., Bressel, M., Neiger, D., Solomon, B., & Mileskin, L. (2013). Suicide in lung cancer: Who is at risk? *Chest*, 144(4), 1245–1252. <https://doi.org/10.1378/chest.12-2986>.
- Vyssoki, B., Gleiss, A., Rockett, I. R. H., Hackl, M., Leitner, B., Sonneck, G., & Kapusta, N. D. (2015). Suicide among 915303 Austrian cancer patients: Who is at risk? *Journal of Affective Disorders*, 175(2015), 287–291. <https://doi.org/10.1016/j.jad.2015.01.028>.
- Walker, J., & Sharpe, M. (2009). Depression care for people with cancer: A collaborative care intervention. *General Hospital Psychiatry*, 31(5), 436–441. <https://doi.org/10.1016/j.genhosppsych.2009.05.010>.
- Wang, B., An, X., Shi, X., & Zhang, J. (2017). Management of Endocrine Disease: Suicide risk in patients with diabetes: A systematic review and meta-analysis. *European Journal of Endocrinology*, 177(4), R169–R181. <https://doi.org/10.1530/EJE-16-0952>.
- Weinberger, M. I., Roth, A. J., & Nelson, C. J. (2009). Untangling the complexities of depression diagnosis in older cancer patients. *The Oncologist*, 14(1), 60–66. <https://doi.org/10.1634/theoncologist.2008-0147>.
- Wilson, K. G., Chochinov, H. M., Graham Skirko, M., Allard, P., Chary, S., Gagnon, P. R., ... Clinch, J. J. (2007). Depression and anxiety disorders in palliative cancer care. *Journal of Pain and Symptom Management*, 33(2), 118–129. <https://doi.org/10.1016/j.jpainsymman.2006.07.016>.
- Wong, G. L., Wong, V. W., Yuen, B. W., Tse, Y., Luk, H. W., Yip, T. C., ... Chan, H. L. (2020). An aging population of chronic hepatitis B with increasing comorbidities: A territory-wide study from 2000 to 2017. *Hepatology*, 71(2), 444–455. <https://doi.org/10.1002/hep.30833>.
- Yeung, S. L. (2015). Assessing and managing potentially suicidal patients: Practical guidelines for doctors. Retrieved from <https://csrp.hku.hk/wp-content/uploads/2015/06/DoctorGuidelines.pdf>.
- Yip, P. S. F., & Zheng, Y. (2020). The impact of suicide on life expectancy: A Hong Kong experience. *Crisis: The Journal of Crisis Intervention and Suicide Prevention*, 1–7. <https://doi.org/10.1027/0227-5910/a000695>.
- Yung, N. C. L., Wong, C. S. M., Chan, J. K. N., Or, P. C. F., Chen, E. Y. H., & Chang, W. C. (2020). Mortality in patients with schizophrenia admitted for incident ischemic stroke: A population-based cohort study. *European Neuropsychopharmacology*, 31, 152–157. <https://doi.org/10.1016/j.euro-neuro.2019.12.107>.
- Zaorsky, N. G., Zhang, Y., Tuanquin, L., Bluethmann, S. M., Park, H. S., & Chinchilli, V. M. (2019). Suicide among cancer patients. *Nature Communications*, 10(1), 1–7. <https://doi.org/10.1038/s41467-018-08170-1>.
- Zhong, B. L., Li, S. H., Lv, S. Y., Tian, S. L., Liu, Z. D., Li, X. B., ... Zhuo, C. J. (2017). Suicidal ideation among Chinese cancer inpatients of general hospitals: Prevalence and correlates. *Oncotarget*, 8(15), 25141–25150. <https://doi.org/10.18632/oncotarget.15350>.
- Zimmermann, C., Swami, N., Krzyzanowska, M., Leighl, N., Rydall, A., Rodin, G., ... Hannon, B. (2016). Perceptions of palliative care among patients with advanced cancer and their caregivers. *Canadian Medical Association Journal*, 188(10), E217–E227. <https://doi.org/10.1503/cmaj.151171>.