

# Conversion of rural abattoir wastes to an organic fertilizer and its application in the field cultivation of tomato in India

Malancha Roy<sup>1</sup>, Rimi Das<sup>1</sup>, Anupam Debsarcar<sup>2</sup>, Pradip Kumar Sen<sup>3</sup> and Joydeep Mukherjee<sup>1\*</sup>

<sup>1</sup>School of Environmental Studies, Jadavpur University, Kolkata 700 032, India

<sup>2</sup>Environmental Engineering Division, Department of Civil Engineering, Jadavpur University, Kolkata 700 032, India

<sup>3</sup>Department of Mathematics, Jadavpur University, Kolkata 700 032, India

\*Corresponding author: [joydeep\\_envstu@school.jdvu.ac.in](mailto:joydeep_envstu@school.jdvu.ac.in)

Accepted 29 June 2015; First published online 12 August 2015

Research Paper

## Abstract

Sophisticated capital intensive waste-recycling technologies are unviable in small rural abattoirs in India due to low volume of wastes (principally blood and rumen digesta) generated and lack of infrastructural facilities. We report recycling of slaughterhouse wastes as an organic fertilizer, ‘bovine-blood-rumen-digesta-mixture’ (BBRDM). Bovine blood and rumen digesta were mixed in 3:1 ratio in a metallic container, boiled and stirred continuously till the mixture was largely free of water. The mass was sun-dried for 3 days to obtain the final product. BBRDM was applied for field cultivation of tomato (*Lycopersicon esculentum* L., local variety ‘Patharkuchi’) in West Bengal state (India) during 2012–13 and 2013–14. We compared tomato yields obtained with BBRDM (N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 30.36:1:5.75) and conventional inorganic fertilizers [diammonium phosphate (DAP), N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 18:46:0 + potash, N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 0:0:44]. BBRDM was applied at a higher rate compared with DAP + potash to meet the farmers’ desire for enhanced yields. 75 kg ha<sup>-1</sup> was applied at the 2nd week while 150 kg ha<sup>-1</sup> was applied at the 8th week after transplantation. Yields (total fruit weight) obtained from BBRDM-treated plants were higher in comparison with DAP + potash-fertilized plants by 46–48% as the former supplied 2.5 times more nitrogen (N) than the latter. The partial factor productivity of DAP + potash was 73–76% higher than BBRDM. Conversely, as BBRDM was produced through local entrepreneurship from slaughterhouse wastes, the cost of this organic product would be expected to be much lower than the commercial inorganic fertilizer. Furthermore, application of BBRDM negates the environmental cost of treating slaughterhouse effluent. Considering the same cost of applying 225 kg fertilizer ha<sup>-1</sup>, higher yield with BBRDM should result in greater potential revenue for the farmer compared with yields with DAP + potash. The C/N ratio of BBRDM is 4.8, having relatively high N content. Accordingly, rapid release of plant-available N was observed in BBRDM-fertilized soils. The temporal increase in soil NH<sub>4</sub><sup>+</sup> may be attributed to lack of soil N immobilization. Local farmers are willing to accept the new fertilizer as a substitute for currently used chemical fertilizers.

**Key words:** waste, slaughterhouse, organic fertilizer, tomato, recycling

## Introduction

India is the world’s fourth largest beef producer and sixth largest domestic consumer. As of October 31, 2012, India is the largest exporter of beef globally and most of the exported beef is buffalo meat (Livestock and Poultry: World Markets and Trade). There are approximately 3600 legal, licensed slaughterhouses (Meat Sector – DSIR) and in parallel, numerous informal slaughterhouses in India (Hiranandani et al., 2010). The majority of the informal slaughterhouses that grew in the villages with traditional skill are more than 75 years old. They

lack modern infrastructural facilities, hygienic practices as well as organized systems of waste disposal (Slaughter House Waste and Dead Animals). Wastewater of small abattoirs in India is generally discharged without any treatment directly into local water bodies or municipal/local sewage systems. This poses a serious threat to human health as well as to surface water quality. Concentrated effluents and the constituent blood rapidly choke the disposal channels (Manual on Municipal Solid Waste Management, 2000). Blood, not recovered from waste streams, contributes to the high chemical oxygen demand (COD). Guidelines of the Indian

environmental regulatory authorities are seldom maintained in traditional rural Indian slaughterhouses. Slaughterhouse wastewater contains high concentrations of organic carbon (C), nitrogen (N) and phosphorus (P) (Zhan et al., 2009). The COD of a typical Indian rural slaughterhouse wastewater is in the range 4400–18,000 mg liter<sup>-1</sup>. The major soluble contaminant in abattoir wastewater is blood having COD 375,000 mg liter<sup>-1</sup> (Satyanarayan et al., 2005). The major solid waste of slaughterhouses is rumen digesta, a semi-solid mass made up of undigested to partially digested grass. The digesta in the rumen is not uniform, but stratified into gas and liquid particles of different sizes and densities (Awodun, 2008). Open dumping of rumen digesta by small informal rural slaughterhouses of India pose a health hazard to humans and adversely affect water sources. On average 20 buffaloes are slaughtered daily in one abattoir and 20 liters of blood as well as 20 kg of rumen digesta are obtained from one buffalo; it is expected that 400 liters of blood and 400 kg of rumen digesta will be generated daily.

Complicated operations to treat this extremely complex waste (blood and rumen digesta) make the wastewater treatment process exceedingly costly. As an estimation, the annual cost for disposal of 58,000 tons of beef slaughter waste in the USA would be approximately US\$10 million (Mittal, 2006). Correspondingly, 11,000 tons of waste are expected to be generated daily from rural Indian abattoirs. Thus, development of alternative processing methodologies for waste of small slaughterhouses (blood and rumen digesta) is imperative to manage the unhygienic waste disposal and to stop numerous informal small-scale slaughterhouses from going out of business. Therefore, this problem has socio-economic connotations as well. We intend to provide a feasible solution to the challenge rather than merely highlighting the problem. For efficient management of slaughterhouse waste bi-methanation and rendering systems have been suggested by the Government of India (Envis Newsletter, 2009). However the high capital costs (INR 38,000,000 for a high rate bi-methanation plant and INR 40,000,000 for a dry rendering plant, to recover animal fat) make them unaffordable for small slaughterhouse owners. Moreover, most small rural slaughterhouses are widely scattered, thus precluding centralized operation. Consequently, sophisticated and capital intensive technologies are unviable in small rural abattoirs due to the low volume of waste generated and lack of infrastructural facilities.

A more pragmatic approach would be to make use of cheap, simple-technology based processes that would be financially and technically viable as well as acceptable to the small slaughterhouses (Envis Newsletter, 2009). Waste products (blood and rumen digesta) may be composted (Pagans et al., 2006) or used as chicken feed (Ekunseitan et al., 2013). However, odorous ammonia emissions (Pagans et al., 2006) in the first recycling method and potential risk of microbial infections to

humans and poultry through the second method (Sapkota et al., 2007) may limit their applications. Soil application of slaughterhouse waste could be an advantageous ecological option to recycle these residues and to mitigate the adverse effects of pollution (Villar et al., 2004). The application of winery and distillery waste to soil as well as reuse of olive solid waste and compost as agricultural fertilizers were demonstrated methods for recycling the organic matter and nutrients (Bustamante et al., 2007; Killi and Kavdir, 2013).

Organic agriculture has the potential to meet the global food demand on one hand and to lessen the damaging environmental effects of conventional agriculture on the other (Badgley et al., 2007). Badgley et al. (2007) suggested application of organic manures in farming practice. Investigations on the application of abattoir waste as an organic fertilizer is therefore an appropriate proposition. Currently half of the nutrient demands for global agriculture are derived from natural and managed N fixation, organic recycling and atmospheric deposition (Smil, 2011). According to the 'Current World Fertilizers Trends and Outlook to 2016' of the Food and Agriculture Organization (FAO), among all Asian countries, the bulk of the increase of world demand for N, P and potash is expected in India (Food and Agricultural Organizations of United Nations, 2012). Essentially, taking into account the domestic and global demands for organic fertilizer, the small abattoir owners may be expected to emerge as important economic actors by selling the organic fertilizer produced through utilization of slaughterhouse waste and earning net profits. Agricultural application of the waste would circumvent the necessity of conventional treatment of slaughterhouse waste, which as observed previously is cost intensive. Therefore, use of slaughterhouse waste for agriculture can provide economic as well as environmental benefits.

Propositions to promote safe disposal of abattoir waste and recommendations to limit the methods of disposal to those internationally allowed were presented in a review (Adeyemi and Adeyemo, 2007). Adeyemi and Adeyemo (2007) suggested rendering of the waste which is a process of cooking that converts the semi-solid waste into a protein-rich substance that appears like sand or soil. Products derived from rendering permit storage for long periods of time. Rendering allows recycling of what would otherwise have been sizeable amounts of waste. Accordingly, we developed a cheap recycling methodology for slaughterhouse waste where a combination of bovine blood and rumen digesta, termed bovine-blood-rumen-digesta-mixture (BBRDM) was converted to a dry powder and applied as an organic fertilizer (Roy et al., 2013). The effectiveness of BBRDM was evaluated against diammonium phosphate (DAP) during pot cultivation of tomato, chili (*Capsicum annum* L.) and brinjal (*Solanum melongena* L.). BBRDM was applied twice at the 2nd and 6th weeks. Fruiting was earlier by 2 weeks in BBRDM-cultivated plants compared with

DAP-treated crops. Yields (total fruit weight) were greater in soils treated with the organic fertilizer. Limited studies have been conducted on the effect of slaughterhouse waste on crop productivity. Agricultural application of processed slaughterhouse waste composts and meat powder were investigated in a field experiment with maize (*Zea mays* L.), mustard (*Sinapis alba* L.) and triticale (*x Triticosecale* Wittm. ex A. Camus.) in Hungary (Ragályi and Kádár, 2012). The hazardous slaughterhouse waste became a nonhazardous product following heat treatment making soil application possible (Ragályi and Kádár, 2012). Slaughterhouse waste compost was applied to sugar beet (*Beta vulgaris* L. var. *saccharifera* Alef.) and spring barley (*Hordeum vulgare* L.) cultivations without any harmful effects on the plants (Petróczki, 2004). The present study, to the best of our knowledge, is the first on the application of abattoir waste for cultivation of tomato.

In continuation of the previous work, we now undertake the translation of the pot study to the agricultural field with the primary objective of evaluating the agronomic performance of BBRDM. To appraise its potential for enhancing tomato yields by substituting conventional chemical fertilizers, comparison of yields obtained by applying BBRDM and DAP + potash in field cultivation was done. Tomato is a comparatively sensitive indicator crop that responds to any mineral nutrient deficiencies, imbalances or phytotoxic properties associated with recycled waste as organic fertilizer (Killi and Kavdir, 2013). As of 2012, tomato featured among the seven most globally important food and agricultural commodities with a world production of 161,793,834 Mt and valued at 59,108,521 (International US\$1000) (FAOSTAT). India produced 17,874,420 tons of tomato and West Bengal (the state where this study was carried out) is the seventh largest producer (1,125,600 tons, 6.3% of total produce) in India (out of 32 states and union territories) as notified by the Agricultural and Processed Food Products Export Development Authority, Ministry of Commerce and Industry, Government of India (APEDA AGRIXCHANGE). Therefore, with regard to the domestic and international market, it should further be of significance to use recycled abattoir waste as an organic fertilizer for tomato cultivation.

## Materials and methods

### *Characterization of slaughterhouse waste, preparation and analysis of BBRDM*

Characterization of waste (bovine blood and rumen digesta) through measurement of wastewater parameters was done as described (Roy *et al.*, 2013). To obtain the C/N ratio of BBRDM, organic C of the fertilizer was determined according to the Walkley and Black's method and organic N was ascertained by estimation of total Kjeldahl N (Bustamante *et al.*, 2007). To prepare

BBRDM, fresh blood was collected in containers immediately following slaughtering of animals. Bovine blood and rumen digesta (ratio 3:1) were weighed in a metallic container. To fulfill the requirements of organic production, 'G.1.3. INDOCERT Organic Standards for Non-EU Country Operators, Version 2, 03/2012' (Indocert) was followed. Accordingly, sodium citrate (anti-coagulant) was not applied. The contents (blood and rumen digesta) were boiled for 90 min by placing the container on a coal-fired earthen stove. The mix (blood and rumen digesta) was continually stirred until the content was almost free of water. Finally the mass was dried under the sun for 3 days to obtain BBRDM. The product is a dry, solid, deep brown-colored powder and is easily spread (Supplementary Fig. 1). BBRDM was stored at ambient temperature during the period of study.

### *Experimental site and location*

The field study was carried out (2012–13) in Kuldiya Baganchi village of Magrahat block (latitude 22°15' 06.19"N, longitude 88°20'52.02"E, altitude 4 m) of South 24 Parganas district, West Bengal, a state in eastern India. In the following year (2013–14) the field experiment was done in Bankipur village of Magrahat block (latitude 22°14'25.24"N, longitude 88°22'42.09"E, altitude 4 m) of South 24 Parganas district, West Bengal state, India. Experimental plots were not cultivated earlier. The landscape had a flat topography without any slope. The total plot size was 500 m<sup>2</sup> (25 × 20 m<sup>2</sup>). In both seasons experiments were laid out as a completely randomized block design with six replications for each treatment. The plot was divided into 12 sub-plots, each covering 5.04 m<sup>2</sup> soil surfaces. The sub-plots were arranged randomly, six of which were treated with BBRDM and another six were treated with DAP + potash. Organic fertilizer (BBRDM) was regarded as the test treatment while chemical fertilizer was considered as control treatment. DAP + potash was selected as controls as these fertilizers are generally applied to tomato plants by local farmers. Tomato plants were planted on November 14, 2012 (season 1) and November 28, 2013 (season 2). Plant seedlings in all sub-plots were arranged in four rows, eight plants in a row at a distance of 40 cm from each other along the single row. The distance between rows in each single sub-plot was 60 cm and the distance between sub-plots was 150 cm. There were 32 tomato plants per sub-plot i.e., 20,000 plants ha<sup>-1</sup> (Agele *et al.*, 1999; Balemi, 2008; Sortino *et al.*, 2012).

### *Soil type, crop, time of study and weather conditions*

The greater part of West Bengal state (India) lies in the Gangetic plain and is covered with alluvium. The soil of the study site in South 24 Parganas district is considered as saline-alkali soils of tidal origin (Thampi and

Mukhopadhyay, 1975). Soil color was ascertained by comparing the soil sample with the standard Munsell Soil Color Charts (Torrent and Barron, 1993). Soil pH was measured potentiometrically (Singh et al., 1999). Soil particle size distribution was determined by the hydrometer method (Department of Sustainable Natural Resources). Soil type was determined following the Triangular Classification Chart of the US Bureau of Soils and Chemistry System (Murthy, 1992). The liquid limit of soil was ascertained following the Casagrande method and the plastic limit of soil was analyzed as per American Society for Testing and Materials (ASTM) Standard D 4318–10e1 (ASTM International, Active standard ASTM D4318). Soil density was determined according to ASTM D7263-09-Standard Test Methods for Laboratory Determination of Density (Unit Weight) of Soil Specimens (ASTM International, Active standard ASTM D7263). Soil water content was determined following ASTM D 2216-10-Standard Test Method for Laboratory Determination of Water (Moisture) Content of Soil and Rock by Mass (ASTM International, Active standard ASTM D2216).

Tomato (*Lycopersicon esculentum* L.), local variety 'Patharkuchi', tolerant to pests and diseases was selected for the field experiment. The study was done in the winter season from November 14, 2012 to February 28, 2013 (total 14 weeks in season 1) and from November 28, 2013 to March 15, 2014 (total 14 weeks in season 2). The mean maximum temperature during field cultivation (2012–13) was  $27.1 \pm 2.3^\circ\text{C}$ , whereas the mean minimum temperature was  $15.7 \pm 2.2^\circ\text{C}$ . The mean maximum humidity during the study was  $86.4 \pm 3.0\%$ , while the mean minimum humidity was  $40.8 \pm 5.8\%$ . The mean maximum temperature during field cultivation (2013–14) was  $28.2 \pm 1.3^\circ\text{C}$ , whereas the mean minimum temperature was  $16.2 \pm 1.9^\circ\text{C}$ . The mean maximum humidity during the study was  $85.7 \pm 2.5\%$ , while the mean minimum humidity was  $41.3 \pm 4.6\%$ . The total rainfall in season 1 was  $0.5 \pm 0.1$  mm and the total rainfall in season 2 was  $0.6 \pm 0.1$  mm.

### Nursery operations and raising of seedlings

Tomato seedlings were grown in nursery beds located near the main experimental plot. The soil was ploughed and seedbed was raised. Seeds were sowed at a depth of 0.5–1 cm, in straight rows at an interval of 5 cm allowing 2 cm gap between successive seeds and watered immediately. The top soil was used as a light cover over the seeds. The seedlings were shaded from direct sunlight and watered every alternate day.

### Preparation of main experimental field, transplantation of seedlings and staking

The experimental field was ploughed twice to make the soil to fine texture. The plots were shaped and ridges

were made for growing tomatoes on the top of the bed. In between the plots, furrows were made to serve as irrigation and drainage channels. The seedlings were thoroughly watered before transplanting to the field. At the end of the 4th week following sowing, seedlings of approximately equal heights and about four or five leaf stage were transplanted to the experimental plot in the late afternoon. The experimental plot was watered after transplantation. Furrow irrigation was used for watering plants and sprinkle irrigation when required. The plants were staked 2 weeks after transplantation using thin bamboo sticks of 2 m length. Pruning, weeding by hand pulling as well as other horticultural operations were done when required (Law-Ogbomo and Egharevba, 2009).

### Application of fertilizer

Well prepared BBRDM (N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 30.36:1:5.75) fertilizer was applied to the plants designated for BBRDM. The control plot was fertilized with DAP (N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 18:46:0) and potash (N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 0:0:44). BBRDM as well as DAP + potash (2:1) were applied in equal amounts. The first dose of  $75 \text{ kg ha}^{-1}$  was applied (evenly distributed along the furrows of each sub-plot by hand) at 2 weeks after transplantation of the plants to the main experimental field and the second dose  $150 \text{ kg ha}^{-1}$  was applied at the 8th week after transplantation (Gupta and Shukla, 1977; Balemi, 2008; Law-Ogbomo and Egharevba, 2009) Considering tomato to be an annual crop, the annual fertilizer application rates were  $68.31 \text{ kg N ha}^{-1}$ ,  $2.25 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ ,  $12.9 \text{ kg K}_2\text{O ha}^{-1}$  for BBRDM and  $27 \text{ kg N ha}^{-1}$ ,  $69 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ ,  $33 \text{ kg K}_2\text{O ha}^{-1}$  for DAP + potash.

### Plant diseases

During this study common pests like aphids, bollworms, leaf miners, thrips, whiteflies, spider mites and nematodes were not significantly noticed and pesticide application was not done. Wilts, blight, leaf spots and mildews were not significantly observed during the cultivation. Cultural methods such as removal of weeds, old leaves and branches as well as overshadowed lower leaves were done regularly by hand. Organic neem oil as pesticide was sprayed twice at the 3rd and 9th week after the transplantation of the tomato plants to the main experimental field as a preventive measure (Hinman et al., 2012).

### Measurement of yield parameters

Number of buds and flowers, numbers of fruits, average fruit weight and yield as mentioned in the Results section (Table 1) were measured or counted every week following transplantation of seedlings. Before harvesting the plants, numbers of ripened tomatoes per cluster were counted and the mean was calculated. Yield from each of the six sub-plots of both treatments were recorded

**Table 1.** Comparative analysis of flowering, fruiting and tomato yield during 14 weeks of cultivation. Yield per sub-plot represents cumulative weights during harvesting period. Details are provided in the Materials and methods section.

Plant growth parameters and yield	Season	Soil treatments	
		BBRDM	DAP + potash
Number of buds	S <sup>1</sup>	30 ± 3.56**	15 ± 2.39
	S <sup>2</sup>	34 ± 3.44**	18 ± 2.45
Number of flowers	S <sup>1</sup>	30 ± 3.56**	18 ± 2.86
	S <sup>2</sup>	39 ± 3.03**	20 ± 3.05
Number of fruits	S <sup>1</sup>	65 ± 4.93**	30 ± 4.59
	S <sup>2</sup>	70 ± 4.5**	40 ± 4.33
Average fruit weight(g)	S <sup>1</sup>	70 ± 4.39**	50 ± 3.54
	S <sup>2</sup>	80 ± 4.03**	58 ± 3.42
Yield per sub plot (kg per subplot)	S <sup>1</sup>	52 ± 6**	35 ± 2.8
	S <sup>2</sup>	57 ± 4.6**	39 ± 5.1
Partial factor productivity for N (PFP <sub>N</sub> ) (kg yield kg <sup>-1</sup> N applied)	S <sup>1</sup>	459 ± 4	809 ± 9**
	S <sup>2</sup>	515 ± 7	892 ± 8**

According to the Student's *t*-test \*\*signifies that the difference between two treatments are highly significant at 1% level; \*signifies that the difference between two treatments are significant at 5% level and no star signifies that the difference between two treatments are not significant. Error ranges indicate one SD from the mean. S<sup>1</sup>, season 1; S<sup>2</sup>, season 2.

separately and expressed in kg per sub-plot. Mean of these six values were calculated for each treatment. Amount of N supplied to each sub-plot (kg ha<sup>-1</sup>) was estimated considering applied fertilizer nutrient ratios (N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O 30.36:1:5.75) and a total application rate of 225 kg ha<sup>-1</sup> (a split-application of 75 + 150 kg fertilizer ha<sup>-1</sup>). Partial factor productivity for N (PFP<sub>N</sub>) in each sub-plot was calculated by dividing the yield obtained in a specific sub-plot (Y) by the N fertilizer rate (F<sub>N</sub>) (Zeng *et al.*, 2012; Deng *et al.*, 2014).

$$PFP_N = Y/F_N$$

PFP<sub>N</sub> values of six sub-plots of each treatment were used to calculate mean PFP<sub>N</sub>.

### Harvesting

First harvest was done at the 10th week after planting. The mature red fruits were harvested from all plants in each treatment. Subsequent harvests were done four additional times at 7 day intervals, for a total of five harvest events per growing season.

### Soil sampling method and measurement of available N

Soil samples were taken at 0–15 cm depth every week spanning the crop production cycle from start to end (Hartz and Johnstone, 2006; Tu *et al.*, 2006; Huang *et al.*, 2007; Walworth, 2010–2011). Six soil samples (six cores, 2.5 cm diameter) were air-dried, homogenized and sieved through 2.0 mm sieve to obtain one composite soil sample for BBRDM and DAP + potash treated plots (Sortino *et al.*, 2012; Zhang *et al.*, 2015). Soil samples

were collected randomly from the space beneath the plant canopy as well as the interspace area of the ridges in the central four rows of each sub-plot and 1 m (approximately) away from the ends of the rows. All samples were immediately stored (8–10°C) in sealed plastic bags and transported to laboratory. The easily mineralizable and available N (NH<sub>4</sub><sup>+</sup>N) of soil samples was estimated by the Subbiah and Asija method using alkaline KMnO<sub>4</sub> which oxidizes and hydrolyses the organic matter present in the soil. The liberated ammonia was condensed and absorbed in boric acid which was titrated against standard acid (Subbiah and Asija, 1956). This method does not measure soil nitrate and is suitable for Indian soils in general. Available soil N was measured before and every week during the cultivation. Analyses were performed in triplicate (Sortino *et al.*, 2012).

### Statistical analysis

All statistical analyses were done applying IBM SPSS Statistics 20 software. The Student's *t*-test compares one variable (yield parameter) between two groups (two treatments). Yield parameters of season 1 were compared using Student's *t*-test which compares the means of two treatments. Same parameters for season 2 were compared separately using Student's *t*-test. Significant difference of any parameter are noted in Table 1. Standard deviations were calculated for the mean values of all determinations and the *n* value is specified for each test in the relevant table/figure.

## Results and discussion

*Soil characteristics* The color of the soil in both field cultivation sites was grey to brownish, fine textured and pH

ranged from 7.0 to 8.0. The liquid limit was 40–45%, plastic limit was 21–23% and plasticity index was 19–23%. Soil particle size distribution was: sand 6–8%, silt 70–76% and clay 24–27%. The pre-tillage soil bulk density of the 0–15 cm surface layer was 1.83–1.91 g cm<sup>-3</sup> and the water content was 29.2–31.7%. The soil type was clayey silt.

### Yield parameters of tomato plants

Mean values of the yield parameters obtained in the two treatments are shown in Table 1. Student's *t*-test indicated that yield obtained from BBRDM-treated plants were higher by 48% (season 1) and 46% (season 2) in comparison with DAP + potash-fertilized plants. The enhancement in yield was, however, lower than that attained in our previous small-scale pot study (Roy et al., 2013). The onset of flowering and fruiting in BBRDM-treated plants were noticed to be 2 weeks in advance than in plants treated with DAP + potash.

To standardize results in a better way, yields based on N partial factor productivity (PFP<sub>N</sub>) were considered. BBRDM supplied 68.31 kg N ha<sup>-1</sup> while DAP + potash (2:1 mix) provided 27 kg N ha<sup>-1</sup>. Compared with DAP + potash, BBRDM supplied 2.5 times more N per unit mass of fertilizer applied. However, BBRDM had lower PFP<sub>N</sub> values than the inorganic fertilizer in both seasons (Table 1). PFP<sub>N</sub> of DAP + potash was 76% higher in S1 and 73% higher in S2 in comparison with PFP<sub>N</sub> of BBRDM. Notwithstanding, as BBRDM was produced through local entrepreneurship from slaughterhouse waste, the cost of this organic product would be expected to be much lower than the commercial inorganic fertilizer. Therefore, the farmer may expect higher revenue by purchasing 225 kg of BBRDM than buying the same amount of DAP + potash. Additionally, the lowered PFP may be compensated by other prospective advantages of applying organic fertilizer such as conservation of soil fertility, nutrient recycling, increased soil resiliency and overall environmental sustainability (Peigné et al., 2014). Furthermore, application of BBRDM negates the environmental cost of treating the slaughterhouse waste. The aim of the present investigation was to enhance the yield by replacing the conventional DAP + potash fertilization by organic fertilizer without altering the traditional fertilization practice. Hence, other rates of fertilizer application were not tested. This could be the subject of future field investigations.

According to the local farmers in our study area, deficient fertilization was a reason for lowered tomato yields. We aimed to enhance the yield obtained by application of the organic fertilizer in comparison with chemical fertilization. Therefore, a higher level of organic fertilizer was purposely selected, considering local farmers' desire for improved productivity and slaughterhouse owners' demand for utilization of the highly polluting effluent. Moreover, tomato is a high fertilizer

demanding crop (Colla et al., 2002; Riahi et al., 2009). Our reasoning is supported by reports of previous workers. There are several reports on enhanced tomato yields following increased N fertilization. Three NP fertilizers rates (urea + DAP) were applied to the cultivation of two tomato cultivars in Ethiopia (Balemi, 2008). Highest fertilization rate produced highest fruit yield. Similarly, two tomato cultivars were grown in Nigeria at three levels of NPK fertilizers (Law-Ogbomo and Egharevba, 2009), with an observed increase in marketable yield when fertilizer rate was enhanced. In another study, three levels of N and P were applied to tomato cultivation in India (Gupta and Shukla, 1977). Each successive increment in N dose resulted in significant increase in plant height and Bartlett's earliness index. Yield was highest at the highest level of N fertilization.

Although our study appears confounded through application of dissimilar levels of N, there are several studies on tomato cultivation where different levels of N were applied similar to our approach. In some experiments enhanced rate of organic fertilization improved tomato yields. For example, total as well as marketable yields of the two tomato cultivars in Tunisia were higher when COMP (mixed compost consisting of olive husk, horse manure, poultry manure, mixed compost extract and humus) and MIX (sheep manure, mixed compost, mixed compost extract and humus) were applied in comparison with humus application (Riahi et al., 2009). Riahi et al. (2009) attributed the improvement in yield to the increased compost rate, similar to our results (enhanced yield with increased N fertilization through BBRDM). The form and rate in which olive solid wastes were applied to soils in Turkey (Killi and Kavdir, 2013) (supplying different soil N levels, similar to our experimental design), played a major role in ascertaining the efficacy of the waste as organic fertilizers and in soil quality improvement.

On the other hand, there are reports where enhanced organic N fertilization did not increase tomato yields. In contrast to our results, tomato yields obtained from the organic and low input systems in the USA were not statistically different from those attained from the conventionally managed system, although the organic system possessed highest soil total C, N, soluble P, exchangeable Ca and K levels (Colla et al., 2002). In contrast to our results, a blend of compost-based fertilizer (CBF) and urea supplying higher amount of N did not consistently produce better tomato yield than the synthetic fertilizer alone (Taiwo et al., 2007) in Nigeria. The total N applied to organically grown tomato in Italy was higher than that applied to the conventionally grown crop. Despite increased fertilization and in contrast to our results, there was no statistical difference between the yields obtained in organic and inorganic systems (Campanelli and Canali, 2012). Three organic fertilizers, municipal bio-refuse (CVD), bio-organic fraction of CVD (SBO), insoluble residue of CVD (IOR) and a

**Table 2.** Comparison of tomato yields obtained with BBRDM with that attained using various organic fertilizers in other cultivation systems.

Amount of nitrogen (kg ha <sup>-1</sup> )	Type of organic fertilizer	Yield (t ha <sup>-1</sup> )	Yield (kg kg <sup>-1</sup> nitrogen applied)	Reference #
237	Poultry manure + cover crops	47–81.9	237–345	Colla <i>et al.</i> (2002)
175	CBF	4.35	25	Taiwo <i>et al.</i> (2007)
588	COMP	87.5 (F)	149 (F)	Riahi <i>et al.</i> (2009)
		82.5 (R)	140 (R)	
702	MIX	92.3 (F)	131 (F)	Riahi <i>et al.</i> (2009)
		87.7 (R)	125 (R)	
176	Off-farm organic fertilizers	62.3	354	Campanelli and Canali (2012)
68.31	BBRDM	33	483	Present study

CBF, compost-based fertilizer, derived from maize stover, cassava peels and poultry manure in ratio (1:3:1 v:v:v); COMP, mixed compost (50% olive husk + 30% horse manure + 20% poultry manure) + mixed compost extract + codahumus 20; MIX, sheep manure + mixed compost + mixed compost extract + codahumus 20; F, Firenze var.; R, Rio Grande var.

reference commercial product (RCP) were applied at different N levels, similar to our study of tomato greenhouse cultivation in Italy (Sortino *et al.*, 2012). SBO, providing the least amount of N to the soil, allowed precocious start of all flowering and fruiting stages and significantly improved plant growth and tomato fruit production compared with CVD and IOR. SBO performed better than RCP although the latter supplied more organic N to soil than the former. Water solubility of the organic fertilizer appeared to be more important than organic N content in enhancing yields (Sortino *et al.*, 2012). As evident from Table 2, BBRDM, applied in the present cultivation system, is a better organic fertilizer for tomato cultivation than other cultivation systems as higher tomato yield was attained supplying lower amount of N. BBRDM application did not require potassium or urea supplementation in the soil as was necessary in previous studies (Taiwo *et al.*, 2007; Javaria and Khan, 2011) by earlier workers with slaughterhouse compost as an organic fertilizer. Table 2 and these reports (Colla *et al.*, 2002; Taiwo *et al.*, 2007; Campanelli and Canali, 2012) indicate that apart from the N fertilization rate, there may be other important factors, for example water solubility of the fertilizer (Sortino *et al.*, 2012), C and N contents of the fertilizer that affect the final tomato yield. This aspect is discussed in the following section, in relation to our present experimental study.

Some researchers comparing the efficacy of organic and inorganic fertilizers during pot and field cultivation of tomatoes have applied equivalent levels of N in both systems. Higher yield of tomato in pots upon application of liquid residue of lipopeptide biosurfactant production compared with application of chemical fertilizers was reported from China (Zhu *et al.*, 2013). Similarly, higher tomato yield in pots by application of municipal solid waste compost and water extract tea in comparison with application of conventional NPK fertilizers was attained in Canada (Radin and Warman, 2011). On the other hand, during field studies, Bulluck and Ristaino (2002)


did not consistently report higher tomato yields in plots fertilized with cotton-gin trash, swine manure and ryevetch in the USA compared with chemical fertilizer. Similarly, higher tomato yields were not observed when compost made up by olive residues, sludge and straw was applied in Italy compared with a chemical fertilizer (Rinaldi *et al.*, 2007). The results of the field studies prompted us to increase the BBRDM application rate in anticipation of a higher yield.

It may be criticized that residual mineral N in soil enhanced the environmental risks due to runoff, leaching or gaseous N release. In India tomato is generally cultivated in the dry winter season hence the chance of excess agricultural runoff is negligible. The possible harmful effects of residual BBRDM are, however, insignificant compared with the catastrophic environmental degradation caused by the direct release of blood and rumen contents into water streams as is being practiced in the rural slaughterhouses. Although BBRDM application may substitute use of chemical fertilizers, further research is required to ascertain if residual soil N from BBRDM is higher than that remaining after chemical fertilizer application.

### Status of soil N during cultivation

Figure 1 illustrates that soil NH<sub>4</sub><sup>+</sup> levels in BBRDM were higher than DAP + potash logically due to the higher input of N. Soil NH<sub>4</sub><sup>+</sup> peaks were observed following the application of BBRDM and DAP + potash on the 2nd and 8th weeks (Fig. 1). The temporal increase in soil NH<sub>4</sub><sup>+</sup> may be attributed to lack of soil N immobilization as supported by reports of previous investigators.

In the trials (Marinari *et al.*, 2010) conducted in Italy, the N inputs in the organic system (guano) were greater than in the conventional one (NH<sub>4</sub>NO<sub>3</sub>), similar to our methodology. The peak of N release during N mineralization (short-term, 200 days) in soils supplied with guano was observed after 4 weeks (Marinari *et al.*,



**Figure 1.** Status of soil available nitrogen ( $\text{NH}_4^+$ ) during tomato cultivation. Fertilizers (BBRDM and DAP + potash) were added at the 2nd week ( $175 \text{ kg ha}^{-1}$ ) and at the 8th week ( $50 \text{ kg ha}^{-1}$ ). Available nitrogen ( $\text{NH}_4^+$ ) was measured as described by Subbiah and Asija (1956). For details of the method, please see the complete article, particulars of which are given in the list of references (Subbiah and Asija, 1956). Error ranges indicate one standard deviation from the mean ( $n = 3$ ). Fx1 Soil treated with DAP ( $\text{S}^1$ ), Fx2 Soil treated with BBRDM ( $\text{S}^1$ ), Fx3 Soil treated with DAP ( $\text{S}^2$ ), Fx4 Soil treated with BBRDM ( $\text{S}^2$ ).

2010), analogous to the N release peak observed upon BBRDM application (Fig. 1). Similar to our observation, soil  $\text{NH}_4^+$  content was drastically increased by slurry addition to soil in Canada (Chantigny et al., 2001). Upon addition of *Sesbania sesban* (Jacq.) W. Wight residues to soil in Kenya, the percentage of inorganic N in the soil peaked and then decreased in a 100 day study (Nyberg et al., 2002) similar to our observations (Fig. 1). Hartz et al. (2010) applied fishery waste, feather meal and seabird guano to agricultural soils of the USA and the products mineralized 60–80% of N within 4–8 weeks, analogous to our results. On the other hand, some investigators noted decrease in soil-available N upon addition of organic fertilizers. For most of the manure treatments to agricultural soils (Bechini and Marino, 2009), immobilization of mineral N occurred in the first weeks, followed by slow remineralization of immobilized N in a study carried out in Italy. This observation contrasts our results (Fig. 1). The report (Bechini and Marino, 2009) was, however, supported by other investigators. Albuquerque et al. (2012) noted initial decreases in inorganic-N during short-term study (56 days) which was ascribed to microbial immobilization. In a study conducted for 120 days in Spain (Bustamante et al., 2007), authors concluded that soil immobilization processes, upon application of winery and distillery waste could have succeeded over N losses by ammonia volatilization or by denitrification. Agricultural soils fertilized with Biolyzer (derived from

plant materials) had appreciably lower N availability (Hartz et al., 2010).

The amount of soil mineral N at the end of the incubation was highly correlated with the total N content of the fertilizers. The C/N ratio of the readily degradable fraction of the fertilizers had an important effect in controlling soil N dynamics in the short-medium period (Martín-Olmedo and Rees, 1999; Bustamante et al., 2007; Galvez et al., 2012). BBRDM having C/N ratio of 4.8 is a high-quality organic fertilizer. High-quality organic fertilizers (Class I) have  $>2.5\%$  N (Gentile et al., 2011) and these fertilizers resulted in net N mineralization alike BBRDM in our study. Lower ratios signify a higher quality manure and faster mineralization of nutrients (Nyberg et al., 2002). Conversely, application of manure with high C/N ratios could cause immobilization of N (Nyberg et al., 2002), which was not observed in our experiment. The C/N ratio of the organic fertilizer was implicated by previous investigators in determining whether soil N immobilization or mineralization would occur upon addition of the organic fertilizer. For example, the C/N ratios of the organic fertilizers applied (Bustamante et al., 2007) that caused immobilization varied from 9.8 to 18.2. Similarly the fertilizers of Bechini and Marino (2009) possessed high C/N ratios. Fishery waste, feather meal and seabird guano showing mineralization (Hartz et al., 2010) had lower C/N ratios compared with Biolyzer (Hartz et al., 2010) showing immobilization. In yet another study authors observed mineralization and nitrification of organic N from sheep wool waste which was richer in organic N (over 5%) than manure and compost (Vončina and Mihelič, 2013). The C/N ratio of tithonia was lower than maize and thus had a quicker mineralization rate (Gentile et al., 2011). High-N waste products from agricultural and fishery industries (for example fish powder) demonstrated N mineralization and were the most practical alternative for in-season N fertilization during vegetable production in the USA (Hartz and Johnstone, 2006). The  $\text{NH}_4^+$  concentration of soils fertilized with pine biochar (labile C content less than 1% of total C) peaked immediately following application without exhibiting immobilization (Angst et al., 2014).

Generally, the present field observations were supportive of our previous pot study. The preceding pot experiment (Vončina and Mihelič, 2013), supported the authors' field observations where sheep wool proved to be a good source of N. Similarly, Galvez et al. (2012) observed that in both field as well as laboratory studies a significant fraction of available N was released in the first 30 days after application.

## Conclusions

Through this study, the exceedingly problematic slaughterhouse waste was converted to an organic fertilizer



which was gainfully utilized for the production of tomatoes. This work embodies a practical and feasible method for recycling of organic waste in agriculture. The produce of tomatoes was appreciable and the local farmers are willing to accept the new fertilizer as a substitute for currently used chemical fertilizers. Soil N mineralization kinetics in BBRDM-treated soil was supportive of the enhanced plant N availability. This waste management approach will make the small-scale slaughter houses compliant with environmental norms. This appropriate, affordable and efficient methodology for recycling of waste generated by small abattoirs can be a model in many developing countries with similar dispersed small-scale traditional slaughtering practices with no hygienic waste disposal system. However, the safety of the fertilizer in terms of presence of pathogens and heavy metals as well as fruit quality in terms of lycopene content, nitrate, heavy metal levels and potential toxicity should be evaluated before large-scale application of BBRDM is practiced. This aspect will be the subject of our next communication.

## Supplementary Material

For supplementary material accompanying this paper, visit <http://dx.doi.org/10.1017/S1742170515000289>

**Acknowledgements.** Financial support to MR through DST Women Scientist B Scheme (SSD/SS/009/2011) and from Jadavpur University is thankfully acknowledged. Authors thank Anarul Gazi for supplying blood and rumen digesta. Valuable suggestions of Dr. Joyashree Roy, Coordinator Global Change Programme and Professor of Economics, Jadavpur University are appreciated. Authors are thankful to the anonymous reviewers whose comments vastly improved the manuscript.

## References

- Adeyemi, N.I.G. and Adeyemo, O.K. 2007. Waste management practices at the Bodija abattoir. *International Journal of Environmental Studies* 64:71–82. doi: 10.1080/00207230601124989.
- Agele, S.O., Iremiren, G.O., and Ojeniyi, S.O. 1999. Effects of plant density and mulching on the performance of late-season tomato (*Lycopersicon esculentum*) in southern Nigeria (Article). *Journal of Agricultural Science* 133:397–402. doi: 10.1017/S0021859699006942.
- Alburquerque, J.A., de la Fuente, C., and Bernal, M.P. 2012. Chemical properties of anaerobic digestates affecting C and N dynamics in amended soils. *Agriculture, Ecosystems and Environment* 160:15–22. doi: 10.1016/j.agee.2011.03.007.
- Angst, T.E., Six, J., Reay, D.S., and Sohi, S.P. 2014. Impact of pine chip biochar on trace greenhouse gas emissions and soil nutrient dynamics in an annual ryegrass system in California. *Agriculture, Ecosystems and Environment* 191: 17–26. doi: 10.1016/j.agee.2014.03.009.

- APEDA AGRIXCHANGE. Indian Production of Tomato (HSCODE-07020000) National Horticulture Board (NHB) Year: 2012–13. APEDA AGRIXCHANGE. Available at Web site [http://agriexchange.apeda.gov.in/India%20Production/India\\_Productions.aspx?hscod=07020000](http://agriexchange.apeda.gov.in/India%20Production/India_Productions.aspx?hscod=07020000) (verified 18 March 2015).
- ASTM International, Active standard ASTM D2216. Standard test methods for laboratory determination of water (moisture) content of soil and rock by mass. ASTM International, Active standard ASTM D2216. Available at Web site <http://www.astm.org/Standards/D2216.htm> (verified 18 March 2015).
- ASTM International, Active standard ASTM D4318. Standard test methods for liquid limit, plastic limit and plasticity index of soils. ASTM International, Active standard ASTM D4318. Available at Web site <http://www.astm.org/Standards/D4318.htm> (verified 16 March 2015).
- ASTM International, Active standard ASTM D7263. Standard test methods for laboratory determination of density (unit weight) of soil specimens. ASTM International, Active standard ASTM D7263. Available at Web site <http://www.astm.org/Standards/D7263.htm> (verified 16 March 2015).
- Awodun, M.A. 2008. Effect of nitrogen released from rumen digesta and cowdung on soil and leaf nutrient content of Gboma (*Solanum macrocarpon* L.). *Journal of Applied Biosciences* 7:202–206. ISSN 1997–5902: <http://www.biosciences.elewa.org> (verified 18 March 2015).
- Badgley, C., Moghtader, J., Quintero, E., Zakem, E., Chappell, M.J., Avile's-Va'zquez, K., Samulon, A., and Perfecto, I. 2007. Organic agriculture and the global food supply. *Renewable Agriculture and Food Systems* 22:86–108. doi: 10.1017/S1742170507001640.
- Balemi, T. 2008. Response of tomato cultivars differing in growth habit to nitrogen and phosphorus fertilizers and spacing on vertisol in Ethiopia. *Acta Agriculturae Slovenica* 91:103–119. doi: 10.2478/v10014-008-0011-8.
- Bechini, L. and Marino, P. 2009. Short-term nitrogen fertilizing value of liquid dairy manures is mainly due to ammonium. *Soil Science Society of America Journal* 73:2159–2169. doi: 10.2136/sssaj2008.0217.
- Bulluck, L.R., III and Ristaino, J.B. 2002. Effect of synthetic and organic soil fertility amendments on southern blight, soil microbial communities, and yield of processing tomatoes. *Phytopathology* 92:181–189. doi: 10.1094/PHYTO.2002.92.2.181.
- Bustamante, M.A., Pe'rez-Murcia, M.D., Paredes, C., Moral, R., Pe'rez-Espinosa, A., and Moreno-Caselles, J. 2007. Short-term carbon and nitrogen mineralisation in soil amended with winery and distillery organic wastes. *Bioresource Technology* 98:3269–3277. doi: 10.1016/j.biortech.2006.07.013.
- Campanelli, G. and Canali, S. 2012. Crop production and environmental effects in conventional and organic vegetable farming systems: The case of a long-term experiment in Mediterranean conditions (Central Italy). *Journal of Sustainable Agriculture* 36:599–619. doi: 10.1080/10440046.2011.646351.
- Chantigny, M.H., Rochette, P., and Angers, D.A. 2001. Short-term C and N dynamics in a soil amended with pig slurry and barley straw: A field experiment. *Canadian Journal of Soil Science* 81:131–137.
- Colla, G., Mitchell, J.P., Poudel, D.D., and Temple, S.R. 2002. Changes of tomato yield and fruit elemental composition

- in conventional low input and organic systems. *Journal of Sustainable Agriculture* 20:53–67. doi: 10.1300/J064v20n02\_07.
- Deng, F., Wang, L., Ren, W.-J., and Mei, X.-F.** 2014. Enhancing nitrogen utilization and soil nitrogen balance in paddy fields by optimizing nitrogen management and using polyaspartic acid urea. *Field Crops Research* 169:30–38. doi: 10.1016/j.fcr.2014.08.015.
- Department of Sustainable Natural Resources.** Soil Survey Standard Test Method Particle Size Analysis. Department of Sustainable Natural Resources. Available at Web site <http://www.environment.nsw.gov.au/resources/soils/testmethods/psa.pdf> (verified 18 March 2015).
- Ekunseitan, D.A., Balogun, O.O., Sogunle, O.M., Yusuf, A.O., Ayoola, A.A., Ebgeyale, L.T., Adeyemi, O.A., Allison, I.B., and Iyanda, A.I.** 2013. Health status of birds fed diets containing three differently processed discarded vegetable-bovine blood-rumen content mixtures. *Pakistan Journal of Biological Sciences* 16:325–331. doi: 10.3923/pjbs.2013.325.331.
- Envis Newsletter.** 2009. ENVIS/PPCC/NL-14. Vol- 4. No. 4. Available at Web site <http://dste.puducherry.gov.in/envis-new/fourteenth-NL-JAN-MARCH-2009.pdf> (verified 18 March 2015).
- FAOSTAT.** Food and Agricultural Organizations of United Nations. Available at Web site <http://faostat.fao.org/site/339/default.aspx> (verified 16 March 2015).
- Food and Agricultural Organizations of United Nations.** 2012. Current World Fertilizer Trends and Outlook to 2016. Food and Agricultural Organizations of United Nations, Rome. Available at Web site <ftp://ftp.fao.org/ag/agp/docs/cwfto16.pdf> (verified 16 March 2015).
- Galvez, A., Sinicco, T., Cayuela, M.L., Mingorance, M.D., Fornasier, F., and Mondini, C.** 2012. Short term effects of bioenergy by-products on soil C and N dynamics, nutrient availability and biochemical properties. *Agriculture, Ecosystems and Environment* 160:3–14. doi: 10.1016/j.agee.2011.06.015.
- Gentile, R., Vanlauwe, B., Chivenge, P., and Six, J.** 2011. Trade-offs between the short- and long-term effects of residue quality on soil C and N dynamics. *Plant and Soil* 338:159–169. doi: 10.1016/j.agee.2011.06.015.
- Gupta, A. and Shukla, V.** 1977. Response of tomato (*Lycopersicon esculentum* mill.) to plant spacing, nitrogen, phosphorus and potassium fertilization. *Indian Journal of Horticulture* 34:270–276.
- Hartz, T.K. and Johnstone, P.R.** 2006. Nitrogen availability from high-nitrogen containing organic fertilizers. *HortTechnology* 16:39–42. Available at Web site <http://www.ucanr.org/sites/nm/files/76755.pdf> (verified 16 March 2015).
- Hartz, T.K., Smith, R., and Gaskell, M.** 2010. Nitrogen availability from liquid organic fertilizers. *HortTechnology* 20: 169–172.
- Hinman, T., Baier, A., and Riesselman, L.** 2012. Organic tomato production. National Sustainable Agriculture Information Service. A project of the National Center for Appropriate Technology. 1-800-346-9140. Available at Web site <http://www.attra.ncat.org> (verified 16 March 2015).
- Hiranandani, A., McCall, R.M., and Shaheen, S.** 2010. India's Holy Cash Cow. *New Internationalist Issue* 434. Available at Web site <http://newint.org/columns/currents/2010/07/01/india-leather-cows/> (verified 16 March 2015).
- Huang, B., Sun, W., Zhao, Y., Zhu, J., Yang, R., Zou, Z., Ding, F., and Su, J.** 2007. Temporal and spatial variability of soil organic matter and total nitrogen in an agricultural ecosystem as affected by farming practices. *Geoderma* 139: 336–345. doi: 10.1016/j.geoderma.2007.02.012.
- Indocert.** Available at Web site <http://www.indocert.org/index.php/en/> (verified 16 March 2015).
- Javaria, S. and Khan, M.Q.** 2011. Impact of integrated nutrient management on tomato yield quality and soil environment. *Journal of Plant Nutrition* 34:140–149. doi: 10.1080/01904167.2011.531605.
- Killi, D. and Kavdir, Y.** 2013. Effects of olive solid waste and olive solid waste compost application on soil properties and growth of *Solanum lycopersicum*. *International Biodeterioration and Biodegradation* 82:157–165. doi: 10.1016/j.ibiod.2013.03.004.
- Law-Ogbomo, K.E. and Egharevba, R.K.A.** 2009. Effects of planting density and NPK fertilizer application on yield and yield components of tomato (*Lycopersicon esculentum* mill) in forest location. *World Journal of Agricultural Sciences* 5:152–158. Available at Web site [http://www.idosi.org/wjas/wjas5\(2\)/4.pdf](http://www.idosi.org/wjas/wjas5(2)/4.pdf) (verified 18 March 2015).
- Livestock and Poultry: World Markets and Trade.** United States Department of Agriculture Foreign Agricultural Service Approved by the World Agricultural Outlook Board/ USDA November 2013. Available at Web site <http://www.thepigsite.com/reports/?id=985> (verified 16 March 2015).
- Manual on Municipal Solid Waste Management.** 2000. Available at Web site [http://www.indiawaterportal.org/sites/indiawaterportal.org/files/Manual%20on%20municipal%20solid%20waste%20management\\_%20MoUD\\_GOI\\_2000.pdf](http://www.indiawaterportal.org/sites/indiawaterportal.org/files/Manual%20on%20municipal%20solid%20waste%20management_%20MoUD_GOI_2000.pdf) (verified 18 March 2015).
- Marinari, S., Lagomarsino, A., Moscatelli, M.C., Di Tizio, A., and Campiglia, E.** 2010. Soil carbon and nitrogen mineralization kinetics in organic and conventional three-year cropping systems. *Soil and Tillage Research* 109:161–168. doi: 10.1016/j.still.2010.06.002.
- Martin-Olmedo, P. and Rees, R.M.** 1999. Short-term N availability in response to dissolved-organic-carbon from poultry manure, alone or in combination with cellulose. *Biology and Fertility of Soils* 29:386–393.
- Meat Sector – DSIR.** Available at Web site [http://www.dsir.gov.in/reports/ittp\\_tedo/agro/AF\\_Animals\\_Meat\\_Intro.pdf](http://www.dsir.gov.in/reports/ittp_tedo/agro/AF_Animals_Meat_Intro.pdf) (verified 18 March 2015).
- Mittal, G.S.** 2006. Treatment of wastewater from abattoirs before land application – a review. *Bioresource Technology* 97:1119–1135. doi: 10.1016/j.biortech.2004.11.021.
- Murthy, V.N.S.** 1992. A Textbook of Soil Mechanics and Foundation Engineering - Third revised and enlarged edition. Kripa Technical Consultants, Bangalore.
- Nyberg, G., Ekblad, A., Buresh, R., and Högborg, P.** 2002. Short-term patterns of carbon and nitrogen mineralization in a fallow field amended with green manures from agroforestry trees. *Biology and Fertility of Soils* 36:18–25. doi: 10.1007/s00374-002-0484-2.
- Pagans, E., Barrena, R., Font, X., and Sánchez, A.** 2006. Ammonia emissions from the composting of different organic wastes. Dependency on process temperature. *Chemosphere* 62:1534–1542. doi: 10.1016/j.chemosphere.2005.06.044.
- Peigné, J., Casagrande, M., Payet, V., David, C., Sans, F.X., Blanco-Moreno, J.M., Cooper, J., Gascoyne, K., Antichi, D., Barberi, P., Bigongiali, F., Surböck, A., Kranzler, A., Beeckman, A., Willekens, K., Luik, A.,**

- Darja, M., Grosse, M., Heß, J., Clerc, M., Dierauer, H., and Mäder, P.** 2014. How organic farmers practice conservation agriculture in Europe. *Renewable Agriculture and Food Systems*, 1–14. doi: 10.1017/S1742170514000477.
- Petróczyki, F.** 2004. Effect of sewage sludge and slaughterhouse waste compost on plant growth. *Acta Agronomica Hungarica* 52:253–261. doi: 10.1556/AAgr.52.2004.3.7.
- Radin, A.M. and Warman, P.R.** 2011. Effect of municipal solid waste compost and compost tea as fertility amendments on growth and tissue element concentration in container-grown tomato. *Communications in Soil Science and Plant Analysis* 42:1349–1362. doi: 10.1080/00103624.2011.571742.
- Ragályi, P. and Kádár, I.** 2012. Effect of organic fertilizers made from slaughterhouse wastes on yield of crops. *Archives of Agronomy and Soil Science* 58(sup 1):S122–S126. doi: 10.1080/03650340.2012.695863.
- Riahi, A., Hdidier, C., Sanaa, M., Tarchoun, N., Ben Kheder, M., and Guezal, I.** 2009. The influence of different organic fertilizers on yield and physico-chemical properties of organically grown tomato. *Journal of Sustainable Agriculture* 33: 658–673. doi: 10.1080/10440040903073800.
- Rinaldi, M., Convertini, G., and Elia, A.** 2007. Organic and mineral nitrogen fertilization for processing tomato in southern Italy. *Acta Horticulturae* 758:241–248.
- Roy, M., Karmakar, S., Debsarcar, A., Sen, P.K., and Mukherjee, J.** 2013. Application of rural slaughterhouse waste as an organic fertilizer for pot cultivation of solanaceous vegetables in India. *International Journal of Recycling Organic Waste in Agriculture* 2:1–11. doi: 10.1186/2251-7715-2-6.
- Sapkota, A.R., Lefferts, L.Y., McKenzie, S., and Walker, P.** 2007. What do we feed to food-production animals? A review of animal feed ingredients and their potential impacts on human health. *Environmental Health Perspectives* 115:663–670.
- Satyanarayan, S., Ramakant, and Vanerkar, A.P.** 2005. Conventional approach for abattoir wastewater treatment. *Environmental Technology* 26:441–447. doi: 10.1080/09593332608618554.
- Singh, D., Chhonkar, P.K., and Pandey, R.N.** 1999. *Soil, Plant and Water Analysis - A Method Manual*. IARI, New Delhi.
- Slaughter House Waste and Dead Animals.** Available at Web site <http://localbodies.up.nic.in/swm/chap5.pdf> (verified 16 March 2015).
- Smil, V.** 2011. Nitrogen cycle and world food production. *World Agriculture* 2:9–1. Available at Web site <http://www.vaclavsmil.com/wp-content/uploads/docs/smil-article-worldagriculture.pdf> (verified 18 March 2015).
- Sortino, O., Dipasquale, M., Montoneri, E., Tomasso, L., Perrone, D.G., Vindrola, D., Negre, M., and Piccone, G.** 2012. Refuse derived soluble bio-organics enhancing tomato plant growth and productivity. *Waste Management* 32:1792–1801. doi: 10.1016/j.wasman.2012.04.020.
- Subbiah, B.V. and Asija, G.L.** 1956. A rapid procedure for determination of available nitrogen in soils. *Current Science* 25: 259–260.
- Taiwo, L.B., Adediran, J.A., and Sonubi, O.A.** 2007. Yield and quality of tomato grown with organic and synthetic fertilizers. *International Journal of Vegetable Science* 13:5–19. doi: 10.1300/J512v13n02\_02.
- Thampi, C.J. and Mukhopadhyay, A.K.** 1975. Studies on the structural stability of some soils of West Bengal. *Proceedings of the Indian National Science Academy - Part B: Biological Sciences* 41:47–53.
- Torrent, J. and Barron, V.** 1993. Laboratory measurement of soil color: Theory and Practice. Available at Web site <http://www.uco.es/organiza/departamentos/decrat/pdf/edaf/Soil%20Colour%201993.pdf> (verified 16 March 2015).
- Tu, C., Ristaino, J.B., and Hu, S.** 2006. Soil microbial biomass and activity in organic tomato farming systems: Effects of organic inputs and straw mulching. *Soil Biology and Biochemistry* 38:247–255. doi: 10.1016/j.soilbio.2005.05.002.
- Villar, M.C., Petrikova, V., Díaz-Raviña, M., and Carballas, T.** 2004. Recycling of organic wastes in burnt soils: Combined application of poultry manure and plant cultivation. *Waste Management* 24:365–370. doi: 10.1016/j.wasman.2003.09.004.
- Vončina, A. and Mihelič, R.** 2013. Sheep wool and leather waste as fertilizers in organic production of asparagus (*Asparagus officinalis* L.). *Acta Agriculturae Slovenica* 101:191–200. doi: 10.2478/acas-2013-0015.
- Walworth, J.L.** 2010–2011. *Soil Sampling and Analysis AZ1412*. Arizona Cooperative Extension. College of Agriculture and Life Sciences. The University of Arizona. Available at Web site <http://extension.arizona.edu/sites/extension.arizona.edu/files/pubs/az1412.pdf> (verified 16 March 2015).
- Zeng, X., Han, B., Xu, F., Huang, J., Cai, H., and Shi, L.** 2012. Effects of modified fertilization technology on the grain yield and nitrogen use efficiency of midseason rice. *Field Crops Research* 137:203–212. doi: 10.1016/j.fcr.2012.08.012.
- Zhan, X., Healy, M.G., and Li, J.** 2009. Nitrogen removal from slaughterhouse wastewater in a sequencing batch reactor under controlled low DO conditions. *Bioprocess and Biosystems Engineering* 32:607–614. doi: 10.1007/s00449-008-0283-8.
- Zhang, A., Luo, W., Sun, J., Xiao, H., and Liu, W.** 2015. Distribution and uptake pathways of organochlorine pesticides in greenhouse and conventional vegetables. *Science of the Total Environment* 505:1142–1147. doi: 10.1016/j.scitotenv.2014.11.023.
- Zhu, Z., Zhang, F., Wang, C., Ran, W., and Shen, Q.** 2013. Treating fermentative residues as liquid fertilizer and its efficacy on the tomato growth. *Scientia Horticulturae* 164: 492–498. doi: 10.1016/j.scienta.2013.10.008.