

Emotion regulation as a transdiagnostic factor in the development of internalizing and externalizing psychopathology: Current and future directions

AMELIA ALDAO,^a DYLAN G. GEE,^b ANDRES DE LOS REYES,^c AND ILANA SEAGER^a

^aOhio State University; ^bYale University; and ^cUniversity of Maryland at College Park

Abstract

In response to rapidly growing rates of comorbidity among psychiatric disorders, clinical scientists have become interested in identifying transdiagnostic processes that can help explain dysfunction across diagnostic categories (e.g., Kring & Sloan, 2009). One factor that has received a great deal of attention is that of emotion regulation, namely, the ability to modulate the intensity and/or duration of emotional states (e.g., Cicchetti, Ackerman, & Izard, 1995; Gross, 1998). Recent theoretical and empirical work has begun to emphasize the role that emotion regulation plays in the temporal comorbidity between internalizing and externalizing conditions (e.g., Aldao & De Los Reyes, 2015; De Los Reyes & Aldao, 2015; Drabick & Kendall, 2010; Jarrett & Ollendick, 2008; Patrick & Hajcak, 2016). However, close inspection of this work reveals two very pertinent areas of growth: (a) this literature is characterized by mixed findings that are likely explained, in part, by methodological heterogeneity; and (b) emotion regulation tends to be studied in relatively narrow terms. To address these issues, we provide a series of recommendations for facilitating cross-study comparisons and leveraging multifaceted approaches to studying emotion regulation processes within a developmental psychopathology framework. We hope that our perspective can enhance the organization and growth of this very important area of inquiry, and ultimately result in more effective prevention and treatment programs.

Prior to the publication of the third edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-III; American Psychiatric Association, 1987), approaches to diagnosing psychopathology assumed that a small number of processes (e.g., neuroses and conditioning) explained most forms of dysfunction (e.g., Beauchaine & McNulty, 2013; Nolen-Hoeksema & Watkins, 2011). Thus, the DSM included diagnostic hierarchies that limited the extent to which an individual could receive more than one diagnosis at a given time. For example, in DSM-III, anxiety disorders and depression were thought to share a common etiology and thus could not be diagnosed in the same individual at the same time (e.g., First, 2005). However, in the 1980s, research on diagnosis and classification shifted toward carving psychopathology into more unique conditions defined by specific behaviors that reflected distinct etiologies (e.g., Beauchaine & Klein, in press; Nolen-Hoeksema & Watkins, 2011). As a result, most diagnostic hierarchies were lifted with the publication of DSM-III (American Psychiatric Association, 1987). In addition, each subsequent edition of the DSM continued to define an increasingly larger number of diagnostic categories, each reflecting more narrow forms of pathology (e.g., Frances, 2013).

With this shift, it became necessary to more frequently diagnose patients with multiple conditions in order to properly characterize their psychological dysfunction. This led to a rampant increase in the rates of comorbidity, to the point that most patients are now diagnosed with many disorders at a time (e.g., Brown, Campbell, Lehman, Grisham, & Mancill, 2001; Kessler, Chiu, Demler, & Walters, 2005), and those who receive one psychiatric diagnosis are likely to receive another one over time (e.g., Kessler et al., 2011). That is, comorbidity, both concurrent and longitudinal, has become the norm in diagnosis and classification (e.g., Barlow, Sauer-Zavala, Carl, Bullis, & Ellard, 2014; Caspi et al., 2014). In this respect, a quick search on PsycINFO reveals that since 1987, the number of articles with the term “comorbid” in their title or abstract has grown impressively from the low hundreds in the early 1990s to over 2500 in 2015 (see Figure 1).

This inflated comorbidity presents a substantial barrier for understanding, and preventing and treating, psychological dysfunction because it conflates three distinct processes: pure comorbidity (i.e., true functional relationships among distinct forms of psychopathology), artifactual comorbidity (i.e., splitting one disorder into two or more), and spurious comorbidity (i.e., stemming from shared diagnostic criteria among disorders; e.g., Beauchaine & McNulty, 2013; First, 2005). One way in which the field has sought to isolate patterns of true comorbidity is by adopting a *transdiagnostic approach* (e.g., Aldao, 2013; Barlow, Allen, & Choate, 2004;

Address correspondence and reprint requests to: Amelia Aldao, Department of Psychology, Ohio State University, 1835 Neil Avenue, Columbus, OH 43210; E-mail: aldao.1@osu.edu.




Figure 1. (Color online) The number of articles with the term “comorbid*” in their title or abstract by year, based on a PsycINFO search in April 2016.

Egan, Wade, & Shafran, 2011; Ehrenreich-May, Queen, Bilek, Remmes, & Marciel, 2013; Ehring & Watkins, 2008; Harvey, Watkins, Mansell, & Shafran, 2004; Kring & Sloan, 2009). The primary goal of this approach is to identify which dysfunctional processes (e.g., cognition, emotion, or physiology) cut across extant diagnostic categories (i.e., are transdiagnostic) and which do not (i.e., are disorder specific). Consequently, it relies on traditionally defined symptom-based diagnostic categories (e.g., DSM criteria). It is important to note that a given process might be transdiagnostic for two conditions, but disorder specific in relation to others (e.g., positive affect is blunted in depression and social anxiety disorder, and within normative levels in the rest of the anxiety disorders; e.g., Brown, 2007). Thus, the transdiagnostic label needs to be understood within relative, rather than absolute, terms.

In line with the transdiagnostic approach, the National Institute of Mental Health has developed the Research Domain Criteria framework (Insel et al., 2010), which proposes dimensional models to capture dysfunction across multiple units of analyses (e.g., subjective reports, physiological reactivity, neural circuitry, and genetic predispositions) and functional domains (e.g., negative valence system and social processes; for a review see Sanislow et al., 2010). These domains represent processes that “cut across” existing diagnostic categories. Therefore, knowledge from Research Domain Criteria informed research might further clarify the mechanisms underlying comorbidity.

One construct that has gained an impressive amount of attention within the transdiagnostic approach is *emotion reg-*

ulation, that is, the processes by which people modify the intensity and/or duration of their emotions in response to contextual demands (e.g., Cicchetti et al., 1995; Cole, Michel, & Teti, 1994; Eisenberg & Fabes, 1992; Gross, 1998, 2015; Thompson, 1994). Difficulties with emotion regulation are not confined to emotional or internalizing disorders (e.g., depression and anxiety). Rather, a growing body of evidence suggests that emotion regulation difficulties are present across conditions as diverse as substance abuse (e.g., Aldao, Nolen-Hoeksema, & Schweizer, 2010; Weiss et al., 2015), eating disorders (e.g., Svaldi, Gripenstroh, Tuschen-Caffier, & Ehring, 2012), attention-deficit/hyperactivity disorder (ADHD; e.g., Steinberg & Drabick, 2015), conduct problems (e.g., Beauchaine, Gatzke-Kopp, & Mead, 2007), and psychotic disorders (e.g., Kring & Caponigro, 2010). In parallel, treatments and prevention programs that explicitly teach clients emotion regulation skills have shown promise in treating many of these disorders (e.g., Barlow et al., 2004; Ehrenreich-May & Bilek, 2012; Fairholme, Boisseau, Ellard, Ehrenreich, & Barlow, 2009; Hayes, Strosahl, & Wilson, 1999; Izard, Trentacosta, King, & Mostow, 2004; Linehan, 1993; Roemer, Orsillo, & Salters-Pedneault, 2008; Webster-Stratton, Jamila Reid, & Stoolmiller, 2008).

Closer inspection of this transdiagnostic work on emotion regulation, however, reveals that a majority of studies have assessed this process and psychopathology cross-sectionally, thus reflecting a conceptualization of these processes as static. Yet, emotion regulation is an extremely dynamic process that

changes as a function of context and development (e.g., Aldao, 2013; Cicchetti et al., 1995; Cole et al., 1994; Eisenberg & Fabes, 1992; Gross, 1998, 2015; Thompson, 1994). For example, infants tend to regulate their emotions by moving their gaze, whereas adults do so by relying on a much broader repertoire of strategies, switching between behavioral (e.g., gaze switching) and cognitive (e.g., reappraisal) techniques depending on the context (e.g., Cole et al., 1994). Moreover, psychopathology also evolves across time and development (e.g., Cicchetti, 1989; Masten & Cicchetti, 2010; Rutter & Sroufe, 2000). In this respect, researchers have sought to identify patterns of developmental comorbidity that can shed light onto whether a given disorder might predispose individuals to develop another disorder over time (e.g., Hettema, Prescott, & Kendler, 2003; Moffitt et al., 2007).

Given that both emotion regulation and psychopathology are constantly evolving, we must integrate the transdiagnostic approach with a *developmental psychopathology* framework in order to understand the role of emotion regulation in comorbidity (e.g., Cicchetti, 1984, 1989; De Los Reyes, Bunnell, & Beidel, 2013; De Los Reyes, Henry, Tolan, & Wakschlag, 2009; Mischel & Shoda, 1995; Rutter & Sroufe, 2000; Sroufe & Rutter, 1984). The basic tenet of this framework is that models of psychological dysfunction must take into account the influence of multiple factors (e.g., environmental and biological) across time and development. Of particular importance is the identification of patterns of continuity and discontinuity in symptom expression, specifically by elucidating mechanisms underlying equifinality (i.e., many paths to developing a given outcome or form of psychopathology), multifinality (i.e., a given process resulting in different psychopathological outcomes), and heterotypic continuity (i.e., different manifestations of the same trait at different times in development; e.g., Achenbach, 2011; De Los Reyes, Thomas, Goodman, & Kundey, 2013; Garner, Hake, & Eriksen, 1956; Hinshaw, 2015; Rutter, Kim-Cohen, & Maughan, 2006).

Three recent reviews have illustrated how developmental psychopathology can elucidate the role of emotion regulation in psychiatric comorbidity. Beauchaine and McNulty (2013) proposed the utilization of an *ontogenic* approach, which posits that psychopathology is the result of complex and bidirectional relationships between neurobiological vulnerabilities (e.g., dopaminergic dysfunction) and environmental factors (e.g., parenting, deviant peers, and substance use) that unfold over time (see Hinshaw, 2015). Using the example of heterotypic continuity within externalizing disorders, the authors presented different developmental trajectories in which heritable trait impulsivity (reflecting mesolimbic dopaminergic dysfunction) could lead, through interactions with various biological and environmental factors over time, to the development of oppositional defiant disorder, conduct disorder, substance dependence, and/or antisocial personality disorder. Difficulties with emotion dysregulation comprise a central contextual factor in this model. Specifically, the au-

thors proposed that such deficits result from the reinforcement of affective states, and also evolve over time to acquire a traitlike quality that sets the tone for poor emotional functioning. In this sense, emotion regulation is conceptualized as a dynamic process that is not only a risk/protective factor for psychopathology but also can be an outcome of it.

Beauchaine (2015a) further elaborated on this framework by reviewing the neurobiological substrates (central and peripheral nervous systems) underlying the generation and regulation of emotions in internalizing and externalizing conditions. He outlined a series of future directions on such neurobiological processes: distinguishing between bottom-up and top-down generation and regulation processes, identifying physiological mechanisms underlying reinforcement, improving validity of psychophysiological measures, identifying molecular and genetic processes, and expanding neuroimaging research among children and adolescents. Furthermore, he acknowledged the importance of integrating neurobiological assessments with behavioral ones.

In a similar vein to Beauchaine, Nolen-Hoeksema and Watkins (2011) developed a heuristic for understanding multifinality and divergent trajectories (i.e., two people with the same risk factors developing different disorders). To do so, they differentiated between distal and proximal factors. The former are removed (e.g., with regard to time and/or probability) and include factors such as parent psychopathology, history of trauma, and congenital biological abnormalities. The latter are linked to disorders via specific mechanisms (i.e., intermediate phenotypes) and include processes such as amygdala reactivity, attentional biases, and emotion regulation. According to this model, distal factors (and moderators) influence proximal factors, which subsequently give rise to symptom expression. The authors illustrated this heuristic by focusing on the case of the putatively maladaptive emotion regulation strategy of rumination, which is a form of repetitive negative thought that entails perseverating on the causes of one's shortcomings, mistakes, and regrets (Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008). Specifically, they proposed that in the context of threat, rumination could lead to the development of anxiety disorders, but when accompanied by high sensitivity to alcohol, it could result in substance abuse. Thus, multifinality could stem from a given emotion regulation process and lead to divergent trajectories as a function of moderators. It is worth mentioning that the authors recognized that their heuristic was oversimplified and that the relationship between distal and proximal factors, moderators, and psychopathology is likely recursive and complex (in the words of Beauchaine & McNulty, 2013, an ontogenic process).

Taken together, these three papers provided thought-provoking perspectives on the study of emotion regulation and developmental comorbidity. However, missing from these accounts was a comprehensive discussion of how emotion regulation, as a multifaceted process spanning behavior *and* neurobiology, might lead to heterotypic continuity across internalizing *and* externalizing conditions. Doing so is critical

because emotion regulation spans multiple units of analysis, and thus activity in each level might have differential relations with distinct forms of symptom expression (e.g., Aldao & De Los Reyes, 2015; De Los Reyes & Aldao, 2015; Insel et al., 2010).

To that end, we review the findings from longitudinal studies that have examined emotion regulation in relation to changes in internalizing and externalizing symptoms over time. We omitted studies that did not account for baseline (or prior time point) assessments of psychopathology and/or emotion regulation in order to focus on developmental trajectories. We included studies assessing emotion regulation at the self-report and physiological levels so that we could offer a multimodal conceptualization of this construct. Following this review, we provide a series of recommendations for future work.

Emotion Regulation and the Development of Internalizing and Externalizing Symptoms

In this section, we provide an overview of the research examining the role of emotion regulation in the development of internalizing and externalizing symptoms. To do so, we carefully searched the literature and included any articles that assessed emotion regulation in relation to changes in both internalizing and externalizing symptoms over time. Before we present these findings, however, a word of caution is warranted. Given that this area of inquiry is relatively new, it yields more questions than answers. This is exemplified in two observations. First, we found a great deal of heterogeneity in study methods and findings, so much so that we could not decipher a consistent pattern of findings across studies. Second, despite this variability, the majority of studies have focused on only a handful of emotion regulation processes, thus falling short of capturing the richness of this construct. Following our detailed review of the literature, we provide suggestions on how to address issues raised by the state of work on these topics.

Emotion regulation strategies: Rumination

A vast number of studies on the temporal comorbidity between internalizing and externalizing conditions have examined the trait (i.e., habitual) use of the putatively maladaptive regulation strategy of depressive rumination. This strategy consists of repetitively dwelling upon one's past mistakes, regrets, and/or shortcomings (Nolen-Hoeksema, 1991). People engage in depressive rumination in futile attempts at reducing negative feelings and solving problems, but paradoxically, this strategy increases negative affect, impairs goal-directed action, and erodes relationships (as reviewed in Nolen-Hoeksema et al., 2008). Although the initial work on depressive rumination focused on its role in depression, more recent work indicates that its use is associated with the development and maintenance of a wider range of conditions, including anxiety disorders, eating disorders, and substance abuse in chil-

dren, adolescents, and adults (e.g., Abela & Hankin, 2011; Aldao et al., 2010; Nolen-Hoeksema & Watkins, 2011; Spasojević & Alloy, 2001).

One study examined the moderating role of peer relations on the associations between depressive rumination and internalizing and externalizing symptoms in adolescents (Hilt, Armstrong, & Essex, 2015). Child-reported habitual depressive rumination in Grade 9 was associated with increases in child-reported internalizing symptoms as well as alcohol use in Grade 11. These associations were moderated by the quality of adolescents' relationships with their peers. Specifically, adolescents who experienced elevated peer rejection in Grade 9 exhibited a significant positive association between depressive rumination in Grade 9 and internalizing symptoms in Grade 11, whereas adolescents who had high exposure to peers who consumed alcohol in Grade 9 demonstrated a positive link between depressive rumination in Grade 9 and alcohol use in Grade 11. The results indicated specificity in these associations, such that peer rejection was not a significant moderator in the model predicting alcohol use, and exposure to peers who consumed alcohol was not a significant predictor in the model predicting internalizing symptoms. Therefore, these findings suggest that different types of peer relations might differentially shape the link between rumination and internalizing and externalizing psychopathology.

Other studies on depressive rumination have examined the role of this strategy in the transition from internalizing to externalizing conditions (and vice versa). For example, in one study of adolescents in Grades 6–8 (McLaughlin, Aldao, Wisco, & Hilt, 2014), child-reported habitual use of depressive rumination predicted increases in child-reported depression, anxiety, and aggression over the course of 7 months. Depressive rumination fully mediated the longitudinal associations between aggression and anxiety as well as anxiety/depression and aggression, but only in boys. This is noteworthy because girls tend to ruminate more than boys (Jose & Brown, 2008; Nolen-Hoeksema, Larson, & Grayson, 1999). Thus, these findings indicate that the reliance on depressive rumination might not be the problem per se, but rather who uses it and how (e.g., Aldao, 2013). In order to explain the link between aggression, depressive rumination, and anxiety in boys, the authors suggested that boys might be experiencing more aggression, which leads them to ruminate more about it, and consequently, to feel more anxious around their peers. Such notions could be explored further by administering questionnaires assessing habitual rumination that is anchored to different stressors (e.g., rumination following physical altercations and rumination following social exclusion; see Hartley, Zakriski, & Wright, 2011).

One multiwave study examined fluctuations in child-reported habitual depressive rumination, internalizing, and externalizing symptoms over the course of 5 months in a sample of 6th–10th graders (Hankin, 2008). Depressive rumination at baseline predicted changes in the levels of depression and internalizing symptoms over time. In addition, depressive rumination predicted trajectories of increasing general inter-

nalizing symptoms, but not depression. The author suggested that depressive rumination might have a stronger impact on overall negative emotionality than on specific syndromes. That is, it might have a more pronounced effect on latent traits that, subsequently, lead to specific symptom expression. Baseline depressive rumination was not associated with mean levels or trajectories of anxious arousal or externalizing symptoms, further suggesting specificity to general negative emotionality.

Another multiwave study of adolescents focused on specific types of externalizing psychopathology, namely, bulimia and substance abuse (Nolen-Hoeksema, Stice, Wade, & Bohon, 2007). Participants were female adolescents aged 11–15, who rated their habitual reliance on depressive rumination and their experiences of symptoms of depression, eating disorders, and substance use/abuse once a year over the course of 5 years. Lagged depression symptoms predicted changes in depressive rumination, but depressive rumination was only marginally associated with changes in depression symptoms. Conversely, lagged depressive rumination predicted changes in bulimic symptoms and vice versa. In terms of substance abuse, only lagged depressive rumination predicted changes in substance abuse. That is, substance abuse did not prospectively predict depressive rumination. One possibility is that this reliance on substances may facilitate the avoidance of the unpleasant thoughts, memories, and emotions that might be triggers for rumination. It is also possible that a frequent state of intoxication might render it quite difficult to engage in the type of effortful thinking that underlies rumination. In all, these findings suggest that the associations between rumination and internalizing symptoms might be bidirectional, whereas that between rumination and externalizing symptoms might be unidirectional.

It is also noteworthy that in a large-scale study of adolescents in Grades 8–12 followed over 5 years (Heleniak, Jenness, Stoep, McCauley, & McLaughlin, 2015), self-reported depressive rumination in Grade 8 was *not* associated with changes in either internalizing or externalizing symptoms from 9th to 12th grades. These findings are in sharp contrast with those from the studies discussed above. It is important to note that the regression models in this study included self-reported emotional reactivity and behavioral dysregulation, which suggests that depressive rumination might not be a strong predictor above and beyond those constructs. This would not be surprising, because emotion reactivity and regulation can be difficult to disentangle (e.g., Gross & Barrett, 2011). These three predictors were moderately correlated with one another.

Taken together, the studies reviewed in this section suggest that depressive rumination is associated with the development of internalizing, and to a lesser extent, externalizing, symptoms over time. In addition, these findings indicate that the relationship between rumination and psychopathology may be moderated by both peer relationships and gender. Beyond this, however, these findings are characterized by a substantial amount of heterogeneity that precludes us from drawing further conclusions.

Trait-level emotion regulation abilities

In addition to assessing the habitual use of strategies such as rumination, the study of trait-level emotion regulation has focused on identifying broadly defined regulation skills. That is, instead of assessing whether a participant frequently uses a given strategy, this type of approach entails asking individuals to report on how effectively they can modify their emotions. For example, a sample item from the widely used Difficulties in Emotion Regulation Scale (Gratz & Roemer, 2004) is “When I am upset, it takes me a long time to feel better.” Similarly, an item from the Emotion Reactivity Scale (Nock, Wedig, Holmberg, & Hooley, 2008) is “When I am angry/upset, it takes me much longer than most people to calm down.” Finally, an item from the parent-reported Emotion Regulation Checklist (Shields & Cicchetti, 1997) is “[Child] is repetitive/rigid when stressed.” It is critical that, in the clinical science literature, impaired emotion regulation skills are frequently referred to as “emotion dysregulation.”

Similar to the study described above on peer relationships (e.g., Hilt et al., 2015), one study examined the link between emotion regulation, peer rejection and victimization, and internalizing and externalizing psychopathology (Bierman, Kalvin, & Heinrichs, 2015). Parent-rated child emotion dysregulation at baseline predicted peer rejection and victimization in middle childhood. The latter, in turn, predicted self-reported depression in adolescence. Parent-reported child internalizing symptoms at baseline were also associated with self-reported depression in adolescence. Another model found that emotion dysregulation was associated with peer rejection and victimization in middle childhood. The former was linked to self-reported delinquent behaviors. In this model, internalizing symptoms were not associated with delinquency. In line with studies described above (e.g., Hilt et al., 2015), these findings underscore the importance of modeling social stressors when seeking to understand the role of emotion regulation in the development of internalizing and externalizing conditions.

In another study, the authors examined associations between emotion regulation abilities and symptoms of internalizing and externalizing conditions over the course of 7 months in a sample of adolescents in Grades 6–8 (McLaughlin, Hatzenbuehler, Mennin, & Nolen-Hoeksema, 2011; see McLaughlin et al., 2014). Adolescent-reported emotional understanding, dysregulated expressions of sadness and anger, and ruminative responses to distress formed a unitary latent emotion dysregulation factor that predicted increases in adolescent-reported anxiety symptoms, aggressive behavior, and eating pathology. However, this emotion dysregulation factor did not predict depressive symptoms. The authors sought to explore the source of this null finding by breaking down the latent factor into its individual indicators and testing regression models predicting depression symptoms with each indicator. They found that dysregulated expressions of sadness and anger as well as depressive rumination predicted increases in depression. However, difficulties with emotional

understanding did not. It is possible that these difficulties understanding emotions might not confer risk for depression to the same extent that they might do so for anxiety, aggression, and eating disorder symptoms. The authors suggested that this could be because teenagers with depression might actually be too aware of their emotions. The findings from this study are in contrast with some of the results described above, suggesting that emotion regulation might have a stronger association with internalizing than externalizing symptoms.

Emotion regulation has also been examined in the context of psychosocial adversity (i.e., low socioeconomic status, unstable living conditions, and relationship instability; Halligan et al., 2013). Pregnant women were classified as experiencing high or low adversity, and then their children's emotion regulation was assessed neonatally (mother report), at 12–18 months (mother report, behavioral), and at 5 years of age (mother report, behavioral); behavioral problems at 12–18 months (mother report); and internalizing and externalizing symptoms at 5 years (mother report). Emotion regulation assessed at 12–18 months (and at 5 years) was the only prospective predictor of externalizing symptoms at age 5 (after controlling for behavioral problems at age 12–18 months, child gender, and risk group). However, there was no association between emotion regulation at 12–18 months and internalizing symptoms at age 5. These findings suggest specificity of emotion regulation to externalizing conditions and highlight the value of assessing emotion regulation at a very young age.

Other studies specifically focused on children experiencing stressors at home. In this respect, one study consisted of maltreated and nonmaltreated children (aged 6–12) who attended a weeklong camp for low-income inner-city families (Kim & Cicchetti, 2010). Maltreatment risk factors, such as neglect, physical and sexual abuse, multiple subtypes, and early onset, were associated with poor counselor-rated adaptive emotion regulation abilities. In addition, counselor-rated emotion regulation abilities were associated with greater peer acceptance at follow-up 1 year later, which was also associated with lower counselor-reported internalizing symptoms at the follow-up time point 1 year later. Moreover, low emotion regulation abilities at baseline were associated with higher co-occurring counselor-rated externalizing symptoms at baseline, peer rejection 1 year later, and externalizing symptoms 1 year later. Therefore, these findings provide evidence for the role of emotion regulation in both internalizing and externalizing conditions.

Another study examined the link between parental marital conflict and adolescent psychopathology over the course of 3 years in adolescents aged 11–14 (Buehler, Lange, & Franck, 2007). Child-reported overall emotion dysregulation and avoidance at year 2 mediated the association between marital hostility at baseline and increases in child-reported internalizing, but not externalizing symptoms, at year 3. These findings provide evidence for the specificity of emotion regulation to internalizing symptoms.

Similar to the literature on rumination reviewed above, these findings suggest that trait-level emotion regulation abil-

ities are broadly associated with the development of internalizing and externalizing conditions. However, given the great variability in the findings, it is not possible at this time to draw nuanced conclusions regarding the role of emotion regulation in the development of these forms of psychopathology.

Biological markers of emotion regulation

In line with a current trend in psychological science consisting of delineating biological processes underlying psychological phenomena (e.g., Blair & Diamond, 2008; Casey et al., 2013; Insel et al., 2010; Sanislow et al., 2010), the study of emotion regulation has been placing great emphasis on identifying biological processes that might reflect emotion regulation processes (for reviews see Aldao & De Los Reyes, 2015, 2016; Beauchaine & Thayer, 2015; De Los Reyes & Aldao, 2015; Etkin, Büchel, & Gross, 2015; Ochsner & Gross, 2005; Patrick & Hajcak, 2016). In this respect, one frequently studied biomarker of emotion regulation capacity is high-frequency heart rate variability, also known as respiratory sinus arrhythmia (RSA). RSA reflects the extent to which the parasympathetic nervous system influences the heart rate. Because the parasympathetic system is faster and more flexible than the sympathetic one, elevated parasympathetic activity is considered an adaptive way of mobilizing resources to respond to environmental demands (e.g., Beauchaine et al., 2007; Porges, 2007; Thayer & Lane, 2000). It is critical that RSA needs to be understood as part of a network that also includes the central nervous system (e.g., Beauchaine, 2001). In this respect, accumulating neuroimaging evidence suggests that RSA might be linked to activity in the prefrontal cortex, and in particular, to prefrontal control of subcortical pathways (e.g., Beauchaine & Thayer, 2015; see meta-analysis by Thayer, Ahs, Fredrikson, Sollers, & Wager, 2012).

A growing literature suggests that elevated RSA during periods of rest is associated with adaptive emotion regulation and good mental health. Low resting RSA has been conceptualized as a transdiagnostic factor that cuts across multiple forms of psychopathology (e.g., Beauchaine, 2015b; Zisner & Beauchaine, 2016). Furthermore, Beauchaine and Thayer (2015) have recently suggested that RSA might underlie heterotypic comorbidity across internalizing and externalizing conditions. It is worth noting, however, that even vanilla baselines can require a certain amount of attentional control that will reduce RSA in healthy controls, but not in those with attention problems (e.g., Beauchaine, 2001). As such, it is of utmost importance that the assessments of resting RSA include assessments of potential attentional confounds (e.g., Zisner & Beauchaine, 2015).

Furthermore, the picture becomes much more nuanced when it comes to phasic RSA. Specifically, some studies have shown that reductions in RSA (i.e., vagal withdrawal) in response to stressors reflects an adaptive response (as reviewed in Graziano & Derefinko, 2013), whereas others have found that it is linked to internalizing psychopathology (e.g., Boyce et al., 2001) and that excessive withdrawal is as-

sociated with externalizing symptoms (see Zisner & Beauchaine, 2016). It is crucial that recent work suggests that in order to understand phasic changes in RSA, it might be essential to take into account resting levels. For example, low resting RSA coupled with excessive RSA withdrawal has been associated with internalizing and externalizing symptoms (as reviewed by Beauchaine & Thayer, 2015). Thus, in order to better understand the functional role of phasic RSA, it will be essential to model it in tandem with resting values.

A few studies have examined the role of RSA in the temporal comorbidity between internalizing and externalizing conditions. One study focused on the interactions between baseline RSA and RSA in response to social stressors (Hinnant & El-Sheikh, 2009). Specifically, in a sample of third graders, baseline RSA interacted with RSA changes in response to a social stressor (i.e., adults arguing) to predict changes in parent-reported internalizing symptoms over the course of 2 years. Children who exhibited low resting RSA and high RSA withdrawal in response to the social stressor developed the highest level of internalizing symptoms over time. This pattern was specific to the social stressor, because RSA changes to a frustrating task (i.e., mirror tracing) were not significant predictors of internalizing symptoms. RSA changes to the frustrating task, but not to the social stressor, were significant predictors of externalizing symptoms. Children with low resting RSA and high RSA augmentation showed the highest level of externalizing symptoms. These findings underscore the importance of examining RSA changes in response to tasks that vary in the emotions they elicit and challenges they pose in order to differentiate affective functioning in internalizing and externalizing conditions.

Combining this sample with a similar one (Hinnant & El-Sheikh, 2013), the authors utilized growth mixture modeling to identify profiles of baseline and reactivity RSA. One group (49%) was considered normative because it comprised a large section of the sample and it was characterized by stabilization of externalizing symptoms and declines in internalizing symptoms over time. This group had moderate levels of baseline RSA and RSA withdrawal to the argument task. Another group (41%) had the lowest levels of externalizing symptoms at baseline and over time, and a more gradual decrease in internalizing symptoms over time (relative to the normative group). This low-externalizing/moderate-internalizing group had the highest baseline RSA and the strongest RSA withdrawal to the social stress and frustration tasks. The third group (10%) had the highest initial levels of internalizing and externalizing symptoms, increases in externalizing symptoms, and the smallest decreases in internalizing symptoms over time. This high-externalizing/high-internalizing group was characterized by the lowest baseline RSA and the weakest withdrawal to the tasks. In all, the findings from this study highlight the importance of utilizing person-centered approaches to identify the link between emotion regulation and changes in symptoms over time.

Another study examined baseline RSA and RSA reactivity to sad film clips in children (Pang & Beauchaine, 2013). They

were 8–12 years old at baseline and were followed for 3 years. Both child-reported depression and conduct symptoms were associated with lower resting baseline RSA and higher RSA withdrawal during the initial assessment. However, the interaction between depression and conduct disorder was also significant, such that participants with elevated scores of both conditions had the lowest baseline RSA. Similar findings were obtained for RSA reactivity to the clip. However, there were no prospective associations between psychopathology and RSA, a finding that the authors suggest might be the result of the specific developmental stage of this sample. In this respect, it will be important to test these models across the developmental spectrum.

In contrast, another study utilized data from a large prospective study of Dutch adolescents to examine the link between resting RSA at age 11 and changes in internalizing and externalizing symptoms by age 13 (Oldehinkel, Verhulst, & Ormel, 2008). In this case, resting RSA was associated with higher externalizing symptoms, but not with internalizing symptoms. Thus, these findings indicate specificity in the relationship between resting RSA and externalizing symptoms.

In all, these studies suggest that baseline RSA and RSA reactivity have potential as biomarkers of emotion regulation within a developmental psychopathology framework. However, as has been the case in the previous two sections, the substantial heterogeneity in the study designs and findings prevents us from drawing additional conclusions at this time.

Future Directions

Recent work utilizing a developmental psychopathology framework has led to important advances in the understanding of the role that emotion regulation plays in the development of both internalizing and externalizing conditions. Specifically, the extant research indicates that different forms of emotion regulation (e.g., habitual rumination, emotion regulation skills, and RSA) are associated with the development of internalizing and externalizing symptoms (e.g., Abela & Hankin, 2011; Aldao et al., 2010; Bierman et al., 2015; Halligan et al., 2013; Hilt et al., 2015; McLaughlin et al., 2011, 2014; Oldehinkel et al., 2008; Spasojević & Alloy, 2001) and that they might underlie transitions from disorders in one domain to the other (e.g., McLaughlin et al., 2014; Nolen-Hoeksema et al., 2007). In all, these findings suggest that emotion regulation might be an important factor in the developmental comorbidity between internalizing and externalizing conditions.

However, a quick glance at this work reveals two important limitations that need to be addressed in order to advance our understanding of emotion regulation in relation to the development of internalizing and externalizing psychopathology. First, when we were conducting our review, it proved difficult to synthesize research on the role of emotion regulation in the temporal comorbidity between internalizing and externalizing symptoms. This was due to the heterogeneity in findings, likely driven by variability in populations, study

Table 1. Recommendations for future research

-
-
1. Facilitation of cross-study comparisons
 - a. Meta-analyses
 - b. Data repositories
 - c. Statistics repositories
 2. A more nuanced approach to emotion regulation
 - a. Is emotion regulation a transdiagnostic factor underlying both internalizing and externalizing conditions or does it mediate transitions between these conditions?
 - b. Which are critical emotion regulation processes for understanding the temporal comorbidity between internalizing and externalizing conditions?
 - c. Does social context make a difference?
 - d. Are there crucial differences across different developmental stages?
 - e. Do developmental changes in brain circuitry matter?
 - f. How important is it to assess and model information from multiple informants?
-
-

designs, and analytic approaches. Thus, our first set of recommendations seeks to facilitate cross-study comparisons. Second, despite the marked variability in methods, it is also the case that the majority of the work has focused on a limited number of emotion regulation processes (e.g., use of strategies, regulation abilities, and RSA), which could result in an overly simplistic understanding of the role of emotion regulation in the development of internalizing and externalizing conditions. Therefore, we provide a series of recommendations for how to comprehensively study emotion regulation as a multifaceted construct within a transdiagnostic developmental framework (see Table 1).

Facilitation of cross-study comparisons

Inherent to the developmental psychopathology approach is the utilization of complex study designs that span multiple assessment points, informants, contexts, and psychological processes. Most data sets contain dozens, if not hundreds, of variables. However, any given study can only focus on a small subset of variables, and consequently, different investigators will analyze different combinations of variables in each study that they publish. For example, let us imagine that two research groups have two comparable data sets that assess trait rumination once a year over the course of 5 years in a sample of adolescents. One research team publishes findings focusing on the link between rumination, anxiety, social stress, and parent psychopathology over the first 3 years, and the other team publishes findings on the covariation between rumination and anxiety symptoms over those entire 5 years. Let us further imagine that these two studies entailed different findings regarding the link between rumination and anxiety: the first study did not find evidence of an association, whereas the second one found a moderate positive correlation. How are we to synthesize these findings? Are the discrepant findings the result of the differential time lapses? Are they a function of the inclusion of additional variables in the first model? Are they stemming from sampling issues? Finding systematic ways of evaluating each of these possibilities is essential for the continuing growth of this area of inquiry.

We can think of a few ways of doing so. The first one consists of adopting a meta-analytic approach. In the example above, an investigator interested in the role of rumination and anxiety over time would e-mail authors and ask for the relevant statistics. With enough studies he or she could examine the role of moderators in the strength and direction of the effect sizes. He or she would then write up a manuscript and submit it for a (usually quite lengthy) peer-review process.

We see great merit in this approach, as we ourselves have published a number of meta-analytic reviews (e.g., Aldao et al., 2010; Chaplin & Aldao, 2013; De Los Reyes et al., 2015). However, it is also the case that this approach places the burden on the potential meta-analyst to collect, process, and analyze all the relevant information. Moreover, it assumes that there would be individuals with enough time, expertise, and dedication to want to carry out enough meta-analysis to cover the extensive work on emotion regulation and the development of internalizing and externalizing symptoms.

Another option that would distribute the workload among investigators in the field would entail the creation of data repositories (see Center for Open Science at <https://cos.io/>). In this sense, if an investigator needs an estimate of the temporal covariation between rumination and anxiety symptoms, he or she could pull the data and run the analyses with relative ease. However, this approach might face several obstacles, including reluctance to share data and massive administrative efforts in order to combine the data sets (e.g., variable naming and formatting, and measure scoring).

Thus, it might be more feasible to generate *summary statistics repositories*. That is, rather than uploading raw data, investigators would upload summary statistics (e.g., descriptive and bivariate correlations). Following up on our example above, this would entail uploading bivariate correlations between rumination and anxiety for each of the 5 years during which data were collected. This would not require the sharing of raw data, and the logistics would be much simpler. It could start as a very simple system, including only descriptive statistics (means and standard deviations) and bivariate

correlations for the emotion regulation and symptom variables at each time point. Over time, it could become a more dynamic system in which investigators could actively request and upload a wider range of summary statistics that would help inform their research. Moreover, circling back to our first point, the availability of these data would facilitate the process of conducting meta-analytic reviews.

A more nuanced approach to emotion regulation

Our second set of recommendations pertains to expanding upon how emotion regulation is studied within a transdiagnostic developmental perspective. We present a series of critical questions to the field and venture suggestions for how to best address them.

Is emotion regulation a transdiagnostic factor underlying both internalizing and externalizing conditions or does it mediate transitions between these conditions? Some of the articles reviewed above indicate that emotion regulation might be a transdiagnostic factor associated with the development of internalizing and externalizing conditions (Abela & Hankin, 2011; Aldao et al., 2010; Bierman et al., 2015; Halligan et al., 2013; Hilt et al., 2015; McLaughlin et al., 2011, 2014; Oldehinkel et al., 2008; Spasojević & Alloy, 2001), although the relative strength of the associations remains to be determined. Other studies found that it mediated transitions between one type of condition and the other one (McLaughlin et al., 2014; Nolen-Hoeksema et al., 2007). Identifying whether specific emotion regulation processes play a stronger role in the transition from health to psychopathology or from one form of psychopathology to another one is critical for prevention and treatment programs. Forms of emotion regulation that are associated with the former would be targeted in large-scale prevention efforts aimed at the general population (e.g., entire school system, retirement homes, community centers, and work-place programs). Conversely, those more strongly linked with transitions from one form of psychopathology to another one would be targeted in treatment and relapse prevention efforts with those already suffering from a given condition (or perhaps even at high risk for it). Overall, reaching this level of specificity in our understanding of emotion regulation and internalizing and externalizing conditions has great potential for the development of more efficient and cost-effective prevention and treatment efforts, and ultimately, the reduction of suffering.

In addition, it would be important to adopt a more nuanced approach by examining emotion regulation in relation to specific forms of internalizing and externalizing disorder. A number of candidates emerge from the literature. First, it is important to examine the “internalizing pathway” to alcohol abuse, whereby alcohol consumption initially serves the function of regulating internalizing emotions, such as anxiety and depression, and over time becomes reinforced and problematic (e.g., Hussong, Jones, Stein, Baucom, & Boeding, 2011; Kashdan, Ferssizidis, Collins, & Muraven, 2010;

Kushner, Sher, & Beitman, 1990; Shadur & Lejuez, 2015; Sher & Grekin, 2007). Second, it will be fruitful to explore the link between anxiety and ADHD (e.g., Falk, Lee, & Chorpita, 2015; Lee, Falk, & Aguirre, 2012; Mennin, Biederman, Mick, & Faraone, 2000; Schatz & Rostain, 2006; Van den Bergh & Marcoen, 2004). In a related vein, it will be important to examine the relationship between anxiety and disruptive behavior disorders, such as oppositional defiant disorder and conduct disorder, especially given the strong commonalities between these disorders in issues with emotion regulation (e.g., Bubier & Drabick, 2009; Drabick, Ollendick, & Bubier, 2010; Forehand, Jones, & Parent, 2013; Fraire & Ollendick, 2013). Third, given the risk of fatality in both depression and substance use disorders, it will be critical for future research to examine the role of emotion regulation in the link between these disorders as well as transitions between them (e.g., Brière, Rohde, Seeley, Klein, & Lewinsohn, 2014; Deykin, Levy, & Wells, 1987; Levy & Deykin, 1989; Lewinsohn, Gotlib, & Seeley, 1995).

Which are critical emotion regulation processes for understanding the temporal comorbidity between internalizing and externalizing conditions? Our survey of the literature revealed that the majority of existing research has focused on trait level self-reports of the use of rumination and perceived regulation skills, and of a particular biomarker of emotion regulation (RSA). Nevertheless, emotion regulation is a complex construct that entails a wide range of targets (e.g., positive and negative emotions), strategies (e.g., reappraisal and avoidance), abilities (e.g., focus on a task in the face of intense emotions), goals (e.g., increased and decreased intensity or duration), and outcomes (e.g., subjective feelings, facial expressivity, motivated behavior, physiological reactivity, and neurobiological activity; Mauss & Robinson, 2009). That is, it unfolds in myriad ways that might have different functional relations with symptom development (e.g., Cicchetti et al., 1995; Cole et al., 1994; Thompson, 1994). Therefore, it is critical to adopt a *multiprocess* approach to the assessment of emotion regulation that entails focusing on multiple processes simultaneously.

A multiprocess approach has two clear advantages. First, it will allow investigators to include multiple emotion regulation processes in the same statistical models and, thus, identify which aspects of emotion regulation are more central to the development of internalizing versus externalizing conditions. That is, this approach might facilitate a more nuanced understanding of which aspects of emotion regulation might be transdiagnostic risk factors versus those that might have specificity to certain conditions. Second, by assessing multiple forms of emotion regulation, investigators open the door to the possibility of understanding how different emotion regulation processes might influence one another. For example, recent work suggests that the use of maladaptive emotion regulation strategies (e.g., avoidance, rumination, and suppression) moderates the association between the use of adaptive strategies (e.g., acceptance and reappraisal) and symptoms

of depression, anxiety, and alcohol abuse (e.g., Aldao, Jazaieri, Goldin, & Gross, 2014; Aldao & Nolen-Hoeksema, 2012; Conklin et al., *in press*). In a similar vein, a number of studies have begun to derive profiles of emotion regulation repertoires and to link them to mental health functioning (e.g., Dixon-Gordon et al., *in press*; Hollenstein, Granic, Stoolmiller, & Snyder, 2004). Further, it will be important to combine the assessment of emotion regulation strategies and abilities (e.g., Tull & Aldao, 2015).

Investigators can adopt this multiprocess approach in a number of ways. First, as we mentioned above, the majority of the work on the habitual use of regulation strategies has focused on depressive rumination. This strategy, albeit a very important one, is only one of many that people have in their repertoires (e.g., Aldao & Dixon-Gordon, 2014). In this vein, it will be important to assess the habitual use of additional putatively maladaptive strategies, such as avoidance, suppression, and worry. Similarly, it will be useful to assess the chronic use of putatively adaptive regulation strategies, such as cognitive reappraisal and acceptance (e.g., Aldao et al., 2010) because it is possible that different strategies have different associations with the development of internalizing and externalizing symptoms. Cross-sectional meta-analytic research on adults and children suggests that maladaptive strategies have a moderate effect size in relation to internalizing conditions (e.g., anxiety and depression; $d = 0.42$), but a small effect size in relation to externalizing ones (e.g., substance abuse and eating disorders, $d = 0.25$). However, the chronic use of adaptive strategies has small effect sizes in relation to both types of psychopathology ($ds = -0.23$ and -0.11 , respectively; see Aldao et al., 2010). However, it is important to keep in mind that the sample sizes for the externalizing cells in this study were considerably smaller than for the internalizing conditions ($k = 30$ vs. 211), reflecting an asymmetry in the study of emotion regulation in relation to internalizing versus externalizing conditions. In all, it will be important to examine how different strategies might simultaneously predict the development of internalizing and externalizing conditions as well as mediate the transitions between these disorders.

Second, it will be useful to assess variations of a given strategy or abilities that might be particularly relevant to certain contexts. A clear example is that of depressive rumination (Nolen-Hoeksema, 1991) and angry rumination (Sukhodolsky, Golub, & Cromwell, 2001). Whereas the former focuses on questioning one's actions and capabilities to deal with stressors, the latter entails perseverating on feelings of having been wronged, slighted, and hurt. Therefore, it is possible that depressive rumination might be more strongly associated with the development of internalizing conditions (Nolen-Hoeksema et al., 1999), whereas angry rumination might be more strongly linked to the development of externalizing problems (Peled & Moretti, 2007). Testing these strategies simultaneously would be of great value for developing a more nuanced understanding of the functional relationship between rumination and different forms of psychopathology. More broadly, a

similar approach can be beneficial by parsing out different types of negative perseverative thinking, including postevent processing, obsessive thinking, and worry.

Third, it might be helpful to anchor trait-level questionnaires of habitual use of strategies and regulation abilities to specific stressors. For example, investigators could ask participants to rate the extent to which they engaged in rumination in response to having been victimized, having felt embarrassed, having underperformed in school or work, or having had romantic setbacks. Moreover, they could assess whether a given regulation ability (e.g., "When I am upset, it takes me a long time to feel better" from the Difficulties in Emotion Regulation Scale; Gratz & Roemer, 2004) is more likely to manifest in response to certain types of stressors. This approach would allow investigators to begin to parse out how the use of regulation strategies and abilities in certain situations might pose different risks for internalizing versus externalizing conditions. In a similar vein, investigators could focus on the regulation of internalizing and externalizing emotions (along the lines of the tasks involving mirror tracing and listening to arguments used by Hinnant and El-Sheikh, 2009, 2013). In all cases, the use of experience sampling methodologies could play a key role in facilitating more precise estimates of the link between stressors, regulation, and outcomes (e.g., Silk, Steinberg, & Morris, 2003).

Does social context make a difference? Our review of the literature suggests that social processes (e.g., peer victimization and delinquent peers) might be important moderators of the link between emotion regulation and the development of internalizing/externalizing symptoms (e.g., Bierman et al., 2015; Buehler et al., 2007; Hilt et al., 2015). Social context is one of the primary moderators outlined in the developmental psychopathology frameworks of Beauchaine and McNulty (2013) and Nolen-Hoeksema and Watkins (2011). This notion is aligned with a growing literature documenting the crucial role that social processes play in emotion regulation (e.g., Aldao, 2013; Christensen et al., *in press*; Hofmann, 2014; Marroquín, 2011; Shallcross, Frazier, & Anders, 2014; Troy, Shallcross, & Mauss, 2013) and with decades of research in developmental psychopathology emphasizing the role that caregivers, siblings, and teachers play in the development of emotion regulation skills (e.g., Cole et al., 1994; Gee et al., 2014; Gunnar & Donzella, 2002; Hofer, 1994; McCoy & Masters, 1985). For these reasons, it can be extremely valuable to systematically model social processes in the study of the association between emotion regulation and the development of internalizing and externalizing conditions. Below, we provide a series of suggestions for how to do so.

The first and most straightforward way to incorporate social processes is by examining regulation in response to social stressors (e.g., peer victimization, aggression, divorce, and job performance). A second method is to examine the extent to which individuals recruit others to regulate their emotions (and also help others regulate theirs). This is known as *interpersonal emotion regulation*, a rapidly growing area of re-

search that used to be primarily confined to infants and younger children (e.g., Cole et al., 1994; Hofmann, 2014; Marroquín, 2011; Zaki & Williams, 2013). For example, when an adolescent is sad and disappointed about a fight with her boyfriend, she might end up ruminating about these emotions by herself or with a friend. The latter is referred to as corumination and has been linked to increases in symptoms of depression in adolescent girls (e.g., Rose, 2002; Rose, Carlson, & Waller, 2007). Thus, studying patterns of coregulation might enhance our understanding of the development of internalizing and externalizing conditions.

In this respect, it would be important to examine whether the use of a given regulation strategy in isolation confers greater risk for, or protection against, internalizing and externalizing conditions than its joint use with other people. That is, is rumination by oneself more or less strongly associated with the development of psychopathology than corumination, or does drinking by oneself to regulate one's anxiety confer more or less of a risk than drinking with others to achieve anxiety reduction? Does self-regulation versus coregulation confer different protection against developing internalizing versus externalizing symptoms (a crucial point)?

In the future, a more fine-grained approach could entail distinguishing whether the consequences of coregulation vary as a function of specific social context. For example, coruminating with individuals who are experiencing the same stressors (e.g., coworkers experiencing problems with their bosses) and therefore have less of a "big picture" perspective might be more problematic than with individuals who are experiencing different stressors and might therefore offer a more distanced perspective (Kross, Ayduk, & Mischel, 2005; Smith & Rose, 2011).

Though it may appear overwhelming to capture interpersonal emotion regulation, a rich amount of information can be collected by adopting the multiple-informant method that is at the cornerstone of developmental psychopathology (e.g., De Los Reyes, 2013; De Los Reyes et al., 2015; De Los Reyes & Kazdin, 2005; Hunsley & Mash, 2007). For example, investigators can assess patterns of interpersonal regulation by asking participants to complete self-report and other-report versions of the use of emotion regulation strategies and abilities. Moreover, with the availability of myriad tools for online data collection, investigators can more easily reach out to friends, family members, significant others, and/or coworkers of participants to request information about their coregulation with the participants. These multiple-informant assessments, when integrated with independent assessments of the social processes described previously, can provide a rich, comprehensive picture of the interplay between social contexts and individual differences in displays of emotion regulation strategies (see De Los Reyes, Bunnell, et al., 2013; De Los Reyes et al., 2009; Mischel & Shoda, 1995).

Are there crucial differences across different developmental stages? The studies we found on emotion regulation as a process related to the development of internalizing and external-

izing symptoms primarily focused on children and adolescents. Consequently, much remains to be done in terms of elucidating how this process might play a role in the development of internalizing and externalizing conditions over the course of the lifespan and in relation to critical developmental periods, such as transition to elementary school, adolescence, college, marriage, parenting, divorce, menopause, and cognitive decline, among others. Adopting this expanded developmental psychopathology approach is particularly important in light of a growing literature documenting marked differences in emotion regulation as a function of aging. This work tends to find that older adults regulate their emotions more effectively than younger adults (e.g., Isaacowitz & Blanchard-Fields, 2012; Livingstone & Isaacowitz, 2015; Nolen-Hoeksema & Aldao, 2011; Prakash, Whitmoyer, Aldao, & Schirda, 2015; Pruzan & Isaacowitz, 2006; Yeung, Wong, & Lok, 2011; Zhang, Ersner-Hersfield, & Fung, 2010). According to socioemotional selectivity theory, these differences might stem from older adults being more selective when building their social networks (e.g., Carstensen, Isaacowitz, & Charles, 1999; Carstensen et al., 2011; Sims, Hogan, & Carstensen, 2015). What remains to be understood is how these, and other, changes in emotion regulation over time might result in differential patterns of internalizing and externalizing symptoms.

Moreover, elucidating the neurodevelopmental mechanisms underlying age-related changes in emotion regulation and their association with risk for the onset of internalizing and externalizing disorders will be critical to informing early intervention and prevention approaches. Given the dynamic changes that occur in frontolimbic circuitry across the life span, mechanisms of illness and treatment are likely to differ depending on developmental stage (Lee et al., 2014). Research that seeks to identify the similarities and differences in neurobiological mechanisms characterizing different psychiatric disorders will play a critical role in better understanding emotion regulation as a transdiagnostic factor.

Do developmental changes in brain circuitry matter? In our review of the literature, we found that, despite growing enthusiasm for identifying the neurobiological underpinnings of emotion regulation and psychopathology (e.g., Aldao & De Los Reyes, 2015; Beauchaine & Thayer, 2015; Etkin et al., 2015; Hinnant & El-Sheikh, 2009, 2013; Ochsner & Gross, 2005; Oldehinkel et al., 2008; Pang & Beauchaine, 2013; Patrick & Hajcak, 2016), much remains to be understood regarding how developmental changes in brain circuitry might play a role in the association between emotion regulation and the development of internalizing and externalizing conditions. Thus, below we provide an overview of developmental research on brain circuitry and provide suggestions for future work.

Interactions between subcortical limbic and cortical prefrontal regions are fundamental to the processing and regulation of emotional reactivity (Banks, Eddy, Angstadt, Nathan, & Phan, 2007; Kim, Somerville, Johnstone, Alexander, &

Whalen, 2003; Lieberman et al., 2007; Ochsner, Bunge, Gross, & Gabrieli, 2002). Emotion regulation and the connections between cortical and subcortical brain regions undergo dynamic changes across the lifespan, which are likely to have important implications for the development of psychiatric disorders. Interactions between bottom-up, subcortical regions supporting emotional reactivity and top-down, cortical regions underlying regulatory control are central to emotion regulation. In general, subcortical regions, such as the amygdala and ventral striatum, facilitate emotional reactivity and motivational processes that may be more automatic and tendency related, whereas top-down control mediated by prefrontal regions may be more involved in the volitional regulation of emotion (see Beauchaine, 2015a).

Findings on the normative maturation of these emotion generation and regulation systems, and the connections between them, can provide an important reference with which to compare deviations from typical development in individuals who develop internalizing and externalizing disorders. Risk for psychiatric disorders increases during development, and many psychiatric disorders have been conceptualized through a neurodevelopmental framework (e.g., Lee et al., 2014; Pine, Cohen, Gurley, Brook, & Ma, 1998). Moreover, dynamic changes in this circuitry may help to explain the influence of environmental and genetic factors on the onset of certain disorders at specific developmental windows (e.g., Burghy et al., 2012; reviewed in Gee & Casey, 2015; Gee et al., 2016). For example, altered cortical–subcortical interactions may be a common possible pathway through which early-life stress and maltreatment influence both internalizing and externalizing behaviors (Burghy et al., 2012; Gee, Gabard-Durnam, et al., 2013; Hanson et al., 2010; Herringa et al., 2013). Understanding the nature of disruptions in the connectivity between bottom-up and top-down regions is also likely to have important implications for the treatment of psychopathology during development.

Subcortical regions involved in emotional reactivity tend to mature earlier in normative development relative to more protracted development of cortical regions. The amygdala undergoes rapid change early in development (reviewed in Tottenham & Sheridan, 2009). Children show heightened amygdala reactivity to fearful faces and other emotional stimuli, with reactivity typically decreasing following childhood (e.g., Decety, Norman, Berntson, & Cacioppo, 2012; Gee, Humphreys, et al., 2013; Silvers, Weber, Wager, & Ochsner, 2014; Swartz, Williamson, & Hariri, 2014; Vink, Derks, Hoogendam, Hillegers, & Kahn, 2014), a finding that parallels decreases in normative fears that occur early in life (Gee, Humphreys, et al., 2013). Activation in the ventral striatum related to motivational cues has been shown to peak during adolescence (e.g., Galvan et al., 2006; Somerville, Hare, & Casey, 2011). By contrast, the prefrontal cortex (PFC) undergoes more protracted development into young adulthood (e.g., Chareyron, Lavenex, Amaral, & Lavenex, 2012; Lenroot & Giedd, 2006; Machado & Bachevalier, 2003; Payne, Machado, Bliwise, & Bachevalier, 2010). The connections

between these subcortical regions and the PFC also show protracted maturation, with these developmental shifts underlying dynamic behavioral changes in regulation during childhood and adolescence (see Casey, Galván, & Somerville, 2016).

Connections between the amygdala and medial PFC that support effective emotion regulation in adults change substantially across the course of childhood and adolescence (e.g., Decety et al., 2012; Gabard-Durnam et al., 2014; Gee, Humphreys, et al., 2013; Lebel et al., 2012; Perlman & Pelphrey, 2011; Swartz et al., 2014; Vink et al., 2014), consistent with behavioral changes in emotion regulation during development (e.g., Bunge, Dudukovic, Thomason, Vaidya, & Gabrieli, 2002; Tottenham, Hare, & Casey, 2011). For example, the regulatory inverse pattern of frontoamygdala functional connectivity that is associated with effective emotion regulation in adults appears to emerge around the transition from childhood to adolescence (Gee, Humphreys, et al., 2013). Prior to adolescence, external sources of emotion regulation, such as parents, have been shown to regulate children's behavior via modulation of frontoamygdala connectivity (Gee et al., 2014). During adolescence, inverse frontoamygdala functional connectivity has been associated with greater amygdala habituation, suggesting a similar function of this circuitry by adolescence (Hare et al., 2008). Similarly, stronger inverse connectivity and medial PFC recruitment correspond to improved reappraisal success with age (McRae et al., 2012; Silvers et al., 2014). Given the role of the medial PFC in regulating amygdala reactivity, evidence of stronger inverse coupling and reduced amygdala reactivity with age may provide a neurobiological basis for developmental improvements in regulation of negative emotion and normative anxiety.

Although cortical–subcortical interactions support emotion regulation and deviations in related neurodevelopment are likely to contribute to the emergence of psychopathology in development, much remains unknown about the specific nature of these deviations for internalizing and externalizing disorders. One possibility is that impairments in specific circuitry might mediate core dimensional processes, such as emotion regulation, which confer risk for psychopathology in general rather than for specific disorders (Buckholtz & Meyer-Lindenberg, 2012), yet there appears to be some distinction between the primary neural mechanisms of regulation in internalizing versus externalizing disorders (see Beauchaine, 2015a). In general, the regulation of internalizing symptoms like anxiety appears to occur through top-down lateral and medial prefrontal control of amygdala reactivity (e.g., Lieberman et al., 2007; Ochsner et al., 2002), whereas the regulation of externalizing symptoms like impulsivity involves dorsolateral prefrontal and orbitofrontal modulation of striatal activity (e.g., Dalley, Mar, Economidou, & Robbins, 2008; Heatherton & Wagner, 2011).

Consistent with its role in the modulation of negative emotion, frontoamygdala circuitry has been strongly implicated in internalizing disorders (e.g., Rauch, Shin, & Wright, 2003). Neuroimaging involving the regulation of negative emotion

in healthy adults has generally underscored the role of lateral regions of the PFC (e.g., ventrolateral and dorsolateral PFC), in addition to medial regions (e.g., ventromedial PFC) that are central to fear reduction processes such as extinction (e.g., Delgado, Nearing, LeDoux, & Phelps, 2008; Erk et al., 2010; Goldin, McRae, Ramel, & Gross, 2008; Lieberman et al., 2007; Ochsner et al., 2002; Phelps, Delgado, Nearing, & LeDoux, 2004; Wager, Davidson, Hughes, Lindquist, & Ochsner, 2008). These same regions have been implicated in pediatric anxiety (e.g., De Bellis et al., 2000; Guyer et al., 2008; Monk et al., 2008; Roy et al., 2013) and depression (e.g., Gaffrey et al., 2011; Hulvershorn, Cullen, & Anand, 2011; Luking et al., 2011; Yang et al., 2010). For example, youth with generalized anxiety disorder had weaker inverse connectivity between the ventrolateral PFC and the amygdala during a task involving masked angry faces, consistent with weaker regulatory control (Monk et al., 2008). However, few studies have examined frontoamygdala circuitry in pediatric psychiatric populations during tasks of emotion regulation, and much remains unknown about the specific associations between risk for emotion regulation difficulties and neural circuitry abnormalities prior to adulthood.

Moreover, disruptions in frontoamygdala neurodevelopment are not specific to internalizing disorders, because they have also been observed in children and adolescents with ADHD (e.g., Hulvershorn et al., 2014; Posner et al., 2011), bipolar disorder (e.g., Passarotti, Ellis, Wegbreit, Stevens, & Pavuluri, 2012; Pfeifer, Welge, Strakowski, Adler, & Delbello, 2008; Rich et al., 2008), psychotic symptoms (e.g., Gee et al., 2012; Wolf et al., 2015), and callous–unemotional traits (e.g., Marsh et al., 2008). While much of the research on frontoamygdala circuitry in adults with psychiatric disorders has focused on anxiety (reviewed in Rauch et al., 2003) and depression (e.g., Siegle, Thompson, Carter, Steinhauer, & Thase, 2007; reviewed in Heller, 2016), altered frontoamygdala circuitry has also been observed in adults with bipolar disorder (e.g., Townsend et al., 2013; Wang et al., 2009), schizophrenia (e.g., Anticevic, Van Snellenberg, & Barch, 2012; Fakra, Salgado-Pineda, Delaveau, Hariri, & Blin, 2008; Taylor et al., 2012), borderline personality disorder (e.g., New et al., 2007; Silbersweig et al., 2007), psychopathy (e.g., Blair, 2008), and ADHD (e.g., Maier et al., 2014; Plichta et al., 2009; Tajima-Pozo et al., 2016). Thus, future research is needed to better understand the nature of frontoamygdala changes and transdiagnostic risk associated with deficits in emotion regulation across many internalizing and externalizing disorders.

By contrast, frontostriatal circuitry is more strongly associated with regulatory difficulties in externalizing disorders. Disrupted interactions between frontal regions (e.g., dorsolateral PFC and orbitofrontal cortex) and the ventral striatum have been observed in adults with substance abuse disorders (e.g., Goldstein & Volkow, 2011; Koob & Le Moal, 2001), ADHD (e.g., Casey et al., 2007), and psychopathy (e.g., Glenn & Yang, 2012). Similarly, frontostriatal circuitry has been implicated in children and adolescents with externaliz-

ing disorders such as conduct disorder (e.g., De Brito et al., 2009), ADHD (e.g., Shaw et al., 2012), ADHD (e.g., Plichta & Scheres, 2014), and psychopathic traits (e.g., Blair, 2013). However, just as frontoamygdala circuitry is not uniquely implicated in internalizing disorders, alterations in frontostriatal circuitry are not specific to externalizing disorders. For example, neuroimaging investigations of altered reward processing and positive affect in depression have led to an increased focus on frontostriatal interactions in depression (Heller, 2016).

Thus, it remains unknown the extent to which the neurobiological disruptions underlying deficits in emotion regulation might be similar or distinct across internalizing and externalizing disorders. Consistent with the association between individual differences in normative anxiety and frontoamygdala connectivity (e.g., Hare et al., 2008; Pezawas et al., 2005), frontoamygdala circuitry is disrupted in pathological anxiety (reviewed in Kim et al., 2011). Specifically, weaker frontoamygdala connectivity, amygdala hyperreactivity, and prefrontal hypoactivity appear to play central roles in the pathophysiology of anxiety disorders (e.g., Shin et al., 2005; reviewed in Rauch et al., 2003). Findings on frontoamygdala alterations are more variable across other disorders, and it may be that the specific nature of emotion regulation and related neural mechanisms differ by symptom type. For example, emerging evidence suggests that, unlike in anxiety disorders, the core mechanism underlying depression may be a failure to upregulate and sustain positive emotion via frontostriatal circuitry (e.g., Heller et al., 2009). Similarly, the nature of neural abnormalities in schizophrenia is markedly different from many other psychiatric disorders, because patients with schizophrenia exhibit hypoactive amygdala activation (Taylor et al., 2012). Thus, although similar circuitry is implicated, neurobiological alterations may relate to excessive or insufficient emotion depending on the form of psychopathology. Moreover, it may be that deficits in emotion regulation differentiate some patients from others with the same disorder (e.g., Hulvershorn et al., 2014; Karalunas et al., 2014). Future research will be essential to delineating the commonalities and differences in the neurobiological substrates of emotion regulation and transdiagnostic risk across internalizing and externalizing disorders.

How important is it to assess and model information from multiple informants? Much like symptom expression, emotion regulation capabilities too change over the course of the life span. Thus, a developmental psychopathology approach encompassing emotion regulation ought to take into account the larger context in which assessments are conducted (e.g., biology, culture, environment, and social processes). In this respect, one of the most robust findings in this field is that these variations in assessment produce substantial inconsistencies in findings regarding psychopathology and its risk and protective factors (Achenbach, 2011; De Los Reyes, 2013; De Los Reyes et al., 2015). Recently, we developed a theoretical model to account for these varia-

tions (De Los Reyes & Aldao, 2015; De Los Reyes, Thomas, et al., 2013).

Specifically, the operations triad model (OTM) accounts for variations among the outcomes of multiple-informant assessments by distinguishing three different kinds of possible outcomes from these assessments. In one instance, two or more informants (e.g., parents and teachers) provide corresponding reports of a target individual's mental health (e.g., child displays externalizing difficulties across reports), thus signaling that the target individual displays the assessed concern consistently across contexts observed by the informants (e.g., home and school). These assessments yield outcomes that reflect *converging operations* (see Garner et al., 1956). Two other instances reflect circumstances in which informants provide inconsistent reports of a target's mental health (e.g., child displays externalizing difficulties based on teacher report and not parent report). When inconsistencies reflect *diverging operations*, the reports meaningfully point to variations in the assessed concerns across contexts (e.g., child displays difficulties at school and not home). When they reflect *compensating operations*, mundane, methodological factors exist to parsimoniously explain the inconsistencies (e.g., different item content across informants' reports). Overall, the OTM presents a hypothesis-driven, theory-guided approach to making sense of consistencies and inconsistencies in the outcomes of multiple-informant mental health assessments.

Recent empirical work supports the OTM, and indicates that variations in measurement do not necessarily pose barriers to our basic understanding of psychopathology. Instead, these variations may inform our understanding of individual differences in mental health functioning (e.g., De Los Reyes, Thomas, et al., 2013). For example, in assessments of disruptive behavior among young children, clinical assessments often incorporate parent and teacher reports, and these reports commonly disagree in estimates of disruptive behavior. In recent work involving parent and teacher reports and independent observations of young children interacting with various adult authority figures (i.e., parents vs. unfamiliar clinical examiners), researchers profiled young children as to whether they behaved disruptively within controlled laboratory interactions with their parents, clinical examiners, neither of these interactions, or both of them (De Los Reyes et al., 2009). Consistent with diverging operations, children who behaved disruptively with parents and not clinical examiners were also those children who were rated as disruptive by parents and not teachers. Further, children who behaved disruptively with clinical examiners and not parents were also those children who were rated as disruptive by teachers and not parents, and children who behaved disruptively with *both* clinical examiners and parents were rated as disruptive by teachers *and* parents. Effects consistent with the OTM have also been observed in assessments of adolescents (De Los Reyes, Alfano, Lau, Augenstein, & Borelli, 2016) and adults (De Los Reyes, Bunnell, et al., 2013).

The work reviewed previously may inform longitudinal research on the role of deficits in emotion regulation and vulnerability to psychopathology. For example, we can posit that adolescents for whom multiple informants' reports reflect converging operations in displays of high degrees of internalizing psychopathology are at particular risk for pervasive emotion regulation difficulties (e.g., parents, teachers, and peers), relative to adolescents for whom only one informant suggests such psychopathology (i.e., diverging operations that reflect context-specific emotion regulation difficulties). Later in adolescence and into emerging adulthood, the *pervasive* adolescents may manifest a particularly high risk for maladaptive strategies for coping with their mental health concerns, namely, risk-taking behaviors (e.g., substance use and risky sexual behavior), thus resulting in development of a comorbid internalizing–externalizing clinical presentation. Conversely, the *context-specific* adolescents might not manifest comorbid externalizing difficulties, but may nonetheless display emotion regulation strategies that maintain internalizing concerns persistently across development. With these examples, a combination of multiple-informant assessments of mental health concerns, and laboratory tasks designed to assess emotion regulation difficulties, might provide a particularly robust paradigm to assess the development of comorbidity in psychopathology, in ways similar to the cross-sectional mental health research described previously (e.g., De Los Reyes et al., 2009, 2016; De Los Reyes, Bunnell, et al., 2013). These paradigms and their implications for emotion regulation research, merit further study.

Concluding Remarks

In the last decade, there has been exponential growth in the study of emotion regulation as a transdiagnostic process that cuts across multiple forms of psychopathology (e.g., Aldao et al., 2010; Gross & Jazaieri, 2014; Kring & Sloan, 2009). However, only a small fraction of this work has adopted a developmental psychopathology approach, and within that, only a few studies have examined the temporal comorbidity between internalizing and externalizing conditions. In this paper, we reviewed this small but promising literature, and we provided a series of recommendations for future research. Specifically, we advocated the putting in place of systems that can facilitate greater cross-communication across laboratories and therefore result in a more systematic growth of this literature. In addition, we provided specific suggestions regarding how to study emotion regulation in more nuanced ways. In all, we hope that these suggestions can lead to a more in-depth delineation of the role of emotion regulation in the temporal comorbidity between internalizing and externalizing conditions, and more broadly, to a more nuanced understanding of comorbidity patterns across psychopathology that can inform research, prevention, and treatment.

References

- Abela, J. R., & Hankin, B. L. (2011). Rumination as a vulnerability factor to depression during the transition from early to middle adolescence: A multiwave longitudinal study. *Journal of Abnormal Psychology, 120*, 259.
- Achenbach, T. M. (2011). Commentary: Definitely more than measurement error: But how should we understand and deal with informant discrepancies? *Journal of Clinical Child and Adolescent Psychology, 40*, 80–86.
- Aldao, A. (2013). The future of emotion regulation research: Capturing context. *Perspectives on Psychological Science, 8*, 155–172.
- Aldao, A., & De Los Reyes, A. (2015). Commentary: A practical guide for translating basic research on affective science to implementing physiology in clinical child and adolescent assessments. *Journal of Clinical Child and Adolescent Psychology, 44*, 341–351.
- Aldao, A., & De Los Reyes, A. (2016). Introduction to the special section: Toward implementing physiological measures in clinical assessments of adult mental health. *Journal of Psychopathology and Behavioral Assessment, 38*, 1–4.
- Aldao, A., & Dixon-Gordon, K. L. (2014). Broadening the scope of research on emotion regulation strategies and psychopathology. *Cognitive Behaviour Therapy, 43*, 22–33.
- Aldao, A., Jazaieri, H., Goldin, P. R., & Gross, J. J. (2014). Adaptive and maladaptive emotion regulation strategies: Interactive effects during CBT for social anxiety disorder. *Journal of Anxiety Disorders, 28*, 382–389.
- Aldao, A., & Nolen-Hoeksema, S. (2012). When are adaptive strategies most predictive of psychopathology? *Journal of Abnormal Psychology, 121*, 276.
- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review, 30*, 217–237.
- American Psychiatric Association. (1987). *DSM-III-R: Diagnostic and statistical manual of mental disorders*. Washington, DC: Author.
- Anticevic, A., Van Snellenberg, J. X., & Barch, D. M. (2012). Neurobiology of emotional dysfunction in schizophrenia: New directions revealed through meta-analyses. *Biological Psychiatry, 71*, e23–e24; author reply e25.
- Banks, S. J., Eddy, K. T., Angstadt, M., Nathan, P. J., & Phan, K. L. (2007). Amygdala–frontal connectivity during emotion regulation. *Social Cognitive and Affective Neuroscience, 2*, 303–312.
- Barlow, D. H., Allen, L. B., & Choate, M. L. (2004). Toward a unified treatment for emotional disorders. *Behavior Therapy, 35*, 205–230.
- Barlow, D. H., Sauer-Zavala, S., Carl, J. R., Bullis, J. R., & Ellard, K. K. (2014). The nature, diagnosis, and treatment of neuroticism back to the future. *Clinical Psychological Science, 2*, 344–365.
- Beauchaine, T. P. (2001). Vagal tone, development, and Gray's motivational theory: Toward an integrated model of autonomic nervous system functioning in psychopathology. *Development and Psychopathology, 13*, 183.
- Beauchaine, T. P. (2015a). Future directions in emotion dysregulation and youth psychopathology. *Journal of Clinical Child & Adolescent Psychology, 44*, 875–896. doi:10.1080/15374416.2015.1038827.
- Beauchaine, T. P. (2015b). Respiratory sinus arrhythmia: A transdiagnostic biomarker of emotion dysregulation and psychopathology. *Current Opinion in Psychology, 3*, 43–47.
- Beauchaine, T. P., Gatzke-Kopp, L., & Mead, H. K. (2007). Polyvagal theory and developmental psychopathology: Emotion dysregulation and conduct problems from preschool to adolescence. *Biological Psychology, 74*, 174–184.
- Beauchaine, T. P., & Klein, D. N. (in press). Classifying psychopathology: The DSM, empirically-based taxonomies, and the Research Domain Criteria. In T. P. Beauchaine & S. P. Hinshaw (Eds.), *Child and adolescent psychopathology* (3rd ed.). Hoboken, NJ: Wiley.
- Beauchaine, T. P., & McNulty, T. (2013). Comorbidities and continuities as ontogenic processes: Toward a developmental spectrum model of externalizing psychopathology. *Development and Psychopathology, 25*, 1505–1528.
- Beauchaine, T. P., & Thayer, J. F. (2015). Heart rate variability as a transdiagnostic biomarker of psychopathology. *International Journal of Psychophysiology, 98*, 338–350.
- Bierman, K. L., Kalvin, C. B., & Heinrichs, B. S. (2015). Early childhood precursors and adolescent sequelae of grade school peer rejection and victimization. *Journal of Clinical Child and Adolescent Psychology, 44*, 367–379.
- Blair, C., & Diamond, A. (2008). Biological processes in prevention and intervention: The promotion of self-regulation as a means of preventing school failure. *Developmental Psychopathology, 20*, 899–911.
- Blair, R. J. R. (2008). The amygdala and ventromedial prefrontal cortex: Functional contributions and dysfunction in psychopathy. *Philosophical Transactions of the Royal Society of London B: Biological Sciences, 363*, 2557–2565.
- Blair, R. J. R. (2013). The neurobiology of psychopathic traits in youths. *Nature Reviews Neuroscience, 14*, 786–799.
- Boyce, W. T., Quas, J., Alkon, A., Smider, N. A., Essex, M. J., Kupfer, D. J., et al. (2001). Autonomic reactivity and psychopathology in middle childhood. *British Journal of Psychiatry, 179*, 144–150.
- Brière, F. N., Rohde, P., Seeley, J. R., Klein, D., & Lewinsohn, P. M. (2014). Comorbidity between major depression and alcohol use disorder from adolescence to adulthood. *Comprehensive Psychiatry, 55*, 526–533.
- Brown, T. A. (2007). Temporal course and structural relationships among dimensions of temperament and DSM-IV anxiety and mood disorder constructs. *Journal of Abnormal Psychology, 116*, 313.
- Brown, T. A., Campbell, L. A., Lehman, C. L., Grisham, J. R., & Mancill, R. B. (2001). Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *Journal of Abnormal Psychology, 110*, 585.
- Bubier, J. L., & Drabick, D. A. G. (2009). Co-occurring anxiety and disruptive behavior disorders: The roles of anxious symptoms, reactive aggression, and shared risk processes. *Clinical Psychology Review, 29*, 658–669.
- Buckholtz, J. W., & Meyer-Lindenberg, A. (2012). Psychopathology and the human connectome: Toward a transdiagnostic model of risk for mental illness. *Neuron, 74*, 990–1004.
- Buehler, C., Lange, G., & Franck, K. L. (2007). Adolescents' cognitive and emotional responses to marital hostility. *Child Development, 78*, 775–789.
- Bunge, S. A., Dudukovic, N. M., Thomason, M. E., Vaidya, C. J., & Gabrieli, J. D. (2002). Immature frontal lobe contributions to cognitive control in children: Evidence from fMRI. *Neuron, 33*, 301–311.
- Burghy, C. A., Stodola, D. E., Ruttle, P. L., Molloy, E. K., Armstrong, J. M., Oler, J. A., et al. (2012). Developmental pathways to amygdala–prefrontal function and internalizing symptoms in adolescence. *Nature Neuroscience, 15*, 1736–1741.
- Carstensen, L. L., Isaacowitz, D. M., & Charles, S. T. (1999). Taking time seriously: A theory of socioemotional selectivity. *American Psychologist, 54*, 165.
- Carstensen, L. L., Turan, B., Scheibe, S., Ram, N., Ersner-Hershfield, H., Samanez-Larkin, G. R., et al. (2011). Emotional experience improves with age: Evidence based on over 10 years of experience sampling. *Psychology and Aging, 26*, 21–33.
- Casey, B. J., Craddock, N., Cuthbert, B. N., Hyman, S. E., Lee, F. S., & Resler, K. J. (2013). DSM-5 and RDoC: Progress in psychiatry research? *Nature Reviews Neuroscience, 14*, 810–814.
- Casey, B. J., Epstein, J. N., Buhle, J., Liston, C., Davidson, M. C., Tonev, S. T., et al. (2007). Frontostriatal connectivity and its role in cognitive control in parent–child dyads with ADHD. *American Journal of Psychiatry, 164*, 1729–1736.
- Casey, B. J., Galván, A., & Somerville, L. H. (2016). Beyond simple models of adolescence to an integrated circuit-based account: A commentary. *Developmental Cognitive Neuroscience, 17*, 128–130.
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., et al. (2014). The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science, 2*, 119–137.
- Chaplin, T. M., & Aldao, A. (2013). Gender differences in emotion expression in children: A meta-analytic review. *Psychological Bulletin, 139*, 735.
- Chareyron, L. J., Lavenex, P. B., Amaral, D. G., & Lavenex, P. (2012). Postnatal development of the amygdala: A stereological study in macaque monkeys. *Journal of Comparative Neurology, 520*, 1965–1984.
- Christensen, K. A., Aldao, A., Sheridan, M., & McLaughlin, K. A. (in press). Habitual reappraisal in context: Peer victimisation moderates its association with physiological reactivity to social stress. *Cognition and Emotion*. doi:10.1080/02699931.2015.1103701
- Cicchetti, D. (1984). The emergence of developmental psychopathology. *Child Development, 55*, 1–7.
- Cicchetti, D. (1989). Developmental psychopathology: Some thoughts on its evolution. *Development and Psychopathology, 1*, 1–4.
- Cicchetti, D., Ackerman, B. P., & Izard, C. E. (1995). Emotions and emotion regulation in developmental psychopathology. *Development and Psychopathology, 7*, 1–10.
- Cole, P. M., Michel, M. K., & Teti, L. O. (1994). The development of emotion regulation and dysregulation: A clinical perspective. *Monographs of the Society for Research in Child Development, 59*, 73–102.

- Conklin, L. R., Cassiello-Robbins, C., Brake, C. A., Sauer-Zavala, S., Farchione, T. J., Ciraulo, D. A., et al. (2015). Relationships among adaptive and maladaptive emotion regulation strategies and psychopathology during the treatment of comorbid anxiety and alcohol use disorders. *Behaviour Research and Therapy*, *73*, 124–130.
- Dalley, J. W., Mar, A. C., Economidou, D., & Robbins, T. W. (2008). Neurobehavioral mechanisms of impulsivity: Fronto-striatal systems and functional neurochemistry. *Pharmacology Biochemistry and Behavior*, *90*, 250–260.
- De Bellis, M. D., Casey, B. J., Dahl, R. E., Birmaher, B., Williamson, D. E., Thomas, K. M., et al. (2000). A pilot study of amygdala volumes in pediatric generalized anxiety disorder. *Biological Psychiatry*, *48*, 51–57.
- De Brito, S. A., Mechelli, A., Wilke, M., Laurens, K. R., Jones, A. P., Barker, G. J., et al. (2009). Size matters: Increased grey matter in boys with conduct problems and callous-unemotional traits. *Brain*, *132*, 843–852.
- Decety, J., Norman, G. J., Berntson, G. G., & Cacioppo, J. T. (2012). A neurobehavioral evolutionary perspective on the mechanisms underlying empathy. *Progress in Neurobiology*, *98*, 38–48.
- Delgado, M. R., Nearing, K. I., LeDoux, J. E., & Phelps, E. A. (2008). Neural circuitry underlying the regulation of conditioned fear and its relation to extinction. *Neuron*, *59*, 829–838.
- De Los Reyes, A. (2013). Strategic objectives for improving understanding of informant discrepancies in developmental psychopathology research. *Development and Psychopathology*, *25*, 669–682.
- De Los Reyes, A., & Aldao, A. (2015). Introduction to the special issue: Toward implementing physiological measures in clinical child and adolescent assessments. *Journal of Clinical Child and Adolescent Psychology*, *44*, 221–237.
- De Los Reyes, A., Alfano, C. A., Lau, S., Augenstein, T. M., & Borelli, J. L. (2016). Can we use convergence between caregiver reports of adolescent mental health to index severity of adolescent mental health concerns? *Journal of Child and Family Studies*, *25*, 109–123.
- De Los Reyes, A., Augenstein, T. M., Wang, M., Thomas, S. A., Drabick, D. A. G., Burgers, D. E., et al. (2015). The validity of the multi-informant approach to assessing child and adolescent mental health. *Psychological Bulletin*, *141*, 858–900.
- De Los Reyes, A., Bunnell, B. E., & Beidel, D. C. (2013). Informant discrepancies in adult social anxiety disorder assessments: Links with contextual variations in observed behavior. *Journal of Abnormal Psychology*, *122*, 376.
- De Los Reyes, A., Henry, D. B., Tolan, P. H., & Wakschlag, L. S. (2009). Linking informant discrepancies to observed variations in young children's disruptive behavior. *Journal of Abnormal Child Psychology*, *37*, 637–652.
- De Los Reyes, A., & Kazdin, A. E. (2005). Informant discrepancies in the assessment of childhood psychopathology: A critical review, theoretical framework, and recommendations for further study. *Psychological Bulletin*, *131*, 483–509.
- De Los Reyes, A., Thomas, S. A., Goodman, K. L., & Kundey, S. M. (2013). Principles underlying the use of multiple informants' reports. *Annual Review of Clinical Psychology*, *9*, 123.
- Daykin, E. Y., Levy, J. C., & Wells, V. (1987). Adolescent depression, alcohol and drug abuse. *American Journal of Public Health*, *77*, 178–182.
- Dixon-Gordon, K. L., Aldao, A., & De Los Reyes, A. (2015). Emotion regulation in context: Examining the spontaneous use of strategies across emotional intensity and type of emotion. *Personality and Individual Differences*, *86*, 271–276.
- Drabick, D. A. G., & Kendall, P. C. (2010). Developmental psychopathology and the diagnosis of mental health problems among youth. *Clinical Psychology: Science and Practice*, *17*, 272–280.
- Drabick, D. A. G., Ollendick, T. H., & Bubier, J. L. (2010). Co-occurrence of ODD and anxiety: Shared risk processes and evidence for a dual-pathway model. *Clinical Psychology*, *17*, 307–318.
- Egan, S. J., Wade, T. D., & Shafran, R. (2011). Perfectionism as a transdiagnostic process: A clinical review. *Clinical Psychology Review*, *31*, 203–212.
- Ehrenreich-May, J., & Bilek, E. L. (2012). The development of a transdiagnostic, cognitive behavioral group intervention for childhood anxiety disorders and co-occurring depression symptoms. *Cognitive and Behavioral Practice*, *19*, 41–55.
- Ehrenreich-May, J., Queen, A. H., Bilek, E. L., Remmes, C. R., & Marciel, K. K. (2013). The unified protocols for the treatment of emotional disorders in children and adolescents. In D. H. Barlow, T. J. Farchione, C. P. Fairholme, K. K. Ellard, C. L. Boisseau, L. B. Allen, et al. (Eds.), *Transdiagnostic treatments for children and adolescents: Principles and practice* (pp. 267–292). Oxford: Oxford University Press.
- Ehring, T., & Watkins, E. R. (2008). Repetitive negative thinking as a transdiagnostic process. *International Journal of Cognitive Therapy*, *1*, 192–205.
- Eisenberg, N., & Fabes, R. A. (1992). Emotion, regulation, and the development of social competence. In M. S. Clark (Ed.), *Review of personality and social psychology: Vol. 14. Emotion and social behavior* (pp. 119–150). Newbury Park, CA: Sage.
- Erk, S., Mikschl, A., Stier, S., Ciaramidaro, A., Gapp, V., Weber, B., et al. (2010). Acute and sustained effects of cognitive emotion regulation in major depression. *Journal of Neuroscience*, *30*, 15726–15734.
- Etkin, A., Büchel, C., & Gross, J. J. (2015). The neural bases of emotion regulation. *Nature Reviews Neuroscience*, *16*, 693–700.
- Fairholme, C. P., Boisseau, C. L., Ellard, K. K., Ehrenreich, J. T., & Barlow, D. H. (2009). Emotions, emotion regulation, and psychological treatment: A unified perspective. In A. M. Kring & D. M. Sloan (Eds.), *Emotion regulation and psychopathology: A transdiagnostic approach to etiology and treatment* (pp. 283–309). New York: Guilford Press.
- Fakra, E., Salgado-Pineda, P., Delaveau, P., Hariri, A. R., & Blin, O. (2008). Neural bases of different cognitive strategies for facial affect processing in schizophrenia. *Schizophrenia Research*, *100*, 191–205.
- Falk, A. E., Lee, S. S., & Chorpita, B. F. (2015). Differential association of youth attention-deficit/hyperactivity disorder and anxiety with delinquency and aggression. *Journal of Clinical Child and Adolescent Psychology*. Advance online publication.
- First, M. B. (2005). Mutually exclusive versus co-occurring diagnostic categories: The challenge of diagnostic comorbidity. *Psychopathology*, *38*, 206–210.
- Forehand, R., Jones, D. J., & Parent, J. (2013). Behavioral parenting interventions for child disruptive behaviors and anxiety: What's different and what's the same. *Clinical Psychology Review*, *33*, 133–145.
- Fraire, M. G., & Ollendick, T. H. (2013). Anxiety and oppositional defiant disorder: A transdiagnostic conceptualization. *Clinical Psychology Review*, *33*, 229–240.
- Frances, A. (2013). *Saving normal: An insider's revolt against out-of-control psychiatric diagnosis, DSM-5, big pharma, and the medicalization of ordinary life* (1st ed.). New York: William Morrow.
- Gabard-Durnam, L. J., Flannery, J., Goff, B., Gee, D. G., Humphreys, K. L., Telzer, E., et al. (2014). The development of human amygdala functional connectivity at rest from 4 to 23 years: A cross-sectional study. *NeuroImage*, *95*, 193–207.
- Gaffrey, M. S., Luby, J. L., Belden, A. C., Hirshberg, J. S., Volsch, J., & Barch, D. M. (2011). Association between depression severity and amygdala reactivity during sad face viewing in depressed preschoolers: An fMRI study. *Journal of Affective Disorders*, *129*, 364–370.
- Galvan, A., Hare, T. A., Parra, C. E., Penn, J., Voss, H., Glover, G., et al. (2006). Earlier development of the accumbens relative to orbitofrontal cortex might underlie risk-taking behavior in adolescents. *Journal of Neuroscience*, *26*, 6885–6892.
- Garner, W. R., Hake, H. W., & Eriksen, C. W. (1956). Operationism and the concept of perception. *Psychological Review*, *63*, 149.
- Gee, D. G., & Casey, B. J. (2015). The impact of developmental timing for stress and recovery. *Neurobiology of Stress*, *1*, 184–194.
- Gee, D. G., Fetcho, R. N., Jing, D., Li, A., Glatt, C. E., Drysdale, A. T., et al. (2016). Individual differences in frontolimbic circuitry and anxiety emerge with adolescent changes in endocannabinoid signaling across species. *Proceedings of the National Academy of Sciences*, *113*, 4500–4505.
- Gee, D. G., Gabard-Durnam, L. J., Flannery, J., Goff, B., Humphreys, K. L., Telzer, E. H., et al. (2013). Early developmental emergence of human amygdala–prefrontal connectivity after maternal deprivation. *Proceedings of the National Academy of Sciences*, *110*, 15638–15643.
- Gee, D. G., Gabard-Durnam, L., Telzer, E. H., Humphreys, K. L., Goff, B., Shapiro, M., et al. (2014). Maternal buffering of human amygdala–prefrontal circuitry during childhood but not during adolescence. *Psychological Science*, *25*, 2067–2078.
- Gee, D. G., Humphreys, K. L., Flannery, J., Goff, B., Telzer, E. H., Shapiro, M., et al. (2013). A developmental shift from positive to negative connectivity in human amygdala–prefrontal circuitry. *Journal of Neuroscience*, *33*, 4584–4593.
- Gee, D. G., Karlsgodt, K. H., van Erp, T. G., Bearden, C. E., Lieberman, M. D., Belger, A., et al. (2012). Altered age-related trajectories of amygdala–prefrontal circuitry in adolescents at clinical high risk for psychosis: A preliminary study. *Schizophrenia Research*, *134*, 1–9.

- Glenn, A. L., & Yang, Y. (2012). The potential role of the striatum in antisocial behavior and psychopathy. *Biological Psychiatry*, *72*, 817–822.
- Goldin, P. R., McRae, K., Ramel, W., & Gross, J. J. (2008). The neural bases of emotion regulation: Reappraisal and suppression of negative emotion. *Biological Psychiatry*, *63*, 577–586.
- Goldstein, R. Z., & Volkow, N. D. (2011). Dysfunction of the prefrontal cortex in addiction: Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, *12*, 652–669.
- Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment*, *26*, 41–54.
- Graziano, P., & Derefinko, K. (2013). Cardiac vagal control and children's adaptive functioning: A meta-analysis. *Biological Psychology*, *94*, 22–37.
- Gross, J. J. (1998). The emerging field of emotion regulation: An integrative review. *Review of General Psychology*, *2*, 271.
- Gross, J. J. (2015). Emotion regulation: Current status and future prospects. *Psychological Inquiry*, *26*, 1–26.
- Gross, J. J., & Barrett, L. F. (2011). Emotion generation and emotion regulation: One or two depends on your point of view. *Emotion Review*, *3*, 8–16.
- Gross, J. J., & Jazaieri, H. (2014). Emotion, emotion regulation, and psychopathology: An affective science perspective. *Clinical Psychological Science*, *2*, 387–401.
- Gunnar, M. R., & Donzella, B. (2002). Social regulation of the cortisol levels in early human development. *Psychoneuroendocrinology*, *27*, 199–220.
- Guyer, A. E., Monk, C. S., McClure-Tone, E. B., Nelson, E. E., Roberson-Nay, R., Adler, A. D., et al. (2008). A developmental examination of amygdala response to facial expressions. *Journal of Cognitive Neuroscience*, *20*, 1565–1582.
- Halligan, S. L., Cooper, P. J., Fearon, P., Wheeler, S. L., Crosby, M., & Murray, L. (2013). The longitudinal development of emotion regulation capacities in children at risk for externalizing disorders. *Development and Psychopathology*, *25*, 391–406.
- Hankin, B. L. (2008). Rumination and depression in adolescence: Investigating symptom specificity in a multiwave prospective study. *Journal of Clinical Child and Adolescent Psychology*, *37*, 701–713.
- Hanson, J. L., Chung, M. K., Avants, B. B., Shirliff, E. A., Gee, J. C., Davidson, R. J., et al. (2010). Early stress is associated with alterations in the orbitofrontal cortex: A tensor-based morphometry investigation of brain structure and behavioral risk. *Journal of Neuroscience*, *30*, 7466–7472.
- Hare, T. A., Tottenham, N., Galvan, A., Voss, H. U., Glover, G. H., & Casey, B. J. (2008). Biological substrates of emotional reactivity and regulation in adolescence during an emotional go-nogo task. *Biological Psychiatry*, *63*, 927–934.
- Hartley, A. G., Zakriski, A. L., & Wright, J. C. (2011). Probing the depths of informant discrepancies: Contextual influences on divergence and convergence. *Journal of Clinical Child and Adolescent Psychology*, *40*, 54–66.
- Harvey, A. G., Watkins, E., Mansell, W., & Shafran, R. (2004). *Cognitive behavioural processes across psychological disorders: A transdiagnostic approach to research and treatment*. New York: Oxford University Press.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (1999). *Acceptance and commitment therapy*. New York: Guilford Press.
- Heatherton, T. F., & Wagner, D. D. (2011). Cognitive neuroscience of self-regulation failure. *Trends in Cognitive Sciences*, *15*, 132–139.
- Heleniak, C., Jenness, J. L., Stoep, A. V., McCauley, E., & McLaughlin, K. A. (2015). Childhood maltreatment exposure and disruptions in emotion regulation: A transdiagnostic pathway to adolescent internalizing and externalizing psychopathology. *Cognitive Therapy and Research*. Advance online publication.
- Heller, A. S. (2016). Cortical–subcortical interactions in depression: From animal models to human psychopathology. *Frontiers in Systems Neuroscience*. Advance online publication. doi:10.3389/fnsys.2016.00020.
- Heller, A. S., Johnstone, T., Shackman, A. J., Light, S. N., Peterson, M. J., Kolden, G. G., et al. (2009). Reduced capacity to sustain positive emotion in major depression reflects diminished maintenance of fronto-striatal brain activation. *Proceedings of the National Academy of Sciences*, *106*, 22445–22450.
- Herringa, R. J., Birn, R. M., Ruttler, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., et al. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences*, *110*, 19119–19124.
- Hettema, J. M., Prescott, C. A., & Kendler, K. S. (2003). The effects of anxiety, substance use and conduct disorders on risk of major depressive disorder. *Psychological Medicine*, *33*, 1423–1432.
- Hilt, L. M., Armstrong, J. M., & Essex, M. J. (2015). Rumination and moderators of multifinality: Predicting internalizing symptoms and alcohol use during adolescence. *Journal of Clinical Child and Adolescent Psychology*. Advance online publication.
- Hinnant, J. B., & El-Sheikh, M. (2009). Children's externalizing and internalizing symptoms over time: The role of individual differences in patterns of RSA responding. *Journal of Abnormal Child Psychology*, *37*, 1049–1061.
- Hinnant, J. B., & El-Sheikh, M. (2013). Codevelopment of externalizing and internalizing symptoms in middle to late childhood: Sex, baseline respiratory sinus arrhythmia, and respiratory sinus arrhythmia reactivity as predictors. *Development and Psychopathology*, *25*, 419–436.
- Hinshaw, S. P. (2015). Developmental psychopathology, ontogenic process models, gene–environment interplay, and brain development: An emerging synthesis. *Journal of Abnormal Psychology*, *124*, 771.
- Hofer, M. A. (1994). Early relationships as regulators of infant physiology and behavior. *Acta Paediatrica*, *83*, 9–18.
- Hofmann, S. G. (2014). Interpersonal emotion regulation model of mood and anxiety disorders. *Cognitive Therapy and Research*, *38*, 483–492.
- Hollenstein, T., Granic, I., Stoolmiller, M., & Snyder, J. (2004). Rigidity in parent–child interactions and the development of externalizing and internalizing behavior in early childhood. *Journal of Abnormal Child Psychology*, *32*, 595–607.
- Hulvershorn, L. A., Cullen, K., & Anand, A. (2011). Toward dysfunctional connectivity: A review of neuroimaging findings in pediatric major depressive disorder. *Brain Imaging and Behavior*, *5*, 307–328.
- Hulvershorn, L. A., Mennes, M., Castellanos, F. X., Di Martino, A., Milham, M. P., Hummer, T. A., et al. (2014). Abnormal amygdala functional connectivity associated with emotional lability in children with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, *53*, 351–361.
- Hunsley, J., & Mash, E. J. (2007). Evidence-based assessment. *Annual Review of Clinical Psychology*, *3*, 29.
- Hussong, A. M., Jones, D. J., Stein, G. L., Baucom, D. H., & Boeding, S. (2011). An internalizing pathway to alcohol use and disorder. *Psychology of Addictive Behaviors*, *25*, 390.
- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., et al. (2010). Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*, *167*, 748–751.
- Isaacowitz, D. M., & Blanchard-Fields, F. (2012). Linking process and outcome in the study of emotion and aging. *Perspectives on Psychological Science*, *7*, 3–17.
- Izard, C. E., Trentacosta, C. J., King, K. A., & Mostow, A. J. (2004). An emotion-based prevention program for Head Start children. *Early Education and Development*, *15*, 407–422.
- Jarett, M. A., & Ollendick, T. H. (2008). A conceptual review of the comorbidity of attention-deficit/hyperactivity disorder and anxiety: Implications for future research and practice. *Clinical Psychology Review*, *28*, 1266–1280.
- Jose, P. E., & Brown, I. (2008). When does the gender difference in rumination begin? Gender and age differences in the use of rumination by adolescents. *Journal of Youth and Adolescence*, *37*, 180–192.
- Karalunas, S. L., Fair, D., Musser, E. D., Aykes, K., Iyer, S. P., & Nigg, J. T. (2014). Subtyping attention-deficit/hyperactivity disorder using temperament dimensions: Toward biologically based nosologic criteria. *JAMA Psychiatry*, *71*, 1015–1024.
- Kashdan, T. B., Ferrisizidis, P., Collins, R. L., & Muraven, M. (2010). Emotion differentiation as resilience against excessive alcohol use: An ecological momentary assessment in underage social drinkers. *Psychological Science*, *21*, 1341–1347.
- Kessler, R. C., Chiu, W. T., Demler, O., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*, 617–627.
- Kessler, R. C., Ormel, J., Petukhova, M., McLaughlin, K. A., Green, J. G., Russo, L. J., et al. (2011). Development of lifetime comorbidity in the World Health Organization world mental health surveys. *Archives of General Psychiatry*, *68*, 90–100.
- Kim, H., Somerville, L. H., Johnstone, T., Alexander, A. L., & Whalen, P. J. (2003). Inverse amygdala and medial prefrontal cortex responses to surprised faces. *NeuroReport*, *14*, 2317–2322.

- Kim, J., & Cicchetti, D. (2010). Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *51*, 706–716.
- Kim, M. J., Loucks, R. A., Palmer, A. L., Brown, A. C., Solomon, K. M., Marchante, A. N., et al. (2011). The structural and functional connectivity of the amygdala: From normal emotion to pathological anxiety. *Behavioural Brain Research*, *223*, 403–410.
- Koob, G. F., & Le Moal, M. (2001). Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology*, *24*, 97–129.
- Kring, A. M., & Caponigro, J. M. (2010). Emotion in schizophrenia where feeling meets thinking. *Current Directions in Psychological Science*, *19*, 255–259.
- Kring, A. M., & Sloan, D. M. (2009). *Emotion regulation and psychopathology: A transdiagnostic approach to etiology and treatment*. New York: Guilford Press.
- Kross, E., Ayduk, O., & Mischel, W. (2005). When asking “why” does not hurt: Distinguishing rumination from reflective processing of negative emotions. *Psychological Science*, *16*, 709–715.
- Kushner, M. G., Sher, K. J., & Beitman, B. D. (1990). The relation between alcohol problems and the anxiety disorders. *American Journal of Psychiatry*, *147*, 685–695.
- Lebel, C., Gee, M., Camicioli, R., Wieler, M., Martin, W., & Beaulieu, C. (2012). Diffusion tensor imaging of white matter tract evolution over the lifespan. *NeuroImage*, *60*, 340–352.
- Lee, F. S., Heimer, H., Giedd, J. N., Lein, E. S., Šestan, N., Weinberger, D. R., et al. (2014). Adolescent mental health—Opportunity and obligation. *Science*, *346*, 547–549.
- Lee, S. S., Falk, A. E., & Aguirre, V. P. (2012). Association of comorbid anxiety with social functioning in school-age children with and without attention-deficit/hyperactivity disorder (ADHD). *Psychiatry Research*, *197*, 90–96.
- Lenroot, R. K., & Giedd, J. N. (2006). Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience & Biobehavioral Reviews*, *30*, 718–729.
- Levy, J. C., & Deykin, E. Y. (1989). Suicidality, depression, and substance abuse in adolescence. *American Journal of Psychiatry*, *146*, 1462–1467.
- Lewinsohn, P. M., Gotlib, I. H., & Seeley, J. R. (1995). Adolescent psychopathology: IV. Specificity of psychosocial risk factors for depression and substance abuse in older adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*, *34*, 1221–1229.
- Lieberman, M. D., Eisenberger, N. I., Crockett, M. J., Tom, S. M., Pfeifer, J. H., & Way, B. M. (2007). Putting feelings into words: Affect labeling disrupts amygdala activity in response to affective stimuli. *Psychological Science*, *18*, 421–428.
- Linehan, M. (1993). *Cognitive-behavioral treatment of borderline personality disorder*. New York: Guilford Press.
- Livingstone, K. M., & Isaacowitz, D. M. (2015). Situation selection and modification for emotion regulation in younger and older adults. *Social Psychological and Personality Science*, *6*, 904–910.
- Luking, K. R., Repovs, G., Belden, A. C., Gaffrey, M. S., Botteron, K. N., Luby, J. L., et al. (2011). Functional connectivity of the amygdala in early-childhood-onset depression. *Journal of the American Academy of Child & Adolescent Psychiatry*, *50*, 1027–1041.
- Machado, C. J., & Bachevalier, J. (2003). Non-human primate models of childhood psychopathology: The promise and the limitations. *Journal of Child Psychology and Psychiatry*, *44*, 64–87.
- Maier, S. J., Szalkowski, A., Kamphausen, S., Feige, B., Perlov, E., Kalisch, R., et al. (2014). Altered cingulate and amygdala response towards threat and safe cues in attention deficit hyperactivity disorder. *Psychological Medicine*, *44*, 85–98.
- Marroquín, B. (2011). Interpersonal emotion regulation as a mechanism of social support in depression. *Clinical Psychology Review*, *31*, 1276–1290.
- Marsh, A. A., Finger, E. C., Mitchell, D. G., Reid, M. E., Sims, C., Kosson, D. S., et al. (2008). Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *American Journal of Psychiatry*, *165*, 712–720.
- Masten, A. S., & Cicchetti, D. (2010). Developmental cascades. *Development and Psychopathology*, *22*, 491–495.
- Mauss, I. B., & Robinson, M. D. (2009). Measures of emotion: A review. *Cognition and Emotion*, *23*, 209–237.
- McCoy, C. L., & Masters, J. C. (1985). The development of children’s strategies for the social control of emotion. *Child Development*, *56*, 1214–1222.
- McLaughlin, K. A., Aldao, A., Wisco, B. E., & Hilt, L. M. (2014). Rumination as a transdiagnostic factor underlying transitions between internalizing symptoms and aggressive behavior in early adolescents. *Journal of Abnormal Psychology*, *123*, 13.
- McLaughlin, K. A., Hatzenbuehler, M. L., Mennin, D. S., & Nolen-Hoeksema, S. (2011). Emotion dysregulation and adolescent psychopathology: A prospective study. *Behaviour Research and Therapy*, *49*, 544–554.
- McRae, K., Gross, J. J., Weber, J., Robertson, E. R., Sokol-Hessner, P., Ray, R. D., et al. (2012). The development of emotion regulation: An fMRI study of cognitive reappraisal in children, adolescents and young adults. *Social Cognitive and Affective Neuroscience*, *7*, 11–22.
- Mennin, D., Biederman, J., Mick, E., & Faraone, S. V. (2000). Towards defining a meaningful anxiety phenotype for research in ADHD children. *Journal of Attention Disorders*, *3*, 192–199.
- Mischel, W., & Shoda, Y. (1995). A cognitive-affective system theory of personality: Reconceptualizing situations, dispositions, dynamics, and invariance in personality structure. *Psychological Review*, *102*, 246.
- Moffitt, T. E., Harrington, H., Caspi, A., Kim-Cohen, J., Goldberg, D., Gregory, A. M., et al. (2007). Depression and generalized anxiety disorder: Cumulative and sequential comorbidity in a birth cohort followed prospectively to age 32 years. *Archives of General Psychiatry*, *64*, 651–660.
- Monk, C. S., Telzer, E. H., Mogg, K., Bradley, B. P., Mai, X., Louro, H. M., et al. (2008). Amygdala and ventrolateral prefrontal cortex activation to masked angry faces in children and adolescents with generalized anxiety disorder. *Archives of General Psychiatry*, *65*, 568–576.
- New, A. S., Hazlett, E. A., Buchsbaum, M. S., Goodman, M., Mitelman, S. A., Newmark, R., et al. (2007). Amygdala-prefrontal disconnection in borderline personality disorder. *Neuropsychopharmacology*, *32*, 1629–1640.
- Nock, M. K., Wedig, M. M., Holmberg, E. B., & Hooley, J. M. (2008). The emotion reactivity scale: Development, evaluation, and relation to self-injurious thoughts and behaviors. *Behavior Therapy*, *39*, 107–116.
- Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology*, *100*, 569.
- Nolen-Hoeksema, S., & Aldao, A. (2011). Gender and age differences in emotion regulation strategies and their relationship to depressive symptoms. *Personality and Individual Differences*, *51*, 704–708.
- Nolen-Hoeksema, S., Larson, J., & Grayson, C. (1999). Explaining the gender difference in depressive symptoms. *Journal of Personality and Social Psychology*, *77*, 1061.
- Nolen-Hoeksema, S., Stice, E., Wade, E., & Bohon, C. (2007). Reciprocal relations between rumination and bulimic, substance abuse, and depressive symptoms in female adolescents. *Journal of Abnormal Psychology*, *116*, 198.
- Nolen-Hoeksema, S., & Watkins, E. R. (2011). A heuristic for developing transdiagnostic models of psychopathology explaining multifinality and divergent trajectories. *Perspectives on Psychological Science*, *6*, 589–609.
- Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspectives on Psychological Science*, *3*, 400–424.
- Ochsner, K. N., Bunge, S. A., Gross, J. J., & Gabrieli, J. D. (2002). Rethinking feelings: An fMRI study of the cognitive regulation of emotion. *Journal of Cognitive Neuroscience*, *14*, 1215–1229.
- Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, *9*, 242–249.
- Oldehinkel, A. J., Verhulst, F. C., & Ormel, J. (2008). Low heart rate: A marker of stress resilience. The TRAILS study. *Biological Psychiatry*, *63*, 1141–1146.
- Pang, K. C., & Beauchaine, T. P. (2013). Longitudinal patterns of autonomic nervous system responding to emotion evocation among children with conduct problems and/or depression. *Developmental Psychobiology*, *55*, 698–706.
- Passarotti, A. M., Ellis, J., Wegbreit, E., Stevens, M. C., & Pavuluri, M. N. (2012). Reduced functional connectivity of prefrontal regions and amygdala within affect and working memory networks in pediatric bipolar disorder. *Brain Connectivity*, *2*, 320–334.
- Patrick, C. J., & Hajcak, G. (2016). Reshaping clinical science: Introduction to the special issue on psychophysiology and the NIMH Research Domain Criteria (RDoC) initiative. *Psychophysiology*, *53*, 281–285.
- Payne, C., Machado, C. J., Bliwise, N. G., & Bachevalier, J. (2010). Maturation of the hippocampal formation and amygdala in *Macaca mulatta*: A volumetric magnetic resonance imaging study. *Hippocampus*, *20*, 922–935.
- Peled, M., & Moretti, M. M. (2007). Rumination on anger and sadness in adolescence: Fueling of fury and deepening of despair. *Journal of Clinical Child and Adolescent Psychology*, *36*, 66–75.

- Perlman, S. B., & Pelphrey, K. A. (2011). Developing connections for affective regulation: Age-related changes in emotional brain connectivity. *Journal of Experimental Child Psychology, 108*, 607–620.
- Pezawas, L., Meyer-Lindenberg, A., Drabant, E. M., Verchinski, B. A., Munoz, K. E., Kolachana, B. S., et al. (2005). 5-HTTLPR polymorphism impacts human cingulate–amygdala interactions: A genetic susceptibility mechanism for depression. *Nature Neuroscience, 8*, 828–834.
- Pfeifer, J. C., Welge, J., Strakowski, S. M., Adler, C., & Delbello, M. P. (2008). Meta-analysis of amygdala volumes in children and adolescents with bipolar disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 47*, 1289–1298.
- Pelphrey, E. A., Delgado, M. R., Nearing, K. I., & LeDoux, J. E. (2004). Extinction learning in humans: Role of the amygdala and vmPFC. *Neuron, 43*, 897–905.
- Pine, D. S., Cohen, P., Gurley, D., Brook, J., & Ma, Y. (1998). The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives of General Psychiatry, 55*, 56–64.
- Plichta, M. M., & Scheres, A. (2014). Ventral–striatal responsiveness during reward anticipation in ADHD and its relation to trait impulsivity in the healthy population: A meta-analytic review of the fMRI literature. *Neuroscience & Biobehavioral Reviews, 38*, 125–134.
- Plichta, M. M., Vasic, N., Wolf, R. C., Lesch, K.-P., Brummer, D., Jacob, C., et al. (2009). Neural hypo-responsiveness and hyper-responsiveness during immediate and delayed reward processing in adult attention-deficit/hyperactivity disorder. *Biological Psychiatry, 65*, 7–14.
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology, 74*, 116–143.
- Posner, J., Nagel, B. J., Maia, T. V., Mechling, A., Oh, M., Wang, Z., et al. (2011). Abnormal amygdalar activation and connectivity in adolescents with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 50*, 828–837.
- Prakash, R. S., Whitmoyer, P., Aldao, A., & Schirda, B. (2015). Mindfulness and emotion regulation in older and young adults. *Aging & Mental Health*. Advance online publication.
- Pruzan, K., & Isaacowitz, D. M. (2006). An attentional application of socio-emotional selectivity theory in college students. *Social Development, 15*, 326–338.
- Rauch, S. L., Shin, L. M., & Wright, C. I. (2003). Neuroimaging studies of amygdala function in anxiety disorders. *Annals of the New York Academy of Sciences, 985*, 389–410.
- Rich, B. A., Fromm, S. J., Berghorst, L. H., Dickstein, D. P., Brotman, M. A., Pine, D. S., et al. (2008). Neural connectivity in children with bipolar disorder: Impairment in the face emotion processing circuit. *Journal of Child Psychology and Psychiatry, 49*, 88–96.
- Roemer, L., Orsillo, S. M., & Salters-Pedneault, K. (2008). Efficacy of an acceptance-based behavior therapy for generalized anxiety disorder: Evaluation in a randomized controlled trial. *Journal of Consulting and Clinical Psychology, 76*, 1083.
- Rose, A. J. (2002). Co-rumination in the friendships of girls and boys. *Child Development, 73*, 1830–1843.
- Rose, A. J., Carlson, W., & Waller, E. M. (2007). Prospective associations of co-rumination with friendship and emotional adjustment: Considering the socioemotional trade-offs of co-rumination. *Developmental Psychology, 43*, 1019–1031.
- Roy, A. K., Fudge, J. L., Kelly, C., Perry, J. S. A., Daniele, T., Carlisi, C., et al. (2013). Intrinsic functional connectivity of amygdala-based networks in adolescent generalized anxiety disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 52*, 290–299.
- Rutter, M., Kim-Cohen, J., & Maughan, B. (2006). Continuities and discontinuities in psychopathology between childhood and adult life. *Journal of Child Psychology and Psychiatry, 47*, 276–295.
- Rutter, M., & Sroufe, L. A. (2000). Developmental psychopathology: Concepts and challenges. *Development and Psychopathology, 12*, 265–296.
- Sanislow, C. A., Pine, D. S., Quinn, K. J., Kozak, M. J., Garvey, M. A., Heinssen, R. K., et al. (2010). Developing constructs for psychopathology research: Research domain criteria. *Journal of Abnormal Psychology, 119*, 631.
- Schatz, D. B., & Rostain, A. L. (2006). ADHD with comorbid anxiety: A review of the current literature. *Journal of Attention Disorders, 10*, 141–149.
- Shadur, J. M., & Lejuez, C. W. (2015). Adolescent substance use and comorbid psychopathology: Emotion regulation deficits as a transdiagnostic risk factor. *Current Addiction Reports, 2*, 354–363.
- Shallcross, S. L., Frazier, P. A., & Anders, S. L. (2014). Social resources mediate the relations between attachment dimensions and distress following potentially traumatic events. *Journal of Counseling Psychology, 61*, 352.
- Shaw, P., Malek, M., Watson, B., Sharp, W., Evans, A., & Greenstein, D. (2012). Development of cortical surface area and gyrification in attention-deficit/hyperactivity disorder. *Biological Psychiatry, 72*, 191–197.
- Sher, K. J., & Grekin, E. R. (2007). Alcohol and affect regulation. In J. G. Gross (Ed.), *Handbook of emotion regulation* (pp. 560–580). New York: Guilford Press.
- Shields, A., & Cicchetti, D. (1997). Emotion regulation among school-age children: The development and validation of a new criterion Q-sort scale. *Developmental Psychology, 33*, 906–916.
- Shin, L. M., Wright, C. I., Cannistraro, P. A., Wedig, M. M., McMullin, K., Martis, B., et al. (2005). A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Archives of General Psychiatry, 62*, 273–281.
- Siegle, G. J., Thompson, W., Carter, C. S., Steinhilber, S. R., & Thase, M. E. (2007). Increased amygdala and decreased dorsolateral prefrontal BOLD responses in unipolar depression: Related and independent features. *Biological Psychiatry, 61*, 198–209.
- Silbersweig, D., Clarkin, J. F., Goldstein, M., Kernberg, O. F., Tiescher, O., Levy, K. N., et al. (2007). Failure of frontolimbic inhibitory function in the context of negative emotion in borderline personality disorder. *American Journal of Psychiatry, 164*, 1832–1841.
- Silk, J. S., Steinberg, L., & Morris, A. S. (2003). Adolescents' emotion regulation in daily life: Links to depressive symptoms and problem behavior. *Child Development, 74*, 1869–1880.
- Silvers, J. A., Weber, J., Wager, T. D., & Ochsner, K. N. (2014). Bad and worse: Neural systems underlying reappraisal of high- and low-intensity negative emotions. *Social Cognitive and Affective Neuroscience*. Advance online publication.
- Sims, T., Hogan, C. L., & Carstensen, L. L. (2015). Selectivity as an emotion regulation strategy: Lessons from older adults. *Current Opinion in Psychology, 3*, 80–84.
- Smith, R. L., & Rose, A. J. (2011). The “cost of caring” in youths' friendships: Considering associations among social perspective taking, co-rumination, and empathetic distress. *Developmental Psychology, 47*, 1792–1803.
- Somerville, L. H., Hare, T., & Casey, B. J. (2011). Frontostriatal maturation predicts cognitive control failure to appetitive cues in adolescents. *Journal of Cognitive Neuroscience, 23*, 2123–2134.
- Spasojević, J., & Alloy, L. B. (2001). Rumination as a common mechanism relating depressive risk factors to depression. *Emotion, 1*, 25.
- Sroufe, L. A., & Rutter, M. (1984). The domain of developmental psychopathology. *Child Development, 55*, 17–29.
- Steinberg, E. A., & Drabick, D. A. G. (2015). A developmental psychopathology perspective on ADHD and comorbid conditions: The role of emotion regulation. *Child Psychiatry and Human Development, 46*, 951–966.
- Sukhodolsky, D. G., Golub, A., & Cromwell, E. N. (2001). Development and validation of the anger rumination scale. *Personality and Individual Differences, 31*, 689–700.
- Svaldi, J., Griepentrog, J., Tuschen-Caffier, B., & Ehring, T. (2012). Emotion regulation deficits in eating disorders: A marker of eating pathology or general psychopathology? *Psychiatry Research, 197*, 103–111.
- Swartz, J. R., Williamson, D. E., & Hariri, A. R. (2014). Developmental change in amygdala reactivity during adolescence: Effects of family history of depression and stressful life events. *American Journal of Psychiatry, 172*, 276–283.
- Tajima-Pozo, K., Yus, M., Ruiz-Manrique, G., Lewczuk, A., Arrazola, J., & Montañes-Rada, F. (2016). Amygdala abnormalities in adults with ADHD. *Journal of Attention Disorders*. Advance online publication. doi:10.1177/1087054716629213
- Taylor, S. F., Kang, J., Brege, I. S., Tso, I. F., Hosanagar, A., & Johnson, T. D. (2012). Meta-analysis of functional neuroimaging studies of emotion perception and experience in schizophrenia. *Biological Psychiatry, 71*, 136–145.
- Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J., & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews, 36*, 747–756.
- Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *Journal of Affective Disorders, 61*, 201–216.

- Thompson, R. A. (1994). Emotion regulation: A theme in search of definition. *Monographs of the Society for Research in Child Development*, 59, 25–52.
- Tottenham, N., Hare, T. A., & Casey, B. J. (2011). Behavioral assessment of emotion discrimination, emotion regulation, and cognitive control in childhood, adolescence, and adulthood. *Frontiers in Psychology*, 2, 2.
- Tottenham, N., & Sheridan, M. A. (2009). A review of adversity, the amygdala and the hippocampus: A consideration of developmental timing. *Frontiers in Human Neuroscience*. Advance online publication. doi:0.3389/neuro.09.068.2009
- Townsend, J. D., Torrisi, S. J., Lieberman, M. D., Sugar, C. A., Bookheimer, S. Y., & Altshuler, L. L. (2013). Frontal–amygdala connectivity alterations during emotion downregulation in bipolar I disorder. *Biological Psychiatry*, 73, 127–135.
- Troy, A. S., Shallcross, A. J., & Mauss, I. B. (2013). A person-by-situation approach to emotion regulation cognitive reappraisal can either help or hurt, depending on the context. *Psychological Science*, 24, 2505–2514.
- Tull, M. T., & Aldao, A. (2015). Editorial overview: New directions in the science of emotion regulation. *Current Opinion in Psychology*, 3, iv–x.
- Van den Bergh, B. R., & Marcoen, A. (2004). High antenatal maternal anxiety is related to ADHD symptoms, externalizing problems, and anxiety in 8- and 9-year-olds. *Child Development*, 75, 1085–1097.
- Vink, M., Derks, J. M., Hoogendam, J. M., Hillegers, M., & Kahn, R. S. (2014). Functional differences in emotion processing during adolescence and early adulthood. *NeuroImage*, 91, 70–76.
- Wager, T. D., Davidson, M. L., Hughes, B. L., Lindquist, M. A., & Ochsner, K. N. (2008). Prefrontal-subcortical pathways mediating successful emotion regulation. *Neuron*, 59, 1037–1050.
- Wang, F., Kalmar, J. H., He, Y., Jackowski, M., Chepenik, L. G., Edmiston, E. E., et al. (2009). Functional and structural connectivity between the perigenual anterior cingulate and amygdala in bipolar disorder. *Biological Psychiatry*, 66, 516–521.
- Webster-Stratton, C., Jamila Reid, M., & Stoolmiller, M. (2008). Preventing conduct problems and improving school readiness: Evaluation of the Incredible Years Teacher and Child Training Programs in high-risk schools. *Journal of Child Psychology and Psychiatry*, 49, 471–488.
- Weiss, N. H., Tull, M. T., Davis, L. T., Searcy, J., Williams, I., & Gratz, K. L. (2015). A preliminary experimental investigation of emotion dysregulation and impulsivity in risky behaviours. *Behaviour Change*, 32, 127–142.
- Wolf, D. H., Satterthwaite, T. D., Calkins, M. E., Ruparel, K., Elliott, M. A., Hopson, R. D., et al. (2015). Functional neuroimaging abnormalities in youth with psychosis spectrum symptoms. *JAMA Psychiatry*, 72, 456–465.
- Yang, T. T., Simmons, A. N., Matthews, S. C., Tapert, S. F., Frank, G. K., Max, J. E., et al. (2010). Adolescents with major depression demonstrate increased amygdala activation. *Journal of the American Academy of Child & Adolescent Psychiatry*, 49, 42–51.
- Yeung, D. Y., Wong, C. K. M., & Lok, D. P. P. (2011). Emotion regulation mediates age differences in emotions. *Aging & Mental Health*, 15, 414–418.
- Zaki, J., & Williams, W. C. (2013). Interpersonal emotion regulation. *Emotion*, 13, 803.
- Zhang, X., Ersner-Hershfield, H., & Fung, H. H. (2010). Age differences in poignancy: Cognitive reappraisal as a moderator. *Psychology and Aging*, 25, 310–320.
- Zisner, A., & Beauchaine, T. (2016). Psychophysiological methods and developmental psychopathology. In D. Cicchetti (Ed.), *Developmental psychopathology: Vol. 2. Developmental neuroscience* (3rd ed., pp. 832–884). Hoboken, NJ: Wiley.
- Zisner, A., & Beauchaine, T. P. (2015). Psychophysiology (peripheral). In S. Lilienfeld (Ed.), *Encyclopedia of clinical psychology*. Hoboken, NJ: Wiley.