

NORMAL STRUCTURE FOR BANACH SPACES WITH SCHAUDER DECOMPOSITION

BY
M. A. KHAMSI

ABSTRACT. We introduce a new constant in Banach spaces which implies, in certain cases, the weak- or weak*-normal structure.

Introduction. M. S. Brodskii and D. P. Mil'man [3] have introduced a geometric property, called normal structure, for convex subsets of Banach spaces. This property was introduced into fixed point theory by W. A. Kirk [10]. In this paper we associate to every Banach space, with Schauder finite decomposition, a constant which may be easily calculated. This constant allows us to decide, in certain cases, if the Banach space has weak-normal structure (alternatively weak*-normal structure in the dual case). For details on normal structure and its generalizations in Banach spaces, we suggest the survey of S. Swaminathan [17].

The author wishes to thank B. Madary for useful discussions regarding this paper.

Definitions and notations. In this paper X will always denote a real Banach space with F.D.D. (see definition 6). For terms not explicitly defined reference may be made to Day's book [6].

DEFINITION 1. A point x of a convex bounded set C is called a non-diametral point for C if

$$\sup\{\|x - y\| : y \in C\} < \text{diam}(C).$$

DEFINITION 2. A sequence (x_n) is called diametral if $c = \text{diam}(x_i) > 0$ and

$$\lim_{n \rightarrow \infty} d(x_n, \text{conv}\{(x_i) : i < n\}) = c$$

It follows that for every $x \in \text{conv}(x_n)$, we have $\lim \|x_n - x\| = c$. Every $x \in \text{conv}(x_n)$ is a diametral point, any subsequence of (x_n) is also diametral and is not convergent.

DEFINITION 3. We say that X has normal structure if every bounded convex subset K of X , which contains more than one point, has a non-diametral point. Restricting

Received by the editors December 3, 1987 and, in revised form, June 30 1988.

AMS Subject Classifications (1980): Primary 46B20, Secondary 47H10.

Key words: Normal structure, Schauder decomposition, fixed point property.

© Canadian Mathematical Society 1988.

K to be weak (respectively weak*) compact, we similarly define weak (respectively weak*) normal structure.

Let us recall the most useful characterization of normal structure [3].

THEOREM 1. *A bounded convex subset K of X contains a non-diametral point if and only if K does not contain a diametral sequence.*

DEFINITION 4. *An ultrapower \tilde{X} of X is defined by $\tilde{X} = \ell_\infty(X)/\mathcal{N}$ with $\ell_\infty(X) = \{(x_n \in X \text{ and } \sup \|x_n\| < \infty)\}$ and $\mathcal{N} = \{(x_n) \in \ell_\infty(X); \lim_{\mathcal{U}} \|x_n\| = 0\}$, where \mathcal{U} is a free ultrafilter over \mathbf{N} . Let (P) be a property on X . We say that X has the property super- (P) if every ultrapower of X has the property (P) .*

For more details about ultrapower spaces, we refer to [15].

DEFINITION 5. *A Banach space F is called a spreading model for X , generated by the bounded sequence (x_n) if the following hold: (a) There exists a sequence (e_n) such that F is the closed linear space of $\{e_n; n \in \mathbf{N}\}$. (b) There exists a subsequence (y_m) of (x_n) so that (y_m) has no norm convergent subsequence and for all $n \in \mathbf{N}$ and scalars $(\alpha_i)_{1 \leq i \leq n}$, we have*

$$\left\| \sum_{i=1}^{i=n} \alpha_i e_i \right\|_F = \lim \left\{ \left\| \sum_{i=1}^{i=n} \alpha_i y_{m_i} \right\| : m_1 < \dots < m_n, m_1 \rightarrow \infty \right\}.$$

Let (P) be a property on X . We say that X has the property $M - (P)$ if every spreading model for X has the property (P) .

For more information about the spreading model we refer to [2,4,7].

DEFINITION 6. *A sequence (X_n) of finite dimensional subspaces of X is called a Schauder finite dimensional decomposition (F.D.D) of X , if every $x \in X$ has unique representation of the form $x = \sum x_i$ with $x_i \in X_i$ for every $i \in \mathbf{N}$.*

Let $x \in X$, we defined $\text{supp}(x)$ to be the set of integers $i \in \mathbf{N}$ such that $x_i \neq 0$. Let A and B be two subsets of \mathbf{N} and $k \in \mathbf{N}$, we will write $A < B$ (resp. $A < B + k$) if for every $(a, b) \in A \times B$, we have: $a < b$ (resp. $a < b + k$).

For more details about F.D.D. and Schauder bases, we suggest [14].

Examples and basic results.

DEFINITION 7. *Let X be a Banach space, with a F.D.D. Define $\beta_p(X)$, for $p \in [1, \infty]$, to be the infimum of the set of numbers λ such that*

$$(\|x\|^p + \|y\|^p)^{1/p} \leq \lambda \|x + y\|$$

for every x and y in X which verify $\text{supp}(x) < \text{supp}(y)$.

EXAMPLES: 1. The space $\ell_{p,q}$ defined in [5], which is simply ℓ_p renormed by:

$$|x| = (\|x^+\|_{\ell_p}^q + \|x^-\|_{\ell_p}^q)^{1/q}$$

where x^+ and x^- are the positive and negative parts of x in the lattice structure of ℓ_p . W. Bynum has shown that $\ell_{p,1}$, using the constant WCS (see definition in [5]) which seems to be more difficult to compute, has normal structure.

2. The James space J which consists of sequences $x = (x_n) \in c_0$ such that

$$\|x\| = \sup \{ (x_{p_1} - x_{p_1})^2 + (x_{p_2} - x_{p_3})^2 + \dots + (x_{p_{n-1}} - x_{p_n})^2 \}$$
 is finite,

where the supremum is taken for every n and every increasing sequence of integers (p_i) . Recall that J fails normal structure. Nevertheless it is shown in [1] that for every u and v which verify $\text{supp}(u) + 1 < \text{supp}(v)$, we have

$$(\|u\|^2 + \|v\|^2)^{1/2} \leq \|u + v\|.$$

We deduce that $\beta_2(J) = 1$.

REMARK. Sometimes the term $(x_{p_m} - x_{p_1})$ is added in the definition of $\|x\|$, and then we obtain a new space J_1 which is isomorphic to J . In [9] the author proves that any weakly-compact convex subset K of J_1 has the fixed point property, i.e. any selfmapping T defined on K has a fixed point, provided that T satisfies $\|T(x) - T(y)\| \leq \|x - y\|$ for every x, y in K . It is not clear whether J_1 has weak-normal structure.

PROPOSITION 1. a) Let $p \in [1, \infty]$ and $q > p$, we have

$$1 \leq \beta_q(X) \leq \beta_p(X) \leq 2^{(q-p)/pq} \beta_q(X).$$

b) We have $\beta_p(\ell_p) = 1$ and $\beta_1(\ell_{p,1}) = 2^{1-1/p}$.

PROOF. a) For every pair of scalars (α, β) we have:

$$(|\alpha|^q + |\beta|^q)^{1/q} \leq (|\alpha|^p + |\beta|^p)^{1/p} \leq 2^{(q-p)/pq} (|\alpha|^q + |\beta|^q)^{1-q}$$

from which we obtain our inequalities on β_p and β_q .

b) Let $(u, v) \in \ell_p$ be such that $\text{supp}(u) < \text{supp}(v)$. Then $\text{supp}(u^+) < \text{supp}(v^+)$ and $\text{supp}(u^-) < \text{supp}(v^-)$. This implies that $(*) \|u + v\| = (\|u\|^p + \|v\|^p)^{1/p}$, $\|u^+ + v^+\|^p = \|u^+\|^p + \|v^+\|^p$ and $\|u^- + v^-\|^p = \|u^-\|^p + \|v^-\|^p$. On the other hand we have, by definition of the norm $|\cdot|$, $|u| + |v| = (\|u^+\| + \|u^-\|) + (\|v^+\| + \|v^-\|)$. Thus $|u| + |v| \leq 2^{1-1/p} [(\|u^+\|^p + \|v^+\|^p)^{1/p} + (\|u^-\|^p + \|v^-\|^p)^{1/p}] = 2^{1-1/p} (\|u^+ + v^+\| + \|u^- + v^-\|)$. As the hypothesis on u and v implies that $(u + v)^+ = u^+ + v^+$ and $(v + u)^- = u^- + v^-$, we have $(**) |u| + |v| \leq 2^{1-1/p} |u + v|$. We conclude by $(*)$ and $(**)$ that $\beta_p(\ell_p) = 1$ and $\beta_1(\ell_{p,1}) \leq 2^{1-1/p}$. If we take $u = e_1$ and $v = e_2$, we obtain $2^{1-1/2} \leq \beta_1(\ell_{p,1})$. \square

The following theorem studies the relation between the constant β_p and the Banach-Mazur distance (for isomorphic Banach spaces X and Y , the Banach-Mazur distance from X to Y , denoted $d(X, Y)$, is defined to be the infimum of $\|U\| \|U^{-1}\|$ taken over all bicontinuous linear operators U from X onto Y).

THEOREM 2. *Let X and Y be two isomorphic Banach spaces. Suppose that X has a F.D.D. Then Y also has a F.D.D. and we have*

$$\beta_p(Y) \leq d(X, Y)\beta_p(X).$$

PROOF. Our theorem will be deduced from the following obvious lemma.

LEMMA 1. *Let $(X, \|\cdot\|)$ be as in Theorem 2, and $|\cdot|$ be an equivalent norm on X such that $\|\cdot\| \leq |\cdot| \leq \lambda\|\cdot\|$, then $\beta_p(X, |\cdot|) \leq \lambda\beta_p(X, \|\cdot\|)$.*

Indeed let $\epsilon > 0$ and $U : X \rightarrow Y$ be an isomorphism such that $\|U^{-1}\| = 1$ and $\|U\| = d_\epsilon$ where $d_\epsilon \leq d + \epsilon$. Therefore we have for every $x \in X$

$$(*) \quad \|x\|_X \leq \|U(x)\|_Y \leq d_\epsilon \|x\|_X.$$

Let (X_i) be the F.D.D. of X , if we put $Y_i = U(X_i)$, we obtain a F.D.D. for Y which satisfies the following property (**) $\text{supp}(y) = \text{supp}(U^{-1}(y))$ for every y in Y . Given (*) and (**) and the lemma the conclusion of Theorem 2 is deduced.

We obtain the following result as an application of Theorem 2.

COROLLARY 1. (a) *Let $|\cdot|$ be an equivalent norm on c_0 , we have $2^{1/p} \leq \beta_p(c_0, |\cdot|)$. (b) *Let X be a Banach space which contains a subspace isomorphic to c_0 , then $2^{1/p} \leq \beta_p(X)$.**

PROOF. Let us first show how (b) can be deduced from (a). Let (e_i) be the basis of c_0 . Then (e_i) converges weakly to 0 in X . By a classical argument, for $\epsilon > 0$ there exists in X a sequence (u_i) of blocks such that $d([u_i], c_0) \leq 1 + \epsilon$, where $[u_i]$ is the closed linear span of $\{u_i; i \in \mathbb{N}\}$. Therefore by (a) we deduce that $2^{1/p} \leq \beta_p(c_0) \leq (1 + \epsilon)\beta_p([u_i])$. As (u_i) are successive blocks in X , we have: $\beta_p([u_i]) \leq \beta_p(X)$. This gives the conclusion of (b), since ϵ is arbitrary. Let us complete the proof by proving (a). In [8] it is shown that if $\|\cdot\|$ is an equivalent norm on c_0 , then given $\epsilon > 0$, there exists u and v in c_0 which satisfy $\text{supp}(u) < \text{supp}(v)$ and $\text{Max}(|\alpha|, |\gamma|) \leq \|\alpha u + \gamma v\| \leq \text{Max}(|\alpha|, |\gamma|)(1 + \epsilon)$ for all scalars α and γ . The definition of β_p implies $(\|u\|^p + \|v\|^p)^{1/p} \leq \beta_p(c_0, \|\cdot\|)\|u + v\|$. Therefore we obtain $2^{1/p}(1 + \epsilon)^{-1} \leq \beta(c_0, \|\cdot\|)$. As ϵ is arbitrary, the conclusion of (a) holds. □

Main result.

THEOREM 3. *Let X be a Banach space, with a finite codimensional subspace Y such that $\beta_p(Y) < 2^{1/p}$ for some $p \in [1, \infty]$. Then X has weak-normal structure.*

PROOF. Suppose that X fails weak-normal structure. Then there exists a weakly convergent diametral sequence (x_n) . Since diametral property is invariant by translation, we may suppose that (x_n) converges weakly to 0. Our hypothesis on Y implies that there exists Z , a finite dimensional subspace of X , such that $X = Y \oplus Z$. Therefore

$x_n = y_n + z_n$ with $y_n \in Y$ and $z_n \in Z$ for every $n \in \mathbf{N}$. Since (x_n) converges weakly to 0, we deduce that both (y_n) and (z_n) converge weakly to 0. Using the fact that Z has a finite dimension, (z_n) converges (in norm) to 0 in X . Therefore there exists a sequence (u_n) of successive blocks and a subsequence (y_{m_n}) such that $\lim \|y_{m_n} - u_n\| = 0$. The definition of $\beta_p(Y)$ implies that

$$(\|u_{n+1}\|^p + \|u_n\|^p)^{1/p} \leq \beta_p(Y)\|u_{n+1} - u_n\|.$$

But $\lim \|u_n\| = \lim \|y_{m_n}\| = \lim \|x_{m_n} - z_{m_n}\| = \lim \|x_{m_n}\| = \text{diam}(x_i) = c$ and $\lim \|u_{n+1} - u_n\| = \lim \|y_{m_{n+1}} - y_{m_n}\| = \lim \|x_{m_{n+1}} - x_{m_n}\| = \text{diam}(x_i) = c$. We deduce that $2^{1/p} \leq \beta_p(Y)$ since $c > 0$. This yields a contradiction to our hypothesis on Y . □

COROLLARY 2. *Let X be a Banach space isomorphic to ℓ_p . Assume that $d(X, \ell_p) < 2^{1/p}$, then X has normal structure.*

PROOF. Since X is reflexive, the weak-normal structure and normal structure are the same. Theorem 2 implies that $\beta_p(X) \leq d(X, \ell_p)\beta_p(\ell_p)$. Since $\beta_p(\ell_p) = 1$ we obtain $\beta_p(X) \leq d(X, \ell_p) < 2^{1/p}$. The conclusion follows from Theorem 2.

REMARK. Let us note that the number $2^{1/p}$ is optimum in the sense that for every $p \in]1, \infty[$ there exists X_p such that $d(X_p, \ell_p) = 2^{1/p}$ and X_p fails normal structure. Indeed let $\epsilon > 0$ and define an equivalent norm $|\cdot|_\epsilon$ on ℓ_p by $|x|_\epsilon = \text{Max}(\|x\|_p, \epsilon\|x\|_1)$. Consider the space $X_\epsilon = (\ell_p, |\cdot|_\epsilon)$. It is clear that $d(X_\epsilon, \ell_p) = \text{Max}(1, \epsilon)$. Let us notice that if $2^{1/p} \geq \epsilon$ then the canonical basis of ℓ_p is diametral. The desired space is $X_{2^{1/p}}$.

The next result gives a partial positive answer to T. Landes' problem [12].

PROPOSITION 2. *Let (X_i) be a sequence of Banach spaces such that*

$$\beta = \sup_{i \in \mathbf{N}} \beta_1(X_i) < 2.$$

Then $X = \otimes_i X_i$ has weak-normal structure.

PROOF. Suppose that X fails weak-normal structure. Then there exists a weakly convergent diametral sequence (x_n) in X . We may suppose, with loss of generality, that (x_n) converges weakly to 0. By definition of X we have $x_n = \Sigma x_i(n)$. Since (x_n) converges weakly to 0, we deduce that $(x_i(n))$ converges weakly to 0 in X_i for every $i \in \mathbf{N}$. By passing to a subsequence, we may assume that $(x_i(n))$ is a sequence of blocks, related to the F.D.D. of X_i , for every $i \in \mathbf{N}$. Our hypothesis on (X_i) implies that

$$\|x_i(n+1)\|_{X_i} + \|x_i(n)\|_{X_i} \leq \beta \|x_i(n+1) - x_i(n)\|_{X_i} \quad \text{for every } i \in \mathbf{N}.$$

Then by definition of the norm of X , we get

$$\|x_{n+1}\|_X + \|x_n\|_X \leq \beta \|x_{n+1} - x_n\|_X.$$

Since (x_n) is diametral and convergent weakly to 0 (this implies that $0 \in \overline{\text{conv}}(x_i)$) and therefore $\lim \|x_n\| = \text{diam}(x_i)$, we get $2 \leq \beta$. This is a contradiction. \square

Our last theorem concerns the M -(normal structure)-property (see definition 5).

THEOREM 4. *Let X be a reflexive Banach space. Assume that $\beta_p(X) < 2^{1/p}$, for some $p \in [1, \infty[$. Then X has M -(weak-normal structure).*

PROOF. Let (x_n) be a bounded sequence and F be the spreading model generated by (x_n) . Since X is reflexive we may suppose that (x_n) is weakly convergent to x . *First case:* $x = 0$. By a classical argument, there exists a sequence (u_i) of successive blocks and a subsequence $(x_{n(i)})$ such that $\lim \|u_i - x_{n(i)}\| = 0$. This implies that the spreading model generated by (u_i) is F . Since (x_n) converges weakly to 0, the sequence (e_i) is a Schauder basis of F , (see [7]). Let u and v be in F which satisfy: $\text{supp}(u) < \text{supp}(v)$. We write

$$u = \sum_{i \leq a} c_i e_i \text{ and } v = \sum_{a < i \leq b} b_i e_i.$$

Therefore

$$\|u+v\|_F = \lim \left\{ \left\| \sum c_i u_{n_i} + \sum b_j u_{m_j} \right\|_X : n_1 < \dots < n_a < m_1 < \dots < m_b, n_1 \rightarrow \infty \right\}.$$

Hence $(\|u\|_F^p + \|v\|_F^p)^{1/p} \leq \beta_p(X) \|u + v\|_F$, which implies that $\beta_p(F) \leq \beta_p(X)$. The hypothesis on $\beta_p(X)$ implies that F has weak-normal structure. *Second case:* $x \neq 0$. It is shown in [7] that in this case F is isometric to a subspace of G with $G = F_0 \oplus \mathbf{R}$, where F_0 is the spreading model generated by the sequence $(x_n - x)$. The first case implies that $\beta_p(F_0) < 2^{1/p}$ and by Theorem 3 we deduce that G has weak-normal structure. Since F is isometric to a subspace of G , we conclude that F has weak normal structure.

COROLLARY 3. *The M -(normal structure) is not a self dual property.*

PROOF. Indeed let $X = \ell_{p,1}$ with $p \in]1, \infty[$. We have shown that $\beta_1(X) = 2^{1/q}$ where q is the conjugate to p (i.e. $1/p + 1/q = 1$). By Theorem 4, X has M -weak-normal structure. Since X is superreflexive, X has M -(normal structure) whereas $X^* = \ell_{q,\infty}$ fails to have normal structure.

REMARK. We don't know if $\ell_{p,1}$ has super-normal structure.

The constant β_p and weak*-normal structure. Let X be a Banach space with a shrinking Schauder basis (x_i) . Recall that X^* has a basis (x_i^*) which is boundedly complete ((x_i^*) are the biorthogonal functionals associated to the basis (x_i)). Let Y be a Banach space, isomorphic to X^* . In this paper, the weak*-topology considered on Y is the weak*-topology defined by the couple (X, X^*) . This implies, in particular, that if (y_i) converges weak* to 0 (in Y), then there exists a sequence (u_i) of successive blocks and a subsequence $(y_{n(i)})$ such that $\lim_{i \rightarrow \infty} \|y_{n(i)} - u_i\| = 0$. The same proof, as in Theorem 3, gives us the following result:

THEOREM 5. *Let X be a Banach space with a shrinking basis. Assume that $\beta_p(X^*) < 2^{1/p}$. Then X^* has weak*-normal structure.*

An application of Theorem 4 is Lim's theorem [13] which says that ℓ_1 has weak*-normal structure. Let us recall that P. Soardi [16] proved that if X is a Banach space which satisfies $d(X, \ell_1) < 2$, then X has the weak*-fixed point property, i.e., any weak*-compact convex subset K of X and any selfmapping T defined on K has a fixed point, provided that T satisfies $\|T(x) - T(y)\| \leq \|x - y\|$ for every x, y in K . Kirk [11] asked whether X has weak*-normal structure.

COROLLARY 4. *Let Y be a Banach space which is isomorphic to ℓ_1 . Assume that $d(Y, \ell_1) < 2$. Then Y has weak*-normal structure.*

REMARK. Here, we consider on Y , isomorphic to ℓ_1 , the weak*-topology given by the couple (c_o, ℓ_1) . Recall that there exists a Banach space X such that X^* is linearly isometric to ℓ_1 and fails the weak*-fixed point property and therefore cannot have weak*-normal structure (in this case the weak*-topology is related to the couple (X, X^*)). The space X is the Banach space c which consists of all convergent sequences.

REFERENCES

1. A. Andrew, *Spreading basic sequences and subspaces of James' quasi reflexive space*. Math. Scad. **48** (1981), 108–118.
2. B. Beauzamy and J. T. Lapreste, *Modèles étalés des espaces de Banach*, Publication du Dept. de Math. Université Claude Barnard-Lyon 1.
3. M. S. Brodskii and D. P. Mil'man, *On the center of a convex set*, Dokl. Akad. Nauk, USSR **59** (1948) 837–840.
4. A. Brunel and L. Sucheston, *On B-convex Banach spaces*, Math. System Theory, **7** (1974) 294–299.
5. W. L. Bynum, *Normal structure coefficients for Banach spaces*, Pac. J. Math. Vol. 86, No. 2, (1980) 427–436.
6. M. M. Day, *Normed linear spaces*, 3rd Edn. Springer-Verlag, 1973.
7. S. Guerre and J. T. Lapreste, *Quelques propriétés des modèles étalés sur un espace de Banach*, Ann. IHP. Section B **16-4** (1980) 339–347.
8. R. C. James, *Bases and reflexivity of Banach spaces*, Ann. of Math. **52** (1950) 518–527.
9. M. A. Khamsi, *James quasi-reflexive space has the fixed point property*, to appear in Bull. Aust. Math. Soc.
10. W. A. Kirk, *A fixed point theorem for mappings which do not increase distance*, Amer. Math. Monthly **72** (1965) 1004–1006. MR 32=6436.
11. ——— *Fixed point theory for non expansive mappings, II*. Contemporary Mathematics, Vol. **18** (1983) 121–140.
12. T. Landes, *Permanence properties of normal structure*, Pac. J. Math. Vol 110, No. 1 (1984) 125–143.
13. T. C. Lim, *Asymptotic centers and nonexpansive mappings in conjugate Banach spaces*, Pac. J. Math. **90** (1980) 134–143.
14. J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces-I-Sequence spaces*, Springer-Verlag 1977.
15. B. Sims, "Ultra"-techniques in Banach space theory, Queen's papers in Pure and Applied Mathematics, no. 60, Queen's University, Kingston, Ontario, Canada.
16. P. M. Soardi, *Schauder bases and fixed points of nonexpansive mappings*, Pac. J. Math. Vol. 101, No. 1 (1982) 193–198.

17. S. Swaminathan, *Normal structure in Banach spaces and its generalizations*, Contemporary Mathematics, Vol. 18 (1983) 201–215.

Department of Mathematics
University of Southern California
Los Angeles, CA.

Current address
Department of Mathematical Sciences
The University of Texas at El Paso
El Paso, Texas, 79968 0514