

## Research Article

**Cite this article:** André WPP, Junior JRP, Cavalcante GS, Ribeiro WLC, Araújo-Filho JV, Morais SM, Oliveira LMB, Abreu FOMS, Bevilaqua CML (2021). Carvacryl acetate nanoencapsulated with chitosan/chichá gum exhibits reduced toxicity in mice and decreases the fecal egg count of sheep infected with gastrointestinal nematodes. *Parasitology* **148**, 1560–1565. <https://doi.org/10.1017/S0031182021001220>

Received: 9 February 2021

Revised: 15 June 2021

Accepted: 23 June 2021

First published online: 9 July 2021










**Keywords:**

Acetylation; biopolymers; carvacrol; *Haemonchus contortus*; toxicity

**Author for correspondence:**

Claudia M. L. Bevilaqua,  
E-mail: [bevilaqua.ucece@gmail.com](mailto:bevilaqua.ucece@gmail.com)

# Carvacryl acetate nanoencapsulated with chitosan/chichá gum exhibits reduced toxicity in mice and decreases the fecal egg count of sheep infected with gastrointestinal nematodes

Weibson P. P. André<sup>1</sup> , José R. P. Junior<sup>2</sup> , Géssica S. Cavalcante<sup>3</sup> , Wesley L. C. Ribeiro<sup>4</sup> , José V. Araújo-Filho<sup>1</sup> , Selene M. Morais<sup>3</sup> , Lorena M. B. Oliveira<sup>1</sup> , Flávia O. M. S. Abreu<sup>2</sup>  and Claudia M. L. Bevilaqua<sup>1</sup> 

<sup>1</sup>Laboratório de Doenças Parasitárias, Programa de Pós-Graduação em Ciências Veterinárias, Universidade Estadual do Ceará, Fortaleza, Brazil; <sup>2</sup>Laboratório de Química Analítica e Ambiental, Programa de Pós-Graduação em Ciências Naturais, Universidade Estadual do Ceará, Fortaleza, Brazil; <sup>3</sup>Laboratório de Química de Produtos Naturais, Programa de Pós-Graduação em Ciências Veterinárias, Universidade Estadual do Ceará, Fortaleza, Brazil and <sup>4</sup>Departamento de Fisiologia e Farmacologia, Faculdade de Medicina, Universidade Federal do Ceará, Fortaleza, Brazil

**Abstract**

The nanoencapsulation of biocomposites with anthelmintic action has been proposed as an alternative for improving their efficiency. Thus, the current study aimed to evaluate the efficacy of carvacryl acetate nanoencapsulated with biopolymers (nCVA) in the control of sheep gastrointestinal nematodes. CVA was nanoencapsulated with chitosan/chichá gum and characterized in terms of its efficacy of encapsulation (EE), yield and zeta potential. The acute toxicity of nCVA was evaluated in mice. For the fecal egg count reduction test, 40 animals were divided into four groups ( $n = 10$ ) and orally administered the following treatments: G1, 250 mg kg<sup>-1</sup> CVA; G2, 250 mg kg<sup>-1</sup> nCVA; G3, chitosan/chichá gum (negative control) and G4, 2.5 mg kg<sup>-1</sup> monepantel (positive control). Feces were collected on days 0 and 16 posttreatment to determine the eggs per gram of feces (epg). The EE and yield of nCVA were 72.8 and 57.5%, respectively. The nanoparticles showed a size of  $764.5 \pm 302.5$  nm, and the zeta potential at pH 3.2 was +22.0 mV. nCVA presented a 50% lethal dose (LD<sub>50</sub>) of 2609 mg kg<sup>-1</sup>. By 16 days posttreatment, CVA, nCVA and monepantel reduced the epg by 52.9.7, 71.5 and 98.7%, respectively, and the epg of sheep treated with nCVA differed from that of the negative control ( $P > 0.05$ ) but did not differ from that of sheep treated with CVA. In conclusion, the nanoencapsulation of CVA reduced its toxicity, and nCVA showed anthelmintic activity.

**Introduction**

Gastrointestinal nematodes are one of the principal sanitary problems of small ruminants in tropical and temperate regions throughout the world (Duarte *et al.*, 2019; Sallé *et al.*, 2019). The control of these nematodes relies almost exclusively on the administration of anthelmintics, such as benzimidazoles, macrocyclic lactones, imidazothiazoles, monepantel (amino-acetonitrile derivative) and derquantel (spiroindole). However, the continuous use of these drugs has promoted the selection of resistant and/or multiresistant nematode populations (Silva *et al.*, 2018; Santos *et al.*, 2019; Nixon *et al.*, 2020).

The control of gastrointestinal nematodes using plant essential oils and their bioactive compounds has been highlighted as a promising alternative (Camurça-Vasconcelos *et al.*, 2007; Katiki *et al.*, 2017; Araújo-Filho *et al.*, 2018; Macedo *et al.*, 2019). Carvacrol is a phenolic monoterpene found in the essential oils of oregano (*Origanum vulgare*), thyme (*Thymus vulgaris*) and other plants (Sharifi-Rad *et al.*, 2018). This monoterpene has antiparasitic activity (Katiki *et al.*, 2017; Novato *et al.*, 2018). Carvacrol presents high toxicity, and acetylation is an alternative that can potentiate its biological activity and reduce toxicity. Carvacryl acetate (CVA) reduced motility and caused cuticle lesions in adult *Haemonchus contortus* and promotes a 65.9% reduction in egg count per gram of feces (epg) of sheep, but the efficacy does not reach the desired therapeutic level (André *et al.*, 2016). Nanoencapsulation has been proposed as an alternative to protect and promote the sustained release of bioactive compounds that show promising anthelmintic action since it can increase their bioavailability and potentiate their nematicidal action (Mesquita *et al.*, 2013; Ribeiro *et al.*, 2017; Katiki *et al.*, 2019; André *et al.*, 2020b). Biopolymers are used for the nanoencapsulation of bioactive compounds *via* polyelectrolyte complexation (Paula *et al.*, 2016). Among the biopolymers, chitosan is a polysaccharide obtained by the deacetylation of chitin. Chitosan possesses a high density of positive charges due to the protonation of amino groups and presents properties such as nontoxicity, biocompatibility and biodegradability (Luo and Wang, 2014). The positively charged amino groups of chitosan react with an anionic group of polysaccharides,

such as chichá gum, leading to the formation of a polyelectrolyte complex (Paula *et al.*, 2016). Chichá gum is extracted from *Sterculia striata*, a tree species belonging to the family Sterculiaceae that is widely found in the northeastern and central regions of Brazil (Mangas *et al.*, 2013). This polymer has been used as a possible polymer for drug nanoencapsulation (Magalhães *et al.*, 2016; Sombra *et al.*, 2019). Thus, in a study evaluating the best reaction conditions for the nanoencapsulation of CVA with biopolymers, it was found that nanoencapsulation with chitosan/chichá gum in a polymeric bilayer presented controlled release and inhibited the motility of *H. contortus* adults *in vitro*. Thus, the use of a biopolymer, such as chichá gum, which has a greater number of carboxylic groups, can trigger greater polyethylene complexation and increase the efficiency of nanoencapsulation (André *et al.*, 2020a).

Thus, the aim of the current study was to evaluate the toxicity and anthelmintic activity of nanoencapsulated carvacryl acetate (nCVA) with chitosan/chichá gum against sheep gastrointestinal nematodes.

## Materials and methods

### Gas chromatography coupled to mass spectrometry (GC-MS)

Carvacrol acetylation was performed according to André *et al.* (2016). The chemical analysis of the CVA was performed with a Shimadzu QP-2010 Ultra instrument with the following conditions: column, Rtx-5 MS (Crossbond 5%, diphenyl/95% dimethyl polysiloxane) with 30 m × 0.25 mm × 0.25 μm df; carrier gas, He (24.2 mL min<sup>-1</sup>, in constant linear velocity mode); injector temperature of 250 °C, split mode (1:100) and detector temperature of 250 °C. The column temperature was programmed as follows: the column increased from 35 to 180 °C at 4 °C min<sup>-1</sup> and then from 180 to 280 °C at 17 °C min<sup>-1</sup> and was subsequently maintained at 280 °C for 10 min. The mass spectra were acquired with an electron impact of 70 eV. Compounds were identified by their relative retention times obtained by GC to those of known compounds and by comparison of their mass spectra with those present in the computer data bank (NIST) and published literature (Adams, 2007).

### Nanoencapsulation of CVA

The nanoparticles were produced from a polyelectrolytic complexation system (André *et al.*, 2020a). Chitosan (1%) (w/v) and Tween 80 were subjected to mechanical stirring; 300 μL of CVA was then added, and the solution was maintained in an ultrasonic bath (Ultra 800, Ciencor Scientific Ltd., São Paulo, Brazil) for 15 min. For formation of the prenucleus of the nanoparticles, 0.1% sodium tripolyphosphate P.A. (STP) (Dinâmica®, São Paulo, Brazil) was added dropwise at a chitosan:STP ratio of 50:1, and the solution was subjected to magnetic stirring for 30 min. Then, 1% chichá gum (w/v) (Dinâmica®, São Paulo, Brazil) was added to the solution at a chitosan:chichá gum ratio of 10:1, and the solution was subjected to magnetic stirring for another 30 min. For the production of particles in bilayers, the monolayer particles were suspended in 20 mL of distilled water and subjected to magnetic stirring. Chitosan (1%) was added dropwise and stirred continuously for an additional 30 min. Subsequently, 1% chichá gum (w/v) was added and stirred continuously for an additional 30 min. The solution was centrifuged at 4000 rpm for 20 min, the supernatant was removed, and the bilayer nanoparticles were subjected to the lyophilization process.

### Encapsulation efficiency

The encapsulation efficiency (EE) of CVA was determined by absorption spectroscopy with a Genesys 10S UV-Vis (Thermo Scientific, Massachusetts, USA) at a wavelength of 271 nm. A solution of 10 mL of 95% ethanol with 10 mg of nanoparticles was stirred constantly for 48 h. Subsequently, 2 mL of the solution was filtered using a 0.25-μm syringe filter, and the concentration of CVA was determined according to the calibration curve using equation (1), with the absorbance (*abs*) and the CVA concentration (*conc*) in mg/mL:

$$abs = 0.0028conc - 0.0368; \quad R^2 = 0.994 \quad (1)$$

### Physicochemical characterization of nanoparticles

The nanoparticles were characterized by Fourier-transform infrared (FTIR) spectroscopy using a Nicolet iS5 spectrophotometer (Thermo Scientific). The samples were prepared as potassium bromide pellets (KBr) at a ratio of 1:20 (m/m) (sample:KBr). The particle size, surface charge at different pH values (3–8) and the point of zero charge were determined through zeta-potential measurements using a Nano ZetaSizer analyzer (Malvern 3600, Worcestershire, UK) with a laser wavelength of 632.8 nm and a fixed dispersion angle of 173 °.

### Acute toxicity

Acute toxicity was assessed according to OECD-425/2008 (The 'Up-and-Down' method for acute toxicity testing) (OECD, 2008). Female Swiss mice ( $n = 11$ ) weighing 25.1 ± 1.1 g were allowed to acclimatize to the laboratory conditions (luminosity: 12 h/12 h, light/dark; temperature: 22 ± 2 °C; relative humidity: 60%) for 7 days and were provided commercial feed (Labina®, Purina, São Paulo, Brazil) and water *ad libitum*. The animals were divided into two groups and received the following treatments orally: G1 ( $n = 8$ ), nCVA at doses of 175, 440, 1110 and 2800 mg kg<sup>-1</sup> and G2 ( $n = 3$ ) and chitosan/chichá gum (negative control). The dose correction factor was 3.2-fold, and the dose was increased or decreased according to the survival and mortality of the animal. Each animal received a single dose and was evaluated for 48 h prior to selection of the dose to be administered to the next animal. An increase or reduction in the dose occurred according to the observation of the survival or mortality of the animals, respectively. The dosage was discontinued when five reversals occurred in six consecutive nCVA-treated animals. All decisions regarding the doses administered and the estimation of the 50% lethal dose (LD<sub>50</sub>) were performed using AOT425StatPgm software.

### Fecal egg count reduction test (FECRT)

We used 40 sheep of both sexes ranging in age from 6 to 18 months, with an average weight of 30 kg, and subjected them *via* semi-extensive management with feeding on native pastures of the semiarid region of northeastern Brazil. The animals selected had over 500 epg, as determined using the McMaster technique with a 50-epg detection limit and sugar flotation (Coles *et al.*, 2006). Sheep were divided into 4 groups ( $n = 10$ ) according to the epg and randomly assigned to the following treatments: G1, 250 mg kg<sup>-1</sup> CVA; G2, nCVA; G3, chitosan/chichá gum (negative control) and G4, 2.5 mg kg<sup>-1</sup> monepantel (Zolvix®, Novartis, New Zealand) (positive control). The treatments were administered orally in a single dose. Fecal samples from each animal were collected on days 0, 8 and 16 posttreatment to determine the epg. According to Coles *et al.* (2006),




Fig. 1. FTIR spectra of CVA (—), nCVA (—), chitosan (—) and chichá gum (—).

when more than one anthelmintic class is being evaluated, a longer period of 14 days should be used, justifying the evaluation of egg on days 8 and 16 posttreatment. Aliquots of 2 g of feces from each animal belonging to the same group were homogenized and divided into three coprocultures containing approximately 20 g each (Roberts and O'Sullivan, 1950), and larvae were identified according to Van Wyk and Mayhew (2013).

### Statistical analysis

The efficacy of the FECRT was calculated using the following formula:  $FECRT = 100 \times (1 - [T2/T1][C1/C2])$ , in which the arithmetic fecal egg count means in controls (C) and treated (T) animals before (T1 and C1) and 8 or 16 days after (T2 and C2) deworming were compared (Dash et al., 1988) and 95% confidence intervals (CIs) were estimated using BootStreat 1.0 software (Cabaret, 2014). The egg values were log transformed ( $\log_{10}[x + 1]$ ), analysed by analysis of variance and compared using Tukey's test with GraphPad Prism® 7.0 software. The CI of the percentage of L3 recovered in coprocultures was calculated according to the following formula:  $2\sqrt{pq/n}$ , where  $p$  is the total number of larvae of a nematode genus,  $q$  is the total number of the other genera of nematodes and  $n$  is the total number of nematodes identified.

### Results

GC-MS analysis of the CVA indicated 99.3% CVA and 0.69% carvacrol. The EE and yield of nCVA were 72.8 and 57.5%, respectively. The FTIR spectra of CVA, chitosan, chichá gum and nCVA are shown in Fig. 1. CVA presented the following bands:  $1590\text{ cm}^{-1}$  (C–H flexion),  $1765\text{ cm}^{-1}$  (acetyl group) and  $816\text{ cm}^{-1}$  (aromatic ring) (André et al., 2016). The  $1086\text{ cm}^{-1}$  (C–O–C stretch) and  $890\text{ cm}^{-1}$  (pyranose ring) bands correspond to chitosan (Keawchaoon and Yoksan, 2011). The bands at  $1728\text{ cm}^{-1}$  indicate stretching and bending deformation out of plane due to the presence of acetyl groups in the chichá gum (Brito et al., 2004). CVA was not detected in the nCVA, possibly due to overlap with other bands of the biopolymer matrix.

The average size of the nanoparticles was  $764.5 \pm 302.5\text{ nm}$  and had a unimodal distribution.

The nanoparticles showed a decrease in the zeta potential with an increase in pH. At pH 3.2, the zeta potential was  $+22.0\text{ mV}$ , and the point of zero charge was situated at approximately pH 6.9 (Fig. 2).


In the acute toxicity test, nCVA presented an  $LD_{50}$  value of  $2609\text{ mg kg}^{-1}$ . The encapsulating matrix showed no toxicity.

The FECRT results of CVA, nCVA and monepantel are expressed as the mean egg on days 0, 8 and 16 posttreatment (Table 1). CVA, nCVA and monepantel reduced the egg by 52.9, 71.5 and 98.7%, respectively, by 16 days posttreatment. The CVA and nCVA results were not significantly different ( $P > 0.05$ ). However, nCVA was significantly different from the negative control ( $P < 0.05$ ).

The prevalence of nematode genera in FECRT is presented in Table 2. After treatment, the frequency of *Trichostrongylus* spp. increased in relation to *Haemonchus* spp. in the CVA- and nCVA-treated groups.

### Discussion

The evaluation of the zeta potential of nCVA at different pH values (3 and 8) showed that the nanoparticles were more stable at pH 3.2, where they had a zeta potential of  $+18.3\text{ mV}$ . The positive zeta potential of nCVA at acidic pH occurred due to the protonation of the amine groups of chitosan, increasing the positive charges on the surface of the nanoparticles. These positively charged nanoparticles can interact electrostatically with the negative charges of components present in the mucus that protect the mucosa of the stomach of monogastric animals, resulting in the mucoadhesion of nanoparticles (Ways et al., 2018). The acidic pH of ovine abomasum may promote the protonation of chitosan, the adhesion of nanoparticles to the abomasal mucosa and a prolonged release of CVA. However, for nanoparticles that do not adhere to the abomasal mucosa and reach the small intestine where the pH is basic, deprotonation of the amine groups of chitosan occurs, and the carboxyl groups of gum chichá are ionized,



**Fig. 2.** Zeta-potential variation with pH for nCVA. Zeta potential (■) and weighted mean zeta potential pH 8.1 (—).

**Table 1.** Fecal egg count reduction (FECR) and egg counts per gram of feces (epg ± standard deviation) of sheep treated with CVA, nCVA and monepantel.

Treatments	Day 0	Day 8	Day 16
<b>CVA</b>			
Mean epg	2085 ± 1078.1 <sup>Aa</sup>	1050 ± 390 <sup>Ba</sup>	945 ± 216 <sup>Ba</sup>
FECR (95% CI)		53.1 (–12 to 79)	52.9 (–2 to 76)
<b>nCVA</b>			
Mean epg	2140 ± 838 <sup>Aa</sup>	1055 ± 686 <sup>Ba</sup>	585 ± 261 <sup>Bb</sup>
FECR (95% CI)	–	54.1 (–11 to 81)	71.5 (41 to 87)
<b>Monepantel</b>			
Mean epg	2145 ± 962.9 <sup>Aa</sup>	85 ± 88.8 <sup>Bb</sup>	27.7 ± 31.2 <sup>Bc</sup>
FECR (95% CI)		96.2 (89 to 100)	98.7 (97 to 100)
<b>Polymer matrix</b>			
Mean epg	2140 ± 658 <sup>Aa</sup>	2300 ± 1460 <sup>Aa</sup>	2060 ± 1102 <sup>Aa</sup>

Capital letters compare means in the rows and small letters compare means in the columns. Different letters indicate significantly different values ( $P < 0.05$ ).

**Table 2.** Frequency (%) and 95% CI of third-stage larvae identified in coprocultures on days 0, 8 and 16 posttreatment treated with CVA, nCVA, monepantel and polymer matrix

Groups	Day 0	Day 8	Day 16
<b>CVA</b>			
<i>Haemonchus</i> spp.	73 (64.2–81.7)	31 (21.9–40.0)	10 (4.1–15.8)
<i>Trichostrongylus</i> spp.	24 (15.6–32.3)	59 (68.6–52.6)	88 (81.6–94.3)
<i>Oesophagostomum</i> spp.	3 (–0.3 to 6.3)	10 (7.2–15.8)	2 (–0.7 to 4.7)
<b>nCVA</b>			
<i>Haemonchus</i> spp.	70 (61.0–78.9)	17 (9.6–24.3)	19 (11.3–26.6)
<i>Trichostrongylus</i> spp.	25 (16.5–33.4)	68 (58.8–77.1)	80 (72.1–87.8)
<i>Oesophagostomum</i> spp.	5 (0.7–9.2)	5 (0.7–9.2)	1 (–0.9 to 2.9)
<b>Monepantel</b>			
<i>Haemonchus</i> spp.	68 (58.8–77.1)	50 (40.2–59.8)	52 (42.2–61.7)
<i>Trichostrongylus</i> spp.	26 (17.4–34.5)	38 (28.4–47.5)	33 (23.7–42.2)
<i>Oesophagostomum</i> spp.	6 (1.3–10.6)	12 (5.6–18.3)	15 (8.0–21.9)
<b>Polymer matrix</b>			
<i>Haemonchus</i> spp.	69 (59.9–78.0)	44 (36.2–55.7)	50 (43.2–62.7)
<i>Trichostrongylus</i> spp.	31 (21.9–40.0)	50 (40.2–59.8)	47 (37.2–56.7)
<i>Oesophagostomum</i> spp.	0	6 (1.3–10.6)	3 (–0.7 to 4.7)

Groups were treated with 250 mg kg<sup>–1</sup> CVA or nCVA. The positive control group was treated with 2.5 mg kg<sup>–1</sup> monepantel (Zolvix®), and the negative control group received polymer matrix.



which reduces the zeta potential of the nanoparticles to approach the isoelectric point, where charge neutralization occurs (Abreu et al., 2008); consequently, the nanoencapsulated biocomposite is rapidly released.

The LD<sub>50</sub> of nCVA was 2609 mg kg<sup>-1</sup>, while the LD<sub>50</sub> of CVA was 1544.5 mg kg<sup>-1</sup>, as obtained from the acute toxicity test in mice (André et al., 2016). The nanoencapsulation of CVA reduced toxicity and increased the toxicology safety of this biocomposite. The reduction in toxicity of a nanoencapsulated essential oil was also verified by an evaluation of the acute toxicity of *Eucalyptus staigeriana* essential oil (LD<sub>50</sub> = 1603.9 mg kg<sup>-1</sup>) and its nanoemulsion (LD<sub>50</sub> = 3495.9 mg kg<sup>-1</sup>) (Ribeiro et al., 2015). The reduction in toxicity may be associated with the sustained release of these biocomposites, which maintains their therapeutic effect and reduces the occurrence of plasma peaks that may trigger toxic effects on animal cells.

In the current study, a significant difference ( $P < 0.05$ ) in the egg values was found between the nCVA-treated group and the negative control group; however, no significant difference ( $P > 0.05$ ) in the egg values was detected between the CVA and nCVA groups. The similarity in the egg values between the treated and negative control groups was also verified when evaluating the effectiveness of the nanoencapsulated CVA with chitosan/gum arabic and the nanoemulsion of the essential oil of *Eucalyptus citriodora* (Ribeiro et al., 2014; André et al., 2020a). There was a reduction in the percentage of *Haemonchus* spp. L3 and a resulting increase in *Trichostrongylus* spp. L3 in posttreatment coprocultures. CVA activity against *Haemonchus* spp. was probably improved by the process of encapsulation with chitosan, which is a bioadhesive polymer used in drug delivery systems, by promoting sustained release at acidic pH, increasing the bioavailability of CVA and therefore potentiating anthelmintic efficacy on abomasal nematodes. Similar results were found when evaluating the anthelmintic activity of *E. staigeriana* essential oil nanoencapsulated with chitosan in the parasitic load of sheep infected with *Haemonchus* spp., *Trichostrongylus* spp. and *Oesophagostomum* spp., where only the parasitic burden of *Haemonchus* spp. has been reduced (Mesquita et al., 2013). In addition, it has been shown that the efficacy varies according to the nematode species, particularly when these inhabit separate digestive organs (Hoste et al., 2008).

The nanoencapsulation system was effective, presenting a nanometric size, zeta potential and sustained release in acidic pH, in addition to reducing the toxicity of the biocomposite. Thus, studies evaluating the anthelmintic action of CVA in experimentally infected animals with gastrointestinal nematodes should be performed. In addition, the bioavailability of biocomposites must be performed to prove that there is a greater release of CVA in the abomasum and to thus justify a greater effectiveness on *Haemonchus* spp. in relation to *Trichostrongylus* spp.

**Acknowledgements.** We acknowledge the Universidade Estadual do Ceará (UECE) and Universidade Federal do Ceará (UFC) for providing the necessary infrastructure and carrying out the experiments.

**Author contributions.** FOMSA and CMLB: conceptualization. WPPA, JRPJ, GSC, WLCR and JVAF: performed experiments. WPPA, GSC, WLCR and JVAF: data analysis, supervision and project administration: FOMSA, LMBO, SMM, CMLB and WPPA: writing the manuscript. All the authors reviewed and approved the manuscript.

**Financial support.** This study was supported by Brazilian funding agencies Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). Dr Bevilaqua has received a researcher fellowship from CNPq (305911/2019-8) and Dr André has received a doctoral scholarship at CAPES (1582708).

**Conflict of interest.** The authors declare that they have no conflicts of interest.

**Ethical standards.** This study was approved by the Ethics Committee on the Use of Animals of the Universidade Estadual do Ceará (Protocol Number: 6511846/2016).

## References

- Abreu FOMS, Bianchini C, Forte MMC and Kist TBL (2008) Influence of the composition and preparation method on the morphology and swelling behavior of alginate–chitosan hydrogels. *Carbohydrate Polymers* **74**, 283–289.
- Adams RP, (2007) *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*. Carol Stream: Allured Publishing Corporation. p. 804.
- André WPP, Ribeiro WLC, Cavalcante GS, Santos JML, Macedo ITF, Paula HCB, Freitas RM, Maia SM, Melo JV and Bevilaqua CML (2016) Comparative efficacy and toxic effects of carvacryl acetate and carvacrol on sheep gastrointestinal nematodes and mice. *Veterinary Parasitology* **218**, 52–58.
- André WPP, Junior JRP, Cavalcante GS, Ribeiro WLC, Araújo-Filho JV, Santos JML, Alves APNN, Monteiro JP, Moraes SM, Silva ING, Oliveira LMB, Abreu FOMS and Bevilaqua CML (2020a) Anthelmintic activity of nanoencapsulated carvacryl acetate against gastrointestinal nematodes of sheep and its toxicity in rodents. *Revista Brasileira de Parasitologia Veterinária* **29**, e013119.
- André WPP, Junior JRP, Cavalcante GS, Ribeiro WLC, Araújo-Filho JV, Cavalcanti BC, Moraes SM, Oliveira LMB, Bevilaqua CML and Abreu FOMS (2020b) Chitosan nanoparticles loaded with carvacrol and carvacryl acetate for improved anthelmintic activity. *Journal of the Brazilian Chemical Society* **31**, 1614–1622.
- Araújo-Filho JV, Ribeiro WLC, André WPP, Cavalcante GS, Guerra MCM, Muniz CR, Macedo ITF, Rondon FCM, Bevilaqua CML and Oliveira LMB (2018) Effects of *Eucalyptus citriodora* essential oil and its major component, citronellal, on *Haemonchus contortus* isolates susceptible and resistant to synthetic anthelmintics. *Industrial Crops and Products* **124**, 294–299.
- Brito ACF, Silva DA, De Paula RCM and Feitosa JPA (2004) *Sterculia striata* exudate polysaccharide: characterization, rheological properties and comparison with *Sterculia urens* (karaya) polysaccharide. *Polymer International* **53**, 1025–1032.
- Cabaret J (2014) Reliable phenotypic evaluations of anthelmintic resistance in herbivores: how and when should they be done? In: Quick W (ed.), *Anthelmintics – Clinical Pharmacology, Uses in Veterinary Medicine and Efficacy*. New York: Nova Science Publisher, pp. 1–26.
- Camurça-Vasconcelos AL, Bevilaqua CM, Moraes SM, Maciel MV, Costa CT, Macedo IT, Oliveira LMB, Braga RR, Silva RA and Vieira LS (2007) Anthelmintic activity of *Croton zehntneri* and *Lippia sidoides* essential oils. *Veterinary Parasitology* **148**, 288–294.
- Coles GC, Jackson F, Pomroy WE, Prichard RK, Von Samson-Himmelstjerna G, Silvestre A, Taylor MA and Vercruyse J (2006) The detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary Parasitology* **136**, 167–185.
- Dash KM, Hall K and Barger IA (1988) The role of arithmetic and geometric mean worm egg counts in faecal egg count reduction tests and in monitoring strategic drenching programs in sheep. *Australian Veterinary Journal* **65**, 66–68.
- Duarte ER, Matias AD, Bastos GA, Maia RC, Júnior VSM and Soares ACM (2019) Anthelmintic efficacy of trichlorfon and blood parameters of young lambs infected with *Haemonchus contortus*. *Veterinary Parasitology* **272**, 40–43.
- Hoste H, Torres-Acosta JF, Alonso-Diaz MA, Brunet S, Sandoval-Castro C and Adote SH (2008) Identification and validation of bioactive plants for the control of gastrointestinal nematodes in small ruminants. *Tropical Biomedicine* **25**, 56–72.
- Katiki LM, Barbieri AME, Araujo RC, Veríssimo CJ, Louvandini H and Ferreira JFS (2017) Synergistic interaction of ten essential oils against *Haemonchus contortus* in vitro. *Veterinary Parasitology* **243**, 47–51.
- Katiki LM, Araujo RC, Ziegelmeier L, Gomes ACP, Gutmanis G, Rodrigues L, Bueno MS, Veríssimo CJ, Louvandini H, Ferreira JFS and Amarante AFT (2019) Evaluation of encapsulated anethole and carvone in lambs artificially- and naturally-infected with *Haemonchus contortus*. *Experimental Parasitology* **197**, 36–42.

- Keawchaoon L and Yoksan R (2011) Preparation, characterization and *in vitro* release study of carvacrol-loaded chitosan nanoparticles. *Colloids and Surfaces B: Biointerfaces* **84**, 163–171.
- Luo Y and Wang Q (2014) Recent development of chitosan-based polyelectrolyte complexes with natural polysaccharides for drug delivery. *International Journal of Biological Macromolecules* **64**, 353–367.
- Macedo ITF, Oliveira LMB, André WPP, Araújo Filho JV, Santos JML, Rondon FCM, Ribeiro WLC, Camurça-Vasconcelos ALF, Oliveira EF, Paula HCB and Bevilaqua CML (2019) Anthelmintic effect of *Cymbopogon citratus* essential oil and its nanoemulsion on sheep gastrointestinal nematodes. *Revista Brasileira de Parasitologia Veterinária* **28**, 522–527.
- Magalhães Jr GA, Neto EM, Sombra VG, Richter AR, Abreu CMWS, Feitosa JPA, Paula HCB, Goycoolea FM and Paula RCM (2016). Chitosan/*Sterculia striata* polysaccharides nanocomplex as a potential chloroquine drug release device. *International Journal of Biological Macromolecules* **88**, 244–253.
- Mangas MBP, Mello VM and Suarez PAZ (2013) Thermal behavior of *Sterculia striata* oil. *Thermochimica Acta* **564**, 39–42.
- Mesquita MA, Silva Júnior JB, Panassol AM, Oliveira EF, Camurça-Vasconcelos ALF, Paula HCB and Bevilaqua CML (2013) Anthelmintic activity of *Eucalyptus staigeriana* encapsulated oil on sheep gastrointestinal nematodes. *Parasitology Research* **112**, 3161–3165.
- Nixon SA, Welz C, Woods DJ, Costa-Junior L, Zamanian M and Martin RJ (2020) Where are all the anthelmintics? Challenges and opportunities on the path to new anthelmintics. *International Journal for Parasitology: Drugs and Drug Resistance* **14**, 8–16.
- Novato T, Gomes GA, Zeringóta V, Franco CT, Oliveira DR, Melo D, Carvalho MG, Daemon E and Monteiro CMO (2018) *In vitro* assessment of the acaricidal activity of carvacrol, thymol, eugenol and their acetylated derivatives on *Rhipicephalus microplus* (Acari: Ixodidae). *Veterinary Parasitology* **260**, 1–4.
- OECD (2008) Organization for Economic Cooperation and Development. Guidelines for the testing of chemicals: *Repeated Dose 28-Day Oral Toxicity Study in Rodents*. Paris: OECD, Guideline 407.
- Paula HCB, Oliveira EF, Carneiro MJM and Paula RCM (2016) Matrix effect on the spray drying nanoencapsulation of *Lippia sidoides* essential oil in chitosan-native gum blends. *Planta Medica* **22**, 1–7.
- Ribeiro JC, Ribeiro WLC, Camurça-Vasconcelos ALF, Macedo ITF, Santos JML, Paula HCB, Araújo-Filho JV, Magalhães RD and Bevilaqua CML (2014) Efficacy of free and nanoencapsulated *Eucalyptus citriodora* essential oils on sheep gastrointestinal nematodes and toxicity for mice. *Veterinary Parasitology* **204**, 243–248.
- Ribeiro WL, Camurça-Vasconcelos AL, Macedo IT, Santos JM, Araújo-Filho JV, Ribeiro JC, Pereira VA, Viana DA, Paula HC and Bevilaqua CM (2015) *In vitro* effects of *Eucalyptus staigeriana* nanoemulsion on *Haemonchus contortus* and toxicity in rodents. *Veterinary Parasitology* **212**, 444–447.
- Ribeiro WLC, Camurça-Vasconcelos ALF, Santos JML, Macedo ITF, Ribeiro JC, Oliveira EF, Paula HCB and Bevilaqua CML (2017) The use of *Eucalyptus staigeriana* nanoemulsion for control of sheep haemonchosis. *Pesquisa Veterinária Brasileira* **37**, 221–226.
- Roberts FHS and O'Sullivan JP (1950) Methods for egg counts and larval cultures for strongyles infesting the gastrointestinal tract of cattle. *Australian Journal of Agricultural Research* **1**, 99–102.
- Sallé G, Doyle SR, Cortet J, Cabaret J, Berriman M, Holroyd N and Cotton JA (2019) The global diversity of *Haemonchus contortus* is shaped by human intervention and climate. *Nature Communications* **10**, 4811.
- Santos JML, Vasconcelos JF, Frota GA, Freitas EP, Teixeira M, Vieira, Bevilaqua CML and Monteiro JP (2019) Quantitative molecular diagnosis of levamisole resistance in populations of *Haemonchus contortus*. *Experimental Parasitology* **205**, 107734.
- Sharifi-Rad M, Varoni EM, Iriti M, Martorell M, Setzer WN, Contreras MM, Salehi B, Soltani-Nejad A, Rajabi S, Tajbakhsh M and Sharifi\_Rad J (2018) Carvacrol and human health: a comprehensive review. *Phytotherapy Research* **32**, 1675–1687.
- Silva FF, Bezerra HMFF, Feitosa TF and Vilela VLR (2018) Nematode resistance to five anthelmintic classes in naturally infected sheep herds in Northeastern Brazil. *Revista Brasileira de Parasitologia Veterinária* **27**, 423–429.
- Sombra FM, Richter AR, Araújo AR, Ribeiro FOS, Mendes JFS, Fontenelle ROS, Silva DA, Paula HCB, Feitosa JPA, Goycoolea FM and Paula RCM (2019) Nanocapsules of *Sterculia striata* acetylated polysaccharide as a potential monomeric amphotericin B delivery matrix. *International Journal of Biological Macromolecules* **130**, 655–663.
- Van Wyk JA and Mayhew E (2013) Morphological identification of parasitic nematode infective larvae of small ruminants and cattle: a practical lab guide. *Onderstepoort Journal of Veterinary Research* **80**, 539.
- Ways TMM, Lau WM and Khutoryanskiy VV (2018) Chitosan and its derivatives for application in mucoadhesive drug delivery systems. *Polymers* **10**, 267.