

# Climate change and the epidemiology of protostrongylid nematodes in northern ecosystems: *Parelaphostrongylus odocoilei* and *Protostrongylus stilesi* in Dall's sheep (*Ovis d. dalli*)

E. J. JENKINS<sup>1\*</sup>, A. M. VEITCH<sup>2</sup>, S. J. KUTZ<sup>1</sup>, E. P. HOBERG<sup>3</sup> and L. POLLEY<sup>1</sup>

<sup>1</sup>Department of Veterinary Microbiology, Western College of Veterinary Medicine, 52 Campus Drive, University of Saskatchewan, Saskatoon, Saskatchewan, Canada, S7N 5B4

<sup>2</sup>Wildlife Management, Environment and Natural Resources, Box 130, Government of the Northwest Territories, Norman Wells, Northwest Territories, Canada, X0E 0V0

<sup>3</sup>United States Department of Agriculture, Agricultural Research Service, US National Parasite Collection, Animal Parasitic Diseases Laboratory, BARC East No. 1180, 10300 Baltimore Avenue, Beltsville, Maryland, 20705, USA

(Received 26 May 2005; revised 9 September 2005; accepted 9 September 2005; first published online 7 December 2005)

## SUMMARY

We describe the epidemiology of the protostrongylid parasites *Parelaphostrongylus odocoilei* and *Protostrongylus stilesi* in Dall's sheep (*Ovis dalli dalli*) from the Mackenzie Mountains, Northwest Territories, Canada (65°N; 128°W). Peak numbers of 1st-stage larvae of both parasites were shed by Dall's sheep on their winter range from March until May. In larval development experiments in the Mackenzie Mountains, peak numbers of infective 3rd-stage larvae of *P. odocoilei* were available in gastropod intermediate hosts in August–September. For both protostrongylids, the majority of transmission likely occurs on the winter range, with infection of gastropods when they emerge from hibernation in spring, and infection of Dall's sheep upon their return in fall. We validated a degree-day model for temperature-dependent development of larval *P. odocoilei* in gastropods, and applied degree-day models to describe and predict spatial and temporal patterns in development of *P. odocoilei* and *P. stilesi* in northern North America. Temperature-dependent larval development may currently limit northward range expansion of *P. odocoilei* into naïve populations of Dall's sheep in the Arctic, but climate warming may soon eliminate such constraints. In Subarctic regions where both *P. odocoilei* and *P. stilesi* are endemic, the length of the parasite 'growing season' (when temperatures were above the threshold for larval development) and amount of warming available for parasite development has increased over the last 50 years. Further climate warming and extension of the seasonal window for transmission may lead to amplification of parasite populations and disease outbreaks in host populations.

Key words: protostrongylid, parasite, epidemiology, climate change, *Parelaphostrongylus odocoilei*, *Protostrongylus stilesi*, *Ovis dalli dalli*, *Deroceras leae*.

## INTRODUCTION

New host and geographical records have recently been established for the protostrongylid parasites *Parelaphostrongylus odocoilei* (muscleworm) and *Protostrongylus stilesi* (lungworm) in wild thimhorn sheep (Dall's sheep, *Ovis d. dalli*, and Stone's sheep, *O. d. stonei*) in Subarctic and Arctic North America (Kutz *et al.* 2001; Hoberg *et al.* 2002; Jenkins *et al.* 2005a). Like all protostrongylid nematodes, these parasites have indirect life-cycles, where 1st-stage larvae (L1) are shed in the faeces of the mammalian definitive host, invade a gastropod intermediate host, and develop to infective 3rd-stage larvae (L3). The

life-cycle of *P. odocoilei* has been described in experimentally infected thimhorn sheep (Jenkins, Hoberg and Polley, 2005b), but the epidemiology (including seasonal patterns of larval development, transmission, prevalence, and intensity) of protostrongylid parasites in populations of wild thimhorn sheep has not been investigated. Seasonal aspects of transmission of *P. odocoilei* in mule deer (*Odocoileus h. hemionus*) and *P. stilesi* in bighorn sheep (*Ovis c. canadensis*) at temperate latitudes in North America have been characterized (Samuel, Platt and Knispel-Krause, 1985; Robb and Samuel, 1990).

Development of protostrongylid larvae within the poikilothermic intermediate host is temperature-dependent and can be predicted using degree-day models, which assume that a fixed amount of heating accumulated above a critical threshold temperature is necessary for development (Saunders, Tompkins and Hudson, 2002; Harvell *et al.* 2002). A degree-day model has been developed and validated in the

\* Corresponding author: Canadian Wildlife Service, 115 Perimeter Road, Saskatoon, Saskatchewan, Canada, S7N 0X4. Tel: 011 1 306 966 7242 or 011 1 306 975 5357. Fax: 011 1 306 966 7242 or 011 1 306 975 4089. E-mail: emily.jenkins@usask.ca or emily.jenkins@ec.gc.ca




Fig. 1. Distribution of protostrongylid parasites in Dall's sheep (extrapolated from Worley *et al.* 2004; Jenkins *et al.* 2005a) and locations of weather stations and study areas in northern Canada: (a) Ivvavik National Park, Yukon Territory (YT) (to the immediate left are the British Mountains, continuous with the Brooks and Baird Mountains in Alaska, USA); (b) Inuvik, Northwest Territories (NT) (to the left are the Richardson Mountains, NT and YT); (c) Norman Wells, NT; (d) Fort Simpson, NT; (e) Palmer Lake study area, Mackenzie Mountains, NT; (f) Katherine Creek study area, Mackenzie Mountains, NT.

field for the protostrongylid *Umingmakstrongylus pallikuukensis*, a lungworm of muskoxen (*Ovibos moschatus*) (Kutz *et al.* 2002), whereas similar models for *P. stilesi* and *P. odocoilei* have been developed but not yet validated (Samson and Holmes, 1985; Jenkins *et al.* 2005c). Such models are powerful tools for comparing development of protostrongylid species across latitudinal gradients, isolating factors that are potentially limiting parasite distribution, and determining the potential for range expansion. A degree-day model for *U. pallikuukensis* has also been used to describe historical patterns of parasite development and to predict responses of protostrongylid-host systems to climate change (Kutz *et al.* 2005). Climate change, especially of the magnitude projected in northern regions, will profoundly influence ecosystems, including the ecological relationships between hosts and parasites (Harvell *et al.* 2002; Dobson *et al.* 2003; Hassol, 2004). The distribution of thimhorn sheep in Canada (see Bowyer and Leslie, 1992) corresponds to the 2 districts (i.e., North British Columbia Mountains/Yukon and Mackenzie District) currently experiencing the strongest warming trends in Canada (<http://www.msc-smc.ec.gc.ca/ccrm/bulletin/>, accessed April 2005).

Our objectives were (i) to describe the epidemiology of *P. odocoilei* and *P. stilesi* in Dall's sheep in the Mackenzie Mountains, Northwest Territories (NT), Canada, (ii) to validate a degree-day model for development of larvae of *P. odocoilei* in an important gastropod intermediate host and (iii) to apply degree-day models to describe and predict trends in development of *P. odocoilei* and *P. stilesi* throughout the geographical range of Dall's sheep in northern North America, and in a future of climate warming.

#### MATERIALS AND METHODS

##### *Patterns of larval shedding*

Every 1–3 months from March 2000 to April 2003, we collected approximately 30 separate faecal piles that appeared to have been deposited within the previous 48 h by Dall's sheep in the Katherine Creek study area, Mackenzie Mountains, NT (65°01'N; 127°35'W; elevation *ca.* 610–1100 m) (Fig. 1). Faecal pellets were sealed in plastic bags and frozen for 2–31 weeks before recovery of L1 from 5 g samples using a modified beaker Baermann technique (Forrester and Lankester, 1997; Jenkins *et al.* 2005a). Samples with high numbers of free-living nematodes, insect larvae,

Table 1. Development trials with larvae of *Parelaphostrongylus odocoilei* in slugs (*Deroceras laeve*) at Palmer Lake, 2002–2003

Exp.	Size ratios of slugs (%)			Start date	Number of enclosures with infective larvae/number with infected slugs (% slug recovery) at days PI*															
	Small	Med.	Large		14	21	28	35	42	56	70	84	98	308	336–357					
1	10	50	40	15-Jul-02	0/3 (60)	0/3 (73)	0/3 (40)	0/3 (63)	0/3 (53)											
2	10	40	50	22-Jul-02	0/3 (80)	0/3 (47)	0/3 (73)	0/3 (73)												
3	10	40	50	29-Jul-02	0/3 (67)	0/3 (63)	0/3 (70)													
4	30	30	40	2-Jun-03	0/3 (57)		0/3 (43)		0/3 (33)	3/3 (23)	2/2 (7)	1/1 (7)								
5	40	10	50	16-Jun-03	0/3 (50)		0/3 (37)		2/2 (13)†	1/1 (17)	3/3 (17)	3/3 (13)								
6	20	40	40	30-Jun-03	0/3 (37)		1/3 (40)†		2/2 (10)	3/3 (13)	1/1 (7)									
7	10	40	50	14-Jul-03	0/3 (70)		0/3 (33)		0/3 (37)	2/3 (33)†										
8	20	30	50	28-Jul-03	0/3 (73)		0/3 (77)		0/3 (60)											
9	60	20	20	11-Aug-03	0/3 (47)		0/3 (40)													

\* Days post-infection.

† Infective L3 observed for the first time on 28 July for Exps 4, 5, and 6, and 8 Sept. for Exp. 7.

or dead L1 were assumed to have been on the ground for longer than 48 h and results were not used in analyses. The prevalence (percentage of samples positive) and mean intensity (mean number of L1 per gram of wet faeces, or LPG) were calculated separately for *P. odocoilei* and *P. stilesi*.

*Development of P. odocoilei in D. laeve*

Using methods adapted from Kutz *et al.* (2002), in 2002 and 2003 we determined the rates of development for larvae of *P. odocoilei* in experimentally infected native slugs (*Deroceras laeve*) at the Palmer Lake study area, Mackenzie Mountains, NT (64°28'N; 129°37'W, elevation 1100 m) (Fig. 1). Slugs were collected from a wet sedge meadow separated by a packed trail from the 20 × 40 m study site, which was enclosed by a battery-powered electric fence to deter animals (e.g., grizzly bears, *Ursus arctos horribilis*, and woodland caribou, *Rangifer tarandus caribou*). The study site was a wet-mesic sedge meadow covered by 5–15% standing water depending on the time of year, and the vegetation was composed primarily of sedge and moss, with mats of mountain aven (*Dryas* sp.) and some willow (*Salix* spp.), dwarf birch (*Betula* sp.), and white spruce seedlings (*Picea glauca*).

Slugs were collected 1–3 times each day for 7–14 days prior to the start of each experiment (Table 1), except for Exp. 4, when slugs were collected for 26 days in May 2003 because the ground was still frozen, and traps were occasionally frozen or obscured by fresh snow. Slugs were measured at full extension and grouped into 1 of 3 size classes: small, up to 9 mm; medium, 10–14 mm; and large, greater than or equal to 15 mm (modified from Samuel *et al.* 1985). Slugs were housed in buckets containing native vegetation at mean temperatures of 6–12 °C.

First-stage larvae of *P. odocoilei* (originally from the Mackenzie Mountains) were recovered from faeces of experimentally infected Stone's sheep (Jenkins *et al.* 2005b). Groups of 12 slugs were exposed to 3000 motile L1 for 3 h (Hoberg *et al.* 1995), except for uninfected control slugs that were otherwise treated in a similar fashion. According to the size ratios at collection in each experiment, infected and uninfected slugs were housed in groups of 10 in separate enclosures randomly placed within the study site. Each enclosure consisted of a turf plug (approximate height 16 cm) inside a 5 litre, high-density polyethylene bucket (21 cm top diameter, 18 cm base diameter, height 19 cm; Ropak Packaging, Fullerton, California, USA) with drainage holes. Each bucket also contained a cut-down aluminum tomato cage (composed of 2 wire circles connected by 3 vertical rods, 24 cm top diameter, 18 cm base diameter, height 28 cm), and the entire enclosure was lined with white nylon/polyester netting. To expel wild slugs, turf plugs were flooded

for at least 24 h prior to addition of experimental slugs.

Starting at 14 days post-infection (p.i.), every 7 days in 2002, and every 14 days in 2003, infected slugs were recovered by visual examination and sequential flooding for 3 days of 3 randomly selected enclosures from each experiment (Table 1). In 2002, starting at 21 days p.i., uninfected control slugs were recovered from 3 enclosures from each experiment every 14 days, and in 2003, starting at 28 days p.i., uninfected slugs were collected from control enclosures every 28 days. In May–June 2003, overwintered enclosures (3 with infected slugs and 3 with uninfected slugs) from each experiment established in 2002 were collected at 308 days p.i., and 3 overwintered enclosures with infected slugs from each experiment were collected on June 16 (Exp. 1, 336 days p.i.), July 7 (Exp. 2, 350 days p.i.), and July 21 (Exp. 3, 357 days p.i.) (Table 1).

Slugs recovered from enclosures were held at approximately 6 °C for up to 7 days before individual digest in a pepsin-hydrochloric acid solution to recover, count, and classify stages of larvae (Hoberg *et al.* 1995; Jenkins *et al.* 2005*c*). In 2003, the seasonal abundance of infective L3 was calculated for slugs from Exps 4–7 by multiplying the mean number of infective L3/slug by the number of slugs recovered in each collection. In 2003, vegetation from enclosures containing infected slugs was examined for emerged infective L3 on July 28 (Exp. 4), August 11 (Exps 4–6), and August 25 (Exps 4–7) (Kutz, Hoberg and Polley, 2000). Any L3 recovered from uninfected control slugs were identified using molecular techniques (Jenkins *et al.* 2005*a*).

To ensure that larvae were viable, 10 infected and 10 uninfected slugs from each experiment (according to the size ratios at collection) were housed indoors at the warmest ambient temperatures possible under field conditions (Exps 1–3 and 6–9, average 19.5 °C, range 9–28 °C; Exps 4 and 5, average 15.5 °C, range 3–24 °C). These slugs were examined for survival and digested at 19 days p.i. in 2002, and 24 days p.i. in 2003, except for Exp. 4 where, because of cooler temperatures, slugs were examined for survival at 24 days p.i. and digested at 36 days p.i.

#### Validation of model

Temperatures 2 cm below and at the soil surface were recorded inside and outside 3 enclosures containing no slugs placed randomly within the study area. Temperatures were recorded every 1 h using external probes from HOBO XT monitors (Onset Computer Corporation, Pocasset, Massachusetts) (15 July to 25 August 2002), every 4 h using HOBO XT monitors (25 August 2002 to 7 May 2003), or every 1 h using HOBO H08 Pro monitors (7 May to 8 September 2003). The means for each temperature measurement for the 3 enclosures were used for

further calculations, except for surface temperatures inside enclosures over the winter of 2002–2003, from which data were available from only 2 enclosures. Air temperatures 1.5 m above ground level were recorded every 0.5 h (15 July to 25 August 2002) or every 1 h (25 August 2002 to 8 September 2003) by a HOBO weather station in the study area.

To predict when infective L3 should have been present in experiments in 2002 and 2003, we used recorded temperatures (soil, surface, and air), the laboratory-derived threshold temperature ( $T_0$ : 8.5 °C) and thermal constant (minimum amount of heating necessary for development of infective larvae, 163 degree days, or DD) (Jenkins *et al.* 2005*c*), and a degree-day model for protostrongylid larval development incorporating slug avoidance of micro-habitat temperatures exceeding 21 °C (Dainton, 1989; Kutz *et al.* 2002). Temperatures above 21 °C were converted to 21 °C, the threshold temperature subtracted from all temperatures, and positive values summed to determine the amount of heating above threshold on a daily basis. These were converted to degree-days (DD) by dividing by 48 for temperatures recorded every 0.5 h, 24 for temperatures recorded every 1 h, or 6 for temperatures recorded every 4 h. For each experiment, we calculated cumulative DD based on soil, surface, and air temperatures, and compared dates of predicted (based on the thermal constant) versus observed development of the first infective L3.

#### Application of model

We obtained hourly air temperature data from Environment Canada weather stations at 4 locations in Subarctic and Arctic Canada, starting in the year that consecutive hourly temperature data were first recorded at each location (Fig. 1). These included the following: Ivvavik National Park, Yukon Territory (69°09'N; 140°09'W; elevation 244 m; 1996–2004); Inuvik, NT (68°18'N; 133°29'W; 68 m; 1961–2004); Norman Wells, NT (65°17'N; 126°48'W; 169 m; 1955–2004); and Fort Simpson, NT (61°52'N; 121°21'W; 132 m; 1955–1963; and 61°45'N; 121°14'W; 74 m; 1963–2004). We used temperature data between 1 April and 31 October for the 3 northernmost locations, and between 1 March and 30 November for Fort Simpson. Similar to Kutz *et al.* (2005), at each location we used degree-day models to describe historical trends and variation for theoretical development of larvae of *P. odocoilei* and *P. stilesi*, for the latter using laboratory-derived parameters ( $T_0$ : 7.8 °C, 305 DD) from Samson and Holmes (1985). To adjust temperatures recorded at the northernmost location (Ivvavik), we used a crude correction factor of –1 °C per 200 m gain in elevation (<http://www.atmosphere.mpg.de/enid/16h.html>, accessed Apr 2005), and an altitude of 1000 m for Dall's sheep winter range, which assumed

that animals winter at altitudes closer to the valley bottoms than the high elevation summer range ([http://www.taiga.net/wmac/consandmanagementplan\\_volume1/landscape.html](http://www.taiga.net/wmac/consandmanagementplan_volume1/landscape.html), accessed Aug 2005).

We used surface temperatures recorded outside enclosures at Palmer Lake in 2003 as the baseline for climate change projections. The last date before 5 consecutive days with average surface temperatures below 0 °C was assumed to be the last day for gastropod activity in each year (see Kutz *et al.* 2005). From a climate change scenario (Canadian Centre for Climate Modeling and Analysis Global Coupled Model 2, A21 Special Report on Emission Scenarios, Economic Regional Focus Simulation 1) yielding mid-range values for the grid box incorporating the study site, we projected mean temperature increases of 1.2 °C by 2020, 3.3 °C by 2050, and 4.9 °C by 2080 (<http://www.cics.uvic.ca/scenarios/>, accessed Feb. 2005). By adding these values to the observed temperatures in 2003, and using degree-day models for development of larvae of *P. odocoilei* and *P. stilesi*, we predicted future trends in development of protostrongylids in the Mackenzie Mountains.

## RESULTS

### *Patterns of larval shedding*

Larvae of *P. odocoilei* were present in 96% (84–100%), and *P. stilesi* in 82% (18–100%), of 650 faecal samples collected from March 2000 to April 2003 (9–47 samples per collection). The mean intensities of L1 of *P. odocoilei* and *P. stilesi* over all collections were, respectively, 568 LPG (141–1350 LPG per collection) and 348 LPG (32–1075 LPG per collection). The prevalence of L1 of *P. stilesi* fell in August of each year, and rose again each fall (Fig. 2). Larval intensity for both parasites peaked in the spring of each year, decreased in summer, and gradually increased throughout fall and winter (Fig. 2).

### *Development of P. odocoilei in D. laeve*

Pre-infective, but not infective, larvae developed under field conditions in Exps 1–3 in 2002, and were recovered from live and dead slugs examined in 2003 after overwintering. In 2003, infective L3 were first observed on 28 July from Exps 4–6, on 8 September from Exp. 7, and were not recovered from Exp. 8 or 9 (for corresponding days p.i., see Table 1). Abundance of infective L3 pooled from all experiments was highest in August and September 2003 (Fig. 3). Emerged protostrongylid larvae were not recovered from any enclosures. One L3 was recovered from each of 2 uninfected control slugs collected in late July from Exp. 6, which were identified as *P. odocoilei* and an unnamed protostrongylid species endemic in caribou. One

unidentified protostrongylid L3 was recovered from an uninfected control slug collected in late August from Exp. 8.

Recovery of uninfected versus infected slugs from enclosures in the field did not differ consistently or substantially for any experiments, but decreased with the passage of time in all experiments. In enclosures from Exps 1–3 that overwintered, slug recoveries were low, and dead slugs were frequently observed. Slug recoveries were also low in Exps 4, 5, 6, and 9, even at 14 and 28 days p.i. (Table 1). Moribund or dead slugs were often observed in collections in May and June 2003, especially those in the medium size class. Ratios of size classes of slugs were similar for experiments starting in July 2002 (Exps 1–3) and 2003 (Exps 6–8) (Table 1). In 2003, ratios were approximately the same for all 3 size classes in May (Exp. 4), the relative proportion of medium slugs dropped to its lowest in early June (Exp. 5), and the largest relative proportion of small slugs (60%) was observed in early August (Exp. 9). Eggs were observed in collection dishes and field enclosures from late May until early September in 2003, most noticeably in June and July. Many newly hatched slugs were observed on traps in late July and early August, and were recovered from field enclosures from mid-July to September 2003.

Survival of both uninfected and infected slugs housed indoors ranged from 80–100% for Exps 1–3 and 7–9, and 40–80% for Exps 4–6. Most (74–95%) larvae recovered from indoor control slugs were infective L3, except in Exp. 5, in which 82% of larvae were early L3. Mean infection intensity in indoor controls ranged from 11–43 larvae/slug for the 9 experiments.

### *Validation of model*

Based on all sources of temperature data (soil, surface, and air), larvae were not predicted to develop to infective stage for experiments starting in 2002 until 19 May 2003 (Exp. 1, 308 days p.i.), 9 June 2003 (Exp. 2, 322 days p.i.), and 23 June 2003 (Exp. 3, 329 days p.i.) at the earliest (Fig. 4). For experiments starting in 2003, both air and surface temperatures adequately predicted when infective larvae were first observed, although air temperatures predicted somewhat slower development (Fig. 5). Degree-days based on soil temperatures did not accumulate rapidly enough to explain observed rates of development.

In the summer of 2003, among the 3 enclosures where hourly soil and surface temperatures were monitored, the mean standard deviation for soil temperature was approximately 0.75 °C (range 0–4.6 °C), while surface temperatures were more variable, with a mean standard deviation of approximately 1.25 °C (range 0.01–8.8 °C). In general, surface temperatures in summer were higher than either soil or air temperatures, except in July 2003,




Fig. 2. (A) Prevalence (%) and (B) mean intensity (larvae per gram of faeces, or LPG) of 1st-stage larvae of *Parelaphostrongylus odocoilei* and *Protostrongylus stilesi* shed by Dall's sheep in the Katherine Creek study area, March 2000–April 2003.

when air and surface temperatures were comparable. In winter, soil temperatures were slightly higher than surface temperatures, and both were less variable and much warmer than air temperatures (Fig. 6). Surface and soil temperatures were marginally lower inside versus outside enclosures, except in summer, when soil temperatures inside enclosures were slightly warmer. Summer temperatures at Palmer Lake were, on average, 5 °C cooler than at Norman Wells.

#### Application of model

For *P. odocoilei*, the length of the 'growing season', when temperatures were above the threshold for larval development, and maximum number of cumulative degree-days had an inverse relationship with latitude (Table 2). As well, moving from north

to south, larvae were predicted to develop to infective stage earlier in the year and could begin development later in the year and still reach infective stage. At Inuvik, Norman Wells (Fig. 7), and Fort Simpson, the maximum number of cumulative degree-days and the length of the growing season for development has increased over the last 4–5 decades. Despite annual variability, this relationship was statistically significant for cumulative degree-days at all 3 locations (Inuvik  $R^2=0.14$ ,  $F=6.8$ ,  $P=0.013$ ; Norman Wells  $R^2=0.15$ ,  $F=8.7$ ,  $P=0.005$ ; Fort Simpson  $R^2=0.18$ ,  $F=10.3$ ,  $P=0.002$ ), but not growing season.

Using recorded air temperatures, larvae of both *P. odocoilei* and *P. stilesi* could have developed within 1 season at all locations in every year, although the thermal constant for *P. stilesi* was only narrowly




Fig. 3. Seasonal abundance (number of slugs  $\times$  mean number of L3/slug) of infective larvae (L3) of *Parelaphostrongylus odocoilei* in the slug *Deroceras laeve* for Exps 4–7 in 2003.




Fig. 4. Predicted cumulative degree-days (Cum DD) for larvae of *Parelaphostrongylus odocoilei* in experiments starting in 2002 using hourly surface temperatures inside enclosures at Palmer Lake. Thermal constant (hatched line) = 163 DD.

exceeded at the 2 Arctic locations (Ivvavik and Inuvik). The two high-elevation data sets (Ivvavik corrected for elevation, and Palmer Lake) yielded approximately half the DD accumulated using uncorrected or nearby low elevation data sets (Norman Wells for Palmer Lake) (Table 2). Using elevation corrected data from Ivvavik, L3 of *P. odocoilei* would not have developed within a single summer in 5 of 9 years (1996–2004), and L3 of *P. stilesi* would not have developed within a single summer in any year.

Under a climate warming scenario, the maximum number of cumulative degree-days at Palmer Lake increased through the 2020, 2050, and 2080 time slices for both *P. odocoilei* and *P. stilesi* (Table 3). As warming progressed, the transmission period lengthened (due to infective larvae becoming available earlier in the year and gastropod activity extending later in the year), and larvae could begin development later in the year and still reach infective stage. In all time slices, infective L3 of *P. stilesi* would be available approximately 3 weeks later in the year than *P. odocoilei*, and the last date that larvae of *P. stilesi* could begin development and reach infective stage larvae occurred approximately 3 weeks earlier in the year (Table 3).

## DISCUSSION

### *Epidemiology of protostrongylid parasites*

*Parasites.* In the Mackenzie Mountains, large numbers of L1 of *P. odocoilei* and *P. stilesi* were deposited by Dall's sheep over winter, with peaks in March–May. Using the mean intensity of larval shedding for November 2000–April 2001 ( $\sim 600$  LPG for each protostrongylid), and an average daily faecal production of  $\sim 300$  grams (Jenkins *et al.* 2005b), each sheep would deposit approximately 30 million L1 of each protostrongylid species on the winter range. Based on laboratory investigation of the effects of temperature and humidity on survival of L1 of *P. odocoilei*, and laboratory and field trials with L1 of a related protostrongylid (*P. tenuis*) as well as *P. stilesi* (Forrester and Senger, 1963; Uhazy, Holmes and Stelfox, 1973; Shostak and Samuel, 1984; Forrester and Lankester, 1998), a proportion (20–60%) of these L1 would survive overwinter in faeces and be available to infect gastropods at snowmelt. Subsequently, gastropod infection may continue throughout summer, although L1 survival decreases with freeze-thaw cycles, increasing temperature, and exposure to sunlight (Rose, 1957;




Fig. 5. Predicted cumulative degree-days (Cum DD) for larvae of *Parelaphostrongylus odocoilei* in experiments starting in 2003 at Palmer Lake using (A) hourly surface temperatures inside enclosures and (B) hourly air temperatures. Circled markers indicate the first dates that infective larvae were observed. Thermal constant (hatched line)=163 DD.




Fig. 6. Mean biweekly air, surface, and soil temperatures (T, °C) outside enclosures at Palmer Lake from September 2002 to May 2003.

Shostak and Samuel, 1984; Lorentzen and Halvorsen, 1986). Larval shedding by sheep in the Mackenzie Mountains declined precipitously over the summer, especially for *P. stilesi*. Although a summer decline in prevalence and intensity of *Protostrongylus* spp. has been reported in some bighorn sheep populations (Uhazy *et al.* 1973; Yde, Brown and Worley, 1988), the marked decline that we observed is unusual, and could be due to seasonal

factors in combination with an interspecific interaction between parasites (Pybus *et al.* 1990; Ball, Lankester and Mahoney, 2001; Lello *et al.* 2004).

In slugs infected on 2–30 June 2003 (Exps 4, 5, and 6), infective larvae of *P. odocoilei* were first present in all 3 experiments on 28 July 2003. As well, 2 uninfected control slugs collected at this time contained L3 of *P. odocoilei* and an unnamed protostrongylid endemic to caribou in the Mackenzie



Table 2. Historical patterns of theoretical development based on recorded air temperatures at locations in Subarctic Canada where *Parelaphostrongylus odocoilei* is present (Fort Simpson, Norman Wells, and Palmer Lake) and Arctic locations where it is absent (Inuvik and Ivvavik)

	Start of season* Range	End of season* Range	Growing season† Mean (range)	Date of first L3‡ Range	Last date for L3§ Range	Cumulative DD Mean (range)
Ivvavik (corrected)	May 10 to Jun 3	Aug 28 to Oct 3	11.4 (8.6–13.3)	Jul 17 to Aug 10	Jun 9 to Jun 30	166 (117–249)
Ivvavik (uncorrected)	Apr 18 to May 31	Sep 6 to Oct 4	13.3 (10.2–16.2)	Jun 28 to Jul 19	Jul 2 to Jul 21	355 (283–510)
Inuvik	Apr 16 to May 30	Sep 5 to Oct 14	13.7 (11.0–16.6)	Jun 20 to Jul 22	Jun 30 to Aug 8	442 (300–674)
Palmer Lake 2003	Apr 24	Oct 5	16.4	Jul 19	Jul 18	320
Norman Wells	Apr 1 to May 15	Sep 10 to Oct 15	16.0 (12.6–19.0)	Jun 4 to Jul 4	Jul 17 to Aug 18	676 (450–906)
Fort Simpson	Mar 3 to Apr 30	Sep 22 to Nov 28	19.3 (15.5–23.6)	May 24 to Jun 29	Jul 23 to Aug 21	767 (540–996)

\* First and last dates that temperatures were above threshold (8.5 °C) and degree-days (DD) were accumulated.

† Number of days between the first and last dates that DD were accumulated in each year.

‡ First date that infective 3rd-stage larvae (L3) could have developed within each year.

§ Last date that 1st-stage larvae could begin development in gastropods and reach infective L3 stage before winter.




Fig. 7. Historical trend and annual variability for development of *Parelaphostrongylus odocoilei* using hourly air temperatures recorded in Norman Wells, NT, 1955–2004. (A) Length of parasite growing season (linear regression, relationship not significant). (B) Maximum number of cumulative degree-days (linear regression,  $F = 8.7$ ,  $P = 0.005$ ).

Mountains, suggesting that L3 are available in naturally infected gastropods by late July. Long periods of minimal development at cooler temperatures are likely interspersed with rapid bursts of development at warmer temperatures, leading to ‘pulses’ of infective larvae. In slugs infected in mid-July 2003 (Exp. 7), but not in those infected in mid-July 2002 (Exp. 1), infective larvae developed in early September. The summer of 2003 was approximately 0.5 °C warmer than the summer of 2002, a normal year for this region of the Mackenzie District ([http://www.msc-smc.ec.gc.ca/ccrm/bulletin/archive\\_e.cfm](http://www.msc-smc.ec.gc.ca/ccrm/bulletin/archive_e.cfm), accessed April 2005). Therefore, even small differences in overall temperature between years led to different seasonal patterns in larval development and availability.

Infective larvae had high abundance in experimentally infected *D. laeve* in August–September 2003, similar to the pattern observed in gastropods naturally infected with *P. odocoilei* and *P. stilesi* in temperate regions (Samuel *et al.* 1985; Robb and Samuel, 1990), as well as *U. pallikuukensis* in Arctic Canada (Kutz *et al.* 2002). Infective larvae in gastropods would remain available to sheep until late fall/early winter, when gastropods enter hibernation. Transmission after this point is theoretically possible, because a small proportion of infective larvae of both *P. odocoilei* and *P. stilesi* emerged from experimentally infected gastropods in the laboratory, and L3 of *P. odocoilei* survived in the environment for months under conditions simulating the subnivean environment (Monson and Post, 1972; Jenkins *et al.*

Table 3. Present and future patterns of development for *Parelaphostrongylus odocoilei* and *Protostrongylus stilesi* at Palmer Lake, Northwest Territories, based on surface temperatures recorded in 2003 and projected warming using a mid-range climate change scenario

	Start of season*	End of season*	Growing season†	Date of first L3§	Last date gastropods¶	Transmission period‡	Last date for L3¥	Cumulative DD
<i>P. odocoilei</i>								
2003	Apr 27	Oct 7	163	Jul 2	Sep 12	72	Jul 22	434
2020	Apr 27	Oct 9	165	Jun 26	Oct 12	108	Jul 26	508
2050	Apr 27	Oct 9	165	Jun 17	Oct 13	118	Aug 6	648
2080	Apr 26	Oct 13	170	Jun 11	Nov 16	158	Aug 12	763
<i>P. stilesi</i>								
2003	Apr 27	Oct 7	163	Jul 23	Sep 12	51	Jul 1	484
2020	Apr 27	Oct 9	165	Jul 16	Oct 12	88	Jul 9	564
2050	Apr 26	Oct 9	166	Jul 6	Oct 13	99	Jul 19	713
2080	Apr 26	Oct 13	170	Jun 30	Nov 16	139	Jul 26	836

\* First and last dates that temperatures were above threshold (8.5 °C) and degree-days (DD) were accumulated.

† Number of days between the first and last dates that DD were accumulated in each year.

§ First date that infective 3rd-stage larvae (L3) could have developed within each year.

¶ Last date before 5 consecutive days with average surface temperatures below 0 °C, assumed to be last day for gastropods.

‡ Number of days between the date that L3 were first available and the last date of gastropod activity.

¥ Last date that 1st-stage larvae could begin development in gastropods and reach infective L3 stage before winter.

2005c). Although we did not observe emergence of larvae of *P. odocoilei* in the field, this may reflect poor slug survival and methods used for examination. Emergence may play a lesser role in transmission of *P. odocoilei* in the Subarctic than for *U. pallikuukensis* at higher latitudes, where emergence significantly extends both the spatial and temporal windows of transmission (Kutz *et al.* 2000). Further investigation of the role of larval emergence and the possibility of overwinter transmission for *P. odocoilei* and other protostrongylids is indicated, as overwinter transmission of L3 of gastrointestinal nematodes occurs even in the High Arctic (Halvorsen *et al.* 1999).

**Gastropods.** The slug *Deroceras laeve* is important in transmission of elaphostrongylines in Canada (Lankester and Anderson, 1968; Samuel *et al.* 1985; Ball *et al.* 2001), and, along with the snails *Catinella* sp. and *Euconulus fulvus*, is probably an important intermediate host for *P. odocoilei* in the Mackenzie Mountains (Jenkins *et al.* 2005c). Based on temperature data and observations in the Mackenzie Mountains, slugs were active from May until September, as compared to April–November in temperate regions (Aitchison, 1979; Rollo and Shibata, 1991; Lankester and Peterson, 1996). In the current study, slugs of all size classes survived overwinter, with mortality of older individuals in May–June and recruitment primarily in July–August, as observed in temperate areas (Lankester and Anderson, 1968; Samuel *et al.* 1985). This pattern, and the appearance of moribund slugs consistent with senescence and shrinkage (Rollo and Shibata, 1991), likely account for low slug recoveries from field and laboratory enclosures for Exps 4, 5,

and 6. In Exp. 9, as 90–100% of indoor control slugs were recovered, low recoveries from field enclosures may have been due to the escape of small slugs (which constituted 60% of slugs in this experiment). Few slugs were recovered from overwintered enclosures (see Kutz *et al.* 2002), which may reflect natural mortality in combination with colder soil and surface temperatures inside enclosures, probably due to decreased snow cover and limited options for microhabitat selection (South, 1989).

We observed a large cohort of slugs hatching in late summer/early fall, as has been reported in temperate areas (Lankester and Anderson, 1968; Boag and Wishart, 1982). This cohort would be immediately available to L1 the following spring and is likely the most important source of infective larvae of *P. odocoilei* later in the summer, because young of the year slugs would not be available until July. This is consistent with the observation that larger, older slugs (versus young of the year) had higher prevalence of infective larvae of *P. odocoilei* in an endemic region at temperate latitudes (Samuel *et al.* 1985). The late-summer cohort may survive a second winter and die the following spring (i.e., a life-span of ~1.5 years). This differs from the conventional view that *D. laeve* is an annual species (Lankester and Anderson, 1968; Boag and Wishart, 1982; Rollo and Shibata, 1991), although longer life-spans for *D. laeve* and terrestrial snails have been described (Getz, 1959; Robb and Samuel, 1990).

**Dall's sheep.** In April–May, Dall's sheep leave their low elevation winter range (subalpine shrubs and boreal meadows), following the melting snow to higher elevations, where lambing occurs in late May (Hoefs and Cowan, 1979; Simmons, 1982). In

June–early August, sheep primarily feed on alpine tundra, where suitable gastropod habitat is thought to be scarce (Hoefs and Cowan, 1979; Boag and Wishart, 1982), although heavily used, naturally-occurring mineral licks and water sources may provide suitable gastropod habitat and thereby serve as foci of protostrongylid transmission. In late August, nursery groups of sheep (lambs, ewes, and yearlings) begin to return to the low elevation winter range, habitat more suitable for hydrophilic terrestrial gastropods like *D. laeve* and *Catinella* sp. (Hoefs and Cowan, 1979; Simmons, 1982). Upon their return, lambs are likely to become infected by consuming infective L3 in gastropods. Therefore, the observed increase in intensity of L1 shedding in November in all 3 years of the study may be due to the exponential rise in larval counts following the 70 day pre-patent period in newly infected lambs (Jenkins *et al.* 2005 *b*), in combination with increased larval shedding by adult males under rutting stress (Halvorsen, Skorping and Hansen, 1985). Intensity of larval shedding continued to rise throughout winter, which may in part reflect immunological compromise due to nutritional stress, and/or increased parasite fecundity due to seasonal triggers (Halvorsen *et al.* 1985; Slomke, Lankester and Peterson, 1995; Grenfell *et al.* 1995).

Based on seasonal overlap of protostrongylid larvae, gastropods, and Dall's sheep in the Mackenzie Mountains, the majority of Dall's sheep likely become infected in the fall on their winter range, as suggested for *P. odocoilei* in mule deer and *Protostrongylus* spp. in bighorn sheep in temperate areas (Samuel *et al.* 1985; Robb and Samuel, 1990). Transmission in spring is also possible, most likely from infective larvae of *P. odocoilei* that have overwintered in gastropods. In the current study, pre-infective larvae of *P. odocoilei* that overwintered in slugs were predicted to develop to L3 in late May and June, by which time sheep have left the winter range, nor did we observe development to L3 in overwintered slugs in Exp. 1 in mid-June (336 days p.i.). This, in combination with overwinter mortality of slugs, suggests that such larvae contribute only minimally to transmission of *P. odocoilei*. In contrast, overwinter survival and resumption of development of larvae of *U. pallikuukensis* in *D. laeve* are important for maintenance of this Arctic parasite in years when L3 do not develop in a single summer (Kutz *et al.* 2002).

#### *Historical and future development of protostrongylid parasites*

**Model validation.** Using a degree-day model, we successfully predicted rates of development of larval *P. odocoilei* in the summers of 2002 and 2003. Infective L3 were present in 1 enclosure in Exp. 6 somewhat earlier than anticipated, possibly due to

heterogeneity of microhabitat within the study area, especially for soil surface temperatures. Despite concerns about a non-linear relationship at lower temperatures and the effects of stochasticity (Saunders *et al.* 2002; Jenkins *et al.* 2005 *c*), degree-day models have been validated for larval development of *U. pallikuukensis* (Kutz *et al.* 2002) and now *P. odocoilei*, which represent different subfamilies (Muellerinae and Elaphostrongylineae) in different hosts (muskoxen and Dall's sheep) and habitats (Arctic tundra and Subarctic alpine). Such models, validated under field conditions, may have broad applicability to other protostrongylids, and perhaps other host-parasite systems (Kutz *et al.* 2005).

We used air temperatures from weather stations located nearest to Dall's sheep habitat to describe trends in protostrongylid development over the last 50 years, and to compare parasite development on a latitudinal gradient. These stations were located at lower elevations than are typically used by Dall's sheep, but long-term data sets from high elevations are unavailable for northern Canada. This overestimated the cumulative degree-days; for example, summer temperatures were  $\sim 5^\circ\text{C}$  cooler, and approximately half the number of degree-days were accumulated, at Palmer Lake in the Mackenzie Mountains than at Norman Wells in the Mackenzie River valley. Likewise, summer temperatures recorded at Ivvavik, when adjusted for elevation, were  $4^\circ\text{C}$  cooler, and the number of degree-days decreased by a half.

Air temperatures, while predictive at Palmer Lake in 2003, generally produce conservative estimates of development rates (Kutz *et al.* 2002). Therefore, we used surface temperatures recorded at Palmer Lake in 2003 as the baseline for climate change projections. This was probably a 'warm start' scenario, based on the 1955–2004 trend line for DD accumulated at Norman Wells and Environment Canada data, which indicated that average temperatures in the summer of 2003 were  $0.4^\circ\text{C}$  above the baseline since 1948 in the Mackenzie District ([http://www.msc-smc.ec.gc.ca/ccrm/bulletin/archive\\_e.cfm](http://www.msc-smc.ec.gc.ca/ccrm/bulletin/archive_e.cfm), accessed April 2005).

**Spatial patterns.** We validated the degree-day model for development of larvae of *P. odocoilei* in gastropods at the northern limits of its range (the northern Mackenzie Mountains in Subarctic Canada), and applied the degree-day model to determine if larvae of *P. odocoilei* could develop within 1 summer at the northern limits of the distribution of Dall's sheep in Canada (the British Mountains, Ivvavik National Park in Arctic Canada). At Ivvavik, using the degree-day model and temperatures corrected for elevation, development to infective larvae could have occurred within 1 summer in only 4 of the last 9 years. Under current temperature conditions at high elevation at Arctic latitudes, larvae of *P. odocoilei* would therefore have to survive

overwinter in slugs and resume development the following summer (a multi-year cycle); however, it is not clear that *P. odocoilei* can undergo a multi-year cycle in *D. laeve*, based on poor slug survival and failure of larvae to resume development after overwintering in the current study. If this, in combination with geographical barriers (isolation by distance), currently excludes *P. odocoilei* from Arctic latitudes, further climate warming may soon release this constraint on range expansion.

Under conditions suitable for larval development, *P. odocoilei* would likely establish at Arctic latitudes, as L1 are resistant to desiccation and freezing (Shostak and Samuel, 1984), and gastropod species suitable as intermediate hosts for *P. odocoilei* are present in Alaska and on the Arctic coast (Pilsbry, 1948; Dau, 1981; Hoberg *et al.* 2002). If introduced, either through translocation or natural movements of definitive hosts (such as cervids), warming could allow *P. odocoilei* to establish in naïve populations of Dall's sheep in the Arctic. Such range expansions have been associated with increased prevalence and severity of clinical disease associated with elaphostrongylines, and even epizootics and population declines (Ball *et al.* 2001).

While *P. odocoilei* is absent in Dall's sheep populations in the Arctic ranges (Richardson, British, Brooks, and Baird), *P. stilesi* is present in all wild sheep populations surveyed in Subarctic and Arctic North America (Jenkins *et al.* 2005a). Based on a degree-day model (using parameters from Samson and Holmes, 1985), larvae of *P. stilesi*, with its higher DD requirements, would not have developed to infective stage in 1 summer at Ivavik, the northernmost location, in any of the last 9 years. This suggests that *P. stilesi* must undergo a multi-year cycle at Arctic latitudes, as described for another Arctic parasite, *U. pallikuukensis* (Kutz *et al.* 2002). Alternatively, transplacental transmission (involving hypobiosis of L3 in ewes, possibly unique to this species of protostrongylid) may allow maintenance of *P. stilesi* in Dall's sheep at these latitudes (Hibler *et al.* 1974). The degree-day model for *P. stilesi* should be validated in the field using a native intermediate host before it is broadly applied; the parameters derived by Samson and Holmes (1985) for *P. stilesi* were based on development in a snail species that may not be a natural intermediate host, potentially accounting for slower development and the higher thermal constant. A longer developmental period for *P. stilesi* may also reflect evolutionary constraints. *Protostrongylus* spp. are thought to have evolved with caprines in Eurasia, and later became distributed at high latitudes in North America (Jenkins, 2005).

Through application of a degree-day model, we have examined temperature as a potentially limiting factor for parasite distribution and identified a wildlife population at risk. This provides the basis for wildlife managers to avoid translocation of

thinhorn sheep, a common practice for bighorn sheep in North America, and eliminates the need for experimental translocation of parasites to determine if establishment is possible in non-endemic regions (Zarnke *et al.* 1990; Hoffmann and Blows, 1994).

*Temporal trends.* Based on our findings and a previous study on *U. pallikuukensis* (Kutz *et al.* 2005), climate warming has already increased the length of the growing season for protostrongylid development and the amount of heating available for larval development in Subarctic and Arctic Canada. Transmission of protostrongylids to Dall's sheep in the fall likely occurs in a narrow window between mid-August, when sheep return to winter range, and mid-September, when gastropods go into hibernation. Climate warming will cause L3 to be available earlier in the year, but this may have little significance if sheep do not return to the winter range before mid-August. Climate warming may have its most significant effects by extending gastropod activity earlier and later in the season, and by increasing the absolute numbers of infective larvae available to infect sheep. As well, at high elevations at Arctic latitudes, climate warming may increase the number of years when development within 1 summer is possible, facilitating a shift from a multi- to a 1-year cycle, similar to that described for *U. pallikuukensis* (Kutz *et al.* 2005).

Climate warming could lead to amplification of parasite populations and increased transmission to sheep, possibly resulting in disease outbreaks in endemic regions (Hoberg, Kocan and Rickard, 2001; Kutz *et al.* 2004). Outbreaks of clinical cerebrospinal elaphostrongylosis (caused by *E. rangiferi*) in reindeer in northern Norway occurred in years when summer temperatures were 1.5 °C above normal (Handeland and Slettbakk, 1994). Recent findings suggest that, in addition to muscular and respiratory pathology, *P. odocoilei* also has the potential to cause neurological disease in thinhorn sheep (Jenkins *et al.* 2005b). Based on its higher DD requirements and distribution at Arctic latitudes, climate warming could have disproportionately greater effects on transmission of *P. stilesi* as compared to *P. odocoilei*. *Protostrongylus stilesi* is a ubiquitous lungworm of bighorn and thinhorn sheep, and in thinhorn sheep, may cause additive or even synergistic pulmonary pathology in conjunction with eggs and larvae of *P. odocoilei* (Uhazy *et al.* 1973; Kutz *et al.* 2001; Jenkins, 2005). Healthy Dall's sheep populations (total 14 000–26 000 animals) in the Mackenzie Mountains (140 000 km<sup>2</sup> area) are valuable resources for subsistence harvest and trophy hunting (Veitch *et al.* 1998).

Although our investigation focused on the positive effects of warming on larval development, warming may also increase mortality of free-living L1 (Levine,

1963), which could in part compensate for amplification of L3 numbers. However, L1 may be produced in sufficient numbers that increased mortality would have little effect on larval abundance in gastropods. Based on our field trials, there was no evidence that infected slugs experienced higher mortality than uninfected controls, suggesting that increased host mortality would not compensate for increased L3 availability.

In addition to temperature, humidity would also influence survival of L1 and the distribution and abundance of gastropods. It was excluded from our model for larval development within gastropods because it was assumed that slugs would buffer larvae from extremes of desiccation through microhabitat selection, behaviour, and obligate maintenance of water homeostasis (Rollo, 1991). Precipitation was the limiting factor for transmission of *Fasciola* sp., another gastropod-transmitted parasite, in temperate England (Ollerenshaw and Smith, 1969), and is important in the Mackenzie Mountains, which receive a desert-like 25–30 cm of rainfall each year, primarily in July and August (Simmons, 1982). However, all climate scenarios project wetter conditions for this region (<http://www.cics.uvic.ca/scenarios/>, accessed Feb. 2005), which favours protostrongylid transmission by increasing gastropod abundance and contact rates with L1 (Uhazy *et al.* 1973; Forrester and Littell, 1976).

In conclusion, the high prevalence of protostrongylid infections in Dall's sheep in Subarctic and Arctic North America bears testament to the success of these parasites at northern latitudes. This may in part reflect abundance and environmental resistance of free-living stages, seasonal concentration of hosts and parasites on winter range, and mobile intermediate hosts capable of microhabitat selection (Inglis, 1965; Dobson *et al.* 2003). Under a regime of climate warming, these parasites may undergo amplification in endemic regions, and range expansion into naïve host populations. Without targeted surveillance programs, such signs may go undetected, especially in remote northern regions. Predictive models are key to determining vulnerabilities of wildlife due to climate warming, and to monitoring and proactively managing wildlife diseases (Kutz *et al.* 2004). Unfortunately, few empirically-derived and field-validated models are available for the broad range of pathogens that will be affected by climate change. As well, such models must be complemented by sound knowledge of the biology and temporal and spatial patterns of availability of both pathogens and hosts (Morgan *et al.* 2004). Such data are critically lacking for northern wildlife, where survey and inventory of pathogen biodiversity (Hoberg *et al.* 2003), as well as baseline data on biologically relevant parameters recorded in appropriate microhabitat types (Danks, 1992), are still desperately needed.

We gratefully acknowledge the financial and logistical support of Bill and Carol McKenzie and the Grinde family (Gana River Outfitters), and the Association of Mackenzie Mountain Outfitters, without whom this research would not have been possible. We also thank Michelle Swallow, Arianna Zimmer, Jenny Skific, Ariel Brunn, Aaron Genest, Richard Popko, Sean Tutcho, Nancey Stevens-Whiteman, James Auld, and Brent Wagner for invaluable assistance. We are grateful to North-Wright Airways, Canadian Helicopters, and Sahtu Helicopters for transporting researchers, mail, and field supplies. Don Ryback at the Meteorological Service of Canada provided weather data. Mathew Herperger and Greg Appleyard performed molecular identification of larvae.

This project was funded by the following organizations: Arctic Institute of North America Grant-in-Aid; Association of Mackenzie Mountain Outfitters; Northern Scientific Training Program; Department of Environment and Natural Resources, Government of the NT; Foundation for North American Wild Sheep, national and regional chapters of Alberta, Iowa, Utah, and Washington; Fraternity of the Desert Bighorn; Safari Club International (Pittsburgh Chapter); Western College of Veterinary Medicine (WCVM) Interprovincial Graduate and Undergraduate Student Scholarships; WCVM Wildlife Health Fund; and the Wild Sheep Society of British Columbia.

#### REFERENCES

- Aitchison, C. W.** (1979). Notes on low temperature activity of oligochaetes, gastropods, and centipedes in southern Canada. *American Midland Naturalist* **102**, 399–400.
- Ball, M. C., Lankester, M. W. and Mahoney, S. P.** (2001). Factors affecting the distribution and transmission of *Elaphostrongylus rangiferi* (Protostrongylidae) in caribou (*Rangifer tarandus caribou*) of Newfoundland, Canada. *Canadian Journal of Zoology* **79**, 1265–1277.
- Boag, D. A. and Wishart, W. D.** (1982). Distribution and abundance of terrestrial gastropods on a winter range of bighorn sheep in southwestern Alberta. *Canadian Journal of Zoology* **60**, 2633–2640.
- Bowyer, R. T. and Leslie, D. M.** (1992). *Ovis dalli*. *Mammalian Species* **393**, 1–7.
- Dainton, B. H.** (1989). Field and laboratory observations on slug and snail behaviour. In *Monograph No. 41 Slugs and Snails in World Agriculture* (ed. Henderson, I.), pp. 201–207. British Crop Protection Council, Thornton Heath, UK.
- Danks, H. V.** (1992). Arctic insects as indicators of environmental change. *Arctic* **45**, 159–166.
- Dau, J.** (1981). *Protostrongylus*, *Muellerius* and *Dictyocaulus*. In *Alaskan Wildlife Diseases* (ed. Dieterich, R. A.), pp. 141–146. University of Alaska, Fairbanks, Alaska, USA.
- Dobson, A., Kutz, S., Pascual, M. and Winfree, R.** (2003). Pathogens and parasites in a changing climate. In *Climate Change and Biodiversity: Synergistic Impacts, Advances in Applied Biodiversity Science* (ed. Hannah, L. and Lovejoy, T. E.), pp. 33–38. Centre for Applied Biodiversity Science, Conservation International, Washington, DC, USA.
- Forrester, D. J. and Littell, R. C.** (1976). Influence of rainfall on lungworm infections in bighorn sheep. *Journal of Wildlife Diseases* **12**, 48–51.

- Forrester, D. J. and Senger, C. M.** (1963). Effect of temperature and humidity on survival of first-stage *Protostrongylus stilesi* larvae. *Experimental Parasitology* **13**, 83–89.
- Forrester, S. G. and Lankester, M. W.** (1997). Extracting protostrongylid nematode larvae from ungulate feces. *Journal of Wildlife Diseases* **33**, 511–516.
- Forrester, S. G. and Lankester, M. W.** (1998). Over-winter survival of first-stage larvae of *Parelaphostrongylus tenuis* (Nematoda: Protostrongylidae). *Canadian Journal of Zoology* **76**, 704–710.
- Getz, L. L.** (1959). Notes on the ecology of slugs: *Arion circumscriptus*, *Deroceras reticulatum*, and *D. laeve*. *American Midland Naturalist* **61**, 485–498.
- Grenfell, B. T., Wilson, K., Isham, V. S., Boyd, H. E. G. and Dietz, K.** (1995). Modelling patterns of parasite aggregation in natural populations: trichostrongylid nematode-ruminant interactions as a case study. *Parasitology* **111** (Suppl.), S135–S151.
- Halvorsen, O., Skorpung, A. and Hansen, K.** (1985). Seasonal cycles in the output of first stage larvae of the nematode *Elaphostrongylus rangiferi* from reindeer, *Rangifer tarandus tarandus*. *Polar Biology* **5**, 49–54.
- Halvorsen, O., Stien, A., Irvine, J., Langvain, R. and Albon, S.** (1999). Evidence for continued transmission of parasitic nematodes in reindeer during the Arctic winter. *International Journal for Parasitology* **29**, 567–579.
- Handeland, K. and Slettbakk, T.** (1994). Outbreaks of clinical cerebrospinal elaphostrongylosis in reindeer (*Rangifer tarandus tarandus*) in Finnmark, Norway, and their relation to climatic conditions. *Journal of Veterinary Medicine Series B* **41**, 407–410.
- Harvell, C. D., Mitchell, C. E., Ward, J. R., Altizer, S., Dobson, A. P., Ostfeld, R. S. and Samuel, M. D.** (2002). Climate warming and disease risks for terrestrial and marine biota. *Science* **296**, 2158–2162.
- Hassol, S. J.** (2004). *Impacts of a Warming Arctic, Arctic Climate Impact Assessment*, Cambridge University Press, New York, USA.
- Hibler, C. P., Metzger, C. J., Spraker, T. R. and Lange, R. E.** (1974). Further observations on *Protostrongylus* sp. infection by transplacental transmission in bighorn sheep. *Journal of Wildlife Diseases* **10**, 39–41.
- Hoberg, E. P., Kocan, A. A. and Rickard, L. G.** (2001). Gastrointestinal strongyles in wild ruminants. In *Parasitic Diseases of Wild Mammals*, 2nd Edn (ed. Samuel, W. M., Pybus, M. J. and Kocan, A. A.), pp. 193–227. Iowa State University Press, Ames, Iowa, USA.
- Hoberg, E. P., Kutz, S. J., Galbreath, K. and Cook, J.** (2003). Arctic biodiversity: from discovery to faunal baseline – revealing the history of a dynamic ecosystem. *Journal of Parasitology* **89** (Suppl.), S84–S95.
- Hoberg, E. P., Kutz, S. J., Nagy, J., Jenkins, E., Elkin, B., Branigan, M. and Cooley, D.** (2002). *Protostrongylus stilesi* (Nematoda: Protostrongylidae): ecological isolation and putative host-switching between Dall's sheep and muskoxen in a contact zone. *Comparative Parasitology* **69**, 1–9.
- Hoberg, E. P., Polley, L., Gunn, A. and Nishi, J. S.** (1995). *Umingmakstrongylus pallikuukensis* gen nov et sp nov (Nematoda: Protostrongylidae) from muskoxen, *Ovibos moschatus*, in the central Canadian Arctic, with comments on biology and biogeography. *Canadian Journal of Zoology* **73**, 2266–2282.
- Hoefs, M. and Cowan, I. M.** (1979). Ecological investigation of a population of Dall sheep (*Ovis dalli dalli* Nelson). *Syesis* **12** (Suppl.), S1–S81.
- Hoffmann, A. A. and Blows, M. W.** (1994). Species borders: ecological and evolutionary perspectives. *Trends in Ecology and Evolution* **9**, 223–227.
- Inglis, W. G.** (1965). Patterns of evolution in parasitic nematodes. In *Evolution of Parasites, Third Symposium of the British Society for Parasitology* (ed. Taylor, A. E. R.), pp. 79–124. Blackwell Scientific Publications, Oxford, UK.
- Jenkins, E. J.** (2005). Ecological investigation of a new host-parasite relationship: *Parelaphostrongylus odocoilei* in thinhorn sheep (*Ovis dalli*), Ph.D. Thesis, University of Saskatchewan, Saskatoon, Saskatchewan, Canada.
- Jenkins, E. J., Appleyard, G. D., Hoberg, E. P., Rosenthal, B. M., Kutz, S. J., Veitch, A. M., Schwantje, H. M., Elkin, B. T. and Polley, L.** (2005a). Geographic distribution of the muscle-dwelling nematode *Parelaphostrongylus odocoilei* in North America, using molecular identification of first-stage larvae. *Journal of Parasitology* **91**, 574–584.
- Jenkins, E. J., Hoberg, E. P. and Polley, L.** (2005b). Development and pathogenesis of *Parelaphostrongylus odocoilei* (Nematoda: Protostrongylidae) in experimentally infected thinhorn sheep (*Ovis dalli*). *Journal of Wildlife Diseases* (in the Press).
- Jenkins, E. J., Kutz, S. J., Hoberg, E. P. and Polley, L.** (2005c). Bionomics of larvae of *Parelaphostrongylus odocoilei* (Nematoda: Protostrongylidae) in experimentally-infected gastropod intermediate hosts. *Journal of Parasitology* (in the Press).
- Kutz, S. J., Hoberg, E. P., Nagy, J., Polley, L. and Elkin, B.** (2004). “Emerging” parasitic infections in arctic ungulates. *Integrative and Comparative Biology* **44**, 109–118.
- Kutz, S. J., Hoberg, E. P., Nishi, J. and Polley, L.** (2002). Development of the muskox lungworm, *Umingmakstrongylus pallikuukensis* (Protostrongylidae), in gastropods in the Arctic. *Canadian Journal of Zoology* **80**, 1977–1985.
- Kutz, S. J., Hoberg, E. P. and Polley, L.** (2000). Emergence of third-stage larvae of *Umingmakstrongylus pallikuukensis* from three gastropod intermediate host species. *Journal of Parasitology* **86**, 743–749.
- Kutz, S. J., Hoberg, E. P., Polley, L. and Jenkins, E. J.** (2005). Global warming is changing the dynamics of Arctic host-parasite systems. *Proceedings of the Royal Society of London, B* (in the Press). DOI: 10.1098/rspb.2005.3285
- Kutz, S. J., Veitch, A. M., Hoberg, E. P., Elkin, B. T., Jenkins, E. J. and Polley, L.** (2001). New host and geographic records for two protostrongylids in Dall's sheep. *Journal of Wildlife Diseases* **37**, 761–774.
- Lankester, M. W. and Anderson, R. C.** (1968). Gastropods as intermediate hosts of *Pneumostrongylus tenuis* Dougherty of white-tailed deer. *Canadian Journal of Zoology* **46**, 373–383.
- Lankester, M. W. and Peterson, W. J.** (1996). The possible importance of wintering yards in the

- transmission of *Parelaphostrongylus tenuis* to white-tailed deer and moose. *Journal of Wildlife Diseases* **32**, 31–38.
- Lello, J., Boag, B., Fenton, A., Stevenson, I. R. and Hudson, P. J.** (2004). Competition and mutualism among the gut helminths of a mammalian host. *Nature, London* **428**, 840–844.
- Levine, N. D.** (1963). Weather, climate, and the bionomics of ruminant nematode larvae. *Advances in Veterinary Science* **8**, 215–261.
- Lorentzen, G. and Halvorsen, O.** (1986). Survival of the first stage larva of the metastrongyloid nematode *Elaphostrongylus rangiferi* under various conditions of temperature and humidity. *Holarctic Ecology* **9**, 301–304.
- Monson, R. A. and Post, G.** (1972). Experimental transmission of *Protostrongylus stilesi* to bighorn-mouflon sheep hybrids. *Journal of Parasitology* **58**, 29–33.
- Morgan, E. R., Milner-Gulland, E. J., Torgerson, P. R. and Medley, G. F.** (2004). Ruminating on complexity: macroparasites of wildlife and livestock. *Trends in Ecology and Evolution* **19**, 181–188.
- Ollerenshaw, C. B. and Smith, L. P.** (1969). Meteorological factors and forecasts of helminthic disease. *Advances in Parasitology* **7**, 283–323.
- Pilsbry, H. A.** (1948). *Land Mollusca of North America (north of Mexico), Volume II, Part 2*. Academy of Natural Sciences Monograph No. 3, Philadelphia, Pennsylvania, USA.
- Pybus, M. J., Samuel, W. M., Welch, D. A. and Wilke, C. J.** (1990). *Parelaphostrongylus andersoni* (Nematoda: Protostrongylidae) in white-tailed deer from Michigan. *Journal of Wildlife Diseases* **26**, 535–537.
- Robb, L. A. and Samuel, W. M.** (1990). Gastropod intermediate hosts of lungworms (Nematoda: Protostrongylidae) on a bighorn sheep winter range: aspects of transmission. *Canadian Journal of Zoology* **68**, 1976–1982.
- Rollo, C. D.** (1991). Endogenous and exogenous regulation of activity in *Deroceras reticulatum*, a weather-sensitive terrestrial slug. *Malacologia* **33**, 199–220.
- Rollo, C. D. and Shibata, D. M.** (1991). Resilience, robustness, and plasticity in a terrestrial slug, with particular reference to food quality. *Canadian Journal of Zoology* **69**, 978–987.
- Rose, J. H.** (1957). Observations on the bionomics of the free-living first stage larvae of the sheep lungworm, *Muellerius capillaris*. *Journal of Helminthology* **31**, 17–28.
- Samson, J. and Holmes, J. C.** (1985). The effect of temperature on rates of development of larval *Protostrongylus* spp. (Nematoda, Metastrongyloidea) from bighorn sheep, *Ovis canadensis canadensis*, in the snail *Vallonia pulchella*. *Canadian Journal of Zoology* **63**, 1445–1448.
- Samuel, W. M., Platt, T. R. and Knispel-Krause, S. M.** (1985). Gastropod intermediate hosts and transmission of *Parelaphostrongylus odocoilei*, a muscle-inhabiting nematode of mule deer, *Odocoileus h. hemionus*, in Jasper National Park, Alberta. *Canadian Journal of Zoology* **63**, 928–932.
- Saunders, L. M., Tompkins, D. M. and Hudson, P. J.** (2002). Stochasticity accelerates nematode egg development. *Journal of Parasitology* **88**, 1271–1272.
- Shostak, A. W. and Samuel, W. M.** (1984). Moisture and temperature effects on survival and infectivity of first-stage larvae of *Parelaphostrongylus odocoilei* and *P. tenuis* (Nematoda: Metastrongyloidea). *Journal of Parasitology* **70**, 261–269.
- Simmons, N. M.** (1982). Seasonal ranges of Dall's sheep, Mackenzie Mountains, Northwest Territories. *Arctic* **35**, 512–518.
- Slomke, A. M., Lankester, M. W. and Peterson, W. J.** (1995). Intrapopulation dynamics of *Parelaphostrongylus tenuis* in white-tailed deer. *Journal of Wildlife Diseases* **31**, 125–135.
- South, A.** (1989). The effect of weather and other factors on numbers of slugs on permanent pasture. In *Monograph No. 41 Slugs and Snails in World Agriculture* (ed. Henderson, I.), pp. 355–360. British Crop Protection Council, Thornton Heath, UK.
- Uhazy, L. S., Holmes, J. C. and Stelfox, J. G.** (1973). Lungworms in the Rocky Mountain bighorn sheep of western Canada. *Canadian Journal of Zoology* **51**, 817–824.
- Veitch, A. M., Simmons, E., Adamczewski, J. and Popko, R.** (1998). Status, harvest, and co-management of Dall's sheep in the Mackenzie Mountains, Northwest Territories. *Biennial Symposium of the Northern Wild Sheep and Goat Council* **11**, 134–153.
- Worley, K., Strobeck, C., Arthur, S., Carey, J., Schwantje, H., Veitch, A. and Coltman, D. W.** (2004). Population genetic structure of North American thimhorn sheep (*Ovis dalli*). *Molecular Ecology* **13**, 2545–2556.
- Yde, C. A., Brown, G. W. and Worley, D.** (1988). Lungworm larvae discharge levels within the Ural-Tweed bighorn sheep population. *Biennial Symposium of the Northern Wild Sheep and Goat Council* **6**, 84–90.
- Zarnke, R. L., Samuel, W. M., Franzmann, A. W. and Barrett, R.** (1990). Factors influencing the potential establishment of the winter tick (*Dermacentor albipictus*) in Alaska. *Journal of Wildlife Diseases* **26**, 412–415.