

# Concentration and retention of *Toxoplasma gondii* surrogates from seawater by red abalone (*Haliotis rufescens*)

KRISTEN C. SCHOTT<sup>1</sup>, COLIN KRUSOR<sup>2</sup>, M. TIM TINKER<sup>3,4</sup>, JAMES MOORE<sup>5</sup>, PATRICIA A. CONRAD<sup>2,6</sup> and KAREN SHAPIRO<sup>2,6\*</sup>

<sup>1</sup> Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544, USA

<sup>2</sup> Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine, University of California-Davis, Davis, CA 95616, USA

<sup>3</sup> Long Marine Laboratory, U.S. Geological Survey, Western Ecological Research Center, 100 Shaffer Road, Santa Cruz, CA 95060, USA

<sup>4</sup> Ecology and Evolutionary Biology, University of California, Santa Cruz, CA 95060, USA

<sup>5</sup> California Department of Fish and Wildlife and UC Davis-Bodega Marine Laboratory, PO Box 247, Bodega Bay, CA 94923, USA

<sup>6</sup> One Health Institute, School of Veterinary Medicine, University of California-Davis, Davis, CA 95616, USA

(Received 7 March 2016; revised 5 June 2016; accepted 1 July 2016; first published online 30 August 2016)

## SUMMARY

Small marine snails and abalone have been identified as high- and low-risk prey items, respectively, for exposure of threatened southern sea otters to *Toxoplasma gondii*, a zoonotic parasite that can cause fatal encephalitis in animals and humans. While recent work has characterized snails as paratenic hosts for *T. gondii*, the ability of abalone to vector the parasite has not been evaluated. To further elucidate why abalone predation may be protective against *T. gondii* exposure, this study aimed to determine whether: (1) abalone are physiologically capable of acquiring *T. gondii*; and (2) abalone and snails differ in their ability to concentrate and retain the parasite. Abalone were exposed to *T. gondii* surrogate microspheres for 24 h, and fecal samples were examined for 2 weeks following exposure. Concentration of surrogates was 2–3 orders of magnitude greater in abalone feces than in the spiked seawater, and excretion of surrogates continued for 14 days post-exposure. These results indicate that, physiologically, abalone and snails can equally vector *T. gondii* as paratenic hosts. Reduced risk of *T. gondii* infection in abalone-specializing otters may therefore result from abalone's high nutritional value, which implies otters must consume fewer animals to meet their caloric needs.

Key words: *Toxoplasma gondii*, transmission, oocysts, surrogates, abalone, marine snails, sea otter.

## INTRODUCTION

*Toxoplasma gondii* is a globally distributed pathogenic parasite with a heteroxenous life-cycle that infects warm-blooded vertebrates, including humans (Tenter *et al.* 2000). The parasite undergoes sexual reproduction in the intestines of infected cats, resulting in the shedding of hundreds of millions of environmentally resistant oocysts in the feces (Dubey and Frenkel, 1972; Dubey, 1998). Although felids are the only known definitive hosts, *T. gondii* infections have been observed in both pelagic and coastal marine mammal species (Dubey *et al.* 2003), indicating the presence of land-to-sea pathogen pollution (Conrad *et al.* 2005). Of particular concern are reports documenting *T. gondii* encephalitis as an important cause of mortality in threatened southern sea otters (*Enhydra lutris nereis*) (Kreuder *et al.* 2003). Sea otters are vital for maintaining a healthy nearshore ecosystem due to their role as keystone predators

(Estes and Palmisano, 1974) and can serve as sentinels for marine water contamination (Conrad *et al.* 2005). Because many sea otter prey species are also harvested as seafood for human consumption, the high prevalence of *T. gondii* in sea otters raises concerns about the risks to human health. Although *T. gondii* infections are usually asymptomatic in otherwise healthy humans, systemic toxoplasmosis can occur in immunocompromised patients, such as people suffering from acquired immune deficiency syndrome (AIDS), and newly acquired infections in pregnant women may cause abortion, stillbirth or serious birth defects due to congenital toxoplasmosis (Luft *et al.* 1993; Dubey and Jones, 2008). In addition, immunocompetent adults have been reported to develop ocular diseases such as retinochoroiditis, and latent *T. gondii* infections may contribute to mental illness (Flegr, 2007; Dubey and Jones, 2008). Gaining an understanding of transmission processes resulting in the infection of sea otters with zoonotic pathogens is therefore important for both conservation of the species and mitigation of a potential public health threat.

One hypothesis that has been proposed as a possible mechanism for transmission of *T. gondii* to sea otters is that oocysts, upon being carried into

\* Corresponding author: Department of Pathology, Microbiology, and Immunology, School of Veterinary Medicine, University of California Davis, One Shields Avenue, Davis, CA 95616, USA. E-mail: [kshapiro@ucdavis.edu](mailto:kshapiro@ucdavis.edu)

the ocean by fresh-water runoff, are concentrated in filter-feeding shellfish commonly preyed upon by sea otters (Miller *et al.* 2002). Bivalve molluscs have been shown to concentrate *T. gondii* oocysts from seawater and retain them for at least 3 days in mussels (Arkush *et al.* 2003) and 85 days in oysters (Lindsay *et al.* 2004) without reducing infectivity. However, epidemiological investigations did not support an association between predation upon bivalves and increased risk of *T. gondii* exposure in otters (Johnson *et al.* 2009). Instead, the same study determined that otters with a diet consisting of  $\geq 10\%$  small marine turban snails (including *Pro-martynia pulligo*, *Chlorostoma brunnea*, *Chlorostoma montereyi* and *Lithopoma undosa*) were 12 times more likely to be seropositive for *T. gondii* than otters that consumed fewer snails, whereas otters with a preference for abalone were significantly less likely to be infected (Johnson *et al.* 2009). Given that both snails and abalone are herbivorous gastropods, this finding raises the question of why species with such similar diets differ so greatly with respect to their epidemiological association with *T. gondii* in sea otters.

A prior study performed to evaluate marine snails as potential paratenic hosts for *T. gondii* showed that snails are capable of concentrating oocysts from seawater by 2–3 orders of magnitude and retaining them for as long as 11 days post-exposure (Krusor *et al.* 2015). However, no such work with abalone has been published to date. As discussed by Johnson *et al.* (2009), the observed reduction in exposure risk associated with a preference for abalone could be either the result of a reduced ability of abalone to acquire, concentrate and retain *T. gondii* oocysts, or a product of the abalone's greater size and the smaller number of individuals therefore needed to satisfy the otter's metabolic needs. The aim of the current study was to examine which of these hypotheses is more plausible by testing whether abalone are physiologically capable of acquiring *T. gondii* from contaminated seawater, and if they are, to compare *T. gondii* concentration and retention ability between abalone and small marine snails.

## MATERIALS AND METHODS

### *Study animals*

This study was conducted using 15 red abalone (*Haliotis rufescens*) maintained at the UC Davis Bodega Marine Laboratory. Average animal weight was 31.6 g. This species was selected because it has been shown to be commonly preyed upon by sea otters in resource-rich systems in central California (Wendell, 1994; Tinker *et al.* 2008), and because it is the most abundant of the abalone species found

within the southern sea otter range (Allen *et al.* 2005).

### *Toxoplasma gondii surrogate microspheres*

To eliminate the biohazardous risk of environmental contamination with *T. gondii* oocysts at the common working space at Bodega Marine Laboratory, Dragon Green (DG) polystyrene microspheres (10.4  $\mu\text{m}$ ) were used as surrogates for *T. gondii* oocysts. These microspheres were validated as appropriate surrogate particles for *T. gondii* oocysts due to their similar physical and chemical surface properties (Shapiro *et al.* 2009) and have been previously shown to accurately mimic oocyst transport in water and passage behaviour through snails (Shapiro *et al.* 2012; Krusor *et al.* 2015). Microspheres were obtained from Bangs Laboratory, Fishers, IN (product number FCO7F/5493).

### *Concentration and retention of T. gondii surrogates by abalone*

Abalone were placed in 4 L tubs, with one control tub (A) containing three abalone and two exposure tubs (B and C) containing six abalone each (Fig. 1). Tub A was filled with continuously aerated filtered (30  $\mu\text{m}$ ) seawater and one 5  $\times$  7.5 cm<sup>2</sup> kelp (*Macrocystis pyrifera*) blade per abalone and maintained in a cold (12–15 °C) seawater bath for 72 h to allow for acclimation. During this period, kelp was replaced, seawater changed, and feces collected using a serological pipette twice daily. At the end of the acclimation period, the exposure tubs (B and C) were spiked with *T. gondii* surrogate microspheres at a final concentration of ten microspheres per mL, while tub A was not spiked and served as a negative control. A fourth tub (designated tub D) containing only kelp and seawater was also spiked with *T. gondii* surrogate microspheres at a final concentration of ten microspheres per mL to determine surrogate distribution in the tub in the absence of abalone. After a 24-h exposure period, fecal samples were collected from tubs A, B and C, and each abalone was placed in an individual 1 L container after being thoroughly rinsed with filtered seawater to remove any surrogates that may have adhered to the shell or foot of the animal. Each container contained aerated filtered seawater and kelp as a source of food. In addition to feces, water samples (50 mL) were also collected from the top, middle, and bottom of the surrogate control tub (D) at the end of the exposure period to determine surrogate distribution in the tub in the absence of abalone. All remaining kelp fragments were also collected and transported to our laboratory at the School of Veterinary Medicine, UC Davis, for surrogate enumeration as described below.




Fig. 1. A schematic diagram representing experimental setup and sample collection. Abalone were placed in 4 L tubs and allowed to acclimate for 72 h, then exposed to *Toxoplasma gondii* oocyst surrogates for 24 h. At the end of the exposure period, feces and kelp were collected from tubs A, B and C, and water and kelp were collected from tub D. Abalone were then placed in individual 1 L containers and monitored for 14 days, during which time daily fecal samples were collected from each animal for surrogate enumeration.

For 14 days following the exposure period, remaining kelp fragments were removed, photographed and discarded, and fecal samples were collected from each 1 L container at 24-h intervals. At the same intervals, individual containers were thoroughly rinsed and fresh filtered, aerated seawater and kelp were provided for each animal. To derive data on the amount of kelp consumed by each abalone, kelp was photographed prior to and following each 24-h interval and subsequently analysed using ImageJ (Schneider *et al.* 2012) to measure the surface areas of each kelp fragment. Aliquots of each fecal sample from days 0, 1, 2 and 3 and complete fecal samples from days 4, 7, 11 and 14 were processed and examined for surrogate enumeration as described below.

*Enumeration of T. gondii surrogates in water and fecal samples*

Detection of *T. gondii* surrogate microspheres in water and abalone feces was conducted following previously published protocols (Shapiro *et al.* 2010a; Krusor *et al.* 2015). Briefly, water samples were filtered through 5 µm mixed cellulose membranes that were mounted on glass slides. Prepared slides were then examined at 100× using a Zeiss Axioskop epifluorescence microscope equipped with a UVA filter. Fecal samples were centrifuged and the volumes of the resulting fecal pellets were estimated. Fecal pellets were then homogenized with a 20 mL syringe equipped with a blunt-tipped 15-gauge needle, diluted with water to

produce a thin slurry, and examined according to the same filtration and microscopy protocols described for water samples.

*Enumeration of T. gondii surrogates from kelp surfaces*

Kelp fragments from each tub were immersed and agitated in a NaPP/guanidinium/Tween (NGT) solution prepared by adding 17.7 g granular sodium polyphosphate (NaPP) to 1 L boiling deionized water, then adding 9.55 g granular guanidine HCl (0.1 M) and 1 mL Tween 20 (Thermo Fisher Scientific, Waltham, MA) to the cooled NaPP solution. Post-immersion, fragments were scraped on both sides with a plastic spoon and rinsed on each side with 10–12 mL NGT solution. The resulting kelp-washed NGT solution was then diluted with deionized water and examined for *T. gondii* surrogates according to the filtration and microscopy protocols described above.

*Data analysis*

Data collected included abalone mass, estimated daily fecal volume, daily surface area of kelp eaten, and daily fecal surrogate counts for each abalone as well as surrogate concentrations for water samples taken from tub D following the 24 h exposure period. Because fecal samples collected on day zero (i.e. immediately following the 24-h exposure period) represent the pooled feces from all animals in each exposure tub, total fecal volume and surrogate counts for these pooled samples were divided

by the number of abalone in the tub ( $n = 3$  for tub A; and  $n = 6$  for tubs B and C). Additionally, estimated fecal volume and surrogate counts were used to calculate the concentration of surrogates per mL feces for each abalone. All data was organized using Microsoft Excel and imported into R (version 3.0.3; <https://www.r-project.org>) for analysis. The Shapiro–Wilk test was used to assess the normality of the surrogate count and concentration data for each day, and based on the results of this test, the Wilcoxon rank sum test was applied to assess the significance of differences between the median surrogate count and concentration values of control and exposed abalone. The Wilcoxon signed rank test was applied to assess the significance of differences between fecal surrogate concentrations and the nominal ‘environmental’ surrogate concentration of ten surrogates per mL of seawater in the exposure tubs. Cumulative fractions of all surrogates that were recovered over the duration of the study were also calculated for each day.

Finally, to evaluate the role of animal size as a potential ‘risk factor’ for oocyst acquisition and retention, correlation analysis (either Pearson’s  $r$  or Kendall  $\tau$ , depending on the normality of the data) was used to test for a relationship between abalone mass and maximum surrogate count, surrogate retention time, fecal volume and amount of kelp eaten. Maximum surrogate count and surrogate retention time were selected as measures of acquisition and retention, while fecal volume and amount of kelp eaten were selected as factors that could potentially influence the number of oocysts an abalone acquires.

#### *Diet-based differences in exposure rates: a feasibility analysis*

We conducted a feasibility analysis to further examine the null hypothesis of constant risk of infection: that is, the possibility that differences in seroprevalence between otters specializing on marine snails (high infection rates) and otters specializing on abalone (low infection rates) could be due simply to differences in prey capture rates, rather than differences in the probability that each prey type contains oocysts. Specifically, assuming that the per-item risk of infection ( $\gamma$ ) was identical for snails and abalone, we calculated expected infection probabilities for a typical abalone specialist and a typical snail specialist using empirical data on individual prey consumption collected from field studies of tagged sea otters in central California (Oftedal *et al.* 2007; Tinker *et al.* 2008, 2012, 2013). We used available data for 17 adult abalone specialists ( $N = 9526$  dives) and 23 adult snail specialists ( $N = 10\,797$  dives) to compute taxa-specific intake rates for each specialist type ( $\pm 1$  standard deviation (S.D.), computed across individual estimated

diets), using the Monte Carlo algorithm described in Tinker *et al.* (2012). We further assumed the likelihood that an adult otter is infected depends on the cumulative exposure risk over the previous 5-year period (1825 days), and thus we used a hazards model to estimate  $P_i$ , the probability of infection for specialist type  $i$ :

$$P_i = 1 - e^{(-1825 \times (Ab_i + Sn_i) \times \gamma)}, \quad (1)$$

where  $Ab_i$  represents the mean number of abalone consumed per day by specialist type  $i$  and  $Sn_i$  represents the mean number of snails consumed per day by specialist type  $i$  (for simplicity we assume only snails and abalone are infective, and thus ignore all other prey). In equation (1), each prey consumption event represents an additive hazard, and thus  $P_i$  is simply  $1 -$  the probability of avoiding infection over the previous 5 years. To adjust for differences in absolute consumption rates between otters of different sex and size, we re-scaled consumption rates to the average observed for a 22 kg female sea otter in Monterey or Big Sur ( $6 \text{ kg d}^{-1}$ ). At present there are no available data with which to estimate  $\gamma$ , the per-item infection risk; however, because our goal was only to calculate relative (and not absolute) rates of infection under the hypothesis of risk-equivalence, the actual value of  $\gamma$  was unimportant. Accordingly, in order to make our analysis relevant to field data on seroprevalence, we arbitrarily adjusted  $\gamma$  to produce a realized infection rate for snail specialists of 0.85 (corresponding to the value reported for snail specialists by Johnson *et al.* 2009), and then calculated the corresponding infection rate for abalone specialists. To provide an indication of within-group variability we also report associated S.D. in computed infection rates, based on the variance in individual diets. We compared the results of this feasibility analysis to the empirically derived infection rates for snail and abalone specialists reported by Johnson *et al.* (2009).

#### RESULTS

Counts and concentrations of *T. gondii* surrogate microspheres in fecal samples were found to be non-normally distributed at all-time points except for the count data from days 1 and 3 (Table S1). Therefore, non-parametric tests were applied for further statistical analyses. Fecal surrogate concentrations were significantly greater than the  $10 \text{ mL}^{-1}$  ambient exposure concentration through 3 days post-exposure (Table 1), and surrogate counts and concentrations in fecal samples from exposed abalone were significantly greater than in samples from control abalone from 0 to 2 days post-exposure (Table S2). In the first 24 h post-exposure, fecal surrogate concentrations were 2–3 orders of magnitude greater than the ambient concentration of surrogates

Table 1. Results of one-sided Wilcoxon signed rank tests employed to assess the significance of differences between the fecal surrogate concentrations at each time point and 10 mL<sup>-1</sup>, which was the nominal ambient surrogate concentration of seawater in the exposure tubs

Day	Median (range) fecal surrogate concentration (mL <sup>-1</sup> )	Number of abalone excreting surrogates/ number tested	V	P value
0	2486 (1564–3408)	12/12	78	<0.001*
1	760 (243–2480)	12/12	78	0.001*
2	105 (43–1150)	12/12	78	0.001*
3	17 (4–240)	12/12	67	0.015*
4	2 (0–11)	9/12	1	0.999
7	1 (0–20)	7/12	11	0.988
11	0 (0–9)	5/12	0	0.999
14	1 (0–20)	6/12	9	0.993

\*Statistically significant ( $P < 0.05$ ).

in seawater during the exposure phase (Fig. 2). Fecal surrogate counts declined rapidly from days 1 to 4, and 99% of all surrogates recovered in fecal samples were detected in samples from this time period. However, six abalone were observed to continue to defecate surrogates through the end of the 14-day post-exposure monitoring period (Table 1).

Unexplained contamination of control abalone occurred approximately 3 days post-exposure, resulting in observation of *T. gondii* surrogates in fecal samples collected from all three control animals on days 3 and 4. On these days, the median fecal surrogate concentration in control abalone was not significantly greater than the 10 mL<sup>-1</sup> nominal exposure concentration (Wilcoxon signed-rank tests,  $V = 3$ ,  $P = 0.625$ ;  $V = 1$ ,  $P = 0.875$ ). On day 7 and beyond, surrogates were no longer detected in fecal samples from the negative control abalone.

Abalone mass, average feces volume and maximum surrogate counts were found to be normally distributed, while average food intake and surrogate retention time were not (Table S3). Significant correlations were found between abalone mass and mean food intake (Fig. S1A; Kendall tau rank sum correlation,  $\tau = 0.4211$ ,  $P < 0.05$ ,  $n = 15$ ), mean fecal productivity (Fig. S1B; Pearson’s correlation,  $r = 0.6999$ ,  $P < 0.01$ ,  $n = 15$ ), and maximum surrogate count (Fig. S1C; Pearson’s correlation,  $r = 0.8033$ ,  $P < 0.01$ ,  $n = 12$ ). However, no correlation was found between abalone mass and retention time (Fig. S1D; Kendall tau rank sum correlation,  $\tau = 0.1551$ ,  $P = 0.1594$ ,  $n = 12$ ).

Surrogate concentrations of water samples taken from the top, middle and bottom of the surrogate control tub (D) measured at 2.6, 3.0 and 7.7 mL<sup>-1</sup>. Assuming even distribution and no loss of microspheres during the spiking procedure, a concentration

of approximately ten surrogates per mL should be measured throughout the tub. All observed values were therefore lower than expected. If each concentration value is assumed to be representative of the concentration of one-third of the water column in the tub, then the total number of microspheres suspended in the water column after 24 h is equal to approximately 50% of the total number added. An additional 10 640 surrogate microspheres, representing about 30% of the total number added, were found to have adhered to the surface of kelp fragments (Table 2). The remaining 20% most likely adhered to the walls of the tubs or were lost during the spiking or sample handling procedures. Kelp surface area, number of surrogates and surrogate density per unit area of kelp were all reduced in the presence of abalone (Table 2).

Our analysis of dietary data showed that a typical 22 kg female sea otter specializing on abalone consumed approximately  $20 \pm 4.6$  abalone and  $11 \pm 32.1$  snails per day, while a similar sea otter specializing on snails consumed approximately  $0.9 \pm 1.6$  abalone and  $705 \pm 251$  snails per day. Assuming the per-item risk of infection ( $\gamma$ ) was identical for snails and abalone, and arbitrarily setting  $\gamma = 1.65E-06$ , we computed expected infection rates of  $0.85 \pm 0.099$  for snail specialists and  $0.09 \pm 0.076$  for abalone specialists. Estimated infection rates were similar to empirically measured rates of 0.85 for snail specialists and 0.07 for abalone specialists (Johnson *et al.* 2009).

DISCUSSION

The physiological ability of abalone to ingest, concentrate, and retain *T. gondii* oocysts from the environment as determined by the present study was similar to that of snails, previously described by Krusor *et al.* (2015). *Toxoplasma gondii* surrogates were detected in abalone feces at significantly higher concentrations than the baseline surrogate concentration of the spiked seawater for three days post-exposure, indicating that abalone are physiologically capable of concentrating oocysts by ingestion. Maximum counts and concentrations of surrogates, which were 2–3 orders of magnitude higher than in the spiked seawater, were observed at 0 and 24 h post-exposure and thereafter declined swiftly through day 4. After day 4, fecal surrogate counts and concentrations did not change significantly as abalone continued to excrete consistently low numbers of surrogates in their feces through the end of the 14-day post-exposure monitoring period. While the infective dose of *T. gondii* oocysts is not known for sea otters, studies with pigs and rodents indicate that susceptible hosts may become infected following ingestion of a single oocyst (Dubey, 1996; Dubey *et al.* 1996). Therefore, the low surrogate counts observed at




Fig. 2. (A) Mean counts and (B) concentrations of *Toxoplasma gondii* surrogates detected in fecal samples from abalone and marine snails on days 0, 1, 2, 3, 4, 7, 11 and 14 following exposure to spiked seawater. Error bars represent the s.e. The dashed line represents the baseline 'environmental' surrogate concentration ( $10 \text{ mL}^{-1}$ ) in the exposure tubs. Snail data were obtained from a previously published investigation (Krusor *et al.* 2015).

Table 2. Surface area of kelp pieces remaining after the 24-h exposure period, the number and density of *Toxoplasma gondii* surrogate microspheres adhered to kelp surfaces, and the percentage of total surrogates added that were recovered from kelp surfaces

Tub <sup>a</sup>	Kelp surface area ( $\text{cm}^2$ )	Number of <i>T. gondii</i> surrogates on kelp	<i>T. gondii</i> surrogate density per $\text{cm}^2$ kelp	% of spiked <i>T. gondii</i> surrogates detected on kelp
B	202·958	3611	18	10
C	169·006	3085	18	9
D	508·902	10 640	21	30

<sup>a</sup> Tubs B and C contained abalone as the exposure animals, while tub D served as a surrogate particle control and did not contain abalone. Tub A served as the negative control and therefore no surrogate particles were added to that treatment.

later time points could still be epidemiologically significant.

Based on recently published findings by Krusor *et al.* (2015), marine snails and abalone demonstrate similar physiological ability to acquire and retain *T. gondii* from surrounding seawater (Fig. 2). Both species concentrated surrogates by 2–3 orders of magnitude compared to the concentration of surrogates in exposure water, and snails continued to excrete surrogates for 11 days (Krusor *et al.* 2015), while abalone excreted them for at least 14. Therefore, the observed epidemiologic association between prey specialization on abalone and the reduced risk of *T. gondii* infection cannot be explained by a reduction in the physiological ability of abalone to concentrate and retain oocysts.

Because the snail study was designed such that six snails were selected for dissection and fecal sample examination each day, the sample size for monitoring fecal excretion of *T. gondii* was effectively half that of the present study. While unlikely, it is possible that the six snails remaining at day 14 were by chance individuals that had shorter retention times and that the observed difference between snails and abalone is an artefact of the smaller snail

population size. In addition, while ingested material should be evident in feces within 24 h based on gut passage time (Garcia-Esquivel and Felbeck, 2009), fecal values of *T. gondii* surrogates on day 0 should be interpreted with caution in both studies due to the possibility of overestimation arising from accidental collection of non-fecal-associated surrogates from spiked exposure water.

Due to the unexplained contamination of fecal samples from negative control abalone, the data from days 4–14 must be interpreted cautiously, as the possibility of surrogate cross-contamination between animals cannot be ruled out. However, the six abalone that had surrogate-positive fecal samples on day 14 were not spatially clustered, and two of them had consistently excreted surrogates at every time point examined. For these results to be explained by contamination rather than retention of surrogates by the abalone, random, inexplicable contamination events would have had to repeatedly affect the same animals without impacting their close neighbours, which is unlikely. Thus, these results suggest that abalone may be capable of retaining *T. gondii* oocysts in their digestive tissues for 13 days or more. This retention time is much higher

than gut passage times reported for abalone fed glass microbeads, which ranged from 20 to 27.5 h (Garcia-Esquivel and Felbeck, 2009). This observation suggests that oocysts may be selectively retained within the abalone digestive system, extending the parasite retention time and risk of its transmission. To adequately measure oocyst retention time in abalone and correct for issues of contamination, a similar experiment could be performed implementing a longer monitoring period to ensure that all surrogates have been excreted prior to termination of the experiment.

In the surrogate control tub (D) containing kelp but no abalone, the concentration of surrogates was highest near the bottom of the tub, indicating that *T. gondii* surrogate microspheres tend to sink in the water column in low turbulence seawater. This result is in agreement with the findings of previous studies that examined the behaviour of *T. gondii* surrogates and oocysts in seawater and supports the hypothesis that oocysts can concentrate in the benthos (Shapiro *et al.* 2009, 2012). After 24 h, an estimated 50% of all surrogates remained suspended in seawater, and an additional 30% had adhered to kelp surfaces at a density of 21 surrogates per cm<sup>2</sup>. This density was slightly lower in the abalone exposure tubs, most likely because abalone shells provided additional surface area for surrogates to adhere to and because some surrogates had been scraped off and consumed by abalone. Furthermore, abalone size was found to be correlated with mean kelp consumption and maximum fecal surrogate count, indicating that large abalone tended to eat more, which in turn resulted in greater likelihood of ingesting oocysts. These findings support the hypothesis that adherence of oocysts to kelp fronds facilitates ingestion by herbivorous invertebrates (Mazzillo *et al.* 2013).

Our finding that abalone can indeed serve as mechanical vectors for *T. gondii* provides indirect support for an alternative explanation for the difference in infection risk observed between snail specialists and abalone specialists: namely, that the former may be at higher risk simply due to differences in the number of prey items consumed. Our feasibility analysis of this hypothesis, informed by extensive data sets on prey consumption rates of snail specialists and abalone specialists, indicated that the discrepancy in reported infection rates (85% of snail specialists *vs* 7% of abalone specialists; Johnson *et al.* 2009) can be explained entirely by differences in the number of prey items consumed, even if each individual snail and abalone conferred the same risk of infection. This remarkable outcome reflects the enormous difference in caloric content between a snail (~4.6 kcal) and an abalone (~200 kcal; Oftedal *et al.* 2007), which means that a typical snail specialist will consume over 700 individual snails per day (in addition to other prey) to meet its energetic

needs, as compared to an abalone specialist that consumes just 20 abalone. Of course the agreement between our feasibility analysis and empirical data on seroprevalence, while striking, does not provide confirmation for this hypothesis. It does, however, demonstrate that differences in infection risk between otters with different diets do not necessarily imply differences in the mechanical abilities of prey to concentrate and retain oocysts, but rather can occur as a by-product of numerical probabilities.

While the model above suggests that *T. gondii* exposure in sea otters can be largely influenced by caloric value and hence number of preferred prey that must be eaten, there may be additional factors that contribute to variation in levels of *T. gondii* contamination among different prey species. For example, one additional factor that may contribute to the probability of abalone *vs* snails encountering *T. gondii* under natural conditions is the difference in the feeding behaviour of these marine gastropods. This study focused on determining the physiological capability of abalone to serve as paratenic hosts for *T. gondii*, and as such the experimental animals were provided with an ample supply of kelp on which to feed. However, wild abalone living within sea otter habitat range are typically found in crevices or other refuges and feed opportunistically on drift algae, or occasionally on fronds of attached macroalgae that sway within their reach (Lowry and Pearse, 1973; Tutschulte and Connell, 1988). As a result, the frequency at which they feed – and therefore the likelihood that they will encounter *T. gondii* oocysts – is governed by chance and the abundance of drift algae. In contrast, marine snails inhabit surfaces of kelp and are therefore able to graze continuously upon fronds or stipes (Watanabe, 1984). The probability of accidental ingestion of *T. gondii* oocysts that become associated with kelp surfaces as described by Mazzillo *et al.* (2013) is therefore likely to be greater for snails than for abalone that opportunistically consume drifting alga.

If abalone and marine snails tend to occupy distinct sites due to variations in habitat preference, then it is also hypothetically possible that the observed differences in infection risk could be explained by abalone tending to occupy coastal sites receiving less freshwater influence, and therefore with reduced likelihood of exposure to *T. gondii* oocysts. To our knowledge, there is no data currently available on the prevalence of *T. gondii* oocyst contamination amongst abalone or snails. However, Johnson *et al.* (2009) emphasize that both abalone and snail specialist otters were observed feeding within the identified high-risk coastal segment with no obvious spatial segregation or difference in foraging microhabitats. It therefore seems unlikely that differences in spatial distribution of prey species are responsible for observed variations in infection risk. Still, field studies quantifying

the prevalence of *T. gondii* oocyst contamination amongst different prey species would be useful to elucidate the dynamics of *T. gondii* oocyst transport in the marine environment. Furthermore, such data could be used as input variables for estimating per-item infection risk used in our feasibility analysis and hence enhance the accuracy of *T. gondii* infection risk modelling.

To test the hypothesis that abalone and snail contamination with *T. gondii* is influenced by differential coastal habitat distribution, species population surveys could be performed to assess whether abalone and snails occupy the distinct nearshore sites. If this hypothesis is supported, prey and water samples could be collected from sites representative of typical habitat for each species and tested for *T. gondii* oocysts. However, one limitation of this approach is that currently, successful detection of *T. gondii* oocysts in seawater has not been reported, despite our team's effort to do so over duration of 3 years. To test for variations in oocyst exposure due to differences in feeding behaviour, animals could also be collected from a high-risk site occupied by both species. Validation of laboratory methods for detection of *T. gondii* oocysts in snails have been recently reported (Krusor *et al.* 2015), though such methods have not been tested to date in abalone. Ideally, samples of invertebrates would be collected during the wet season as well as the dry season to assess seasonal fluctuations in oocyst contamination rate due to rainfall and surface runoff. However, the practicality of conducting such a study is highly challenging. Although Staggs *et al.* (2015) reported *T. gondii* prevalences of 33 and 54% for small samples of mussels, Miller *et al.* (2008) detected only one *T. gondii*-positive mussel out of 1396 tested invertebrates collected from central coastal California, while Krusor *et al.* (2015) reported no positives out of 227 sampled marine snails. Low *T. gondii* contamination prevalences suggest that a sample size of several thousand abalone and snails would be necessary to adequately measure and compare *T. gondii* contamination in these species, which could be particularly problematic given the protected status of central California abalone populations.

Another factor that may impact the likelihood of abalone acting as paratenic hosts for *T. gondii* under natural conditions in California is the presence of withering syndrome (WS), an important abalone pathogen in this state. Clinical signs of WS include reduced feeding and withering of the foot muscle, which reduces the animal's ability to attach firmly to the substrate (Moore *et al.* 2000; Crosson *et al.* 2014). Reduced feeding would diminish the probability that abalone will obtain kelp and thus ingest *T. gondii* oocysts, while an inability to firmly attach to substrates would increase its vulnerability to predation, creating a scenario in which

abalone that suffer from WS are less likely to contain oocysts and are more likely to be preyed upon by sea otters. While WS-induced mass mortality has only been recorded in warmer waters of southern California, particularly the Channel Islands, clinical disease has been reported as far north as San Francisco, and the infectious agent of WS is distributed from Baja California to Sonoma County, California (Crosson *et al.* 2014), an area that encompasses the entirety of the southern sea otter's current range.

One limitation in the current study is that due to utilization of *T. gondii* surrogates to mitigate the biohazard nature of working with viable parasites in large aquaria, the effects of passage through abalone digestive tract on oocyst viability could not be evaluated. It is likely that the robust nature of oocysts renders them resistant to inactivation within abalone, especially as oocysts have been shown to remain infectious after passing through the digestive systems of other marine invertebrates such as mussels and oysters (Arkush *et al.* 2003; Lindsay *et al.* 2004). However, experiments using viable *T. gondii* oocysts and mouse bioassays would be needed to definitively evaluate the infectivity of oocysts after passage through the abalone gut.

The ability of abalone to concentrate and retain *T. gondii* also has important implications for public health, as it demonstrates that this commonly consumed seafood species is capable of harbouring an important zoonotic parasite of humans. Consumption of undercooked meat containing *T. gondii* tissue cysts has been a major source of human toxoplasmosis in the past, but studies of seroprevalence in livestock have demonstrated that *T. gondii* infections in meat-producing animals have been declining (Dubey and Jones, 2008). However, consumption of raw oysters, mussels and clams has been shown to be a significant risk factor for human *T. gondii* infection (Jones *et al.* 2009). If abalone can serve as transport hosts for *T. gondii* oocysts, then raw abalone could also be an important source of *T. gondii* infection for humans in spite of its apparent 'protective' effect in sea otters, particularly when harvested in large numbers by commercial fisheries.

Considering food web-driven disease transmission mechanisms in the ocean is important for conservation policy and management of both sea otters and abalone. In addition to both being threatened, sea otters and abalone have strong predator-prey interactions. Modelling-based studies have demonstrated that these interactions must be taken into account to set and achieve reasonable recovery goals for both species (Chadès *et al.* 2012). However, Raimondi *et al.* (2015) emphasized the importance of pairing such studies with field-collected data to avoid oversimplifying complex trophic relationships when devising multispecies



management plans. The role of abalone in reducing sea otters' risk of exposure to *T. gondii* adds an additional dimension to their predator–prey interaction and could be incorporated into simulations used to evaluate the effectiveness of wildlife and fisheries management decisions. Because this study indicates that the protective effect of an abalone-rich diet is more likely due to their size and caloric content, rather than a species-specific inability to concentrate *T. gondii* oocysts, supporting stocks of abalone as well as other large, high-value prey items may reduce the proportion of sea otters relying on low-quality prey animals, such as snails.

Given that even low-risk prey may harbour oocysts, results suggest that focusing on the underlying issue of coastal water pollution is vitally important for sea otter conservation efforts. Wetland vegetation has been shown to remove *T. gondii* oocysts from runoff water (Hogan *et al.* 2013; Daniels *et al.* 2014), and coastal development and degradation of estuarine wetlands have been associated with increased transport of *T. gondii* oocysts from land to sea (Shapiro *et al.* 2010b). Management measures aimed at reducing delivery of contaminants such as *T. gondii* oocysts to the marine environment via overland runoff could therefore include protection of estuarine wetland habitats, as well as encouraging sustainable 'green' urban planning in coastal areas that are slated for development (Cook, 2007; Shapiro, 2012). An additional management strategy for reducing environmental contamination with *T. gondii* could include effective education of cat owners regarding proper fecal waste disposal and improved control of feral cat populations (Shapiro, 2012). These measures are especially important because the emergence of *T. gondii* in marine mammals is indicative of broader conservation and public health issues associated with land-to-sea pathogen pollution. Other disease-causing microorganisms, including *Sarcocystis neurona* – another encephalitis-inducing coccidian parasite – and several zoonotic enteric bacterial pathogens, have recently been observed in sea otters (Miller *et al.* 2010a, b). Findings of this study suggest that continued research into disease transmission processes in the ocean can inform implementation of policies aimed at reducing pollution of marine ecosystems with terrestrial pathogens, protecting the health of nearshore habitats and mitigating potential public health threats.

#### SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/S0031182016001359>.

#### ACKNOWLEDGEMENTS

We would like to thank Dr James Gould for his advising and support. We thank David Dann for his logistical

assistance with collection of kelp for study animals. We thank Michelle Staedler, Joe Tomoeoni, Gena Bentall, Jess Fujii, and dozens of other employees and volunteers from the Monterey Bay Aquarium and U.S. Geological Survey for contributing data on sea otter diets and foraging behaviour. Any use of trade, product or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. government.

#### FINANCIAL SUPPORT

Funding for this work was provided by the National Science Foundation (NSF) Ecology of Infectious Disease programme (grant no. OCE-1065990) (to K. S., C. K. and P. A. C.); the Princeton University Office of the Dean of the College (to K. C. S.); U.S. Geological Survey Western Ecological Research Center; and the Department of Ecology and Evolutionary Biology John T. Bonner Senior Thesis Fund (to K. C. S.).

#### REFERENCES

- Arkush, K. D., Miller, M. A., Leutenegger, C. M., Gardner, I. A., Packham, A. E., Heckerroth, A. R., Tenter, A. M., Barr, B. C. and Conrad, P. A. (2003). Molecular and bioassay-based detection of *Toxoplasma gondii* oocyst uptake by mussels (*Mytilus galloprovincialis*). *International Journal for Parasitology* **33**, 1087–1097.
- Allen, B., Callaway, M., Haaker, P., Kalvass, P., Karpov, K., Kashiwada, J., Laueremann, A., Moore, J., O'Leary, J., O'Reilly, K., Patyten, M., Ramsay, J., Rogers-Bennett, L., Skeen, C., Taniguchi, I., Tillman, T., Watters, D. and Wine, V. (2005). *Abalone Recovery and Management Plan*. Final Report to the California Fish and Game Commission. California Department of Fish and Game Marine Region Report. 363 pages. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=29494&inline>.
- Chadès, I., Curtis, J. M. R. and Martin, T. G. (2012). Setting realistic recovery targets for two interacting endangered species, sea otter and northern abalone. *Conservation Biology* **26**, 1016–1025.
- Conrad, P. A., Miller, M. A., Kreuder, C., James, E. R., Mazet, J., Dabritz, H., Jessup, D. A., Gulland, F. and Grigg, M. E. (2005). Transmission of *Toxoplasma*: clues from the study of sea otters as sentinels of *Toxoplasma gondii* flow into the marine environment. *International Journal for Parasitology* **35**, 1155–1168.
- Cook, E. A. (2007). Green site design: strategies for storm water management. *Journal of Green Building* **2**, 46–56.
- Crosson, L. M., Wight, N., VanBlaricom, G. R., Kiryu, I., Moore, J. D. and Friedman, C. S. (2014). Abalone withering syndrome: distribution, impacts, current diagnostic methods and new findings. *Diseases of Aquatic Organisms* **108**, 261–270.
- Daniels, M. E., Hogan, J., Smith, W. A., Oates, S. C., Miller, M. A., Hardin, D., Shapiro, K., Huertos, M. L., Conrad, P. A., Dominik, C. and Watson, F. G. R. (2014). Estimating environmental conditions affecting protozoal pathogen removal in surface water wetland systems using a multi-scale, model-based approach. *Science of the Total Environment* **493**, 1036–1046.
- Dubey, J. P. (1996). Pathogenicity and infectivity of *Toxoplasma gondii* oocysts for rats. *Journal of Parasitology* **82**, 951–956.
- Dubey, J. P. (1998). Advances in the life cycle of *Toxoplasma gondii*. *International Journal for Parasitology* **28**, 1019–1024.
- Dubey, J. P. and Frenkel, J. K. (1972). Cyst-induced toxoplasmosis in cats. *Journal of Protozoology* **19**, 155–177.
- Dubey, J. P. and Jones, J. L. (2008). *Toxoplasma gondii* infection in humans and animals in the United States. *International Journal for Parasitology* **38**, 1257–1278.
- Dubey, J. P., Lunney, J. K. and Shen, S. K. (1996). Infectivity of low numbers of *Toxoplasma gondii* oocysts to pigs. *Journal of Parasitology* **82**, 438–443.
- Dubey, J. P., Zarnke, R., Thomas, N. J., Wong, S. K., Van Bonn, W., Briggs, M., Davis, J. W., Ewing, R., Mense, M., Kwok, O. C. H., Romand, S. and Thulliez, P. (2003). *Toxoplasma gondii*, *Neospora caninum*, *Sarcocystis neurona*, and *Sarcocystis canis*-like infections in marine mammals. *Veterinary Parasitology* **116**, 275–296.
- Estes, J. A. and Palmisano, J. F. (1974). Sea otters: their role in structuring nearshore communities. *Science* **185**, 1058–1060.

- Flegel, J.** (2007). Effects of *Toxoplasma* on human behavior. *Schizophrenia Bulletin* **33**, 757–760.
- Garcia-Esquivel, Z. and Felbeck, H.** (2009). Comparative performance of juvenile red abalone, *Haliotis rufescens*, reared in laboratory with fresh kelp and balanced diets. *Aquaculture Nutrition* **15**, 209–217.
- Hogan, J. N., Daniels, M. E., Watson, F. G., Oates, S. C., Miller, M. A., Conrad, P. A., Shapiro, K., Hardin, D., Dominik, C., Melli, A., Jessup, D. A. and Miller, W. A.** (2013). Hydrologic and vegetative removal of *Cryptosporidium parvum*, *Giardia lamblia*, and *Toxoplasma gondii* surrogate microspheres in coastal wetlands. *Applied and Environmental Microbiology* **79**, 1859–1865.
- Johnson, C. K., Tinker, M. T., Estes, J. A., Conrad, P. A., Staedler, M., Miller, M. A., Jessup, D. A. and Mazet, J. A. K.** (2009). Prey choice and habitat use drive sea otter pathogen exposure in a resource-limited coastal system. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 2242–2247.
- Jones, J. L., Dargelas, V., Roberts, J., Press, C., Remington, J. S. and Montoya, J. G.** (2009). Risk factors for *Toxoplasma gondii* infection in the United States. *Clinical Infectious Diseases* **49**, 878–884.
- Kreuder, C., Miller, M. A., Jessup, D. A., Lowenstine, L. J., Harris, M. D., Ames, J. A., Carpenter, T. E., Conrad, P. A. and Mazet, J. A. K.** (2003). Patterns of mortality in southern sea otters (*Enhydra lutris nereis*) from 1998–2001. *Journal of Wildlife Diseases* **39**, 495–509.
- Krusor, C., Smith, W. A., Tinker, M. T., Silver, M., Conrad, P. A. and Shapiro, K.** (2015). Concentration and retention of *Toxoplasma gondii* oocysts by marine snails demonstrate a novel mechanism for transmission of terrestrial zoonotic pathogens in coastal ecosystems. *Environmental Microbiology* **17**, 4527–4537.
- Lindsay, D. S., Collins, M. V., Mitchell, S. M., Wetch, C. N., Rosypal, A. C., Flick, G. J., Zajac, A. M., Lindquist, A. and Dubey, J. P.** (2004). Survival of *Toxoplasma gondii* oocysts in Eastern oysters (*Crassostrea virginica*). *Journal of Parasitology* **90**, 1054–1057.
- Lowry, M. L. F. and Pears, J. S.** (1973). Abalones and sea urchins in an area inhabited by sea otters. *Marine Biology* **9**, 213–219.
- Luft, B. J., Hafner, R., Korzun, A. H., Lepore, C., Antoniskis, D., Bosler, E. M., Bourland, D. D., III, Uttamchandani, R., Fuhrer, J., Jacobson, J., Morlat, P., Vilde, J., Remington, J. S. and Members of the ACTG 077p/ANRS 009 Study Team** (1993). Toxoplasma encephalitis in patients with the acquired immunodeficiency syndrome. *New England Journal of Medicine* **329**, 995–1000.
- Mazzillo, F. F. M., Shapiro, K. and Silver, M. W.** (2013). A new pathogen transmission mechanism in the ocean: the case of sea otter exposure to the land-parasite *Toxoplasma gondii*. *PLoS ONE* **8**, e82477.
- Miller, M. A., Gardner, I. A., Kreuder, C., Paradies, D. M., Worcester, K. R., Jessup, D. A., Dodd, E., Harris, M. D., Ames, J. A., Packham, A. E. and Conrad, P. A.** (2002). Coastal freshwater runoff is a risk factor for *Toxoplasma gondii* infection of southern sea otters (*Enhydra lutris nereis*). *International Journal for Parasitology* **32**, 997–1006.
- Miller, M. A., Miller, W. A., Conrad, P. A., James, E. R., Melli, A. C., Leutenegger, C. M., Dabritz, H. A., Packham, A. E., Paradies, D., Harris, M., Ames, J., Jessup, D. A., Worcester, K., and Grigg, M. E.** (2008). Type X *Toxoplasma gondii* in a wild mussel and terrestrial carnivores from coastal California: new linkages between terrestrial mammals, runoff and toxoplasmosis of sea otters. *International Journal for Parasitology* **38**, 1319–1328.
- Miller, M. A., Byrne, B. A., Jang, S. S., Dodd, E. M., Dorfmeier, E., Harris, M. D., Ames, J., Paradies, D., Worcester, K., Jessup, D. A. and Miller, W. A.** (2010a). Enteric bacterial pathogen detection in southern sea otters (*Enhydra lutris nereis*) is associated with coastal urbanization and freshwater runoff. *Veterinary Research* **41**, 1–13.
- Miller, M. A., Conrad, P. A., Harris, M., Hatfield, B., Langlois, G., Jessup, D. A., Magargal, S. L., Packham, A. E., Toy-Choutka, S., Melli, A. C., Murray, M. A., Gulland, F. M. and Grigg, M. E.** (2010b). A protozoal-associated epizootic impacting marine wildlife: mass-mortality of southern sea otters (*Enhydra lutris nereis*) due to *Sarcocystis neurona* infection. *Veterinary Parasitology* **172**, 183–194.
- Moore, J. D., Robbins, T. T. and Friedman, C. S.** (2000). Withering syndrome in farmed red abalone *Haliotis rufescens*: thermal induction and association with a gastrointestinal rickettsiales-like prokaryote. *Journal of Aquatic Animal Health* **12**, 26–34.
- Oftedal, O., Ralls, K., Tinker, M. T. and Green, A.** (2007). *Nutritional constraints on the southern sea otter in the Monterey Bay National Marine Sanctuary, and a comparison to sea otter populations at San Nicholas Island, California and Glacier Bay, Alaska*. Final Report to the Monterey Bay National Marine Sanctuary and the Marine Mammal Commission. [http://sanctuariesimon.org/regional\\_docs/monitoring\\_projects/100263\\_report.pdf](http://sanctuariesimon.org/regional_docs/monitoring_projects/100263_report.pdf)
- Raimondi, P., Jurgens, L. J. and Tinker, M. T.** (2015). Evaluating potential conservation conflicts between two listed species: sea otters and black abalone. *Ecology* **96**, 3102–3108.
- Schneider, C. A., Rasband, W. S. and Eliceiri, K. W.** (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* **9**, 671–675.
- Shapiro, K.** (2012). Climate and coastal habitat change: a recipe for a dirtier ocean. *Marine Pollution Bulletin* **64**, 1079–1080.
- Shapiro, K., Largier, J., Mazet, J. A. K., Bernt, W., Ell, J. R., Melli, A. C. and Conrad, P. A.** (2009). Surface properties of *Toxoplasma gondii* oocysts and surrogate microspheres. *Applied and Environmental Microbiology* **75**, 1185–1191.
- Shapiro, K., Mazet, J. A. K., Schriewer, A., Wuertz, S., Fritz, H., Miller, W. A., Largier, J. and Conrad, P. A.** (2010a). Detection of *Toxoplasma gondii* oocysts and surrogate microspheres in water using ultrafiltration and capsule filtration. *Water Research* **44**, 893–903.
- Shapiro, K., Conrad, P. A., Mazet, J. A. K., Wallender, W. W., Miller, W. A. and Largier, J. L.** (2010b). Effect of estuarine wetland degradation on transport of *Toxoplasma gondii* surrogates from land to sea. *Applied and Environmental Microbiology* **76**, 6821–6828.
- Shapiro, K., Silver, M. W., Largier, J. L., Conrad, P. A. and Mazet, J. A. K.** (2012). Association of *Toxoplasma gondii* oocysts with fresh, estuarine, and marine macroaggregates. *Limnology and Oceanography* **57**, 449–456.
- Staggs, S. E., Keely, S. P., Ware, M. W., Schable, N., See, M. J., Gregorio, D., Zou, X., Su, C., Dubey, J. P. and Villegas, E. N.** (2015). The development and implementation of a method using blue mussels (*Mytilus* spp.) as biosentinels of *Cryptosporidium* spp. and *Toxoplasma gondii* contamination in marine aquatic environments. *Parasitology Research* **114**, 4655–4667.
- Tenter, A. M., Heckeroth, A. R. and Weiss, L. M.** (2000). *Toxoplasma gondii*: from animals to humans. *International Journal for Parasitology* **30**, 1217–1258.
- Tinker, M. T., Bentall, G. and Estes, J. A.** (2008). Food limitation leads to behavioral diversification and dietary specialization in sea otters. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 560–565.
- Tinker, M. T., Guimarães, P. R., Novak, M., Marquitti, F. M. D., Bodkin, J. L., Staedler, M., Bentall, G. and Estes, J. A.** (2012). Structure and mechanism of diet specialisation: testing models of individual variation in resource use with sea otters. *Ecology Letters* **15**, 475–483.
- Tinker, M. T., Tomoleoni, J., Weitzman, B., Staedler, M., Jessup, D., Murray, M., Miller, M., Burgess, T., Bowen, L., Miles, K., Thometz, N., Tarjan, L., Golson, E., Batac, F., Dodd, E., Berberich, E., Kunz, J., Bentall, G., Fujii, J., Nicholson, T., Newsome, S., Melli, A., LaRoche, N., MacCormick, H., Johnson, A., Henkel, L., Kreuder-Johnson, C. and Conrad, P.** (2013). *Sea otter population biology at Big Sur and Monterey California: investigating the consequences of resource abundance and anthropogenic stressors for sea otter recovery*. Final Report to California Coastal Conservancy and U.S. Fish and Wildlife Service. US Geological Survey Project Report. 242 pages. <https://www.fws.gov/ventura/docs/species/ss0/Big%20Sur%20Monterey%20Sea%20Otter%20Study%20Draft%20Report.pdf>.
- Tutschulte, T. C. and Connell, J. H.** (1988). Feeding behavior and algal food of three species of abalones (*Haliotis*) in southern California. *Marine Ecology Progress Series* **49**, 57–64.
- Watanabe, J. M.** (1984). Food preference, food quality and diets of three herbivorous gastropods (Trochidae: *Tegula*) in a temperate kelp forest habitat. *Oecologia* **62**, 47–52.
- Wendell, F.** (1994). Relationship between sea otter range expansion and red abalone abundance and size distribution in central California. *California Fish and Game* **80**, 45–56.