

## Standard Paper

# Verrucariaceae from Nepal

Alan Orange<sup>1</sup>  and Som G. Chhetri<sup>2</sup>

<sup>1</sup>Honorary Research Fellow, Natural Sciences, Amgueddfa Cymru - National Museum Wales, Cardiff CF10 3NP, Wales, UK and <sup>2</sup>Ratnanagar 6, Bachhauri, Chitwan, Nepal

### Abstract

Twenty-eight species of *Verrucariaceae* are reported from Nepal. One genus and nine species are described as new: *Nesothele* gen. nov., sister to *Staurothele* s. lat., with a crustose to squamulose thallus, hymenial algae and 4–8 colourless muriform ascospores per ascus; *Nesothele glebulosa* sp. nov., resembling *N. rugulosa* but with smaller perithecia; *Thelidium uvidulum* sp. nov., producing a thin thallus with soralia, prominent perithecia, and 1-septate ascospores; *Verrucaria antepotens* sp. nov., having a well-developed thallus with dark-sided areoles, immersed perithecia, and small ascospores 12.5–16.5 µm long; *V. lactea* sp. nov., resembling *V. praetermissa* but with larger ascospores and a strongly deviating ITS sequence; *V. parvipeltata* sp. nov., with brown, basally constricted areoles on an extensive dark prothallus; *V. senta* sp. nov., with a brown cracked thallus and prominent naked perithecia; *Willeya eminens* sp. nov., with prominent perithecia, and differing from *W. protrudens* in its ITS sequence; *W. irrigata* sp. nov. with a thallus cracked into dark-sided areoles, and relatively large ascospores 28.5–40 µm long; *W. nepalensis* sp. nov. with a cracked thallus, immersed perithecia and a distinctive ITS sequence. Three species complexes might contain new taxa, but wider geographical sampling is necessary before delimiting species: *Thelidium minutulum*, *Verrucaria elaeomelaena* and *V. hydrophila*. Five new combinations are made: *Willeya honghensis* comb. nov. (for *Staurothele honghensis*), *Nesothele globosa* comb. nov. (for *Endocarpon globosum*), *N. hymenogonia* comb. nov. (for *Staurothele hymenogonia*), *N. rugulosa* comb. nov. (for *Staurothele rugulosa*) and *N. succedens* comb. nov. (for *Staurothele succedens*). Four species are unidentified.

**Key words:** floristics, ITS, lichens, mtSSU, taxonomy

(Accepted 22 April 2022)

### Introduction

The family *Verrucariaceae* comprises *c.* 43 genera and 943 species (Lücking *et al.* 2016), the vast majority of which are lichen-forming. Important characteristics defining the family include the perithecioid ascomata, with periphyses and periphysoids but no interascal filaments, and the hemiamyloid hymenium. A wide range of growth forms is exhibited, including crustose, squamulose and foliose (Gueidan *et al.* 2009). In two species the form is determined by the blade-like macroscopic growth form of the alga, which in these species is little altered by lichenization (Pérez-Ortega *et al.* 2010, 2018). Photobiont diversity in the family is high (Thüs *et al.* 2011). The family occurs worldwide, and the species inhabit a wide range of substrata, including siliceous rock and limestone (on the seashore and in freshwater), soil, and less frequently bark. The traditional generic delimitation relied heavily on spore septation and thallus growth form. Gueidan *et al.* (2007, 2009) used molecular data from 83 taxa to propose a more natural classification but the phylogenetic position of many taxa is still unknown, including most species of the largest genus in the family, *Verrucaria*. There have been recent revisions of substantial species groups (e.g. Prieto *et al.* 2010, 2012; Savić & Tibell 2012), but no further attempts to produce a revised phylogeny for the whole family.

The crustose members of the family, especially in the genus *Verrucaria*, have a relatively simple morphology. This, combined with plasticity in response to environmental conditions, often makes difficult the identification and delimitation of the species using morphological characteristics alone. Despite these difficulties, or more likely because of them, a large number of species of currently uncertain status have been described. For instance, A. Zahlbruckner described 84 species, of which perhaps only 37 are currently accepted; H. Zschacke described 133 species, of which perhaps only 50 are currently accepted; M. Servit described 424 species, of which perhaps only 60 are currently accepted (unpublished data). Some recent molecular studies have revealed species which are difficult or impossible to separate using morphological characteristics alone (Savić & Tibell 2009; Orange 2012, 2020; Thüs *et al.* 2015). These issues provide a serious challenge to taxonomic and floristic work.

The lichens of Nepal are still underexplored. A provisional checklist contained 792 taxa (Olley 2011), but it is estimated that at least 2000 species occur in the country (Olley & Sharma 2013). Before 2022, only 12 species of *Verrucariaceae* had been reported (Table 1); of these, 10 are mostly squamulose or foliose species which are relatively conspicuous or easy to collect.

Crustose *Verrucariaceae*, in particular, are likely to be under-represented in checklists and lichen collections because they are often inconspicuous, mostly grow on rock, and the taxonomy is still poorly known in many regions. In East Asia the family is well studied in Japan but poorly known elsewhere. The recent checklist of Japan (Ohmura & Kashiwadani 2018) lists 77 species,

**Author for correspondence:** Alan Orange. E-mail: [alanbiosurveys@outlook.com](mailto:alanbiosurveys@outlook.com)

**Cite this article:** Orange A and Chhetri SG (2022) *Verrucariaceae* from Nepal. *Lichenologist* 54, 139–174. <https://doi.org/10.1017/S0024282922000160>

**Table 1.** *Verrucariaceae* previously reported from Nepal.

Taxon	Reference
<i>Agonimia tristicula</i>	Olley (2011)
<i>Catapyrenium cinereum</i>	Awasthi (1991), Olley (2011)
<i>C. daedaleum</i>	Awasthi (1991), Olley (2011)
<i>Dermatocarpon miniatum</i>	Sharma (1995), Olley (2011)
<i>D. moulinsii</i>	Sharma (1995)
<i>D. vellereum</i>	Sharma (1995), Olley (2011)
<i>Endocarpon subrosettum</i>	Rai et al. (2016)
<i>Endocarpon</i> sp.	Olley (2011)
<i>Placidiopsis pseudocinerea</i>	Breuss (1996), Olley (2011)
<i>Placidium squamulosum</i>	Awasthi (1991), Olley (2011)
<i>Verrucaria acrotella</i>	Rai et al. (2016)
<i>V. margacea</i>	Rai et al. (2016)

of which 31 have been described from Japan since 1990 by Harada (1991, 1992a, 1993, 1994, 1995a, b, c, 1996a, b, 1998, 2000, 2003, 2012a, b, 2013a, b). There is no checklist for China, but at least 46 taxa of *Verrucariaceae* have been described from this country, most in or before 1940, for example by Zahlbruckner (1930) or Magnusson (1940). In recent years, Harada & Wang (1996, 2006a, b, 2008) described eight species from freshwater habitats in Yunnan. Only a small number of *Verrucariaceae* are reported from, for instance, Bhutan (Aptroot & Feijen 2002), Taiwan (Wang-Yang & Lai 1973), Thailand (Buaruang et al. 2017) and Vietnam (nine species by Aptroot & Sparrus 2006, with four more reported by Gueidan et al. 2014).

Between 2–17 October 2009, specimens of *Verrucariaceae* were collected in mid-altitude regions of Nepal, mostly in the Kaski District of Gandaki Pradesh, with a small number in the Kathmandu District of Bagmati Pradesh. In this short period c. 28 species were collected, suggesting that Nepal will be a rich source of *Verrucariaceae*. Two collections made in 2007 by an expedition from the Royal Botanic Garden Edinburgh are also treated below.

## Materials and Methods

### Morphology

Descriptions are of the collected material from Nepal, unless otherwise stated. Sections were cut by hand. In the descriptions, the number of spores measured and the number of specimens from which measurements were taken are given in square brackets. In muriform ascospores, the degree of septation is expressed both as the number of cells visible in optical section, and as the number of cells occurring at the apparent periphery of the spore in optical section, which is easier to determine. Where there are sufficient numbers of measurements, spore sizes are cited as: (minimum–) mean minus one standard deviation – mean – mean plus one standard deviation (–maximum).

### Sequence acquisition

DNA was extracted from recently collected or frozen specimens, using the Qiagen DNeasy Plant Mini Kit; the manufacturer's instructions were followed except that warm water was used for

the final elution. PCR amplification was carried out using Bioneer AccuPower PCR Premix 50 µl reaction tubes. The two internal transcribed spacer regions and the 5.8S region (ITS1-5.8S-ITS2) of the nuclear ribosomal genes, and sometimes part of the 28S ribosomal gene (LSU) and part of the small sub-unit of the mitochondrial ribosomal DNA (mtSSU), were amplified, using the primers ITS1F (Gardes & Bruns 1993), ITS4 (White et al. 1990), nu-LSU-155-5' (Döring et al. 2000), LR3, LR5, LR7 (White et al. 1990), mrSSU1 and mrSSU3R (Zoller et al. 1999). The PCR thermal cycling parameters followed Orange (2018) for all gene regions. Sequencing was performed by The Sequencing Service, College of Life Sciences, University of Dundee ([www.dnaseq.co.uk](http://www.dnaseq.co.uk)).

### Phylogenetic analysis

Sequences were assembled in BioEdit 7.0 (Hall 1999) and aligned using PRANK (Löytynoja & Goldman 2010) (<http://wasabiapp.org/software/prank/>) with the online interface at <https://www.ebi.ac.uk/goldman-srv/webprank/>. Gaps in the alignment were coded using FastGap (Borchsenius 2009).

Phylogenetic relationships and support values were investigated using maximum likelihood (ML), as implemented in RaxML (Stamatakis 2006; Stamatakis et al. 2008), hosted on the CIPRES Science Gateway (Miller et al. 2010). Analyses with RAXML used rapid bootstrapping with 1000 iterations and the GTRGAMMA substitution model; a search for the best-scoring ML tree was carried out with the bootstrap analysis in a single run. The resulting tree was visualized using MEGA v.4 (Tamura et al. 2007). Support values ≥ 70% ML bootstrapping were regarded as significant. Sequences used in the analyses are shown in Table 2.

Sequences were assigned to groups of putatively related species following initial BLAST searches and broad analyses. Separate analyses were carried out for each of the following groups:

*Endocarpon* group (Gueidan et al. 2009). A clade comprising *Endocarpon*, *Willeya* and a number of species of *Verrucaria* s. lat.

*Staurothele* group (Gueidan et al. 2009). A clade comprising *Catapyrenium*, *Placidiopsis*, *Staurothele* (including the type species *S. clopima* (Wahlenb.) Th. Fries) and a number of species of *Verrucaria* s. lat.

*Thelidium* s. lat. in part. An informal group of species with colourless, 1–3-septate ascospores, including *Thelidium minutulum* Körb., *T. pluvium* Orange, *T. rehmi* Zschacke and *T. zwackhii* (Hepp) A. Massal. (Thüs & Nascimbene 2008). The phylogenetic position of these within the family is unknown.

*Verrucaria elaeomelaena* group (Thüs et al. 2015). A clade comprising *V. elaeomelaena* (A. Massal.) Arnold, *V. alpicola* Zschacke, *V. funckii* (Spreng.) Zahlbr. and *V. humida* Orange. Thüs et al. (2015) showed that it was a sister clade to the *Endocarpon* group.

*Verrucaria hydrophila* group (Pykälä et al. 2018). An informal group comprising *Verrucaria hydrophila* Orange, *V. dolosa* Hepp, *V. lignicola* Zschacke, *V. placida* Orange and *V. tenebrosa* Pykälä et al.

*Verrucaria praetermissa* group. Part of the *Staurothele* group of Gueidan et al. (2009), comprising *Verrucaria praetermissa* (Trevisan) Anzi, *V. devensis* (G. Salisb.) Orange, *V. elaeina* Borrer and *V. lapidicola* Orange.

Some taxa were not closely related to any other sequenced taxon and were not included in detailed analyses.

**Table 2.** *Verrucariaceae* specimens generated for this study or used in the phylogenetic analyses. New sequences are in bold. \* = out group taxa in Fig. 2 (*Herpotrichiellaceae*).

Species	Locality	Voucher	GenBank Accession number		
			ITS-LSU	LSU only	mtSSU
* <i>Capronia pilosella</i>	?	WUC28	DQ826737	DQ823099	FJ225725
* <i>C. semiimmersa</i>	?	WUC244	-	FJ358226	FJ225726
<i>Catapyrenium cinereum</i>	Sweden	Wedin 7152a	-	AY853364	AY853316
<i>C. daedaleum</i>	?	Gueidan 115	-	EF643748	FJ225672
<i>Dermatocarpon minutum</i>	Nepal	Olley L9	<b>OM228840</b>		
<i>D. vellereum</i>	Nepal	Orange & Chhetri 18504 (NMW)	<b>OM228795</b>	-	-
<i>Endocarpon adscendens</i>	Switzerland	Gueidan 671 (DUKE)	KF959777	-	-
<i>E. adsurgens</i>	?	-	KX758063	-	-
<i>E. adsurgens</i>	China	Yang & Huang GS158	KX538741	-	-
<i>E. adsurgens</i>	China	Yang & Huang QH014	KX538742	-	-
<i>E. crystallinum</i>	?	-	HM237332	-	-
<i>E. crystallinum</i>	Greece	Timdal	KY682866	-	-
<i>E. deserticola</i>	China	Zhang Z07090 (HMAS-L)	KX538747	-	-
<i>E. deserticola</i>	China	Zhang Z10010 (HMAS-L)	KX538750	-	-
<i>E. nigromarginatum</i>	China	Yang & Zhang SPT268	KX538752	-	-
<i>E. pallidulum</i>	?	-	MN103141	-	-
<i>E. pallidulum</i>	USA	Joneson 4028 (DUKE)	DQ826735	-	-
<i>E. petrolepideum</i>	?	-	MN103138	-	-
<i>E. petrolepideum</i>	?	-	MN103140	-	-
<i>E. petrolepideum</i>	USA	U-492F (DUKE)	KF959778	-	-
<i>E. psorodeum</i>	?	-	MN103137	-	-
<i>E. psorodeum</i>	?	-	MN103142	-	-
<i>E. psorodeum</i>	Estonia	Gueidan 684 (DUKE)	KF959779	-	-
<i>E. pusillum</i>	?	-	HM237334	-	-
<i>E. pusillum</i>	?	?	MN103139	-	-
<i>E. pusillum</i>	?	Gueidan 470 (DUKE)	JQ927447	-	-
<i>E. pusillum</i>	China	Yang & Zhang SPT294	KX538754	-	-
<i>E. sinense</i>	China	Yang & Zhang GS030	KX538755	-	-
<i>E. sinense</i>	China	Yang & Zhang GS031	KX538756	-	-
<i>E. unifoliatum</i>	China	Zhang SPT10047 (HMAS-L)	KX538759	-	-
<i>E. unifoliatum</i>	China	Zhang Z10020 (HMAS-L)	KX538762	-	-
<i>Endocarpon</i> Species A	Nepal	Orange & Chhetri 18476 (NMW)	<b>OM228776</b>	-	-
<i>Endocarpon</i> Species B	Nepal	Cross AC18	<b>OM228838</b>	<b>OM179487</b>	<b>OM331703</b>
<i>Endocarpon</i> sp.	?	-	JF831049	-	-
<i>Hydropunctaria reitrophila</i>	Nepal	Orange & Chhetri 18530 (NMW)	<b>OM228809</b>	-	-
<i>Nesothele glebulosa</i>	Nepal	Orange & Chhetri 18502 (KATH, NMW)	<b>OM228793</b>	<b>OM179484</b>	<b>OM331699</b>
<i>N. globosa</i>	Nepal	Orange & Chhetri 18498 (KATH, NMW)	<b>OM228790</b>	-	<b>OM331698</b>
<i>N. globosa</i>	Nepal	Orange & Chhetri 18505 (KATH, NMW)	<b>OM228796</b>	-	<b>OM331700</b>
<i>N. hymenogonia</i>	England	Orange 25438 (hb. A. Orange)	<b>OM228836</b>	-	<b>OM331702</b>
<i>N. rugulosa</i>	England	Hill [E393] (NMW)	<b>OM228839</b>	-	<b>OM331704</b>
<i>N. succedens</i>	Wales	Orange 16104 (NMW)	-	-	<b>OM331693</b>

(Continued)

Table 2. (Continued)

Species	Locality	Voucher	GenBank Accession number		
			ITS-LSU	LSU only	mtSSU
<i>N. succedens</i>	Wales	Orange 16748 (NMW)	<b>OM228769</b>	-	<b>OM331695</b>
<i>N. succedens</i>	Germany	Orange 19390 (hb. A. Orange)	<b>OM228833</b>	-	-
<i>Nesothele</i> sp.	Wales	Orange 18403 (hb. A. Orange)	<b>OM228772</b>	-	<b>OM331696</b>
<i>Placidiopsis cinerascens</i>	?	Gueidan 585	-	EF643759	FJ225686
<i>Staurothele areolata</i>	?	Gueidan 3781	-	EF643772	FJ225699
<i>S. clopima</i>	Norway	Orange 19167 (NMW)	<b>OM228832</b>	-	-
<i>S. drummondii</i>	?	Gueidan 831	-	EF643774	FJ225700
<i>S. fissa</i>	Wales	Orange 16310 (NMW)	<b>OM228768</b>	-	<b>OM331694</b>
<i>S. fissa</i>	Sweden	Palice s.n.	-	DQ329028	DQ329003
<i>S. frustulenta</i>	Greenland	Heiðmarsson 2066	KY697133	KY773278	KY773553
<i>S. frustulenta</i>		Heiðmarsson 268	KY697134	KY773279	KY773554
<i>S. pulvinata</i>	Iceland	-	-	KY773276	KY773546
<i>S. pulvinata</i>	USA	RR 3686	-	KY769550	KY769564
<i>S. rufa</i>	Germany	Lumbsch s.n.	-	DQ329029	DQ329004
<i>Thelidium</i> aff. <i>minutulium</i>	Nepal	Orange & Chhetri 18522 (NMW)	<b>OM228804</b>	-	-
<i>T.</i> aff. <i>minutulium</i>	Nepal	Orange & Chhetri 18529 (NMW)	<b>OM228808</b>	-	-
<i>T.</i> aff. <i>minutulium</i>	Nepal	Orange & Chhetri 18539 (KATH)	<b>OM228814</b>	-	-
<i>T.</i> aff. <i>minutulium</i>	Nepal	Orange & Chhetri 18544 (NMW)	<b>OM228819</b>	-	-
<i>T.</i> aff. <i>minutulium</i>	Nepal	Orange & Chhetri 18548 (NMW)	<b>OM228822</b>	-	-
<i>T. minutulum</i>	Wales	Orange 18319 (NMW)	<b>OM228770</b>	-	-
<i>T. minutulum</i>	Wales	Orange 18333 (NMW)	<b>OM228771</b>	-	-
<i>T. minutulum</i>	England	Orange 21643 (NMW)	<b>OM228834</b>	-	-
<i>T. minutulum</i>	Austria	Thüs W1386	EU249477	-	-
<i>T. minutulum</i>	Germany	Thüs W1387	EU249478	-	-
<i>T. papulare</i>	Nepal	Orange & Chhetri 18499 (NMW)	<b>OM228791</b>	-	-
<i>T. papulare</i>	Nepal	Orange & Chhetri 18500 (KATH, NMW)	<b>OM228792</b>	-	-
<i>T. pluvium</i>	Scotland	Orange 15019 (NMW)	FJ645269	-	-
<i>T. pluvium</i>	Austria	Thüs & Kison (SMNS-STU-F-635)	MT146895	-	-
<i>T. rehmii</i>	Wales	Orange 22738 (NMW)	<b>OM228835</b>	-	-
<i>T. rehmii</i>	Germany	Thüs W1646	EU249483	-	-
<i>T. rehmii</i>	Germany	Thüs W1648	EU249485	-	-
<i>T. rehmii</i>	Germany	Thüs W1276	EU249482	-	-
<i>T. uvidulum</i>	Nepal	Orange & Chhetri 18485 (KATH, NMW)	<b>OM228781</b>	-	-
<i>T. uvidulum</i>	Nepal	Orange & Chhetri 18543 (NMW)	<b>OM228818</b>	-	-
<i>T. uvidulum</i>	Nepal	Orange & Chhetri 18545 (KATH, NMW)	<b>OM228820</b>	-	-
<i>T. zwackhii</i>	Wales	Orange 25510 (hb. A. Orange)	<b>OM228837</b>	-	-
<i>T. zwackhii</i>	Germany	Thüs W1660	EU249479	-	-
<i>Thelidium</i> sp.	Sweden	Savic 3157 (UPS)	EU559732	-	-
<i>Verrucaria alpicola</i>	Germany	Orange 20669 (NMW)	KM243183	-	-
<i>V. alpicola</i>	Wales	Orange 17468 (NMW)	KM243179	-	-
<i>V. alpicola</i>	Wales	Orange 17485 (NMW)	KM243180	-	-

(Continued)

Table 2. (Continued)

Species	Locality	Voucher	GenBank Accession number		
			ITS-LSU	LSU only	mtSSU
<i>V. alpicola</i>	Norway	Orange 19315 (NMW)	KM243181	-	-
<i>V. alpicola</i>	Norway	Orange 19317 (NMW)	KM243182	-	-
<i>V. alpicola</i>	Italy	Nascimbene 1654 (hb. Nascimbene)	KM243208	-	-
<i>V. alpicola</i>	Italy	Nascimbene 1995a (hb. Nascimbene)	KM243210	-	-
<i>V. alpicola</i>	Italy	Nascimbene 2129 (hb. Nascimbene)	KM243212	-	-
<i>V. alpicola</i>	Italy	Nascimbene 2130 (BM)	KM243213	-	-
<i>V. alpicola</i>	Italy	Nascimbene 2138a (hb. Nascimbene)	KM243214	-	-
<i>V. alpicola</i>	Italy	Nascimbene 2138a (hb. Nascimbene)	KM243215	-	-
<i>V. alpicola</i>	Italy	Nascimbene 4293 (hb. Nascimbene)	KM243216	-	-
<i>V. alpicola</i>	Switzerland	Thüs W1757 (BM)	KM243247	-	-
<i>V. alpicola</i>	Austria	Türk 34321 (BM)	KM243249	-	-
<i>V. antepotens</i>	Nepal	Orange & Chhetri 18488 (KATH, NMW)	<b>OM228784</b>	-	-
<i>V. antepotens</i>	Nepal	Orange & Chhetri 18492 (NMW)	<b>OM228788</b>	-	-
<i>V. bella</i>	Nepal	Orange & Chhetri 18486 (NMW)	<b>OM228782</b>	-	-
<i>V. bella</i>	Nepal	Orange & Chhetri 18483 (KATH, NMW)	<b>OM228780</b>	-	-
<i>V. bella</i>	Nepal	Orange & Chhetri 18490 (KATH, NMW)	<b>OM228786</b>	-	-
<i>V. bella</i>	Nepal	Orange & Chhetri 18532 (KATH, NMW)	<b>OM228810</b>	-	-
<i>V. caerulea</i>	?	Gueidan 507	-	EF643787	FJ225703
<i>V. caerulea</i>	Wales	Orange 16451 (NMW)	FJ664837	FJ664837	-
<i>V. craterigera</i>	Nepal	Orange & Chhetri 18517 (KATH, NMW)	<b>OM228801</b>	-	<b>OM331701</b>
<i>V. devensis</i>	Wales	Orange 18804 (NMW)	KF819516	KF819516	-
<i>V. devensis</i>	Wales	Orange 18814 (NMW)	KF819517	-	-
<i>V. devensis</i>	Wales	Orange 18869 (NMW)	KF819518	KF819518	KF819526
<i>V. elaeina</i>	England	Orange 16628 (NMW)	FJ664849	-	-
<i>V. elaeina</i>	Germany	Thüs W1269	-	EF105140	EF105146
<i>V. elaeomelaena</i>	Wales	Orange 16295 (NMW)	FJ664824	-	-
<i>V. elaeomelaena</i>	Wales	Orange 16297 (NMW)	FJ664825	-	-
<i>V. elaeomelaena</i>	England	Orange 16614 (NMW)	FJ664826	-	-
<i>V. elaeomelaena</i>	Wales	Orange 16784 (NMW)	FJ664827	-	-
<i>V. elaeomelaena</i>	Estonia	Orange 18087 (NMW)	KM243184	-	-
<i>V. elaeomelaena</i>	Wales	Orange 18854 (NMW)	KM243185	-	-
<i>V. elaeomelaena</i>	Wales	Orange 20491 (NMW)	KM243189	-	-
<i>V. elaeomelaena</i>	Germany	Orange 19371 (NMW)	KM243190	-	-
<i>V. elaeomelaena</i>	England	Orange 20541 (NMW)	KM243191	-	-
<i>V. elaeomelaena</i>	Wales	Orange 17402 (NMW)	KM243192	-	-
<i>V. elaeomelaena</i>	Wales	Orange 20546 (NMW)	KM243193	-	-
<i>V. elaeomelaena</i>	Wales	Orange 20876 (NMW)	KM243194	-	-
<i>V. elaeomelaena</i>	Wales	Orange 19358 (NMW)	KM243196	-	-
<i>V. elaeomelaena</i>	Germany	Thüs W0959 (BM)	KM243232	-	-
<i>V. elaeomelaena</i>	Germany	Thüs W1411 (BM)	KM243240	-	-
<i>V. elaeomelaena</i>	Germany	Thüs W1545 (BM)	KM243241	-	-

(Continued)

Table 2. (Continued)

Species	Locality	Voucher	GenBank Accession number		
			ITS-LSU	LSU only	mtSSU
<i>V. elaeomelaena</i>	Germany	Thüs W1624 (BM)	KM243244	-	-
<i>V. elaeomelaena</i>	Germany	Thüs W1645 (BM)	KM243245	-	-
<i>V. elaeomelaena</i>	Germany	Thüs W1658 (BM)	KM243246	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18497 (KATH, NMW)	<b>OM228789</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18503 (NMW)	<b>OM228794</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18520 (NMW)	<b>OM228803</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18526 (NMW)	<b>OM228806</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18528 (NMW)	<b>OM228807</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18536 (NMW)	<b>OM228812</b>	-	-
<i>V. elaeomelaena</i>	Nepal	Orange & Chhetri 18560 (NMW)	<b>OM228825</b>	<b>OM179486</b>	-
<i>V. funckii</i>	Wales	Orange 16298 (NMW)	FJ664862	-	-
<i>V. funckii</i>	Germany	Thüs W1174 (BM)	KM243237	-	-
<i>V. humida</i>	Poland	Krzewicka 2719a (KRAM)	KM261814	-	-
<i>V. humida</i>	Germany	Orange 21178 (NMW)	KM243203	-	-
<i>V. humida</i>	Wales	Orange 16783 (NMW)	KM243199	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18479 (NMW)	<b>OM228778</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18515 (NMW)	<b>OM228800</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18537 (NMW)	<b>OM228813</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18561 (KATH, NMW)	<b>OM228826</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18562 (NMW)	<b>OM228827</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18563 (NMW)	<b>OM228828</b>	-	-
<i>V. hydrophila</i> s. lat.	Nepal	Orange & Chhetri 18564 (NMW)	<b>OM228829</b>	-	-
<i>V. lactea</i>	Nepal	Orange & Chhetri 18572 (KATH, NMW)	<b>OM228831</b>	-	-
<i>V. lapidicola</i>	England	Orange 16690 (NMW)	FJ664850	-	-
<i>V. lapidicola</i>	Estonia	Orange 18112 (NMW)	KF819512	-	-
<i>V. lapidicola</i>	Wales	Orange 18790 (NMW)	KF819513	-	-
<i>V. luchunensis</i>	Nepal	Orange & Chhetri 18489 (KATH, NMW)	<b>OM228785</b>	-	-
<i>V. luchunensis</i>	Nepal	Orange & Chhetri 18491 (KATH, NMW)	<b>OM228787</b>	-	-
<i>V. luchunensis</i>	Nepal	Orange & Chhetri 18569 (KATH, NMW)	<b>OM228830</b>	-	-
<i>V. macrostoma</i>	Wales	Orange 17560 (NMW)	JX848567	-	-
<i>V. macrostoma</i>	Wales	Orange 17825 (NMW)	JX848568	-	-
<i>V. nigrescens</i>	Estonia	Orange 18097 (NMW)	JX848569	-	-
<i>V. nigrescens</i>	Germany	Thüs W1164 (BM)	KM243235	-	-
<i>V. nigrescens</i>	Germany	Thüs W0934 (BM)	KM243231	-	-
<i>V. parvipeltata</i>	Nepal	Orange & Chhetri 18508 (KATH, NMW)	<b>OM228797</b>	-	-
<i>V. parvipeltata</i>	Nepal	Orange & Chhetri 18512 (NMW)	<b>OM228799</b>	-	-
<i>V. parvipeltata</i>	Nepal	Orange & Chhetri 18555 (KATH, NMW)	<b>OM228824</b>	-	-
<i>V. praetermissa</i>	Wales	Orange 16257 (NMW)	FJ664881	-	-
<i>V. praetermissa</i>	Germany	Orange 19382 (NMW)	KF819524	-	-
<i>V. praetermissa</i>	Germany	Thüs W1196	-	EF105142	EF105158
<i>V. praetermissa</i>	Nepal	Orange & Chhetri 18518 (KATH, NMW)	<b>OM228802</b>	-	-

(Continued)

Table 2. (Continued)

Species	Locality	Voucher	GenBank Accession number		
			ITS-LSU	LSU only	mtSSU
<i>V. praetermissa</i>	Nepal	Orange & Chhetri 18524 (NMW)	<b>OM228805</b>	-	-
<i>V. praetermissa</i>	Wales	Orange 18815 (NMW)	KF819522	-	-
<i>V. praetermissa</i>	Wales	Orange 18853 (NMW)	KF819523	-	-
<i>V. rosula</i>	Wales	Orange 20542 (NMW)	JX848577	-	-
<i>V. senta</i>	Nepal	Orange & Chhetri 18541 (NMW)	<b>OM228816</b>	-	-
<i>V. senta</i>	Nepal	Orange & Chhetri 18542 (KATH, NMW)	<b>OM228817</b>	-	-
<i>V. submersella</i>	Switzerland	Gueidan 726 (DUKE)	KF959776	-	-
<i>V. viridula</i>	France	Gueidan 587b (MARSSJ)	KF959786	-	-
<i>V. viridula</i>	Scotland	Orange 15145 (NMW)	FJ664822	-	-
<i>V. viridula</i>	Ireland	Orange 17893 (NMW)	JX848582	-	-
<i>Verrucaria</i> Species A	Nepal	Orange & Chhetri 18546 (NMW)	<b>OM228821</b>	<b>OM179485</b>	-
<i>Verrucaria</i> sp.	Wales	Orange 17295 (NMW)	FJ664877	-	-
<i>Willeya</i> cf. <i>japonica</i>	Nepal	Orange & Chhetri 18510 (KATH, NMW)	<b>OM228798</b>	-	-
<i>W. diffractella</i>	?	Lendemmer 13548	KM371609	-	-
<i>W. diffractella</i>	?	Lendemmer 6513	KM371604	-	-
<i>W. eminens</i>	Nepal	Orange & Chhetri 18533 (NMW)	<b>OM228811</b>	-	-
<i>W. fusca</i>	Vietnam	Gueidan 1877 (BM)	KF959805	-	-
<i>W. fusca</i>	Vietnam	Gueidan 1912 (BM)	KF959806	-	-
<i>W. honghensis</i>	Nepal	Orange & Chhetri 18487 (NMW)	<b>OM228783</b>	-	-
<i>W. irrigata</i>	Nepal	Orange & Chhetri 18475 (KATH, NMW)	<b>OM228775</b>	-	-
<i>W. irrigata</i>	Nepal	Orange & Chhetri 18540 (KATH, NMW)	<b>OM228815</b>	-	-
<i>W. laevigata</i>	Vietnam	Gueidan 1852 (BM)	KF959807	-	-
<i>W. nepalensis</i>	Nepal	Orange & Chhetri 18472 (NMW)	<b>OM228773</b>	-	-
<i>W. nepalensis</i>	Nepal	Orange & Chhetri 18473 (NMW)	<b>OM228774</b>	-	-
<i>W. nepalensis</i>	Nepal	Orange & Chhetri 18480 (KATH, NMW)	<b>OM228779</b>	-	<b>OM331697</b>
<i>W. pallidipora</i> s. lat.	Vietnam	Gueidan 1908 (BM)	KF959790	-	-
<i>W. pallidipora</i> s. lat.	Vietnam	Gueidan 1926 (BM)	KF959791	-	-
<i>W. pallidipora</i> s. lat.	Vietnam	Gueidan 1927 (BM)	KF959792	-	-
<i>W. pallidipora</i> s. str.	Vietnam	Gueidan 1865 (BM)	KF959789	-	-
<i>W. pallidipora</i> s. str.	Vietnam	Gueidan 1940b (BM)	KF959793	-	-
<i>W. pallidipora</i> s. str.	Vietnam	Gueidan 1941 (BM)	KF959794	-	-
<i>W. pallidipora</i> s. str.	Nepal	Orange & Chhetri 18478 (NMW)	<b>OM228777</b>	-	-
<i>W. pallidipora</i> s. str.	Australia	PMC 2546 (CANB)	KF959797	-	-
<i>W. protrudens</i>	Vietnam	Gueidan 1885 (BM)	KF959798	-	-
<i>W. protrudens</i>	Vietnam	Gueidan 1945 (BM)	KF959802	-	-
<i>W. protrudens</i>	Vietnam	Gueidan 1957b (BM)	KF959804	-	-
<i>Willeya</i> Species A	Nepal	Orange & Chhetri 18554 (KATH, NMW)	<b>OM228823</b>	-	-


## Results

### Morphology

Mature ascospores could not be found in many of the collections; this is possibly due to poor drying conditions during the expedition, which took place in unseasonably damp weather.


### Phylogenetic analysis

**Endocarpon group, ITS analysis.** ITS sequences were newly prepared for 18 specimens and an additional 50 sequences from GenBank were included in the analysis. The tree resulting from the analysis of the ITS1-5.8S-ITS2 region is shown in Fig. 1.



**Fig. 1.** Phylogenetic relationships of the *Endocarpon* group *sensu* Gueidan *et al.* (2009), based on a maximum likelihood (ML) analysis of the ITS region. Thickened lines indicate ML support values  $\geq 70\%$ . Information for all taxa is presented in Table 2. The tree is rooted using *Verrucaria rosula* and *V. submersella*.






**Fig. 2.** Phylogenetic relationships of the *Staurothele* group *sensu* Gueidan *et al.* (2009), based on a maximum likelihood (ML) analysis of concatenated data of the ITS, LSU and mtSSU regions. Information for all taxa is presented in Table 2. Thickened lines indicate ML support values  $\geq 70\%$ . The tree is rooted using *Capronia pilosella* and *C. semiimmersa*.

*Endocarpon* is recovered as a monophyletic group with good support. The two available *Endocarpon* sequences from Nepal are not very closely related to other sequences in the tree. Most species of *Willeya* are recovered as a monophyletic group with good support, but *W. irrigata* is basal to a poorly supported clade comprising the remainder of *Willeya* together with *Endocarpon*, while *W. honghensis* is basal in a well-supported clade otherwise comprising *Endocarpon* species. Of the seven species of *Willeya* collected in Nepal, only one (*W. pallidipora* s. str.) is apparently conspecific with existing sequenced species. *Verrucaria bella* is recovered as basal to *Endocarpon* and *Willeya* species in a well-supported clade, and *Verrucaria senta* is shown to be not closely related to any previously sequenced species.

*Staurothele* group. A phylogenetic analysis of 76 selected *Verrucariaceae* was carried out based on mtSSU sequences (not shown). This recovered many of the clades shown in Gueidan *et al.* (2009), including the *Staurothele* group, but with low support. This group comprises *Staurothele* s. str., members of the *Verrucaria praetermissa* group, *V. caerulea*, and the genera *Catapyrenium* and *Placidiopsis* (Gueidan *et al.* 2009). In addition, a new clade not sampled by Gueidan *et al.* (2009) was recovered in the group. Taxa in the *Staurothele* group were reanalyzed using mtSSU, LSU and ITS. The resulting trees exhibited a similar topology, so the data were concatenated and analyzed together (Fig. 2). *Staurothele* s. str. was recovered with good support, including the type species *S. clopima*. *Staurothele* s. str. was sister



**Fig. 3.** Phylogenetic relationships of the *Thelidium minutulum* group, based on a maximum likelihood (ML) analysis of the ITS region. Thickened lines indicate ML support values  $\geq 70\%$ . Information for all taxa is presented in Table 2. The tree is rooted using *Verrucaria rosula*.

to a well-supported clade of species also possessing muriform ascospores and hymenial algae but differing in the ascospores being colourless rather than (mostly) brown, and asci that are 4–8-spored rather than 2-spored. This clade is described as the new genus *Nesothele* below.


***Thelidium* s. lat. in part, ITS analyses.** Two sequences of *Thelidium papulare* from Nepal are nested within sequences of *T. papulare* from Europe (not shown). Specimens from Nepal which are morphologically similar to *Thelidium minutulum* are recovered as two entities: a single sequence basal to four specimens of *T. minutulum* from Europe, and a clade of four identical sequences sister to a clade comprising *T. pluvium* and *T. rehmii*, but with low support (Fig. 3). European specimens of *T. minutulum* are heterogeneous, occurring in two clades.

***Verrucaria elaeomelaena* group, ITS analysis.** Specimens from Nepal form three well-supported clades (Fig. 4). One is nested within sequences of *V. alpicola* from Europe, the other two within specimens from Europe named as *V. elaeomelaena*. There is great variation within the ITS region of this group of species, with little


morphological variation, and additional gene regions will be necessary to determine species boundaries.

***Verrucaria hydrophila* group, ITS analysis.** An ITS tree was constructed comprising 24 publicly available and 13 unreleased sequences of *V. hydrophila* Orange s. lat., together with *V. dolosa* Hepp, *V. placida* Orange and *V. tenebrosa* Pykälä *et al.* (not shown). Four Nepalese specimens (Orange & Chhetri 18479, 18561, 18562 and 18563) form a well-supported clade with sequences of *Verrucaria hydrophila* s. lat. from Europe; another (Orange & Chhetri 18537) is basal in a well-supported clade comprising *V. hydrophila* s. lat., while another (Orange & Chhetri 18515) is basal to sequences of *Verrucaria dolosa* from Europe. Specimens attributed to *V. hydrophila* in Europe are morphologically uniform but there is great variation in ITS sequence, some of which may be correlated with substratum preference. Additional gene regions are needed to investigate species boundaries in this aggregate.

***Verrucaria praetermissa* group (part of the *Staurothele* group of Gueidan *et al.* 2009).** Two identical ITS sequences from Nepal (Orange & Chhetri 18518 and 18524) are basal to a well-



**Fig. 4.** Phylogenetic relationships of the *Verrucaria elaeomelaena* group, based on a maximum likelihood (ML) analysis of the ITS region. Thickened lines indicate ML support values  $\geq 70\%$ . Information for all taxa is presented in Table 2. The tree is rooted using *Verrucaria nigrescens*.



**Fig. 5.** Phylogenetic relationships of the *Verrucaria praetermissa* group, based on a maximum likelihood (ML) analysis of the ITS region. Thickened lines indicate ML support values  $\geq 70\%$ . Information for all taxa is presented in Table 2. The tree is rooted using *Verrucaria rosula*.

supported clade of four sequences of *Verrucaria praetermissa* from Europe (Fig. 5).

## Taxonomy

### *Dermatocarpon miniatum* (L.) W. Mann

*Thallus* foliose, monophyllous, up to 23 mm diam., attached by a single central holdfast; margin shallowly lobed, sometimes thallus split to holdfast; thallus *c.* 280–400  $\mu\text{m}$  thick. *Upper surface* grey to locally pale brown, pruinose; *lower surface* light brown in centre, dark brown or blackish near margin, slightly undulating, smooth, matt.

*Ascospores* 10.5–13  $\times$  6–6.5  $\mu\text{m}$ .

*Specimen examined. Nepal:* Bagmati Pradesh: Langtang National Park, near Ghoda Tabela, 28°12'18.8"N, 85°29'13.5"E, on boulder, upland temperate mixed broad-leaved forest, 2007, L. Olley L9 (E).

### *Dermatocarpon vellereum* Zschacke

(Fig. 6A & B)

*Thallus* foliose, unifoliate, simple or somewhat lobed, up to 28 mm diam., attached by a single, central holdfast. *Upper surface* grey-brown, thinly to densely pruinose; *lower surface* dark brown, mostly obscured by densely branched rhizinomorphs.

*Specimen examined. Nepal:* Gandaki Pradesh: Kaski District, Himalaya, just south-west of lodge, 28.4845°N, 83.8885°E, alt. 2840 m, on unshaded boulder by path, 2009, Orange & Chhetri 18504 (NMW.C.2012.002.161).

### *Endocarpon* Species A

(Fig. 6C & D)

*Thallus* of squamules attached directly to rock, arising singly as convex buttons, sometimes contiguous, later becoming lobed, but not forming mats; squamules up to 900  $\mu\text{m}$  wide, underside

pale brown. *Pseudocortex* an alga-free zone *c.* 25  $\mu\text{m}$  thick, of isodiametric cells, dilute brown above. *Lower cortex* absent.

*Exciple* 210  $\mu\text{m}$  wide, dilute to dense brown in upper part, colourless to dilute brown below. *Hymenial algal cells* present, 3–3.5  $\times$  2.5–3  $\mu\text{m}$ , 1.2 times as long as wide. *Asci* 2-spored. *Ascospores* colourless, muriform, 24.5–30.5  $\times$  10–15.5  $\mu\text{m}$ .

*Notes.* The single sequence available is not closely related to any sequenced species (Fig. 1).

*Specimen examined. Nepal:* Gandaki Pradesh: Kaski District, 0.5 km south of Ghandruk, 28°22.358'N, 83°48.560'E, alt. 1940 m, on stones at edge of lawn in garden of guest house, 2009, Orange & Chhetri 18476 (NMW).

### *Endocarpon* Species B

*Thallus* squamulose, growing over moss; squamules repeatedly lobed; lobes 1.1–1.5 mm wide, to 300  $\mu\text{m}$  thick, upper side light to mid-brown, slightly glossy, smooth; underside pale buff at margin, eventually blackish; no rhizines seen, but probably attached to moss by rhizohyphae. *Lower cortex* absent. *Medulla* white above, lower medulla becoming yellow, K–.

*Perithecia* numerous, but apparently immature; no spores seen.

*Notes.* An ITS sequence clusters with a sequence of *Endocarpon adscendens*, but it is unlikely that the two are conspecific (Fig. 1).

*Specimen examined. Nepal:* Bagmati Pradesh: Rasuwa District, Kyanjin Birch Wood, 28.20513°N, 85.56161°E, alt. 3810 m, boulder in juniper-*Rhododendron* pasture, large boulder 10 m high, 2007, A. Cross AC18 (E - E00305720).

### *Hydropunctaria rheitrophila* (Zschacke) C. Keller, Gueidan & Thüs

(Fig. 6E & F)


*Thallus* light brownish green, thin, 35–75  $\mu\text{m}$  thick, uncracked, rough with very numerous projecting black punctae *c.* 10–50  $\mu\text{m}$  wide; cortex absent, thallus cells with no air spaces between them; thallus containing discrete, darkly pigmented patches (punctae) *c.* 15–40  $\mu\text{m}$  wide.

*Perithecia* forming low conical-hemispherical mounds 260–335  $\mu\text{m}$  wide, black at apex, covered by thallus below. *Involucrellum* well developed, surrounding the apex of the exciple, upper surface very irregular, grading into the black punctae of the thallus. *Exciple* *c.* 165  $\mu\text{m}$  wide, dark brown. *Ascospores* (8.5–)9.5–10.6–11.5(–12)  $\times$  (5.5–)6–6.5–7  $\mu\text{m}$ , (1.4–)1.5–1.6–1.8 (–1.9) times as long as wide [12/1].

*Ecology and distribution.* One collection on a stone in a shaded stream. Reported from Europe, North America, China (Harada & Wang 2008), Japan (Harada 1996c), Australia (McCarthy 2003) and New Zealand (McCarthy 1991).

*Notes.* The specimen fits the morphological concept of this species but the ITS sequence shows significant differences to material collected in Europe.

*Specimen examined. Nepal:* Gandaki Pradesh: Kaski District, north-west of Ghandruk, south-east of Tadapani, 28.37866°N, 83.78715°E, alt. 2275 m, on stable stones in rivulet beside stream



**Fig. 6.** A & B, *Dermatomocarpon vellereum* (Orange & Chhetri 18504). C & D, *Endocarpon* Species A (Orange & Chhetri 18476). E & F, *Hydropunctaria reitrophila* (Orange & Chhetri 18530). B, section of thallus showing rhizinomorphs. D & F, sections through perithecia. Scales: A = 5 mm; B, C & E = 500  $\mu$ m; D & F = 100  $\mu$ m. In colour online.

in forest, shaded, with *Hildenbrandia* sp., 2009, Orange & Chhetri 18530 (NMW.C.2002.002.167).

#### *Nesothele* Orange gen. nov.

Mycobank No.: MB 842601

Thallus crustose, usually superficial, to squamulose; ascospores hyaline, muriform, 4–8 per ascus; photobiont cells present in the hymenium.

Type species: *Nesothele succedens* (Rehm ex Arnold) Orange.

Thallus endolithic to more usually superficial, grey-green to brown, of lobed goniocyst-like units, or of convex areoles, or of tightly adnate squamules. Photobiont chlorococcoid.

Ascomata perithecioid, sometimes partly immersed in thallus but without a discrete thalline covering, in one species immersed in convex squamules; 300–560  $\mu$ m wide, black; ostiolar region often pale and conspicuous in several species. *Involucrellum* present or indistinct. Hymenial gel I+ red; hamathecium of periphyses; interascal filaments absent. Hymenial algal cells 3–8.5  $\mu$ m long, 1.0–4.5 times as long as wide. Ascus verrucarioid, I–,

4–8-spored. *Ascospores* colourless, ellipsoid to narrowly ellipsoid, 25–59.5 × 12.5–31.5 µm, muriform, without a perispore.

*Conidiomata* unknown.

**Phylogenetic placement.** *Ascomycota*, *Verrucariales*, *Verrucariaceae*.

**Etymology.** From the Classical Greek nouns *Nēsos* (νήσος; island) and *Thele* (θηλή; nipple or teat), for the tendency of some species to have naked perithecia amongst an uneven or discontinuous thallus, both resembling islands; the name is also a reference to the morphologically similar *Staurothele*. The name is feminine.

**Ecology and distribution.** On rock, often where damp or calcareous.

**Notes.** In recent decades, *Verrucariaceae* with algae in the hymenium have usually been placed in one of two genera *Staurothele* for crustose species, and *Endocarpon* for squamulose species. In a family-level multigene analysis, Gueidan *et al.* (2009) showed that some *Staurothele* species, including the type species *S. clopima*, were recovered in a well-supported ‘*Staurothele* group’, distant from *S. immersa* and *S. rupifraga* which were recovered in a well-supported ‘*Thelidium* group’ together with some species currently placed in *Polyblastia* and *Thelidium*. Squamulose species attributed to *Endocarpon* were recovered in a well-supported ‘*Endocarpon* group’ which also contained a number of crustose species currently placed in *Verrucaria*. However, Heiðmarsson *et al.* (2017) showed that the squamulose *Endocarpon pulvinatum* belonged in *Staurothele* s. str. Gueidan *et al.* (2014) showed that further crustose species attributed to *Staurothele* formed a well-supported sister group to *Endocarpon*, for which the generic name *Willeya* was available. *Nesothele* represents another lineage of species with hymenial algae which was not sampled by Gueidan *et al.* (2009).

Crustose species formerly attributed to *Staurothele* s. lat. are now placed in four clades: *Staurothele* s. str. (*Staurothele* group), with 2-spored asci and colourless to usually brown ascospores, thallus epilithic; *Nesothele* (*Staurothele* group), with 4–8-spored asci and hyaline ascospores, thallus endolithic to epilithic, sometimes with discrete areolae, or squamulose; *Staurothele* s. lat. (*Thelidium* group), with 2–8-spored asci and hyaline to brown ascospores, thallus often endolithic; and *Willeya* (*Endocarpon* group), with (4–)8-spored asci and hyaline ascospores, thallus epilithic, perithecia often immersed in the thallus. *Staurothele caesia* (Arnold) Arnold and *S. guestphalica* (J. Lahm ex Körb.) Arnold also belong to the *Thelidium* group, based on BLAST searches and alignments of ITS and mtSSU sequences, but some species of ‘*Staurothele*’ are not yet confidently assigned to a clade and more sampling is necessary to determine the morphological limits of the four clades.

### *Nesothele glebulosa* Orange sp. nov.

Mycobank No.: MB 842604

Thallus lumpy, of convex areoles; perithecia naked, ascospores 29.5–41 µm long. Resembles the European *Nesothele rugulosa* (see below), differing in the smaller perithecia and the ITS and mtSSU sequences.

Type: Nepal, Gandaki Pradesh, Kaski District, shortly north-west of Dobhan, 28.47253°N, 83.8731°E, alt. 2530 m, on unshaded

bedrock in stream, 6 October 2009, Orange & Chhetri 18502 (KATH—holotype; NMW.C.2013.001.189—isotype).

(Fig. 7A & B)

*Thallus* of convex grey-brown to brown areoles 170–235 µm diam., areole surface becoming uneven.

*Perithecia* forming conical-hemispherical mounds 300–360 µm in diam., naked or with a small number of areoles at base, ostiole inconspicuous or visible as a convex white mound 115 µm wide. *Involucrellum* 50–60 µm thick at sides of exciple, appressed to exciple but spreading somewhat at base, dark brown, yellowish brown where dilute, K–, densely pigmented on the outside, cell outlines visible within; ostiolar region with a large unpigmented area when young, dark in older perithecia. *Exciple* 260–275 µm wide, colourless to later brown. *Periphyses* c. 30 µm. *Hymenial algae* very poorly preserved, possibly ellipsoid to rod-shaped, c. 3.5–4 × 2–3 µm. Mature asci not seen. *Ascospores* colourless, muriform, 29.5–41 × 12.5–15 µm [6/1], with 14–16 cells at the periphery in optical section.

**Etymology.** From the Latin *glebulosus* (lumpy), referring to the form of the thallus.

**Ecology and distribution.** Known from one specimen on a rock in a stream at 2530 m altitude.

**Notes.** Resembles the European species *Nesothele rugulosa* (see below) in the areolate thallus but differs in the ITS and mtSSU sequence, in the smaller perithecia and perhaps the more strongly pigmented thallus; however, more material needs to be examined to establish the morphological differences between the two species. A specimen from Great Britain (Orange 18403) is closely related (Fig. 2), but more material is needed to determine whether it is conspecific.

The following four combinations are made into *Nesothele*:


### *Nesothele globosa* (H. Harada & Li S. Wang) Orange comb. nov.

Mycobank No.: MB 842606

*Endocarpon globosum* H. Harada & Li S. Wang, *Lichenologist* 28, 303 (1996); type: China, Yunnan, Deqin County, Meili Snow Mountain (E side), 28°24'N, 98°45'E, 3600 m alt., on non-calcareous stones submerged in a stream, 28 September 1994, H. Harada & L.-S. Wang 14563 (HKAS—holotype; CBM, TNS —isotypes[ *non vidi*]).

(Fig. 7C–E)

*Prothallus* not apparent. *Thallus* arising as circular squamules, becoming shallowly lobed, dispersed, or contiguous in small groups, or confluent to produce a thallus of crustose appearance; squamules more or less plane to gently convex when sterile, to 630 µm wide, tightly adnate or margin free from the substratum; upper surface smooth, light brown, extreme margin black; squamules pseudoparenchymatous throughout, or algal layer with more or less vertically oriented hyphae. *Upper cortex* formed of the swollen apices of these hyphae, 2.5–3.5 µm wide, or sometimes appearing wider than tall, 5–7 × 2.5–4 µm, colourless to brown. *Lower cortex* dark brown;



**Fig. 7.** A & B, *Nesothele glebulosa* (Orange & Chhetri 18502). C–E, *Nesothele globosa* (C & E, Orange & Chhetri 18498; D, Orange & Chhetri 18505). A, C & D, thallus with perithecia. B & E, sections of thallus and perithecia. Scales: A, C & D = 500  $\mu$ m; B = 100  $\mu$ m; E = 300  $\mu$ m. In colour online.

rhizohyphae sparse, brown, 4 µm wide. Fertile squamules convex to strongly convex, 630–900 µm wide.

*Perithecia* one per areole, almost completely immersed in convex areoles, or forming moderate to prominent mounds 380–670 µm wide when measurable, with only the ostiolar region visible as a dark ring and/or a pale spot, very rarely exposed as a conspicuous black ring. *Involucrellum* absent, or present as a slight pigmented thickening of the upper exciple. *Centrum* 360–600 µm wide. *Exciple* 30–40 µm thick, dark brown throughout. *Periphyses* with free ends c. 20–35 µm long. *Hymenial algae* subglobose to shortly rectangular, 3–5 × 2.5–3 µm. *Asci* (4?–)8-spored (4 mature asci seen). *Ascospores* 45–59.5 × 21.5–31.5 µm, 1.6–2.7 times as long as wide [*n* = 8], colourless, muriform, with c. 32–42 cells visible in optical section, 19–27 cells at the periphery.

*Conidiomata* not seen.

**Ecology and distribution.** On stones on a wet slope or beside a stream.

**Notes.** The thallus in this species can comprise distinct squamules with raised margins, or squamules which are tightly adnate to the margin, or confluent adnate squamules giving the appearance of a crustose thallus. The pigmented lower cortex and the presence of rhizohyphae suggest that it is convenient to regard the thallus units as squamules. The two collections differ somewhat in appearance and in ITS sequence, but they are regarded as conspecific until further material can be examined. *Orange & Chhetri* 18505 agrees with the protologue in the large, convex fertile squamules, but in *Orange & Chhetri* 18498 most squamules are sterile and thin. The protologue of *Endocarpon globosum* differs from the Nepalese material in the very dark brown to black squamules and the smaller ascospores, 32–42 µm long. Despite these differences, they are regarded here as conspecific until more material can be examined.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, immediately south-west of Dobhan, 28.4697°N, 83.86805°E, alt. 2510 m, on rock in stream, more or less unshaded, 2009, *Orange & Chhetri* 18498 (KATH, NMW); Kaski District, just south-west of Himalaya, 28.48406°N, 83.88765°E, alt. 2810 m, on stones on wet slope beside stream, 2009, *Orange & Chhetri* 18505 (KATH, NMW.C.2013.001.192).

***Nesothele hymenogonia* (Nyl.) Orange comb. nov.**

Mycobank No.: MB 844555

*Verrucaria hymenogonia* Nyl., *Actes de la Société Linnéenne de Bordeaux* 21, 430 (1856).—*Staurothele hymenogonia* (Nyl.) Th. Fr., *Botaniska Notiser* 1865, 40 (1865).

***Nesothele rugulosa* (A. Massal.) Orange comb. nov.**

Mycobank No.: MB 842607

*Polyblastia rugulosa* A. Massal., *Memorie Lichenografiche*, 139–140, fig. 171 (1855).—*Staurothele rugulosa* (A. Massal.) Arnold, *Verhandl. zool.-bot. Gesellsch. Wien* 47, 389 (1897).

***Nesothele succedens* (Rehm ex Arnold) Orange comb. nov.**

Mycobank No.: MB 842608

*Polyblastia succedens* Rehm, in Arnold, *Flora* 53, 17 (1870).—*Staurothele succedens* (Rehm) Arnold, *Verhandl. zool.-bot. Gesellsch. Wien* 30, 149 (1880).

***Thelidium* aff. *minutulum* Körb.**

(Fig. 8A & B)

*Prothallus* not apparent. *Thallus* very thin, c. 25 µm, translucent, pale green-brown, uncracked, smooth, apparently homogeneous within and not composed of gonocyst-like units.

*Perithecia* prominent, dark brown, 190–290 µm wide, with thalline covering only at base, sometimes appearing thin-walled and with sunken sides when dry; ostiole inconspicuous, visible as a small pale or dark plane dot or papilla. *Involucrellum* thin, appressed to exciple above, diverging slightly near base, or developed only in upper part of perithecium, 24–50 µm thick in upper part. *Centrum* c. 235 µm. *Periphyses* c. 20 µm. *Asci* 8-spored. *Ascospores* hyaline, 1-septate, (20–)21.5–23.0–24.5(–25) × (10–)10.5–11.1–12 µm, (1.9–)2–2.1–2.2 times as long as wide [11/2].

*Conidiomata* not seen.


**Ecology and distribution.** On stones in streamlets or on irrigated rocks in shade. *Thelidium minutulum* is reported from China (Harada & Wang 2006b), Europe and North America.

**Notes.** Important features of this species are the very thin thallus, small prominent perithecia with a scarcely developed involucrellum, and 1-septate ascospores. Morphologically the material resembles *Thelidium minutulum* (Harada & Wang 2006b; Thüs & Nascimbene 2008), but an ITS tree suggests that the material is heterogeneous and is not conspecific with either *Thelidium minutulum* or the rather similar *T. rehmii*. The specimens from Nepal might represent one or two new species but formal description must await more sequenced material. *Thelidium minutulum* also appears to be heterogeneous in Europe.

The occurrence of an involucrellum is difficult to confirm in this and some other small *Thelidium* species. The exciple is a thin, hyaline layer; pigmented tissue outside this could be termed an involucrellum, but where it is closely appressed and does not diverge from the exciple it has been customary to conclude that an involucrellum is absent. In the present species the pigmented layer is variable in extent, either closely appressed to the upper part of the exciple only or extending to the base-level of the exciple and diverging slightly.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, north-west of Ghandruk, south of Tadapani, 28°23.15'N, 83°46.85'E, alt. 2450 m, on stones in stream (c. 1 m wide) in forest, shaded, 2009, *Orange & Chhetri* 18522 (NMW.C.2013.001.183); Kaski District, north-west of Ghandruk, south of Tadapani, 28°23.02'N, 83°46.85'E, alt. 2535 m, on irrigated rocks in ravine in forest, shaded, 2009, *Orange & Chhetri* 18529 (NMW.C.2013.001.181); Kaski District, north-west of Ghandruk, east of Sitkyu, 28°22.705'N, 83°48.02'E, alt. 2140 m, on stone by streamlet in scrub, *Orange & Chhetri* 18539 (KATH); Kaski District, south of Landruk, 0.5 km north of Tolka, 28.35°N, 83.82448°E, alt. 1775 m, on slightly damp rocks by path in scrubby forest, 2009, *Orange & Chhetri* 18544 (NMW); Kaski District, south of Landruk, immediately south of Tolka, alt. 1795 m, 28°20.68'N, 83°49.43'E, on stone in streamlet by path amongst fields and scrubby forest, 2009, *Orange & Chhetri* 18548 (NMW.C.2013.001.182).





**Fig. 8.** A & B, *Thelidium* cf. *minutulum* s. lat. (Orange & Chhetri 18522). C & D, *Thelidium papulare* (Orange & Chhetri 18500). E & F, *Thelidium uvidulum* (Orange & Chhetri 18485). Scales: A = 500  $\mu$ m; B, D & F = 100  $\mu$ m; C = 1 mm; E = 200  $\mu$ m. In colour online.

### *Thelidium papulare* (Fr.) Arnold

(Fig. 8C & D)

*Thallus* well developed, to c. 100  $\mu$ m thick, pale grey-brown, cracks numerous but not delimiting discrete areoles, sides of cracks unpigmented.

*Perithecia* forming moderate projections 530–670  $\mu$ m wide, not covered by thallus, black, apex somewhat flattened or slightly depressed, ostiolar area visible as a brown or pale disc 50–200  $\mu$ m wide. *Involucrellum* well developed, c. 75–100  $\mu$ m thick in upper

two-thirds of the perithecium. *Centrum* c. 300–355  $\mu$ m. *Exciple* hyaline to brown. *Ascospores* transversely 3-septate, with 0–1 longitudinal septum, 36–39.6–43(–48)  $\times$  (11–)14–16.2–18.5(–19.5)  $\mu$ m, 2.1–2.5–2.9(–3.4) times as long as wide [12/2].

**Ecology and distribution.** Two collections were made from more or less unshaded rocks in streams.

**Notes.** *Thelidium papulare* belongs to the *Verrucaria aethiobola* group (Orange 2020) rather than to *Thelidium* s. str., but transfer to another genus must wait for a wider study of *Verrucaria* s. lat.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, immediately south-west of Dobhan, 28.4696°N, 83.868°E, alt. 2510 m, on rock in stream, more or less unshaded, 2009, *Orange & Chhetri* 18499 (NMW.C.2013.001.188); Kaski District, just north-west of Dobhan, 28.47253°N, 83.8731°E, alt. 2530 m, on unshaded bedrock in stream, 2009, *Orange & Chhetri* 18500 (KATH, NMW.C.2013.001.189).

***Thelidium uvidulum* Orange sp. nov.**

Mycobank No.: MB 842609

Thallus thin, with soralia; perithecia small, prominent, with appressed involucrellum, ascospores 1-septate, (13.5–)15.5–17.0–18.5(–19.5)  $\mu\text{m}$  long.

Type: Nepal, Gandaki Pradesh, Kaski District, c. 2 km south-west of Chhomrong, 28.41293°N, 83.80365°E, alt. 2175 m, on stone embedded in damp bank by path in forest, 4 October 2009, *Orange & Chhetri* 18485 (KATH—holotype; NMW.C.2012.002.147—isotype).

(Fig. 8E & F)

*Prothallus* not seen. *Thallus* thin, 40–70  $\mu\text{m}$  thick, light grey-green to dull grey-brown, either very thin and scurfy, or thicker and composed of indistinct subunits c. 80–140  $\mu\text{m}$  diam., forming an uneven crust; cracks absent, or a small number in thicker areas. *Soralia* usually present, 60–180  $\mu\text{m}$  diam., discrete, plane, pale brownish green; *soredia* c. 15–29  $\times$  15–20  $\mu\text{m}$ , but difficult to measure, comprising repeating units.

*Perithecia* immersed only in lower 0.2–0.3 of height, forming moderately projecting mounds 200–300  $\mu\text{m}$  diam., naked or with a few flecks of thallus in lower part; ostiolar region inconspicuous, or a pale dot 20–40  $\mu\text{m}$  diam. *Involucrellum* appressed to exciple above, spreading somewhat in lower part, dark brown, K+ dark green. *Centrum* 185–260  $\mu\text{m}$  diam. *Exciple* dark-pigmented or only slightly browned. *Periphyses* c. 20  $\mu\text{m}$  long. *Asci* 8-spored. *Ascospores* colourless, 1-septate, slightly browned when overmature, (13.5–)15.5–17.0–18.5(–19.5)  $\times$  (6–)6.5–7.2–7.5(–8)  $\mu\text{m}$ , (2.1–)2.2–2.4–2.6(–3.1) times as long as wide [37/3].

*Conidiomata* not seen.

**Etymology.** From the Latin *uvidulus* (damp), referring to the habitat of damp, shady rocks.

**Ecology and distribution.** Three collections from moist rocks in forest.

**Notes.** Important features include the thin thallus, the presence of soralia, prominent perithecia with a well-developed involucrellum, and small 1-septate ascospores. The specimen *Orange & Chhetri* 18543 differs slightly in ITS sequence from the others. There are no strongly similar sequences in a BLAST search and this species is apparently not closely related to the group of species including *Thelidium minutulum*. *Thelidium sinense* (Harada & Wang 2006b) differs in the slightly smaller ascospores and absence of soralia. *Verrucaria craterigera* H. Harada is similar in having small soralia, small perithecia and small and rather narrow ascospores, but the ascospores are slightly larger and are septate only when overmature. The two taxa are not closely related.

**Additional specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, south of Landruk, 0.5 km north of Tolka, 28.35°N, 83.82448°E, alt. 1775 m, on slightly damp rocks by path in scrubby forest, 2009, *Orange & Chhetri* 18543 (NMW.C.2012.002.148), 18545 (NMW.C.2012.002.149).

***Verrucaria antepotens* Orange sp. nov.**

Mycobank No.: MB 842610

Thallus well developed, cracked into dark-sided areoles, growing margin thick; perithecia immersed in thallus, with weak involucrellum; ascospores (12.5–)13.5–14.6–16(–16.5)  $\mu\text{m}$  long.

Type: Nepal, Gandaki Pradesh, Kaski District, north-east of Chhomrong, south-west of Sinuwa, 28.43256°N, 83.83416°E, alt. 2200 m, on irrigated siliceous rocks, unshaded, overgrowing *Willeya honghensis*, 5 October 2009, *Orange & Chhetri* 18488 (KATH—holotype; NMW.C.2012.002.153—isotype).

(Fig. 9A–D)

*Prothallus* not apparent. *Thallus* well developed, 200–500  $\mu\text{m}$  thick, non-gelatinous, pale to mid brown; growing margin of thallus thick, well delimited, often slightly lobed, continuous, but cracks rapidly appearing with age; mature thallus with extensive wide cracks, these often surrounding discrete areoles; areoles plane or with slightly ascending margins, smooth, matt, the sides black; thallus cells in vertical columns, air spaces numerous between cells; upper surface a pseudocortex with brown pigment; lower part of thallus occupied by a thick layer of strongly pigmented cells, which is already present at the young margin of the thallus.

*Perithecia* completely immersed in thallus, apex visible as a brown dot or a black disc 60–160  $\mu\text{m}$  diam. *Involucrellum* weakly developed, comprising ill-defined areas of pigmented cells flanking the upper part of the exciple. *Centrum* 220–260  $\mu\text{m}$  diam. *Exciple* pigmented throughout, dark brown below, partly replaced by dark green pigment in thickened upper part. *Periphyses* c. 30  $\mu\text{m}$  long. *Asci* 8-spored. *Ascospores* (12.5–)13.5–14.6–16(–16.5)  $\times$  (6–)6.5–7.1–7.5(–8)  $\mu\text{m}$ , (1.8–)1.9–2.1–2.2(–2.4) times as long as wide [14/1; in rather poor condition].


*Pycnidia* numerous, immersed in thallus, the wall colourless to brown, adjacent pycnidia sometimes with confluent loculi. *Conidia* rod-shaped, 5.7–6.2  $\times$  1  $\mu\text{m}$ .

**Etymology.** From the Latin *antepotens* (superior in power, strongest), referring to the thallus overgrowing other species.

**Ecology and distribution.** On irrigated or non-irrigated rocks, sometimes overgrowing other crustose *Verrucariaceae*, including *Willeya honghensis*.

**Notes.** Critical features include the thick thallus, the abrupt thallus margin, the immersed perithecia with a weakly developed involucrellum, and the small ascospores. The two specimens are overgrowing other crustose *Verrucariaceae* but there is no evidence that they are parasitic.

*Verrucaria gongshanensis* H. Harada & Li S. Wang (Harada & Wang 2008) sometimes has largely immersed perithecia, but the involucrellum merges more completely with the basal layer, the ascospores are larger (15–20  $\mu\text{m}$  long), and pycnidia have



**Fig. 9.** A–D, *Verrucaria antepotens* (A, C & D, Orange & Chhetri 18488; B, Orange & Chhetri 18492). E & F, *Verrucaria bella* (Orange & Chhetri 18483). G & H, *Verrucaria craterigera* (Orange & Chhetri 18517). Scales: A, B & E = 1 mm; C, D, F & H = 100  $\mu$ m, G = 500  $\mu$ m. In colour online.

not been reported. *Verrucaria luchunensis* (see below) and *V. nipponica* are similar in having a cracked thallus, immersed perithecia and pycnidia, but they differ in the much larger ascospores. BLAST searches do not suggest any closely related species.

**Additional specimen examined.** **Nepal:** Gandaki Pradesh: Kaski District, north-east of Chhomrong, north-west of Sinuwa in direction of Kuldhigar, 28.4433°N, 83.84415°E, alt. 2390 m, on low siliceous bedrock near stream in forest, shaded, 2009, Orange & Chhetri 18492 (NMW.C.2012.002.154).

### *Verrucaria bella* Zahlbr.

In Handel-Mazzetti, *Symbolae Sinicae* 3, 10 (1930); type: China, NW Yunnan, Diabasfelsen der kalttemperierte (subalpine) Stufe im Bache auf dem Nguka-la zwischen Dschungdien und Djitsung, 4125 m, 24 August 1915, Handel-Mazzetti Iter Sinense 1914–1918 no. 7769 (BM M001107027—isotype!).

(Fig. 9E & F)

*Thallus* well developed, 60–200 µm thick, pale brownish grey in shade to mid grey-brown when lightly shaded, non-gelatinous; marginal sterile areas sparsely cracked, fertile areas with numerous cracks, often dividing the thallus into discrete areoles; areoles mostly 160–400 µm wide (sterile areoles), plane or slightly concave, smooth, matt; thallus sometimes developing a dark basal layer; thallus cells often in vertical columns, often with air spaces between cells; upper surface a pseudocortex without pigment (in shade) or brown.

*Perithecia* forming low or moderately projecting mounds *c.* 540–800 µm diam., covered by thallus to apex, or a small area exposed at the perithecial apex; ostiole inconspicuous or appearing as a pale dot 20–40 µm diam. *Involucrellum* well developed, spreading basally and laterally, more weakly pigmented below, merging with dark basal layer of thallus when present. *Centrum* 370–435 µm diam. *Exciple* colourless or brown-pigmented on the outside throughout. *Periphyses* long, *c.* 60–100 µm, and branched. *Ascospores* (23.5–)27–31.7–43 × (12.5–)14–15.0–16(–17) µm, (1.6–)1.8–2.1–2.4(–2.8) times as long as wide [25/2], perispore not seen.

*Pycnidia* numerous, conspicuous, scattered over the thallus, isodiametric to elongated, 80–300 × 80–300 µm, brown, plane or slightly projecting; ostiole brown, locule often invaginated. *Conidia* rod-shaped, 5–6.5 × *c.* 0.8 µm.

**Ecology and distribution.** On siliceous rock; beside streams and occasionally to frequently submerged, or on shaded rock beside a path, at altitudes of 2055–2205 m.

**Notes.** Characteristic features of this species include the well-developed, cracked thallus, perithecia forming projecting mounds, a well-developed and spreading involucrellum, large ascospores, and the presence of numerous pycnidia. The contrast between the large fertile and small sterile areoles is a distinctive feature of some specimens, but it is not clear in others. LSU and ITS analyses suggest that this belongs in the *Endocarpon* group of Gueidan *et al.* (2009) (Fig. 1). The identification as *Verrucaria bella* remains provisional, as the protologue of *V. bella* does not mention pycnidia, although they are conspicuous in the Nepalese specimens, and also suggests that the ascospores are

eventually brownish (Zahlbruckner 1930). *Verrucaria honghensis* H. Harada & Li S. Wang (Harada & Wang 2008) is similar, but it lacks the conspicuous pycnidia of this species and has slightly smaller ascospores. *Verrucaria gongshanensis* H. Harada & Li S. Wang (Harada & Wang 2008) has a brown, cracked thallus but much smaller ascospores, while *V. nipponica* H. Harada (Harada 1992b, 2012b) differs in the involucrellum being weakly developed above and merging with a black basal layer.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, Kyumnu, north of Kyumnu Khola, 28.4088°N, 83.79945°E, alt. 2055 m, on siliceous boulder by path in forest, 2009, Orange & Chhetri 18483 (KATH, NMW.C.2012.002.132); Kaski District, north-east of Chhomrong, near Tilche, 28.43208°N, 83.83268°E, alt. 2100 m, on siliceous rock in dry streamlet, in light shade, 2009, Orange & Chhetri 18486 (NMW.C.2012.002.133); Kaski District, north-east of Chhomrong, north-west of Sinuwa in direction of Kuldhigar, 28.4433°N, 83.84415°E, alt. 2390 m, on low siliceous bedrock near stream in forest, shaded, 2009, Orange & Chhetri 18490 (KATH, NMW.C.2013.001.173); Kaski District, west of Ghandruk, 28.37845°N, 83.79056°E, alt. 2205 m, on rock in small stream in forest, shaded, frequently submerged, 2009, Orange & Chhetri 18532 (KATH, NMW.C.2012.002.134).

### *Verrucaria craterigera* H. Harada

*Lichenology* 10, 119 (2012); type: Japan, Honshu, Chiba-ken, Kimitsu-shi, 2002, Harada 20424 (CBM-FL 15028—holotype [*non vidi*]).

(Fig. 9G & H)

*Prothallus* not seen. *Thallus* thin, 60–105 µm thick, non-gelatinous, pale grey-green to pale grey-brown, surface slightly uneven; uncracked or locally with short cracks; thallus cells more or less irregularly arranged (but not in gonocyst-like units). *Soralia* present, discrete, 140–200 µm diam., plane, pale brownish green, often with slightly raised, ragged rim; *soredia* 16–40 × 16–22 µm, outer layer slightly irregular, but with more or less compact layer of isodiametric cells *c.* 3.7–4.5 µm diam., some cells projecting slightly; free hyphae absent or few and very short.

*Perithecia* 0.3–0.5-immersed in the thallus, forming moderately projecting mounds 200–300 µm diam., black or partly grey, sometimes with a slight covering of thallus at the base, but mostly naked; ostiolar region a pale spot. *Involucrellum* well developed, not much or slightly diverging from exciple below, occupying upper 0.66 or the entire height of the exciple, dark brown, K+ dark green-brown. *Centrum* 165–200 µm diam. *Exciple* colourless or slightly browned at sides and base. *Periphyses* *c.* 15 µm long. *Asci* 8-spored, *c.* 80 × 20 µm. *Ascospores* (16.5–)18.5–20.2–22(–24.5) × (6.5–)7–7.4–8(–8.5) µm, (2.2–)2.5–2.7–3(–3.2) times as long as wide [20/1; poor condition], some spores 1-septate when overmature; perispore not seen.

*Conidiomata* not seen.

**Ecology and distribution.** One collection from rock in forest.

**Notes.** Significant characters include the thin thallus with well-defined soralia, small perithecia, a well-developed involucrellum, and small ascospores. The material from Nepal resembles

the protologue of *Verrucaria craterigera* (Harada 2012b), which was described from a single specimen from Japan, differing only in the slightly thicker thallus and slightly longer ascospores (16–18 × 6–8 µm *vide* Harada). The identification is provisional, and further material collected from Nepal and elsewhere is required. Vegetative propagules are uncommon in *Verrucariaceae*, especially well-defined soralia as occur in this species; these were termed gonocystangia by Harada (2012b). The single specimen is very similar in appearance to *Thelidium uvidulum*, especially in the presence of soralia, but the ascospores are smaller in that species and the ITS sequence differs considerably.

**Specimen examined. Nepal: Gandaki Pradesh:** Kaski District, north-west of Ghandruk, near Tadapani, 28.39333°N, 83.77066°E, alt. 2640 m, on boulder in forest, 2009, *Orange & Chhetri* 18517 (KATH, NMW.C.2012.002.155).

### *Verrucaria elaeomelaena* (A. Massal.) Arnold

(Fig. 10A & B)

*Prothallus* not apparent. *Thallus* very thin, up to 30 µm thick, smooth, uncracked, pale greenish brown in herbarium, green and translucent when fresh and wet.

*Perithecia* forming irregular, shallowly to rather strongly conical projections 220–560 µm wide, black or concolorous with the thallus, mostly covered by thallus to the apex, but the covering may be patchy or very thin and inconspicuous. *Involucrellum* conical, reaching to substratum, *c.* 50 µm thick in upper part of perithecium. *Centrum* *c.* 200–330 µm. *Exciple* *c.* 13 µm thick at sides and base, hyaline or lightly browned, or dilute dull green near the ostiole. *Periphyses* *c.* 20–25 µm long. *Ascospores* (20.5–)23–26.0–29(–29.5) × (9–)10.5–12.0–13.5(–16.5) µm, (1.6–)2–2.2–2.4(–2.5) times as long as wide [34/7; poor condition].

*Conidiomata* not seen.

**Ecology and distribution.** On stones in small streams, rarely on irrigated rocks. This aggregate species was reported from Japan by Harada (2012b) (as *Verrucaria andesiatica*).

**Notes.** The species is treated here as an aggregate due to the unresolved taxonomy in this group (Thüs *et al.* 2015) and the small quantities of material available from Nepal. Important features of the aggregate species include the very thin, uncracked thallus and the conical involucrellum. The Nepalese specimens are recovered in three clades within an ITS analysis of the *V. elaeomelaena* group (Fig. 4). Two specimens (18503 and 18528) are nested within sequences from Europe identified as *V. alpicola* Zschacke, and the relatively large ascospores, 25.5–29.5 µm long, approach this taxon in size. However, specimen 18497 also has relatively large ascospores but it is nested within *V. elaeomelaena* s. lat. Collections 18536 and 18560 are also nested within *V. elaeomelaena* s. lat. and the ascospores are comparatively small, 20.5–24.5 µm long. However, further analysis is not possible due to the small quantity of material and the poor condition of the spores. It will not be possible to resolve this complex using the ITS region alone, and a careful analysis using several gene regions is needed.

**Specimens examined. Nepal: Gandaki Pradesh:** Kaski District, north-east of Bamboo, 28.46646°N, 83.86521°E, alt. 2460 m, on

stone in intermittent streamlet in forest, 2009, *Orange & Chhetri* 18497 (KATH, NMW.C.2013.001.178); Kaski District, just north-west of Dobhan, 28.47306°N, 83.8735°E, alt. 2540 m, on stone in streamlet by path in forest, in light shade, 2009, *Orange & Chhetri* 18503 (NMW.C.2013.001.176); Kaski District, north-west of Ghandruk, south of Tadapani, 28.38366°N, 83.78083°E, alt. 2535 m, on irrigated rocks in ravine in forest, shaded, 2009, *Orange & Chhetri* 18528 (KATH, NMW.C.2013.001.177); Kaski District, north-west of Ghandruk, south of Tadapani, 28.3885°N, 83.77833°E, alt. 2450 m, on stones in stream (*c.* 1 m wide) in forest, shaded, 2009, *Orange & Chhetri* 18520 (NMW.C.2013.001.179); Kaski District, north-west of Ghandruk, south of Tadapani, 28.38518°N, 83.78018°E, alt. 2640 m, on stone in streamlet in forest, shaded, not submerged at present, 2009, *Orange & Chhetri* 18526 (NMW.C.2013.001.180); Kaski District, north-west of Ghandruk, east of Sitkyu, 28.37833°N, 83.80026°E, alt. 2140 m, on stone by streamlet in scrub, 2009, *Orange & Chhetri* 18536 (NMW.C.2012.002.157); Kaski District, south of Pothana, 28.31003°N, 83.8291°E, alt. 1940 m, on stone beside small stream in open forest, shaded, 2009, *Orange & Chhetri* 18560 (NMW.C.2012.002.158).

### *Verrucaria hydrophila* Orange s. lat.

(Fig. 10C–F)


*Thallus* thin, up to 20 µm thick, often extensive, pale grey-green to mid grey-brown, uncracked, without dark basal layer; thallus often bright green and translucent when wet and fresh.

*Perithecia* forming low to moderately projecting conical to conical-hemispherical mounds 300–400 µm diam., covered by thin layer of thallus to apex, or covering sometimes patchy, or lost in upper part; apex of perithecium rounded, the ostiolar region inconspicuous or visible as a pale dot or papilla 20–60 µm wide. *Involucrellum* conical, densely pigmented at upper surface, in lower part weakly pigmented or grading into colourless tissue at base of exciple. *Exciple* 140–200 µm diam., colourless at sides and base. *Periphyses* *c.* 18–22 µm long. *Ascospores* (17.5–)21–24.0–27(–29.5) × (9–)9.5–10.3–11(–11.5) µm, (1.9–)2.1–2.3–2.6(–2.8) times as long as wide [31/4], lacking a perispore.

**Ecology and distribution.** On rocks and stones in or beside shaded streams.

**Notes.** Important characters include the very thin uncracked thallus and the more or less conical involucrellum. *Verrucaria elaeomelaena* is very similar in morphology, though not closely related; it tends to have larger perithecia and spores, but some specimens can be impossible to place without sequencing. *Verrucaria hydrophila* as defined here is probably a species complex, with much variation in the ITS region but very little morphological variation. In Europe, some ITS clades are associated with specimens from non-aquatic rock or bark by streams. The specimens from Nepal occur in at least two clades within the complex. An analysis using several gene regions is needed to resolve this group.

**Specimens examined. Nepal: Gandaki Pradesh:** Kaski District, *c.* 1 km north-west of Ghandruk, Chharnami, 28.3820°N, 83.7987°E, alt. 1975 m, on rock by stream, unshaded, 2009, *Orange & Chhetri* 18479 (NMW.C.2013.001.191); Kaski District, north-west of Ghandruk, east of Sitkyu, 28.37833°N, 83.80026°E, alt. 2140 m, on stone by streamlet in scrub, 2009, *Orange & Chhetri* 18537



**Fig. 10.** A & B, *Verrucaria elaeomelaena* s. lat. (Orange & Chhetri 18503). C–F, *Verrucaria hydrophila* s. lat. (C & D, Orange & Chhetri 18561; E & F, Orange & Chhetri 18479). A, C & E, thallus and perithecia. B, D & F, sections of perithecia. Scales: A, C & E = 500 µm; B, D & F = 100 µm. In colour online.

(NMW.C.2012.002.159); Kaski District, south of Pothana, 28.31003°N, 83.8291°E, alt. 1940 m, on stone in small stream in open forest, frequently submerged, shaded, 2009, Orange & Chhetri 18561 ([KATH, NMW.C.2013.001.175), 18562 (NMW), 18563 ([NMW.C.2013.001.174), 18564 (NMW.C.2012.002.160).

*Verrucaria lactea* Orange sp. nov.

Mycobank No.: MB 842611

Thallus very pale grey, very sparsely cracked; perithecia immersed, involucrellum shallowly conical. Resembles the European *Verrucaria praetermissa* but differs in the larger ascospores and strongly deviating ITS sequence.


Type: Nepal, Bagmati Pradesh, Kathmandu District, Shivapuri, 27.78933°N, 85.37666°E, alt. 1740 m, on shaded rock by stream in forest, probably inundated during high flows, 17 October 2009,

Orange & Chhetri 18572 (KATH—holotype; NMW.C.2013.001.193—isotype).

(Fig. 11A–C)

Thallus crustose, extensive, 35–175 µm thick, very pale grey, smooth, matt; cracks sparse, not delimiting areoles, sides of cracks unpigmented; thallus locally with a black basal layer.

*Perithecia* mostly immersed in the thallus, not projecting, or forming low and indistinct projections in the thallus; apex visible as a black dot up to 90 µm wide, or projecting as a small black, often rough mound up to 220 µm wide; ostiolar region inconspicuous or visible as a plane pale dot up to 65 µm wide. *Involucrellum* shallowly conical, densely pigmented at surface, paler and sometimes colourless below, cells with large oil droplets; involucrella confluent, forming a dark basal layer locally in the thallus. *Exciple* colourless or lightly browned, 180 µm wide.



**Fig. 11.** A–C, *Verrucaria lactea* (Orange & Chhetri 18572). D & E, *Verrucaria lachunensis* (Orange & Chhetri 18569). A & D, thallus. B, C & E, sections of thallus and perithecia. Scales: A & D = 500  $\mu$ m; B, C & E = 100  $\mu$ m. In colour online.

*Ascospores* simple, colourless, (23.5–)26.5–29.1–32(–33)  $\times$  (9–)9.5–10.5–11.5  $\mu$ m, (2.4–)2.6–2.8–3.0(–3.1) times as long as wide [18/1], but measured spores possibly a little immature; gelatinous perispore present, 4–5  $\mu$ m thick in water.

*Conidiomata* not found.

**Etymology.** From the Latin adjective *lacteus* (milk-white), referring to the extensive, whitish thallus.

**Ecology and distribution.** A single collection from shaded rock near a stream.

**Notes.** This species resembles *Verrucaria praetermissa* in the pale thallus with an unpigmented cortex, immersed perithecia and a shallowly conical involucrellum, but the very pale and scarcely cracked thallus would be unusual for that species. The ITS sequence of the holotype differs widely from *V. praetermissa*, and a BLAST search does not return any closely related species. The ascospores of the new species are larger than in the two specimens of *V. praetermissa*

from Nepal (although the latter were in poor condition) and are larger than the size of (16–)18–25(–28)  $\times$  (6.5–)7–10(–10.5)  $\mu$ m given for European material by Orange (2000).

#### *Verrucaria lachunensis* H. Harada & Li S. Wang

*Lichenology* 7, 17 (2008); type: China, Yunnan, 2005, Harada 21401 (CBM-FL-16560—holotype; KUN-L—isotype [*non vidi*]).

(Fig. 11D & E)

*Prothallus* not seen. *Thallus* well developed, thick, 100–370  $\mu$ m, non-gelatinous, very pale brownish grey to pale brown; margin entire, soon becoming cracked, mature thallus extensively cracked, the cracks sometimes completely surrounding discrete areoles; areoles with sides pale or dark brown, the surface smooth, matt; thallus cells irregularly arranged or in indistinct vertical columns, air spaces between cells numerous; upper surface a pseudocortex with or without brown pigment, cells sometimes

broken but not forming an epinecral layer; lower part of thallus with extensive dark brown pigment, forming a dark basal layer, cells containing large oil droplets.

*Perithecia* completely immersed in the thallus, the ostiolar region visible as a pale grey-brown spot or disc 40–160 µm, often lying in a small depression; dark-pigmented tissue not reaching thallus surface. *Involucrellum* irregular in shape, comprising areas of brown-pigmented tissue flanking the upper exciple, and merging below with the dark basal layer of the thallus. *Centrum* 175–285 µm diam. *Exciple* colourless when young, later brown to greenish brown at the sides and base, pigment K+ brown or dark green. *Asci* 8-spored. *Ascospores* (22.5–)24.5–27.1–29.5(–33) × (8–)9–10.9–12.5(–15.5) µm, (1.8–)2.3–2.5–2.8 (–3.0) times as long as wide [30/3], with a well-defined gelatinous perispore 0.8–1.6 µm thick (in K); overmature spores light brown. *Conidiomata* not seen.

**Ecology and distribution.** Three collections on shady rocks near streams, but never inundated.

**Notes.** Important features include the thick, cracked thallus, completely immersed perithecia, large ascospores, and presence of a perispore. The perispore is easily visible as a smooth gelatinous coating in most spores; when thin, it can be mistaken for a thickened spore wall. The exciple is unusual amongst other *Verrucaria* species described here, in that it is more deeply pigmented than the adjacent involucrellum. The green pigment usually present in parts of the exciple can be striking, especially in K, though traces of a similar pigment can occur in other species. The upper exciple is weakly pigmented and the involucrellum is weakly developed, so that no dark pigments are visible in surface view of the thallus. In the field this species could be mistaken for an unrelated, sterile, crustose lichen from another family.

The protologue of *V. luchunensis* H. Harada & Li S. Wang (Harada & Wang 2008) from Yunnan agrees with the present species in the immersed perithecia, weakly developed involucrellum, rather large ascospores, the presence of a perispore and the absence of pycnidia. It differs in the slightly smaller ascospores and in the dark tissue of the exciple reaching the thallus surface. This last feature may simply be absent in the specimens from Nepal because of the shady habitat. Distinctly halonate ascospores, as found in the Nepalese specimens, are uncommon in *Verrucariaceae*. The similarities outlined here are sufficient to provisionally identify the Nepalese material as this species, but confirmation by molecular methods would be desirable. *Verrucaria nipponica* Zahlbruckner grows on periodically inundated rocks in Japan; it has a similar ascospore size but differs in the presence of pycnidia and the consistently pale sides of the areoles (Harada 1992b). In addition, the occurrence of a perispore is not mentioned (Harada 1992b).

BLAST searches do not suggest any closely related species.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, north-east of Chhomrong, north-west of Sinuwa in direction of Kuldhigar, 28.44091°N, 83.8423°E, alt. 2340 m, on boulder in forest, shaded, near a stream but never inundated, 2009, Orange & Chhetri 18489 (KATH, NMW.C.2012.002.150); *ibid.*, 28.4433°N, 83.84415°E, alt. 2390 m, on low siliceous bedrock near stream in forest, shaded, 2009, Orange & Chhetri 18491 (KATH, NMW.C.2012.002.151). Bagmati Pradesh: Kathmandu District, Shivapuri, 27.79216°N, 85.37508°E, alt. 1750 m, shady rocks near

stream in forest, 2009, Orange & Chhetri 18569 (KATH, NMW.C.2012.002.152).

*Verrucaria parvipeltata* Orange sp. nov.

Mycobank No.: MB 842612

Thallus comprising basally constricted areoles distributed on an extensive dark prothallus; perithecia sometimes partly surrounded by areoles, otherwise free of thallus.

Type: Nepal, Gandaki Pradesh, south-east of Tolka, Bhedi Kharka, 28.33616°N, 83.8302°E, alt. 1890 m, on rock in retaining wall of stone platform by path in forest, shaded, 12 October 2009, Orange & Chhetri 18555 (KATH—holotype; NMW.C.2013.001.186—isotype).

(Fig. 12A–D)

*Prothallus* conspicuous, dark brown. *Areoles* arising singly on the prothallus, grey to pale brown, convex, enlarging and generally becoming flattened and shallowly lobed, up to 330–500 µm wide, constricted below, sometimes the surface becoming nodular; areoles paraplectenchymatous throughout; upper surface with a pseudocortex of brown cells, lower cortex absent.

*Perithecia* brown, forming convex mounds 265–400 µm wide, without a thalline covering, sometimes partly surrounded by areoles, otherwise free of thallus. *Involucrellum* conical-hemispherical, reaching down to the substratum, dark brown, K+ dull dark green or dull brownish green. *Centrum* 175–185 µm wide. *Exciple* hyaline. *Ascospores* 20.5–26 × 8.5–10 µm, 2.1–3 times as long as wide [13/2; rather poorly developed].

*Conidiomata* not seen.

**Etymology.** From the Latin *parvus* (small) and *peltatus* (peltate), referring to the basally constricted areoles.

**Ecology and distribution.** On rock, shaded or unshaded.

**Notes.** This is a striking species which will be readily recognized in the field by its extensive dark prothallus incompletely covered by brown areoles of subsquamulose appearance. The perithecia sometimes occur separated from the areoles and arising on the prothallus. BLAST searches of ITS and LSU do not return any closely related species.


**Additional specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, south of Ghandruk, by Chane Kola, 28.36446°N, 83.80403°E, alt. 1710 m, on stone in vegetated block scree, 2009, Orange & Chhetri 18474 (NMW.C.2013.001.187); Kaski District, south-west of Chhomrong, 28.41066°N, 83.797°E, alt. 2200 m, on damp boulder by path in deciduous forest, 2009, Orange & Chhetri 18508 (KATH, NMW.C.2013.001.184); Kaski District, north-west of Ghandruk, north of Chuile, just south of Kyumnu Khola, 28.40773°N, 83.77653°E, alt. 2045 m, on unshaded stone by path, 2009, Orange & Chhetri 18512 (NMW.C.2013.001.185).

*Verrucaria praetermissa* (Trevis.) Anzi

(Fig. 12E & F)

*Prothallus* white, non-fimbriate. *Thallus* well developed, 60–200 µm thick, non-gelatinous, pale grey-green, continuous or usually





**Fig. 12.** A–D, *Verrucaria parvipeltata* (A & B, Orange & Chhetri 18555; C & D, Orange & Chhetri 18512). E & F, *Verrucaria praetermissa* (Orange & Chhetri 18518). G & H, *Verrucaria senta* (Orange & Chhetri 18542). A, E & G, thallus and perithecia. B, D, F & H, sections of thallus and perithecia. C, section of areole. Scales: A & E = 500  $\mu$ m; B, D, F & H = 100  $\mu$ m; C = 50  $\mu$ m; G = 1 mm. In colour online.

with sparse to numerous cracks, these locally extensive but only rarely completely surrounding discrete areoles; cracks pale-sided (except when adjacent to the basal layer); thallus cells irregularly arranged or in very weakly delimited vertical columns; upper surface an unpigmented pseudocortex; a darkly pigmented basal layer locally present, sometimes probably part of a widely spreading involucrellum.

*Perithecia* immersed in the thallus, at most producing very low mounds that are too poorly delimited to measure; perithecium visible at first by grey ostiolar region, later the apex exposed as a black ring or disc 100–200 µm diam.; ostiolar region visible as a pale dot usually 20–40 µm diam., later sometimes concave, to 80 µm. *Involucrellum* well developed, conical, often wide-spreading, densely pigmented at the surface, in lower part weakly pigmented with dark cell outlines visible, or locally colourless near the base of the exciple; involucrella often coalescing and merging with the concolorous basal layer. *Centrum* 170–210 µm wide. *Exciple* colourless. *Periphyses* c. 20 µm long, in a common gel. *Asci* 8-spored. *Ascospores* c. 18–20 × 8–8.5 µm, but only a small number of old or immature spores seen.

*Conidiomata* not seen.

**Ecology and distribution.** Two collections on stones in shaded streams. Reported from Europe, North America, Japan (Harada 2012b), China (Yunnan; Harada & Wang 2008), Australia (McCarthy 1995b) and New Zealand (Malcolm & Galloway 1997).

**Notes.** Significant characteristics include the pale, well-developed thallus, cracks with unpigmented sides, immersed perithecia and conical involucrellum. The collections are very similar in morphology to European specimens, although ascospore size could not be determined accurately in the Nepalese material. However, the two ITS sequences from Nepal form a sister clade to sequences from Europe (Fig. 5). Further related collections should be studied before any new species are described in this group.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, north-west of Ghandruk, south of Tadapani, 28.3885°N, 83.77833°E, alt. 2450 m, on stones in stream (c. 1 m wide) in forest, shaded, 2009, Orange & Chhetri 18518 (KATH, NMW.C.2012.002.145); *ibid.*, 28.3885°N, 83.77866°E, alt. 2570 m, submerged in stream in forest (water pH 8.3, conductivity 223 µS cm<sup>-2</sup>), shaded, 2009, Orange & Chhetri 18524 (NMW.C.2012.002.146).

#### *Verrucaria senta* Orange sp. nov.

Mycobank No.: MB 842613

Thallus brown, cracked into dark-sided areoles; perithecia prominent, naked, ascospores (21.5–)25.5–31(–33) µm long. Differing from superficially similar species with a brown thallus by the ITS sequence and occurrence of pycnidia.

Type: Nepal, Gandaki Pradesh, Kaski District, 1 km south of Landruk, 28.36833°N, 83.826166°E, alt. 1650 m, on siliceous rock on unshaded, occasionally irrigated face, 11 October 2009, Orange & Chhetri 18542 (KATH—holotype; NMW C.2012.002.144—isotype).

(Fig. 12G & H)

*Prothallus* not seen. *Thallus* well developed, 80–220 µm thick, greenish mid brown to dark brown, extensively cracked, the cracks

often delimiting discrete areoles; sterile areoles mostly 140–500 µm wide, fertile areoles mostly 350–900 µm wide, thallus occasionally with a marked division into sterile and fertile areoles, sometimes most areoles fertile and the distinction not obvious; areoles with upper surface matt, uneven, marked with faint lines (at least in lightly shaded specimen) denoting subunits mostly 90–220 µm wide; sides of areoles black; upper surface a pseudocortex with brown pigment; lower part of thallus often with extensive darkly pigmented tissue, which ascends the areole margin.

*Perithecia* mostly 1–3(–6) per fertile areole, forming moderately projecting mounds 240–500 µm diam., black, more or less smooth to rough, often with thin patches of thallus in lower half, but mostly naked; ostiole mostly inconspicuous, sometimes visible as a plane dot 20–40 µm diam. *Involucrellum* well developed, spreading laterally and downwards, merging with the dark basal layer of the thallus. *Centrum* 295–355 µm diam. *Exciple* slightly to strongly pigmented in outer layers. *Periphyses* c. 30–36 µm long. *Asci* 8-spored. *Ascospores* (21.5–)25.5–28.2–31(–33) × (10.5–)12–12.8–13.5(–14.5) µm, (1.8–)1.9–2.2–2.4(–2.7) times as long as wide [30/2; mostly in poor condition].

*Pycnidia* scattered, mostly immersed, apex projecting, dark brown, of irregular outline, 50–120 µm diam.; wall brown throughout. *Conidia* rod-shaped, 4.1–4.5 × c. 0.8 µm.

**Etymology.** From the Latin *sentus* (rough, rugged, uneven), a reference to the appearance of the thallus.

**Ecology and distribution.** Two collections from rocks that are dry, or irrigated only in wet weather.

**Notes.** Important features include the well-developed thallus, black-sided areoles subdivided into (inconspicuous) smaller units, and the projecting naked perithecia. Pycnidia were conspicuous in one specimen but not seen in the other. LSU and ITS analyses suggest that this belongs in the *Endocarpon* group of Gueidan *et al.* (2009) (Fig. 1). This is a relatively nondescript species, superficially similar to a number of others with a brown thallus and medium to large ascospores. *Verrucaria macrostoma* Dufour ex DC. has a brown thallus and similar ascospore size but differs in the larger areoles. *Verrucaria nigrescens* Pers. has smaller ascospores, (17–)19–27(–30) × 8–14 µm, and *V. fusconigrescens* Nyl. also has smaller ascospores (17–)19.5–23.5(–26) µm long. These three species also differ in ITS sequence and in the absence of pycnidia.


**Additional specimen examined.** **Nepal:** Gandaki Pradesh: west-south-west of Ghandruk, north-west of Landruk, east side of Modi Khola, 28.37463°N, 83.82185°E, alt. 1390 m, on rock amongst rice terraces, unshaded, 2009, Orange & Chhetri 18541a (NMW.C.2012.002.143).

#### *Verrucaria* Species A

(Fig. 13A & B)

*Prothallus* not seen. *Thallus* very thin, c. 20–30 µm thick, mid greenish brown, smooth, uncracked, apparently composed of inconspicuous, indistinct, coalescing units c. 100–200 µm wide; upper surface a pseudocortex with brown pigment.

*Perithecia* forming low to moderately projecting mounds 200–460 µm diam., sometimes covered to the apex by the thallus, but



**Fig. 13.** A & B, *Verrucaria* Species A (Orange & Chhetri 18546). C & D, *Willeya honghensis* (Orange & Chhetri 18487). E & F, *Willeya* cf. *japonica* (Orange & Chhetri 18510). G & H, *Willeya* Species A (Orange & Chhetri 18554). A, C, E & G, thallus and perithecia. B, D, F & H, sections of thallus and perithecia. Scales: A, E & G = 500 µm; B, D, F & H = 100 µm; C = 1 mm. In colour online.

usually the thalline cover patchy, no more than 50%. *Involucrellum* well developed, shallowly conical, densely pigmented at the surface, internal parts more weakly pigmented with dark cell outlines visible. *Centrum* 110–250 µm diam. *Exciple* colourless. *Periphyses* c. 15 µm long. *Asci* 8-spored. *Ascospores* (24.5–)27–29.2–31.5(–33.5) × (11–)11.5–12.7–14(–15) µm, (2.1–) 2.2–2.3–2.5(–2.6) times as long as wide [16/1]; perispore not seen. *Conidiomata* not seen.

**Ecology and distribution.** One collection from a stone in a shady stream.

**Notes.** Important features include the very thin, smooth, uncracked thallus, the conical involucrellum and the large ascospores. The thallus is composed of small, obscurely delimited sub-units, most easily seen in heavily shaded parts of the specimen, or on the perithecia, where they contribute to the patchy appearance of the partial thalline cover. This is in contrast to species including the superficially similar *Verrucaria hydrophila*, where the thallus appears to be continuous. However, the difference is slight and needs to be confirmed in additional specimens. It resembles three species described from Europe: *V. hydrophila* and *V. elaeomelaena* also have a very thin thallus and conical involucrellum, and *V. margacea* (Wahlenb.) Wahlenb. agrees in the slightly patchy thallus, conical involucrellum and large ascospores. However, LSU and ITS sequences do not suggest a close relationship with any of these taxa. Due to the sparse material, the single specimen is not formally described here.

**Specimen examined.** **Nepal:** Gandaki Pradesh: south of Landruk, 1 km south-east of Tolka, 28.3405°N, 83.83033°E, alt. 1775 m, on stone in small stream in forest (water pH 7.4, conductivity 34 µS cm<sup>-1</sup>), shaded, 2009, Orange & Chhetri 18546 (NMW.C.2012.002.156).

#### *Willeya* Müll. Arg.

The genus *Willeya* was resurrected by Gueidan *et al.* (2014) for a number of species with pale ascospores formerly placed in *Staurothele* s. str. *Staurothele* s. str. was restricted to those species with two dark brown ascospores per ascus. Twelve species have been accepted in *Willeya*, and further species of *Staurothele* s. lat. are likely to belong there. The thallus is epilithic (partly endolithic in one taxon), cracked and often distinctly areolate; the perithecia are typically immersed in the thallus but may protrude to some extent. There are 2–8 spores per ascus, and in most species ascospores are in the size range 18–32 µm long (rarely to 52 µm). The described species differ in relatively minor features such as thallus colour (which seems to be rather variable even in one species), ascospore size and the degree of protrusion of the perithecia. Pycnidia are reported for only two species. Because of the small amount of material available, the lack of firm morphological differences and the lack of DNA data, the status of some of the earlier described species is uncertain.

#### *Willeya eminens* Orange sp. nov.

Mycobank No.: MB 842614

Perithecia prominent, with an uneven thallus cover. *Willeya protrudens* Gueidan differs in ITS sequence.

Type: Nepal, Gandaki Pradesh, Kaski District, south of Syaui Bajar, 28.34706°N, 83.8036°E, alt. 1135 m, on siliceous stone in

drystone retaining wall by path, 2 October 2009, Orange & Chhetri 18533 (NMW.C.2012.002.137—holotype).

(Fig. 14G & H)

*Thallus* thin, c. 40 µm thick, light brown, with occasional cracks that do not delimit areoles; dark basal layer absent, except near perithecia; epinecral layer absent.

*Perithecia* forming irregularly conical-hemispherical projections 560–670 µm wide, with an uneven cover of thallus below, but with the black apex visible above; ostiolar area forming a pale dot 85–105 µm wide. *Involucrellum* conical. *Centrum* c. 325 µm wide. *Exciple* brown. *Asci* 8-spored. *Ascospores* colourless, 22.5–29 × 9–12.5 µm, 2.1–2.4 times as long as wide [14/1], muriform, c. 17–25 cells visible in optical section. *Hymenial algae* 5–10 × 1.5 µm, oblong to cylindrical.

**Etymology.** From the Latin *eminens* (projecting), referring to the perithecial mounds.

**Ecology and distribution.** One collection from a wall.

**Notes.** The rather large and prominent perithecia distinguish this species from all other *Willeya* species in Nepal, but *W. protrudens* also has projecting perithecia. The ITS tree (Fig. 1) suggests they are not closely related, but more material needs to be collected before the range of variation in each can be established. The thallus of the type specimen grows immediately adjacent to one of *W. nepalensis* (the remainder of that species separated as Orange & Chhetri 18472). The latter differs in the smaller, less prominent perithecia, although without close examination the two thalli might be considered part of the same colony.

#### *Willeya honghensis* (H. Harada & Li S. Wang) Orange comb. nov.


Mycobank No.: MB 842615

*Staurothele honghensis* H. Harada & Li S. Wang, *Lichenology* 5, 15 (2006); type: China, Yunnan, Hekou-xian, Manhao, 22°59'23"N, 103°24'19"E, 13 January 2005, Harada 21262 (CBM-FL-16423—holotype; KUN-L—isotype [*non vidi*]).

(Fig. 13C & D)

*Prothallus* not seen. *Thallus* well developed, 200–230 µm thick, non-gelatinous, mid grey-brown, cracks numerous, often enclosing discrete areoles; upper surface of areoles more or less plane, smooth, matt, with black sides; thallus margin apparently of discrete areoles that rapidly become confluent (but little free margin was seen); thallus cells in weak vertical columns, with numerous air spaces between cells; upper surface a pseudocortex with dull brown pigment; lower parts of thallus often without living algae, with brown-pigmented areas. *Photobiont* cells 4.5–9 × 5–7.5 µm.

*Perithecia* almost completely immersed in the thallus, not forming mounds, the naked apex forming a low black protrusion 70–140 µm diam.; ostiolar region inconspicuous, sometimes visible as a small pale dot. *Involucrellum* rather weakly developed, scarcely diverging from exciple, sometimes merging with pigmented basal layer of thallus. *Centrum* 160–255 µm diam. *Periphyses* c. 25 µm long. *Hymenial algal cells* broadly ellipsoid, 3.3–4.1 × 2.5–3.3 µm, solitary or dividing by a transverse septum. *Asci* not



**Fig. 14.** A & B, *Willeya irrigata* (Orange & Chhetri 18540). C & D, *Willeya pallidipora* (Orange & Chhetri 18478). E & F, *Willeya nepalensis* (Orange & Chhetri 18480). G & H, *Willeya eminens* (Orange & Chhetri 18533). Scales: A, C & E = 500  $\mu$ m; B, D, F & H = 100  $\mu$ m; G = 1 mm. In colour online.

observed. *Ascospores* *c.* 32–38 × 10–13 µm, colourless, muriform (but only a small number of poorly developed spores seen).

*Pycnidia* immersed in thallus, ostiole brown, wall colourless. *Conidia* rod-shaped, 5.3–7.0 × *c.* 1.2 µm.

**Ecology and distribution.** One collection from irrigated siliceous rocks. Also found in China (Yunnan).

**Notes.** Important features of this species include the well-developed, cracked thallus, black-sided areoles, and immersed perithecia with only the extreme apex visible. The specimen from Nepal is identified provisionally as *Willeya honghensis* (Harada & Wang 2006a). In fig. 1G of the protologue of that species, the perithecia are shown as forming rather distinct projecting mounds, though a section shown in fig. 3A suggests this character is not always well developed. Although pycnidia are present in the Nepalese collection, they were not found in the type specimen. The few overmature ascospores in the Nepal collection are larger than in *W. honghensis*. The significance of all these features is difficult to assess since *W. honghensis* was described from only a single collection. The Nepalese specimen also resembles *Staurothele kwapiensis* Zahlbr. (Zahlbruckner 1930) in the short and broad hymenial algae and in the ascospore length but, according to Harada & Wang (2006a), an involucrellum is absent in *S. kwapiensis*. *Staurothele yunnana* H. Harada & Li S. Wang differs in the perithecia forming mounds (Harada & Wang 2006a). These two species probably belong in *Willeya* but as material has not been examined in the course of the present study, the transfer is not made. *Willeya tetraspora* Aptroot (Aptroot 2016) has ascospores (27–)30–32–35(–37) µm long, and it possesses pycnidia, but it grows on non-irrigated limestone.

In the ITS tree (Fig. 1), *W. honghensis* is recovered as basal to a well-supported clade comprising *Endocarpon* species, and not amongst other *Willeya* species, but more gene regions are needed to recover a reliable phylogeny.

**Specimens examined.** **Nepal:** Gandaki Pradesh: Kaski District, north-east of Chhomrong, south-west of Sinuwa, 28.43256°N, 83.83416°E, alt. 2200 m, on irrigated siliceous rocks, unshaded, 2009, Orange & Chhetri 18487 (NMW.C.2012.002.135).

#### *Willeya irrigata* Orange sp. nov.

Mycobank No.: MB 842616

Thallus cracked into black-sided areoles; perithecia immersed, ascospores (28.5–)29.5–36(–40) µm, hymenial algal cells oblong.

Type: Nepal, Gandaki Pradesh, Kaski District, west-south-west of Ghandruk, west side of Modi Khola, 28.374°N, 83.82145°E, alt. 1360 m, on irrigated rock face beside fields, 10 October 2009, Orange & Chhetri 18540 (KATH—holotype).

(Fig. 14A & B)

*Prothallus* not seen. *Thallus* well developed, 195–560 µm thick, non-gelatinous, pale to mid brown; thallus margin entire, soon becoming cracked; the cracks later extensive, and often delimiting discrete areoles, areoles more or less plane, 330–670 µm wide, with black sides; thallus cells in weakly delimited vertical columns, air spaces present between some cells; upper surface a pseudocortex with dull brown pigment; lower part of thallus

often darkly pigmented, sometimes forming a thick dark basal layer, cells containing large oil droplets.

*Perithecia* almost completely immersed in the thallus, apex visible as a black disc or black convex structure 120–300 µm diam. *Involucrellum* fairly well developed, more or less conical, densely pigmented at the surface, more weakly coloured below, merging with dark basal layer of thallus. *Centrum* 150–280 µm diam. *Exciple* weakly pigmented on outer side. *Periphyses* *c.* 15–26 µm long, unbranched, enclosed in a common gel. *Hymenial algal cells* oblong, 3.3–7 × 2.1–2.5 µm, 1.3–3.4 times as long as wide, often dividing by a transverse septum, cells occasionally forming chains of several cells. *Asci* not seen when mature. *Ascospores* colourless, muriform, (28.5–)29.5–32.8–36(–40) × (9–)11.5–12.7–14(–14.5) µm, (2.1–)2.2–2.7–3.3 (–4.0) times as long as wide [12/2; in poor condition].

*Pycnidia* immersed in thallus, visible as a small brown dot. *Conidia* rod-shaped, 7–7.8 × *c.* 1 µm.

**Etymology.** From the Latin *irrigatus* (watered, irrigated), referring to the habitat.

**Ecology and distribution.** Two collections from occasionally irrigated rock faces.

**Notes.** Important features of this species include the well-developed thallus which is extensively cracked into black-sided areoles, the mostly immersed perithecia, oblong hymenial algae and the comparatively large ascospores (most species of *Willeya* have much smaller ascospores). *Willeya tetraspora* has globose hymenial algae. *Staurothele kwapiensis* also has globose hymenial algae and it lacks pycnidia, while *Willeya honghensis* differs in its ellipsoid hymenial algae and ITS sequence.

**Additional specimen examined.** **Nepal:** Gandaki Pradesh: Kaski District, south of Ghandruk, 28.3680°N, 83.8098°E, 2009, Orange & Chhetri 18475 (KATH, NMW.C.2012.002.141).

#### *Willeya cf. japonica* (B. de Lesd.) Gueidan

In Gueidan *et al.*, *Lichenologist* **46**, 529 (2014).—*Staurothele japonica* B. de Lesd., *Bull. Soc. Bot. Fr.* **68**, 494 (1921); type: Japan, Hokkaido, 1904, *Faurie* (KYO—lectotype [non vidi]).

(Fig. 13E & F)

*Prothallus* brown (but little thallus margin present in specimen). *Thallus* well developed, 130–290 µm thick, non-gelatinous, light greenish brown; initially of discrete patches, these soon coalescing; becoming strongly secondarily cracked, the cracks often delimiting discrete areoles; surface of areoles slightly uneven, more or less plane to slightly convex; sides of areoles black; thallus cells in weakly delimited vertical columns, with numerous air spaces between cells; upper surface of thallus a pseudocortex with dilute brown pigment; lower part of thallus with dark pigment, which ascends the sides of the areoles.

*Perithecia* almost completely immersed, not forming mounds, or these very low and indistinct; initially visible only by the pale brown ostiolar region (visible as a pale dot up to 60 µm wide), eventually apex exposed as a more or less plane, black disc up to 200 µm wide, not or scarcely projecting from the thallus. *Involucrellum* flanking the upper exciple, confluent with a dark basal layer of the thallus. *Exciple* dark brown. *Centrum* 190–273 µm diam. *Periphyses* short, *c.* 21 µm long. *Hymenial algal cells*

broadly ellipsoid,  $2.5\text{--}3.3 \times 2.5\text{--}2.9$   $\mu\text{m}$ . *Asci* not seen when mature. *Ascospores* colourless, muriform, *c.*  $21\text{--}24 \times 9\text{--}14$   $\mu\text{m}$  (but only a small number of dead and immature spores seen).

*Conidiomata* not seen.

**Ecology and distribution.** A single collection on rock in forest in light shade.

**Notes.** Important characteristics include the well-developed thallus with dark-sided areoles, the immersed perithecia, a well-developed and more or less conical involucrellum, and ellipsoid hymenial algae. ITS sequences confirm that this species is distinct from *W. honghensis*, from which it differs in the thallus initially being composed of discrete areoles, the more uneven surface, and the absence of pycnidia. This species matches the published descriptions of *Willeya japonica* (Harada 1992a; Gueidan *et al.* 2014: 529) in the brown, areolate thallus with more or less discrete marginal areoles, almost completely immersed perithecia and comparatively short hymenial algae. However, *W. japonica*, which is known only from the type collection, usually has an epinecral layer (Harada 1992a). In the absence of molecular data, it is not possible to be certain that the Nepalese specimen is conspecific.

**Specimen examined. Nepal: Gandaki Pradesh:** Kaski District, south-west of Chhomrong,  $28.41066^\circ\text{N}$ ,  $83.797^\circ\text{E}$ , alt. 2200 m, on rock beside path in deciduous forest, in light shade, 2009, Orange & Chhetri 18510 (KATH, NMW.C.2012.002.139).

#### *Willeya nepalensis* Orange sp. nov.

Mycobank No.: MB 842617

Thallus cracked into dark-sided areoles; perithecia immersed, *ascospores* (19.5–)20–23.5(–25)  $\mu\text{m}$  long, hymenial algal cells broadly ellipsoid.

Type: Nepal, Gandaki Pradesh, Kaski District, *c.* 1 km north-west of Ghandruk, Chharnami,  $28.3844^\circ\text{N}$ ,  $83.7973^\circ\text{E}$ , alt. 1990 m, on level surface of schist stones on and beside path in forest, in light shade, 4 October 2009, Orange & Chhetri 18480 (KATH—holotype; NMW.C.2012.002.136—istotype).

(Fig. 14E & F)

*Prothallus* inconspicuous, or dark brown and fimbriate. *Thallus* well developed, 80–180  $\mu\text{m}$  thick, non-gelatinous, dull greyish brown or greenish brown; margin entire or of coalescing patches, soon secondarily cracked, cracks numerous, often enclosing discrete areoles, sides of cracks black; surface of areoles plane to slightly concave, smooth, matt; thallus cells in weak vertical columns, with numerous air spaces between cells; upper surface a pseudocortex with dull brown pigment, epinecral layer not seen.

*Perithecia* forming low, ill-defined mounds *c.* 300–340  $\mu\text{m}$  diam.; apex of perithecium flattened, exposed only as a black disc, or as a low black mound 100–240  $\mu\text{m}$  diam.; ostiolar region plane or slightly depressed, often rather conspicuous as a pale dot 20–80  $\mu\text{m}$  diam. *Involucrellum* well developed, more or less conical, densely pigmented at the surface, more weakly pigmented below. *Centrum* 195–220  $\mu\text{m}$  diam. *Exciple* externally brown. *Periphyses* *c.* 30–35  $\mu\text{m}$  long. *Hymenial algal cells* broadly ellipsoid,  $3.7\text{--}5.3 \times 3.7\text{--}4$   $\mu\text{m}$ , solitary or divided by a transverse septum. *Asci* 8-spored,  $55\text{--}65 \times 29\text{--}36$   $\mu\text{m}$ . *Ascospores* colourless, muriform, (19.5–)20–21.6–23.5(–25)  $\times$  (9–)10–10.9–12  $\mu\text{m}$ , (1.7–)1.8–2.0–2.2

(–2.3) times as long as wide [18/2], with 10–16 cells visible in optical section, 9–13 cells on margin in optical section.

*Conidiomata* not seen.

**Etymology.** Referring to the country of origin.

**Ecology and distribution.** Three specimens, from drystone retaining walls and flagstones in a path.

**Notes.** This species is characterized by the cracked thallus with dark-sided areoles, the weakly prominent perithecia, a well-developed involucrellum, ellipsoid hymenial algae, and the small *ascospores*. Amongst the species with short hymenial algae and relatively small *ascospores*, *W. microlepis* (Zahlbr.) Gueidan from southern China differs in the 2-spored *asci* (Gueidan *et al.* 2014). *Willeya japonica* differs in the involucrellum appressed to the upper exciple (Gueidan *et al.* 2014) and the frequent occurrence of an epinecral layer (Harada 1992a), although these could prove to be unimportant features once a wider range of material is available for study. However, the two are not synonymized here as although *Willeya nepalensis* is very similar to *W. cf. japonica* (Orange & Chhetri 18510) in morphology, it is distinct in ITS sequence, highlighting the difficulty of identifying species by morphology alone in this genus. *Willeya eminens* (see above) differs in the more prominent perithecia and the oblong hymenial algae.

**Additional specimens examined. Nepal: Gandaki Pradesh:** Kaski District, south of Syaui Bajar,  $28.3378^\circ\text{N}$ ,  $83.8003^\circ\text{E}$ , alt. 1135 m, on siliceous stone in drystone retaining wall by path, 2009, Orange & Chhetri 18472 (NMW.C.2012.002.137); Kaski District, NNW of Syaui Bajar in the direction of Kimche,  $28^\circ20.82'\text{N}$ ,  $83^\circ48.21'\text{E}$ , alt. 1405 m, on rock beside path, 2009, Orange & Chhetri 18473 (NMW).

#### *Willeya pallidipora* (P. M. McCarthy) Gueidan

In Gueidan *et al.*, *Lichenologist* **46**, 530 (2014).—*Staurothele pallidipora* P. M. McCarthy, *Muelleria* **8**, 275 (1995) as ‘pallidipora’; type: Australia, Queensland, P. M. McCarthy 768 (MEL—holotype; BRI—istotype [non vidi]).

(Fig. 14C & D)

*Prothallus* not seen. *Thallus* well developed, 50–195  $\mu\text{m}$  thick, non-gelatinous, pale grey-green; cracks more or less numerous in mature thalli, but not delimiting discrete areoles; thallus cells in vertical columns, with numerous air spaces between cells; upper surface a pseudocortex, pigment absent; with a dark basal layer formed by confluent involucrella.

*Perithecia* almost completely immersed in thallus, not forming mounds, or the mounds very low and poorly delimited; at first only the pale brown ostiolar region visible, later apex visible as a more or less plane black disc up to 180  $\mu\text{m}$  diam.; ostiolar region visible as a pale dot 20–40  $\mu\text{m}$  diam. *Involucrellum* conical, densely pigmented at surface, less densely below; upper surface sometimes rough; cells in lower part of involucrellum with pigment confined to cell wall (cell outlines easily visible in thin section), containing large oil drops. *Centrum* *c.* 260  $\mu\text{m}$  diam. (one measurement). *Periphyses* short, *c.* 20  $\mu\text{m}$ . *Hymenial algal cells* rod-shaped,  $3.7\text{--}6.6 \times 2.1$   $\mu\text{m}$ , 1.8–3.1 times as long as wide. *Asci* 8-spored. *Ascospores* colourless,  $20.5\text{--}21.6\text{--}22.5(–23) \times 9.5\text{--}$

10.3–11 µm, 2.0–2.1–2.2(–2.3) times as long as wide [7/1], muriform, with c. 16–21 cells visible in optical section.

*Pycnidia* not seen.

**Ecology and distribution.** One collection from a rock beside a stream. Reported from Vietnam (Gueidan *et al.* 2014) and Australia (McCarthy 1995a).

**Notes.** Important characters in the specimen from Nepal include the pale thallus (pigment apparently absent even in unshaded situation) which shows few cracks, the immersed perithecia and rod-shaped hymenial algae. The single specimen forms a well-supported clade with three specimens of *W. pallidipora* from Vietnam and two from Australia (Fig. 1). Three additional sequences from Vietnam were named *W. pallidipora* s. lat. by Gueidan *et al.* (2014), and these appear to represent a distinct species. The single specimen from Nepal agrees well with the protologue of *W. pallidipora*, including the cracked and pale but not strongly areolate thallus; the absence of pigment on the sides of the thallus cracks is not mentioned by McCarthy (1995a) but is suggested by his fig. 2B. *Willeya malayensis* (Zahlbr.) Gueidan, *Staurothele paraguayensis* Malme, *Willeya japonica* and *W. microlepis* differ in the globose hymenial algae; *W. rimosa* Müll. Arg. in the projecting perithecia; *W. laevigata* Gueidan and *W. protrudens* differ in ITS sequence (Gueidan *et al.* 2014). *Willeya australis* (Groenh.) Gueidan from Java appears to be very similar in morphology (Gueidan *et al.* 2014).

**Specimen examined.** **Nepal:** Gandaki Pradesh: Kaski District, c. 1 km north-west of Ghandruk, Chharnami stream, 28°22.92'N, 83°47.93'E, alt. 1975 m, on rock by stream, unshaded, 2009, Orange & Chhetri 18478 (NMW.C.2012.002.138).

### Willeya Species A

(Fig. 13G & H)

*Prothallus* not seen (no free thallus margin present). *Thallus* well developed, 100–260 µm thick, non-gelatinous, very pale grey-brown, cracks numerous and extensive but not delimiting discrete areoles; sides of areoles not pigmented except adjacent to the dark basal layer of thallus, which is absent in young areas of thallus, becoming visible in older thallus areas as a result of confluent involucrella; thallus cells in weakly defined vertical columns,

with numerous air spaces between cells; upper surface a non-pigmented pseudocortex; lower parts of thallus usually with a dark basal layer.

*Perithecia* completely immersed in thallus, not forming mounds, at first visible only as a pale brown disc c. 100 µm diam., later the apex sometimes exposed as a plane black disc up to 180 µm diam.; ostiolar region comprising a small pale depression c. 40 µm diam. surrounded by a pale brown circle to 100 µm wide. *Involucrellum* scarcely developed in the upper half of the exciple, hardly spreading, merging with the dark basal layer when present, densely pigmented at the surface, weakly and patchily pigmented below. *Centrum* 180–230 µm diam. *Exciple* colourless or brown on the outside. *Periphyses* c. 18–33 µm long, enclosed in a common gel. *Hymenial algal cells* oblong, 5–9 × 2.5 µm, 2–3.7 times as long as wide, often dividing by a transverse septum, or forming short chains of several cells. *Asci* 8-spored, c. 74 × 26 µm. *Ascospores* immature.

*Conidiomata* not seen.

**Ecology and distribution.** A single collection from shaded siliceous rock.

**Notes.** Important features include the well-developed, pale, extensively cracked but non-areolate thallus, the completely immersed perithecia and the oblong hymenial algae. Unfortunately, the ascospores are immature. It is not possible to identify this with any previously described species; amongst those with oblong to elongate hymenial algae, *Staurothele fauriei* B. de Lesd., *S. paraguayensis*, *Willeya malayensis* and *W. rimosa* differ in the more or less emergent perithecia (Gueidan *et al.* 2014). *Willeya iwatsukii* (H. Harada) Gueidan agrees in the immersed perithecia and the in-part bacilliform hymenial algae (Harada 1992a), but these features are not distinctive enough to allow an identification in the absence of sequence data. *Willeya fusca* Gueidan, *W. laevigata* and *W. pallidipora* differ in ITS sequence (Fig. 1). The type specimen of *W. australis* is similar in the immersed perithecia and elongate hymenial algae but differs in the more strongly areolate, grey-brown thallus and the pigmented sides to the areoles.

**Specimen examined.** **Nepal:** Gandaki Pradesh: Kaski District, south-east of Tolka, Bhedi Kharka, 28.33616°N, 83.8302°E, alt. 1890 m, on small boulder on slope by platform, by path in forest, shaded, 2009, Orange & Chhetri 18554 (NMW.C.2012.002.140).

### Key to Verrucariaceae reported from Nepal

The records of *Verrucaria acrotella* Ach. and *V. margacea* (Wahlenb.) Wahlenb. published by Rai *et al.* (2016) are not included in the key because the descriptions provided are insufficient to characterize these collections.

1	Thallus foliose or squamulose .....	2
	Thallus crustose .....	11
2(1)	Thallus foliose, attached by usually one central holdfast, underside smooth or with rhizinomorphs .....	3
	Thallus squamulose, underside attached by its underside or by rhizohyphae .....	5
3(2)	Lower surface smooth .....	<b>Dermatocarpon miniatum</b>
	Lower surface with rhizinomorphs .....	4
4(3)	Rhizinomorphs coralloid, with dense, congested branches .....	<b>Dermatocarpon vellereum</b>
	Rhizinomorphs simple to weakly branched, pointed or thickened at apex .. .	<b>Dermatocarpon moulinii</b> (Mont.) Zahlbr.



- 5(2) Thallus of small grey-green squamules up to 1 mm long, lacking rhizohyphae; perithecia amongst the squamules, not immersed in thallus; hymenial algae absent; ascospores muriform, (42–)57–120(–150)  $\mu\text{m}$  long . . . . . **Agonimia tristicula**  
Thallus of squamules that are sometimes attached by rhizohyphae; perithecia immersed in the thallus . . . . . 6
- 6(5) Ascospores muriform, hymenial algae present . . . . . 7  
Ascospores simple or 1-septate, hymenial algae absent . . . . . 8
- 7(6) Ascospores 45–59  $\mu\text{m}$  long, (4?–)8 per ascus . . . . . **Nesothele globosa**  
Ascospores 24–30  $\mu\text{m}$  long, 2 per ascus . . . . . **Endocarpon** spp.
- 8(6) Ascospores 1-septate . . . . . **Placidiopsis pseudocinerea** Breuss  
Ascospores simple . . . . . 9
- 9(8) Rhizohyphae colourless . . . . . **Placidium squamulosum** (Ach.) Breuss  
Rhizohyphae dark . . . . . 10
- 10(9) Lower cortex present, paraplectenchymatous, blackish; with blackish rhizohyphae; squamules usually white-pruinose . . . . .  
. . . . . **Catapyrenium cinereum** (Pers.) Körb.  
Lower cortex absent, medulla merging into dark rhizohyphae; squamules not or weakly pruinose . . . . .  
. . . . . **Catapyrenium daedaleum** (Kremp.) Stein.
- 11(1) Ascospores 9.5–11.5  $\mu\text{m}$  long, simple; thallus with embedded black punctae that can be visible at the surface . . . . .  
. . . . . **Hydropunctaria rheitrophila**  
Ascospores longer; thallus without punctae . . . . . 12
- 12(11) Ascospores simple; hymenial algae absent . . . . . 13  
Ascospores septate or muriform (or ascospores apparently simple, but hymenial algae present) . . . . . 21
- 13(12) Thallus with small soralia; thallus thin (to 105  $\mu\text{m}$ ) with few or no cracks; perithecia forming moderately projecting mounds  
200–300  $\mu\text{m}$  diam., mostly without a thalline covering; ascospores 18.5–24.5  $\mu\text{m}$  long . . . . . **Verrucaria craterigera**  
Thallus without soralia . . . . . 14
- 14(13) Thallus of areoles arising singly on a dark prothallus, later discrete or contiguous, narrowed at base; perithecia forming convex  
mounds without a thalline covering, but sometimes surrounded by areoles . . . . . **Verrucaria parvipeltata**  
Thallus continuous or cracked, areoles not narrowed at base . . . . . 15
- 15(14) Thallus brown, divided into areoles with an uneven surface; perithecia forming moderately projecting mounds 240–500  $\mu\text{m}$   
diam., at most with thin patches of thallus in lower half, mostly naked . . . . . **Verrucaria senta**  
Thallus various; perithecia immersed, or forming projecting mounds with a thalline covering to near apex . . . . . 16
- 16(15) Thallus very thin, to 30  $\mu\text{m}$  thick, sometimes translucent when fresh and wet, uncracked; perithecia forming low to rather  
strong projections, at first with a thin thalline covering; involucrellum conical . . . . .  
. . . . . **Verrucaria elaeomelaena**, **V. hydrophila**, **Verrucaria** Species A  
Note: these three taxa are not closely related, but they can look very similar. *Verrucaria elaeomelaena* tends to have larger perithecia and  
ascospores than *V. hydrophila* but material is too sparse and ascospores often too poorly developed for a meaningful comparison.  
Thallus thicker, not translucent when fresh and wet, cracks sparse to numerous . . . . . 17
- 17(16) Perithecia forming low to moderately projecting mounds more or less covered to the apex by the thallus; ascospores (23.5–)27–43  $\mu\text{m}$   
long; pycnidia numerous . . . . . **Verrucaria bella**  
Perithecia immersed in the thallus, forming at most very low and indistinct projections . . . . . 18
- 18(17) Thallus extensively cracked, often into discrete areoles with pigmented sides; ascospores small, 13.5–16.5  $\mu\text{m}$  long; pycnidia  
numerous . . . . . **Verrucaria antepotens**  
Thallus extensively cracked or cracks sparse, sides of areoles pigmented or not; ascospores 18–33  $\mu\text{m}$  long; pycnidia . . . . .  
not seen . . . . . 19
- 19(18) Involucrellum irregular in shape; exciple well pigmented when mature, often darker than adjacent involucrellum; ascospores  
with perispore, dilute brown when overmature . . . . . **Verrucaria luchunensis**  
Involucrellum more or less conical, typically dark at surface and paler below, often confluent with other involucrella; exciple  
colourless or only lightly pigmented; ascospores with or without perispore, not becoming brown with age . . . . . 20

- 20(19) Ascospores *c.* 18–20  $\mu\text{m}$  long ((16.0–)18.0–25.0(–28.0)  $\times$  (6.5–)7.0–10.0(–10.5)  $\mu\text{m}$  in material from Europe); perispore absent  
 ..... **Verrucaria praetermissa**  
 Ascospores (23.5–)26.5–32(–33)  $\mu\text{m}$  long; perispore present ..... **Verrucaria lactea**
- 21(12) Ascospores 1- or 3-septate ..... 22  
 Ascospores muriform ..... 24
- 22(21) Ascospores 3-septate, 36–43  $\mu\text{m}$  long; perithecial mounds 530–670  $\mu\text{m}$  diam.; involucrellum thick, *c.* 75–100  $\mu\text{m}$  in upper  
 part ..... **Thelidium papulare**  
 Ascospores 1-septate, not exceeding 25  $\mu\text{m}$  long ..... 23
- 23(22) Involucrellum appressed to exciple or slightly diverging at the base, sometimes the two scarcely discernible as separate struc-  
 tures; ascospores 21.5–25  $\mu\text{m}$  long ..... **Thelidium aff. minutulum**  
 Involucrellum distinct, spreading from the exciple in lower part; ascospores 15.5–19.5  $\mu\text{m}$  long ... **Thelidium uvidulum**
- 24(21) Thallus comprising closely adnate squamules with rhizohyphae below; perithecia immersed in large thalline mounds 380–670  
 $\mu\text{m}$  diam.; ascospores 45–59.5  $\mu\text{m}$  long ..... **Nesothele globosa**  
 Thallus of discrete areoles or extensively cracked, never squamulose; perithecia variously immersed or forming projections;  
 ascospores smaller ..... 25
- 25(24) Thallus of discrete convex areoles; perithecia projections without a thalline covering or with a few areoles below; ascospores  
 29.5–41  $\mu\text{m}$  long ..... **Nesothele glebulosa**  
 Thallus a cracked crust but without discrete primary areoles; perithecia immersed or forming projection with a partial thalline  
 covering below; ascospores often smaller ..... 26
- 26(25) Hymenial algal cells oblong to cylindrical, some at least 3 times as long as wide ..... 27  
 Hymenial algal cells broadly ellipsoid ..... 29
- 27(26) Ascospores (28.5–)29.5–36(–40)  $\mu\text{m}$  long (but measured in poor condition); thallus cracks with black sides; pycnidia present  
 ..... **Willeya irrigata**  
 Ascospores 19–29  $\mu\text{m}$  long; pycnidia not seen ..... 28
- 28(27) Perithecia forming conical-hemispherical projections; black basal thallus layer absent ..... **Willeya eminens**  
 Perithecia immersed in thallus, not forming projections ..... **Willeya pallidipora** and **Willeya** Species A  
 Note: these two species are similar in the pale, weakly cracked thallus.
- 29(26) Ascospores *c.* 32–38  $\mu\text{m}$  long; pycnidia present ..... **Willeya honghensis**  
 Ascospores 19.5–25  $\mu\text{m}$  long; pycnidia not seen ..... **Willeya cf. japonica** and **W. nepalensis**  
 Note: these two species are distinct in ITS sequence but no consistent morphological differences can be found in the sparse material studied.


## Discussion

Although most of the material documented in this paper has come from a single collecting trip, the results are a significant advance in our knowledge of the family *Verrucariaceae* in Nepal. However, there is little doubt that the taxa described here are only a fraction of the diversity present in Nepal, especially since collecting for the present study was confined to moderate altitudes due to unseasonably wet weather experienced during the expedition. Unfortunately, habitats above the forest zone were not sampled.

The small number of collections available for each taxon means that morphological and genetic variability cannot be fully determined. Species of *Verrucariaceae* are often difficult to distinguish, being separated by subtle and overlapping characters. This makes it particularly difficult to match Nepalese collections with previously published species, as there is little sequenced material available from the region. In this study, some collections have been identified as previously known taxa because of either ITS sequences or clear-cut morphological features. In some instances, although morphology might suggest an identification,

ITS sequences show differences from previously published sequences, indicating that more than one taxon may be present (for example with *Hydropunctaria rheitrophila*). Reassessment of species complexes, including *Verrucaria elaeomelaena* and *V. hydrophila*, would certainly benefit from a geographically broader study incorporating additional gene regions. Where it has not been possible to identify a collection with total confidence, such taxa have mostly been described here as new, since the only alternative would have been an unhelpful list of unidentified collections. All newly published taxa are associated with an ITS sequence which provides a reliable, though not infallible, way of fixing the application of a name in the future. Although it is likely that most of the species can be reliably identified by morphological characters, the limited material available means that the extent of variation in each species is unknown. Due to the frequently subtle differences between species, DNA sequencing is recommended in floristic surveys of *Verrucariaceae* in other underexplored regions. Indeed, if sequencing is currently impractical, freezing of some material for future study might be helpful.

**Acknowledgements.** We would like to gratefully acknowledge permission to collect specimens, granted by the Department of National Parks and Wildlife Conservation (DNPWC), Ministry of Forests and Environment, Government of Nepal. Hem Sagar Baral is warmly thanked for his assistance in obtaining an export licence and other matters.

**Author ORCID.**  Alan Orange, 0000-0002-3181-5435.

**Competing interests.** The authors declare none.

## References

- Aptroot A** (2016) The first European *Willeya* (Verrucariaceae) on limestone brought from China. *Herzogia* **29**, 688–691.
- Aptroot A and Feijen FJ** (2002) Annotated checklist of the lichens and lichenicolous fungi of Bhutan. *Fungal Diversity* **11**, 21–48.
- Aptroot A and Sparrus LB** (2006) Additions to the lichen flora of Vietnam, with an annotated checklist and bibliography. *Bryologist* **109**, 358–371.
- Awasthi DD** (1991) A key to the microlichens of India, Nepal and Sri Lanka. *Bibliotheca Lichenologica* **40**, 1–337.
- Borchsenius F** (2009) *FastGap 1.2*. Department of Biosciences, Aarhus University. [WWW resource] URL [http://www.aubot.dk/FastGap\\_home.htm](http://www.aubot.dk/FastGap_home.htm).
- Breuss O** (1996) Revision der Flechtengattung *Placidopsis* (Verrucariaceae). *Österreichische Zeitschrift für Pilzkunde* **5**, 65–94.
- Buaruang K, Boonpragob K, Mongkolsuk P, Sangvichien E, Vongshewarat K, Polyiam W, Rangsiruji A, Saipunkaew W, Naksuwankul K, Kalb J, et al.** (2017) A new checklist of lichenized fungi occurring in Thailand. *MycKeys* **23**, 1–91.
- Döring H, Clerc P, Grube M and Wedin M** (2000) Mycobiont-specific PCR primers for the amplification of nuclear ITS and LSU rDNA from lichenized ascomycetes. *Lichenologist* **32**, 200–204.
- Gardes M and Bruns TD** (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113–118.
- Gueidan C, Roux C and Lutzoni F** (2007) Using a multigene phylogenetic analysis to assess generic delineation and character evolution in Verrucariaceae (Verrucariales, Ascomycota). *Mycological Research* **111**, 1145–1168.
- Gueidan C, Savić S, Thijs H, Roux C, Keller C, Tibell L, Prieto M, Heiðmarsson S, Breuss O, Orange A, et al.** (2009) Generic classification of the Verrucariaceae (Ascomycota) based on molecular and morphological evidence: recent progress and remaining challenges. *Taxon* **58**, 184–208.
- Gueidan C, Van Do T and Lu NT** (2014) Phylogeny and taxonomy of *Staurothele* (Verrucariaceae, lichenized ascomycetes) from the karst of northern Vietnam. *Lichenologist* **46**, 515–533.
- Hall TA** (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**, 95–98.
- Harada H** (1991) *Thelidium japonicum*, a new species of the lichen family Verrucariaceae from Japan. *Natural History Research* **1**, 9–11.
- Harada H** (1992a) A taxonomic study on the lichen genus *Staurothele* (Verrucariaceae) in Japan. *Natural History Research* **2**, 39–42.
- Harada H** (1992b) Taxonomic notes on the lichen family Verrucariaceae in Japan (I). Five species of *Verrucaria* previously reported from Japan. *Journal of Japanese Botany* **67**, 218–226.
- Harada H** (1993) Taxonomic notes on the lichen family Verrucariaceae in Japan (IV). *Verrucaria minuscula* sp. nov. *Hikobia* **11**, 231–233.
- Harada H** (1994) Taxonomic notes on the lichen family Verrucariaceae in Japan (V). *Verrucaria iwatsukii* Harada sp. nov. *Hikobia* **11**, 519–522.
- Harada H** (1995a) Taxonomic notes on the lichen family Verrucariaceae in Japan (VII). *Verrucaria marinomuralis* Harada sp. nov. *Natural History Research* **3**, 111–114.
- Harada H** (1995b) Two new species of maritime lichens in the genus *Verrucaria* (Lichenes, Verrucariaceae) from Japan. *Nova Hedwigia* **60**, 73–78.
- Harada H** (1995c) *Endocarpon minutum* Harada sp. nov. (lichenized Ascomycotina, Verrucariaceae) from western Japan. *Bryologist* **98**, 385–388.
- Harada H** (1996a) *Verrucaria igii*, a new freshwater species of Verrucariaceae (lichenized Ascomycotina) from Japan. *Bryologist* **99**, 343–344.
- Harada H** (1996b) Taxonomic notes on the lichen family Verrucariaceae in Japan (X). *Thelidium radiatum* Harada sp. nov. *Hikobia* **12**, 133–136.
- Harada H** (1996c) Taxonomic notes on the lichen family Verrucariaceae in Japan (IX). *Verrucaria rheitrophila* Zsch., new to Japan. *Journal of Japanese Botany* **71**, 317–322.
- Harada H** (1998) Taxonomic notes on the lichen family Verrucariaceae in Japan (XI). *Thelidium pacificum*, a new maritime species. *Hikobia* **12**, 289–291.
- Harada H** (2000) Three new maritime species of *Verrucaria* (lichenized Ascomycota, Verrucariaceae) from Japan. *Bryologist* **103**, 555–562.
- Harada H** (2003) *Psoroglaena japonica* (lichenized Ascomycota, Verrucariaceae), a new species from Chiba-ken, central Japan, with notes on *Psoroglaena*. *Lichenology* **2**, 5–10.
- Harada H** (2012a) Taxonomic study on the freshwater species of Verrucariaceae of Japan (1). *Verrucaria praetermissa* and *V. yoshimurae* sp. nov. *Lichenology* **10**, 33–41.
- Harada H** (2012b) Taxonomic study on the freshwater species of Verrucariaceae of Japan (2). Genus *Verrucaria*. *Lichenology* **10**, 97–135.
- Harada H** (2013a) *Agonimia deguchii* (lichenized Ascomycota, Verrucariaceae), a new saxicolous species from central Japan. *Hikobia* **16**, 307–310.
- Harada H** (2013b) The lichen genus *Thelidium* (Verrucariaceae) in Japan. *Lichenology* **11**, 53–66.
- Harada H and Wang L-S** (1996) Two new freshwater species of Verrucariaceae from Yunnan, China. *Lichenologist* **28**, 297–305.
- Harada H and Wang L-S** (2006a) Taxonomic study on the freshwater species of Verrucariaceae (lichenized Ascomycota) of Yunnan, China (2). Genus *Staurothele*. *Lichenology* **5**, 13–22.
- Harada H and Wang L-S** (2006b) Taxonomic study on the freshwater species of Verrucariaceae (lichenized Ascomycota) of Yunnan, China (3). Genus *Thelidium*. *Lichenology* **5**, 23–30.
- Harada H and Wang L-S** (2008) Taxonomic study on the freshwater species of Verrucariaceae (lichenized Ascomycota) of Yunnan, China (4). Genus *Verrucaria*. *Lichenology* **7**, 1–24.
- Heiðmarsson S, Gueidan C, Miadłlikowska J and Lutzoni F** (2017) Multi-locus phylogeny supports the placement of *Endocarpon pulvinatum* within *Staurothele* s. str. (lichenized ascomycetes, Eurotiomycetes, Verrucariaceae). *Phytotaxa* **306**, 37–48.
- Löytynoja A and Goldman N** (2010) webPRANK: a phylogeny-aware multiple sequence aligner with interactive alignment browser. *BMC Bioinformatics* **11**, 579.
- Lücking R, Hodkinson B and Leavitt SD** (2016) The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota – approaching one thousand genera. *Bryologist* **119**, 361–416.
- Magnusson AH** (1940) Lichens from central Asia I. In Hedin S (ed.), *Reports from the Scientific Expedition to the North-Western Provinces of China Under the Leadership of Dr. Sven Hedin. The Sino-Swedish Expedition* **13**, XI. Botany, 1. Stockholm: Aktiebolaget Thule.
- Malcolm WM and Galloway DJ** (1997) *New Zealand Lichens. Checklist, Key, and Glossary*. Wellington: Museum of New Zealand Te Papa Tongarewa.
- McCarthy PM** (1991) A new species and new records of *Verrucaria* Schrader (lichenized Ascomycotina, Verrucariaceae) from New Zealand. *New Zealand Journal of Botany* **29**, 283–286.
- McCarthy PM** (1995a) Notes on Australian Verrucariaceae (lichenized Ascomycotina). 5. *Staurothele pallidopora* sp. nov. from south-eastern Queensland. *Muelleria* **8**, 275–277.
- McCarthy PM** (1995b) Aquatic species of *Verrucaria* in eastern Australia. *Lichenologist* **27**, 105–126.
- McCarthy PM** (2003) Additional lichen records from Australia 51. Some aquatic pyrenolichens from Kosciuszko National Park, Snowy Mountains, New South Wales. *Australasian Lichenology* **52**, 16–18.
- Miller MA, Pfeiffer W and Schwartz T** (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In *Proceedings of the Gateway Computing Environments Workshop (GCE), 14 November 2010, New Orleans, Louisiana*, pp. 1–8.
- Ohmura Y and Kashiwadani H** (2018) Checklist of lichens and allied fungi of Japan. *National Museum of Nature and Science Monographs* **49**, 1–140.

- Olley L** (2011) *A provisional checklist to the lichens of Nepal*. [WWW document] URL [https://rbg-web2.rbge.org.uk/lichen/staff\\_profiles/olley/nepal\\_checklist.html](https://rbg-web2.rbge.org.uk/lichen/staff_profiles/olley/nepal_checklist.html).
- Olley L and Sharma LR** (2013) A provisional checklist of the lichens of Nepal. *Journal of the Department of Plant Resources Nepal* **35**, 18–21.
- Orange A** (2000) *Verrucaria elaeina*, a misunderstood European lichen. *Lichenologist* **32**, 411–422.
- Orange A** (2012) Semi-cryptic marine species of *Hydropunctaria* (Verrucariaceae, lichenized Ascomycota) from north-west Europe. *Lichenologist* **44**, 299–320.
- Orange A** (2018) A new species-level taxonomy for *Trapelia* (Trapeliaceae, Ostropomycetidae) with special reference to Great Britain and the Falkland Islands. *Lichenologist* **50**, 3–42.
- Orange A** (2020) The *Verrucaria aethiobola* group (lichenised Ascomycota, Verrucariaceae) in North-west Europe. *Phytotaxa* **459**, 1–15.
- Pérez-Ortega S, de los Ríos A, Crespo A and Sancho LG** (2010) Symbiotic lifestyle and phylogenetic relationships of the biotents of *Mastodia tessellata* (Ascomycota, incertae sedis). *American Journal of Botany* **97**, 738–752.
- Pérez-Ortega S, Miller KA and de los Ríos A** (2018) Challenging the lichen concept: *Turgidosculum ulvae* (Verrucariaceae) represents an independent photobiont shift to a multicellular blade-like alga. *Lichenologist* **50**, 341–356.
- Prieto M, Martínez I, Aragón G and Otálora MAG** (2010) Phylogenetic study of *Catapyrenium* s. str. (Verrucariaceae, lichen-forming Ascomycota) and related genus *Placidopsis*. *Mycologia* **102**, 291–304.
- Prieto M, Martínez I, Aragón G, Gueidan C and Lutzoni F** (2012) Molecular phylogeny of *Heteroplacidium*, *Placidium*, and related catapyrenioid genera (Verrucariaceae, lichen-forming Ascomycota). *American Journal of Botany* **99**, 23–35.
- Pykälä J, Launis A and Myllys L** (2018) *Verrucaria tenebrosa* (Verrucariaceae), a new lichen species from Finland and Norway, and notes on the taxonomy of epiphytic taxa belonging to the *V. hydrophila* complex. *Phytotaxa* **361**, 211–221.
- Rai H, Nag P, Khare R, Upreti DK and Gupta RK** (2016) Twenty-eight new records of lichenized fungi from Nepal: a signature of undiscovered biodiversity in central Himalaya. *Proceedings of the National Academy of Sciences, India, Section B Biological Sciences* **87**, 1363–1376.
- Savić S and Tibell L** (2009) Taxonomy and species delimitation in *Sporodictyon* (Verrucariaceae) in Northern Europe and the adjacent Arctic – reconciling molecular and morphological data. *Taxon* **58**, 585–605.
- Savić S and Tibell L** (2012) *Polyblastia* in Northern Europe and the adjacent Arctic. *Symbolae Botanicae Upsalienses* **36**, 1–69.
- Sharma LR** (1995) *Enumeration of the Lichens of Nepal*. Nepal Bio-diversity profile project. Kathmandu: Department of National Parks & Wildlife Conservation.
- Stamatakis A** (2006) RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688–2690.
- Stamatakis A, Hoover P and Rougemont J** (2008) A fast bootstrapping algorithm for the RAXML web servers. *Systematic Biology* **57**, 758–771.
- Tamura K, Dudley J, Nei M and Kumar S** (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**, 1596–1599.
- Thüs H and Nascimbene J** (2008) Contributions toward a new taxonomy of Central European freshwater species of the lichen genus *Thelidium* (Verrucariales, Ascomycota). *Lichenologist* **40**, 499–521.
- Thüs H, Muggia L, Pérez-Ortega S, Favero-Longo SE, Joneson S, O'Brien H, Nelsen MP, Duque-Thüs R, Grube M, Friedl T, et al.** (2011) Revisiting photobiont diversity in the lichen family Verrucariaceae (Ascomycota). *European Journal of Phycology* **46**, 399–415.
- Thüs H, Orange A, Gueidan C, Pykälä J, Ruberti C, Lo Schiavo F and Nascimbene J** (2015) Revision of the *Verrucaria elaeomelaena* species complex and morphologically similar freshwater lichens (Verrucariaceae, Ascomycota). *Phytotaxa* **197**, 161–185.
- Wang-Yang JR and Lai MJ** (1973) A checklist of the lichens of Taiwan. *Taiwania* **18**, 83–104.
- White TJ, Bruns T, Lee S and Taylor J** (1990) Amplification and direct sequencing of fungal ribosomal genes for phylogenetics. In Innis M, Gelfand D, Sninsky J and White T (eds), *PCR Protocols: a Guide to Methods and Applications*. New York: Academic Press, pp. 315–322.
- Zahlbruckner A** (1930) Teil 3. Lichens. In Handel-Mazzetti HRE (ed.), *Symbolae Sinicae: Botanische Ergebnisse der Expedition der Akademie der Wissenschaften in Wein nach Südwest-China, 1914–1918*. Wien: J. Springer
- Zoller S, Scheidegger C and Sperisen C** (1999) PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. *Lichenologist* **31**, 511–516.