


Short Communication

Unravelling the hidden inter and intra-varietal diversity of durum wheat commercial varieties used in Portugal

Miguel Bento, Sónia Gomes Pereira, Wanda Viegas and Manuela Silva* 

Linking Landscape, Environment, Agriculture and Food (LEAF), Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda 1349-017 Lisboa, Portugal

Received 19 December 2018; Accepted 5 April 2019 – First published online 6 May 2019

Abstract

Assessing durum wheat genomic diversity is crucial in a changing environment particularly in the Mediterranean region where it is largely used to produce pasta. Durum wheat varieties cultivated in Portugal and previously assessed regarding thermotolerance ability were screened for the variability of coding sequences associated with technological traits and repetitive sequences. As expected, reduced variability was observed regarding low molecular weight glutenin subunits (LMW-GS) but a specific LMW-GS allelic form associated with improved pasta-making characteristics was absent in one variety. Contrastingly, molecular markers targeting repetitive elements like microsatellites and retrotransposons – Inter Simple Sequence Repeat (ISSR) and Inter Retrotransposons Amplified Polymorphism (IRAP) – disclosed significant inter and intra-varietal diversity. This high level of polymorphism was revealed by the 20 distinct ISSR/IRAP concatenated profiles observed among the 23 individuals analysed. Interestingly, median joining networks and PCoA analysis grouped individuals of the same variety and clustered varieties accordingly with geographical origin. Globally, this work demonstrates that durum wheat breeding strategies induced selection pressure for some relevant coding sequences while maintaining high levels of genomic variability in non-coding regions enriched in repetitive sequences.

Keywords: genetic diversity, glutenins, microsatellites and retrotransposons

Introduction

Durum wheat (*Triticum turgidum* L.) represents the major ingredient of such an important food as pasta (Sissons, 2008). In the context of environmental changes, it is crucial to evaluate the genomic diversity of nowadays commercial varieties. Durum wheat varieties used in Portugal and previously assessed for heat stress response (Bento *et al.*, 2017) were evaluated regarding the variability of coding sequences associated with technological traits and non-coding repetitive sequences.

Glutenins are polymeric gluten proteins determinant to dough elasticity (Sissons, 2008) among which low molecular weight glutenin subunits (LMW-GS) form inter-molecular disulphide bonds creating polymers responsible for gluten structure and properties (D'Ovidio and Masci, 2004). Sequences coding for LMW-GS, a diverse protein family, were previously characterized in bread wheat (Zhang *et al.*, 2011). LMW-1 and LMW-2 subunits were related to durum wheat gluten viscoelasticity, being LMW-2 associated with superior dough quality (Payne *et al.*, 1984).

Genomic characterization using Inter Retrotransposons Amplified Polymorphism (IRAP) and Inter Simple Sequence Repeat (ISSR) techniques has been successfully used to identify several crop species and varieties (Smykal,

*Corresponding author. E-mail: manuelasilva@isa.ulisboa.pt

2006; Bento *et al.*, 2011; Carvalho *et al.*, 2012; Khan *et al.*, 2015; Tomás *et al.*, 2016a, b). Both methodologies uncover insertional polymorphisms: IRAP methodology using outward primers to a long terminal repeat of LTR-retrotransposons (Kalendar and Schulman, 2006) and ISSR anchored primers to SSR (Zietkiewicz *et al.*, 1994).

In this work, we assessed LMW-GS variability and used IRAP and ISSR fingerprinting techniques to characterize the diversity of *T. durum* varieties grown in Portugal.

Experimental

Six durum wheat varieties used in Portugal and previously assessed regarding tolerance to heat stress treatments in distinct developmental stages (Bento *et al.*, 2017) were used: Celta, Hélio and Marialva (Portuguese); Saragolla and Severo (Italian); and DonDuro (Greek). DNA of at least three individual plants from each variety was isolated from leaves using Citogene® DNA Purification Kit. LMW-GS analysis used primers for conserved regions (Zhang *et al.*, 2011) and specific for LMW-1 and LMW-2 subunits (D'Ovidio, 1993). IRAP used primers targeting the LTR region of retrotransposons *Angela* and *Fatima* and ISSR used the primer (AAG)₆C (online Supplementary Table S1).

PCR products were separated through electrophoresis and at least three profiles per individual plant were analysed with GelAnalyzer 2010a. Concatenated data was used to calculate similarity matrixes with simple matching (SM) coefficient and to produce PCoA scatter plot using NTSYSpc version 2.02i (Rohlf, 2000), and NETWORK 5.0.0.0 (Fluxus Technology Ltd., Clare, Suffolk, England) was used to construct median joining networks.

Discussion

Previous studies showed the importance of LMW-GS genes variability in durum wheat end-use quality (D'Ovidio and Masci, 2004; Sissons, 2008). The primer pair designed to characterize bread wheat varieties targeting the conserved regions of LMW-GS genes and flanking the variable ones (Zhang *et al.*, 2011) yielded informative profiles in durum wheat due to the phylogenetic proximity between these two wheat species (online Supplementary Fig. S1a). No intra-varietal diversity and a marked lack of inter-varietal diversity were observed since all varieties except Hélio showed identical banding profiles. On the other hand, the LMW-2 subunit related with superior quality (Payne *et al.*, 1984; D'Ovidio, 1993) was identified in all individuals of Hélio, DonDuro, Saragolla, Severo and three out of four individuals from the variety Marialva as expected in commercial varieties (online Supplementary Fig. S1b). Contrastingly, one individual from the variety Marialva and all Celta individuals showed the presence of LMW-1

allelic form. The median joining network obtained from LMW-GS and LMW-1/LMW-2 concatenated data (Fig. 1 (a)) highlighted the low variability observed since besides Hélio, which is the variety that accumulated more mutational steps, only variety DonDuro was distinguished. Low variability detected among coding regions in durum wheat varieties is not surprising and was previously reported in HMW-GS and LMW-GS composition of Spanish, Italian and Moroccan cultivars (De Vita *et al.*, 2007; Subira *et al.*, 2014; Henkrar *et al.*, 2017). This lack of diversity regarding end-use traits can be a direct consequence of breeding programme strategies generally used (De Vita *et al.*, 2007).

To further access genomic variability, we used ISSR and IRAP molecular markers targeting repetitive elements. These markers highlighted 57% polymorphic amplicons allowing the discrimination between varieties and even between individuals of the same variety (online Supplementary Tables S2 and S3). Even though the polymorphism level detected was lower than previously reported for durum wheat landraces using ISSR or IRAP (Pujar *et al.*, 2002; Khan *et al.*, 2015) it was significant considering only commercial varieties. Moreover, the separated characterization of multiple individuals of each variety, contrarily to previous studies (Pujar *et al.*, 2002; Carvalho *et al.*, 2008; Carvalho *et al.*, 2012; Shimasabian *et al.*, 2014; Khan *et al.*, 2015), allowed the determination of similarity coefficient values higher (0.60–0.92, online Supplementary Table S4) than those previously reported using ISSR in durum wheat landraces (Carvalho *et al.*, 2009) or advanced genotypes (Zamanianfard *et al.*, 2015). Additionally, these multiple individual analyses disclosed a much more obvious intravarietal diversity assessed through IRAP and ISSR in comparison with LMW.

Intravarietal diversity in repetitive genome regions was expected, but the ISSR/IRAP banding profiles and the presence of LMW-1 in one Marialva individual plant similar to Celta variety were not expected. Regarding the Marialva outlier genotype, we cannot completely discard the hypothesis of seed admixture, although all varieties were maintained by controlled self-fertilization during at least four consecutive years.

Importantly, combining ISSR and IRAP median joining network (Fig. 1(b)) produced almost a unique profile for each individual (20 distinct profiles among 23 individuals). However, different individuals of the same variety were clustered, allowing inference of phylogenetic relationship between varieties. PCoA scatter plot (Fig. 2) with the first two principal components (51.9% variation) also grouped individuals by variety and separates varieties in three groups accordingly with their geographical origin.

Overall, our study revealed opposing scenarios coexisting in durum wheat genotypes regarding variability levels of coding sequences associated with important




Fig. 1. Median-joining networks obtained in *Triticum durum* cultivars Celta, Hélivio, Marialva, DonDuro, Saragolla and Severo from (a) LMW-GS genes and LMW-1/LMW-2 subunits concatenated profiles and from (b) IRAP and ISSR concatenated profiles. Branches are proportional to the number of differences between haplotypes and nodes are proportional to frequencies of haplotypes. Black dots on branches indicate the mutational steps when more than one and white dots represent median vectors.




Fig. 2. Principal coordinate analysis of IRAP and ISSR profiles based on the geographic origin of six durum wheat varieties: Celta, Hélivio and Marialva (Portugal); Saragolla and Severo (Italy); and DonDuro (Greece).

technological traits and repetitive genome domains. The first, being a target of artificial selection, showed low variability in durum wheat commercial varieties, favouring the

presence of particular alleles, while more labile and less scrutinized repetitive sequences retrieved high variability levels between and within the varieties studied.

Acknowledgements

We would like to thank Eng^o Benvido Maças (INRB/ INIAV Portugal) and Eng^a Joana Aleixo (ANSEME Portugal) for the seeds used in the present study. Miguel Bento was funded by an FCT (Fundação para a Ciência e a Tecnologia, Portugal) Post-doctoral research fellowship (SFRH/BPD/80550/2011) and Manuela Silva by the FCT Investigator Program (IF/00834/2014). This work was supported by the FCT LEAF Unit (UID/AGR/04129/2013).

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262119000133>

References

- Bento M, Gustafson JP, Viegas W and Silva M (2011) Size matters in Triticeae polyploids: larger genomes have higher remodeling. *Genome* 54: 175–183.
- Bento M, Pereira SG, Viegas W and Silva M (2017) Durum wheat diversity for heat stress tolerance during inflorescence emergence is correlated to TdHSP101C expression in early developmental stages. *PLoS ONE* 12: e0190085.
- Carvalho A, Lima-Brito J, Macas B and Guedes-Pinto H (2008) Genetic variability analysis of a collection of old Portuguese bread wheat using ISSRs. *Options Méditerranéennes Series A* 81: 35–38.
- Carvalho A, Lima-Brito J, Macas B and Guedes-Pinto H (2009) Genetic diversity and variation among botanical varieties of Old Portuguese wheat cultivars revealed by ISSR assays. *Biochemical Genetics* 47: 276–294.
- Carvalho A, Guedes-Pinto H and Lima-Brito JE (2012) Genetic diversity in Old Portuguese durum wheat cultivars assessed by retrotransposon-based markers. *Plant Molecular Biology Reporter* 30: 578–589.
- De Vita P, Nicosia OL, Nigro F, Platani C, Riefolo C, Di Fonzo N and Cattivelli L (2007) Breeding progress in morpho-physiological, agronomical and qualitative traits of durum wheat cultivars released in Italy during the 20th century. *European Journal of Agronomy* 26: 39–53.
- D'Ovidio R (1993) Single-seed PCR of LMW glutenin genes to distinguish between durum-wheat cultivars with good and poor technological properties. *Plant Molecular Biology* 22: 1173–1176.
- D'Ovidio R and Masci S (2004) The low-molecular-weight glutenin subunits of wheat gluten. *Journal of Cereal Science* 39: 321–339.
- Henkrar F, El-Haddoury J, Iraqi D, Bendaou N and Udupa SM (2017) Allelic variation at high-molecular weight and low-molecular weight glutenin subunit genes in Moroccan bread wheat and durum wheat cultivars. *3 Biotech* 7: 287–297.
- Kalendar R and Schulman AH (2006) IRAP and REMAP for retrotransposon-based genotyping and fingerprinting. *Nature Protocols* 1: 2478–2484.
- Khan MK, Pandey A, Thomas G, Akkaya MS, Kayis SA, Ozsensoy Y, Hamurcu M, Gezgin S, Topal A and Hakki EE (2015) Genetic diversity and population structure of wheat in India and Turkey. *Aob Plants* 7, plv083.
- Payne PI, Jackson EA and Holt LM (1984) The association between gamma-gliadin 45 and gluten strength in durum wheat varieties: a direct causal effect or the result of genetic linkage? *Journal of Cereal Science* 2: 73–81.
- Pujar S, Tamhankar SA, Gupta VS, Rao VS and Ranjekar PK (2002) Diversity analysis of Indian tetraploid wheat using inter-simple sequence repeat markers reveals their superiority over random amplified polymorphic DNA markers. *Biochemical Genetics* 40: 63–69.
- Rohlf FJ (2000) NTSYS-pc: numerical taxonomy and multivariate analysis system, version 2.1.
- Shirnasabian S, Etminan A, Mohammadi R and Shoostari L (2014) Molecular variation of improved durum wheat genotypes based on inter-simple sequence repeats fingerprinting. *International Journal of Biosciences* 5: 222–228.
- Sissons MJ (2008) Role of durum wheat composition on the quality of pasta and bread. *Food* 2: 75–90.
- Smykal P (2006) Development of an efficient retrotransposon-based fingerprinting method for rapid pea variety identification. *Journal of Applied Genetics* 47: 221–230.
- Subira J, Pena RJ, Alvaro F, Ammar K, Ramdani A and Royo C (2014) Breeding progress in the pasta-making quality of durum wheat cultivars released in Italy and Spain during the 20th century. *Crop & Pasture Science* 65: 16–26.
- Tomás D, Dias AL, Silva M, Oliveira HR, Suso MJ, Viegas W and Veloso MM (2016a) Genetic diversity assessment of Portuguese cultivated *Vicia faba* L. through IRAP markers. *Diversity* 8: 8–19.
- Tomás D, Rodrigues J, Varela A, Veloso MM, Viegas W and Silva M (2016b) Use of repetitive sequences for molecular and cytogenetic characterization of *Avena* species from Portugal. *International Journal of Molecular Sciences* 17: 203–217.
- Zamanianfard Z, Etminan A, Mohammadi R and Shoostari L (2015) Evaluation of molecular diversity of durum wheat genotypes using ISSR markers. *Biological Forum* 7: 214–218.
- Zhang XF, Liu DC, Yang WL, Liu KF, Sun JZ, Guo XL, Li YW, Wang DW, Ling HQ and Zhang AM (2011) Development of a new marker system for identifying the complex members of the low-molecular-weight glutenin subunit gene family in bread wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics* 122: 1503–1516.
- Zietkiewicz E, Rafalski A and Labuda D (1994) Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20: 176–183.