

Article

Cite this article: Liljas A, Sanyal S (2018). The enigmatic ribosomal stalk. *Quarterly Reviews of Biophysics* **51**, e12, 1–10. <https://doi.org/10.1017/S0033583518000100>

Received: 16 September 2018

Revised: 15 October 2018

Accepted: 31 October 2018

Key words:

Ribosomal stalk; bL12; P1/P2; translational GTPases (trGTPases)

Author for correspondence:

Anders Liljas, E-mail: anders.liljas@biochemistry.lu.se

© The Author(s) 2018. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

The enigmatic ribosomal stalk

Anders Liljas¹ and Suparna Sanyal²

¹Department of Biochemistry and Structural Biology, Center for Chemistry and Chemical Engineering, Lund University, Lund, Sweden and ²Department of Cell and Molecular Biology, Biomedical Center, Uppsala University, Uppsala, Sweden

Abstract

The large ribosomal subunit has a distinct feature, the stalk, extending outside the ribosome. In bacteria it is called the L12 stalk. The base of the stalk is protein uL10 to which two or three dimers of proteins bL12 bind. In archaea and eukarya P1 and P2 proteins constitute the stalk. All these extending proteins, that have a high degree of flexibility due to a hinge between their N- and C-terminal parts, are essential for proper functionalization of some of the translation factors. The role of the stalk proteins has remained enigmatic for decades but is gradually approaching an understanding. In this review we summarise the knowhow about the structure and function of the ribosomal stalk till date starting from the early phase of ribosome research.

Introduction

The ribosome translates a messenger RNA (mRNA) into a chain of amino acid residues to produce proteins. In order to proceed at a speed that is required to sustain life translation factors are needed to catalyze the process. Several of these factors bind in complex with GTP molecules and dissociate after GTP hydrolysis. Non-cleavable GTP analogues inhibit the participation of these factors in protein synthesis. These GTP-depending factors can be called translational GTPases (referred as trGTPases). The induction of GTP hydrolysis has for a long time been an important question.

The two subunits of the ribosome in bacteria are usually composed of two large and one smaller RNA molecule and a large number of ribosomal proteins (See the book by Liljas & Ehrenberg, 2013). The early electron microscopic studies of the ribosomal large subunit identified a prominent stalk-like extension composed of the protein bL12 (Fig. 1) (Boublik *et al.*, 1976; Lake, 1976; Strycharz *et al.*, 1978). Removing protein bL12 has detrimental effects on the functionality of the trGTPases (Kischa *et al.*, 1971; Hamel *et al.*, 1972; Koteliansky *et al.*, 1978; Pettersson and Kurland, 1980; Mohr *et al.*, 2002; Huang *et al.*, 2010).

The organization of the ribosomal stalk proteins in different kinds of ribosomes and their role in relation to the GTP hydrolysis has remained enigmatic for decades but is gradually approaching an understanding.

The L7/L12 stalk

The protein bL12

The protein L7/L12 is an acidic protein that was first characterized in *Escherichia coli*, where its N-terminus can be acetylated (Terhorst *et al.*, 1972). The acetylated form of L12 is referred to as L7; this is rarely seen in other species. A new naming system has been introduced to accentuate whether ribosomal proteins are universal (and get the prefix u) or occur only in bacteria (prefix b), archaea (a) or (e) for uniquely eukaryotic proteins (Ban *et al.*, 2014). According to this naming system, L7/L12 is now called bL12. For reviews on bL12 see Liljas, 1982, 1991; Gudkov, 1997; Sanyal and Liljas, 2000; Wahl and Möller, 2002; Gonzalo and Reboud, 2003.

The stoichiometry of bL12

The primary analysis of the stoichiometry of ribosomal proteins demonstrated that all proteins are present in one copy per ribosome except bL12, which was found in more than two copies per ribosome (Hardy, 1975; Subramanian, 1975). Later chemical crosslinking and small angle X-ray scattering studies showed that bL12 forms dimers (Österberg *et al.*, 1976). Thus the stoichiometry of the protein on the ribosome should always be in multiples of two. In *E. coli* there are two bL12 dimers (Österberg *et al.*, 1977), but some species (e.g. *Deinococcus radiodurans*, *Thermus thermophilus*) can have three dimers of bL12 (Ilag *et al.*, 2005). The dimers of bL12 bind to the ribosome through protein uL10 (Diaconu *et al.*, 2005). The tight pentameric complex (bL12)₄.uL10 was initially identified as a unique protein (called L8) not only in *E. coli* (Kaltschmidt and Wittmann, 1969, 1970; Stöffler, 1974) but also in *Bacillus stearothermophilus*




Fig. 1. A symbolic representation of the bacterial ribosome with its small (brown) and large (orange) subunits labelled as the 30S and 50S, respectively. Four copies of the protein bL12 are seen on the right-hand side (blue). Each monomer is composed of two domains (bL12 NTD and bL12 CTD) joined by a flexible linker. The bL12 NTD bind to the protein uL10 of the 50S subunit while the bL12 CTDs are free to move held only by the linker.

(Cohlberg and Nomura, 1976). It was shown that by mixing bL12 with uL10 *in vitro*, the ‘unique protein’, L8, could be formed (Pettersson *et al.*, 1976).

Ribosomal location and basic structure

Electron microscopy showed that bL12 (formerly L7/L12) forms a stalk-like extension, the so-called L7/L12 stalk on the right-hand side of the large subunit when viewed from the subunit interface (Fig. 1; Boublik *et al.*, 1976; Strycharz *et al.*, 1978). This elongated protein has two structured domains: N-terminal domain (NTD) and C-terminal domain (CTD) interconnected by a highly flexible ‘hinge’ segment. The NTDs form the tight dimer and can be seen closely associated with the C-terminal long $\alpha 8$ helix of the protein uL10 (Diaconu *et al.*, 2005). Due to the flexible nature of the hinge, the CTDs could be crosslinked to various locations on the ribosome (Traut *et al.*, 1995; Dey *et al.*, 1998). The flexibility explains why the bL12 stalk is poorly, if at all, visible in cryo-EM or crystallographic structures (Agrawal *et al.*, 1999; Ban *et al.*, 2000; Yusupov *et al.*, 2001; Valle *et al.*, 2003; Harms *et al.*, 2008; Gao *et al.*, 2009; Schmeing *et al.*, 2009; Voorhees *et al.*, 2010; Chen *et al.*, 2013; Lin *et al.*, 2015).

The structure of the bL12 dimer

The bL12 dimer structures have been solved using X-ray crystallography and NMR (Fig. 2) (Wahl *et al.*, 2000; Bocharov *et al.*, 2004; Christodoulou *et al.*, 2004). These structures clearly depict the structural features of different domains of bL12 and the basis of their dimerization. Additionally, they also demonstrate how the flexible hinge leads to different extended and compact conformations of the bL12 dimer.

The NTD of the bL12 monomer is 35 amino acid long in *Escherichia coli* and composed of two α -helices ($\alpha 1$ and $\alpha 2$; Bocharov *et al.*, 1996; 1998). The helices of the NTDs form a hook around its corresponding partner to form the bL12 dimer. In a bL12 dimer, the monomers are arranged in an anti-parallel fashion with respect to each other (Fig. 2; Sanyal and Liljas, 2000; Wahl *et al.*, 2000; Bocharov *et al.*, 2004).

The CTD of bL12 (residue number 52–120 in *E. coli*) was the first component of the ribosome for which the crystal




Fig. 2. Ribbon diagrams illustrating different conformations of the bL12 dimer. *a, b* – The conformations of the bL12 dimer from NMR (Bocharov *et al.*, 2004; PDB ID: 1RQT and 1RQS) showing both hinges extended (*a*) and one in extended and one in a compact state (*b*). The different domains are marked as CTD (C-terminal domain), NTD dimer (N-terminal domain dimer) and hinge.

structure was determined at atomic resolution (Leijonmarck and Liljas, 1987; Leijonmarck *et al.*, 1980). It has a globular structure with a conserved, positively charged surface, which interacts with the trGTPases (Leijonmarck and Liljas, 1987; Wahl *et al.*, 2000; Bocharov *et al.*, 2004). Two α -helices ($\alpha 4$ and $\alpha 5$) of the bL12 CTDs constitute the universal translation factor interaction sites (Savelsbergh *et al.*, 2005; Helgstrand *et al.*, 2007).

NTD and CTD of bL12 are joined by a flexible link or hinge involving residues 36–51 (Bushuev *et al.*, 1989; Bocharov *et al.*, 2004; Mulder *et al.*, 2004). The length and composition of the bL12 hinge, rich in alanyl and glycyl residues, varies considerably between different organisms (Liljas *et al.*, 1986; Bushuev *et al.*, 1989). The hinge is the source of the flexibility of bL12 proteins. From NMR studies it has been suggested that this hinge may be fully extended. Alternatively, it may also transiently form an α -helix (Fig. 2*a, b*) (Bocharov *et al.*, 2004). In the x-ray structure of bL12 from *T. maritima* (Sanyal and Liljas, 2000; Wahl *et al.*, 2000), where the crystallographic asymmetric unit contains two complete bL12 and two NTDs, both the hinges ($\alpha 3$) were seen to adopt the compact helical conformations.

NMR spectra of large subunits or whole ribosomes show that bL12 CTD behaves, as if it would be essentially free in solution (Tritton, 1980; Gudkov *et al.*, 1982; Cowgill *et al.*, 1984; Bushuev *et al.*, 1989; Mulder *et al.*, 2004; Bernadó *et al.*, 2010). Furthermore, NMR spectra of bL12 in 50S subunits or 70S ribosomes indicate that only two of the four hinge regions are mobile (Christodoulou *et al.*, 2004; Mulder *et al.*, 2004). It is not known whether the flexible bL12 hinges belong to the same or separate dimers (Sanyal and Liljas, 2000).

The interactions of bL12 with uL10

The two or three NTD dimers of bL12 (Ilag *et al.*, 2005) all bind to the C-terminal helix ($\alpha 8$) of protein uL10 (Fig. 3; Diaconu *et al.*, 2005). This helix has a short sequence motif that repeats twice or three times. Each such motif can bind a dimer of bL12. From investigations of the bL12 dimer (Bocharov *et al.*, 1996; Mulder *et al.*, 2004) or the pentameric L8 complex (Bocharov *et al.*, 1998) it is clear that the bL12 CTDs neither interact with each other nor with their NTDs.




Fig. 3. The structure of three dimers of bL12 NTDs bound to $\alpha 8$ of uL10 (blue) extracted from PDB: 1ZAX.

The 'stalk' on archaeal or eukaryotic ribosomes

The P proteins

The proteins corresponding to bacterial bL12 and uL10 in eukaryotes and archaea can be phosphorylated and are therefore called P-proteins (Zinker and Warner, 1976). The protein P0 corresponds to uL10 (Wool *et al.*, 1991). Its N-terminal part is an orthologue of the bacterial form of uL10 (Shimmin *et al.*, 1989). In eukarya there are two related proteins, eP1 and eP2, which functionally correspond to bL12 in that they bind to the long C-terminal helix of uL10 as two heterodimers (Tchorzewski *et al.*, 2000a; Maki *et al.*, 2007) to form the bL12 corresponding stalk of the large subunit (Uchiyama *et al.*, 1987; for a review see Liljas, 1991). eP1/eP2 have no sequence similarity with bL12 but share the acidic nature. In archaea, there is only one protein related to eP1 and eP2. It is called aP1. Like in eukaryotes, the archaeal uL10 can bind three dimers of aP1. In addition, the C-terminal region of P0 in both archaea and eukaryotes contains an extension of about 30 residues identical with the C-terminal residues of P1 and P2 (Santos and Ballesta, 1995).

The eP1 and eP2 have NTDs of about 70 residues followed by a flexible region of about 30 amino acid residues (Wool *et al.*, 1991; Bailey-Serres *et al.*, 1997; Tchorzewski, 2002). This hinge-region is acidic and highly variable in length and amino acid sequence but is rich in alanyl and glycyl residues. The C-termini of all P proteins has a conserved stretch of 10–13 residues. Several protein kinases can phosphorylate a conserved seryl residue in this region (Ballesta *et al.*, 1999). Like for bL12, the C-terminal regions of the eP1/eP2-proteins interact with trGTPases (Bargis-Surgey *et al.*, 1999). Similar to bL12, the NTDs of aP1 in archaea and eP1 and eP2 in eukaryotes are involved in the dimerization (Fig. 4) as well as binding to uL10 (Ballesta and Remacha, 1996). The mode of dimerization

involving helical NTDs is more complex in archaea and eukaryotes than in bacteria and involves four α -helices from each molecule than two α -helices in bacteria (Fig. 4).

In yeast, there are four variants of the P proteins: P1A, P1B, P2A and P2B. P1A forms a heterodimer with P2B and P2A forms a heterodimer with P1B (Tchorzewski *et al.*, 2000b; Guarinos *et al.*, 2001). In plants, there is one additional P protein, P3 (Bailey-Serres *et al.*, 1997). If all P-proteins are deleted, the ribosomes do not function. If only the P1 and P2 proteins are missing, the ribosome is able to support protein synthesis with the C-terminally extended uL10 alone (Santos & Ballesta, 1994; Remacha *et al.*, 1995), but a different subset of mRNAs is translated. However, if the ~ 30 amino acid long C-terminal residues of uL10 corresponding to the C-terminus of P1/P2 are also removed from uL10, the ribosome is no more functional.

The structure of the P-protein stalk

The structures of uL10 and the stalk complex from the archaeal species have been determined at high resolution (Kravchenko *et al.*, 2010; Naganuma *et al.*, 2010). The latter structure contains uL10 with three dimers of the N-termini of aP1. The structure of archaeal uL10 is closely similar to bacterial uL10, whereas the NTDs of aP1 are each composed of four α -helices in two layers. The dimer interface is formed by the two N-terminal helices (Fig. 4).

In all species, this stalk of the ribosome is long and flexible. The archaeal uL10 C-terminal helix binds over the 2-fold symmetry axis of the dimer of aP1, like in the uL10–bL12 interaction. The first and the fourth helix of aP1 interact almost orthogonally with the long helix of uL10 (Fig. 5). The possible arrangement of the archaeal P proteins in the ribosome will not be much different from the situation in the bacterial ribosome. One difference is that the C-terminal domain of the P proteins is smaller and that there is a C-terminal extension of uL10 of the same nature as the C-terminal domain of the P1 and P2 proteins.

The trGTPases and their interactions with bL12 on the ribosome

The trGTPases

There are a number of trGTPases involved in protein synthesis (Table 1).

The trGTPases catalyze different steps of translation. These proteins, in their GTP-bound states catalyze different steps of translation on the ribosome. Further, they hydrolyze the bound GTP molecule and dissociate from the ribosome. The different roles of the trGTPases necessitate their different structures (Fig. 6), but they all have two domains in common, the GTP binding domain (G domain), normally at the N-terminus and the




Fig. 4. Comparison of the structure of the NTD dimer of bacterial bL12 with archaeal protein aP1 and eukaryotic eP1/P2 proteins adopted from the PDBs 1RQT, 3A1Y, and 2LBF, respectively. While the archaeal and eukaryotic domains are similar, in comparison with the bacterial domain it is difficult to identify a structural similarity apart from the helical structure.




Fig. 5. Structure of the archaeal stalk complex (PDB: 3A1Y). uL10 is colored blue, while three aP1 dimers bound to the spine helices are colored in salmon, light blue and lemon. The aP1 dimers bind to the long C-terminal helix of uL10.

Table 1. Some trGTPases involved in protein synthesis on the ribosome

Bacteria	Archaea and Eukarya	Functional role
bIF2	eIF5B	Association of large and small subunits during initiation
bEF-Tu	eEF-1A	Carrier of aminoacyl-tRNA to the decoding site
bSelB	eEF-Sec	Carrier of Se-Met-tRNA to the decoding site
bEF-G	eEF2	Translocation of peptidyl-tRNA from A-site to P-site and recycling of ribosomal subunits together with the ribosome recycling factor, RRF
bRF3	eRF3	Removal of release factors 1 and 2 from the ribosome. The bacterial protein is completely different from the archaeal and the eukaryotic proteins.
bLepA, EF4	mtLepA, cpLepA	Back translocation? Ribosome biogenesis?
BipA	TypA	Ribosome biogenesis?
Tet-O/M		Removal of tetracycline causing resistance

subsequent domain (domain II). Two of the trGTPases, EF-G and RF3, have an insert in the G-domain called G' domain (Fig. 6). EF4 and BipA are paralogs of EF-G, but among other differences, they lack the G' domain. For further structural and functional comparisons see Liljas and Ehrenberg (2013); Heller *et al.* (2017); Gibbs and Fredrick (2018).

The binding of trGTPases to the ribosome

The trGTPases in complex with GTP bind to overlapping sites on the ribosome (Fig. 7) (Heimark *et al.*, 1976). The two common domains of the factors, G and II, interact with the ribosome in similar ways. These interactions, identified by cryo-EM and

crystallography, show that the G-domains interact primarily with the sarcin-ricin loop (SRL) and the GTPase activating region (GAR) of the large subunit at the base of the bL12 stalk. The binding sites include RNA-helices H43 and H44 as well as proteins uL11 and uL6 (Gao *et al.*, 2009; Schmeing *et al.*, 2009). The anticodon-stem-loop of the tRNA in the ternary complex, EF-Tu-GTP-aa-tRNA interacts with the decoding region of the small subunit and so does the domain IV of EF-G. IF2, on the contrary, interacts initially only with the 30S subunit. When the subunits associate IF2 makes substantial contacts with the GAR of the 50S subunit. RF3 interacts mostly with the 50S subunit in a similar fashion as EF-Tu (Gao *et al.*, 2007; Pallesen *et al.*, 2013).




Fig. 6. The structure of the major trGTPases showing G and G' domains in green and cyan. The structures are adopted from the following PDBs: IF2 (PDB: 3JCN), EF-Tu (PDB: 4PC7), EF-G (PDB: 4V90) and RF3 (PDB: 4V89).




Fig. 7. Left - Elongation factor Tu (EF-Tu) in complex with an aminoacyl tRNA and a GTP analogue bound to the ribosome. Right - Elongation factor G with a GTP analogue bound to the ribosome. The similarity in binding is striking. The structures are adopted from the following PDBs: EF-Tu (PDB: 4V5R), EF-G (PDB: 4V5F).

What induces GTPase activity in the trGTPases?

The induction of GTPase activity in the trGTPases by the ribosome has been a challenging question. The trGTPases are part of a large family of GTPases where generally a GTPase activating protein (GAP) participates at the right stage and induces GTP hydrolysis. Furthermore, a residue in a loop of the GTPase, the so-called Switch II, has been thoroughly discussed. In most GTPases, this residue is a glutamine. Its ability to function as a general base is severely limited (Langen *et al.*, 1992). The corresponding residue in the trGTPases is a histidine, which would normally be able to function as a general base. However, since the GTPases have a common evolution it is most likely that the mechanism for GTP hydrolysis is the same, and that the histidine has a different role.

Several investigations have focused on the question of whether bL12 could be the ribosomal GAP and whether the histidine in Switch II of the trGTPases could function as a general base. Previously, it was proposed that isolated bL12 could independently stimulate GTP hydrolysis of EF-G on the ribosome unlike EF-Tu (Savelsbergh *et al.*, 2000). Subsequent kinetic studies revealed that depletion of bL12 from the 70S ribosome selectively affects stimulation of GTP hydrolysis in both EF-Tu and EF-G under multiple turnovers (Mohr *et al.*, 2002; Savelsbergh *et al.*, 2005) or single turnover condition (Diaconu *et al.*, 2005). However, it was demonstrated that for GTPase stimulation on IF2, bL12 does not play any role (Huang *et al.*, 2010). Also, high-resolution structures of EF-Tu and EF-G on the ribosome demonstrated that bL12 is not in contact with the G-domains of the trGTPases (Voorhees *et al.*, 2010; Carlson *et al.*, 2017). Therefore, the precise role of bL12 in the context of GTP hydrolysis by trGTPases remains elusive and awaits further investigation.

For a considerable time, the SRL of the large subunit has also been considered to be an important region of the ribosome for factor interactions. Chemical footprinting showed that both EF-Tu and EF-G protect this loop (Moazed *et al.*, 1988). Crystallographic experiments on EF-Tu bound to the ribosome have made it clear that the phosphate of adenine 2662 in the SRL is critically involved in GTP hydrolysis (Voorhees *et al.*, 2010). It was shown that the phosphate of A2662 hydrogen bonds to His84 in EF-Tu and brings it up against the nucleophilic water molecule that is moved into close proximity of the γ -phosphate of the GTP molecule. His84 was suggested to act like a general base (Berchtold *et al.*, 1993; Voorhees *et al.*, 2010). However, insightful mechanistic analysis supported by *in*




Fig. 8. A possible mechanism for GTP hydrolysis by the trGTPases. The phosphate of A2662 of SRL accepts a hydrogen bond from His 84 of the switch II of EF-Tu and positions it next to the water molecule close to the γ -phosphate. Both side chain nitrogens of the histidine are protonated making the histidine positively charged. The water molecule (red) donates a proton to the γ -phosphate leading to an in-line attack by the hydroxyl ion on the γ -phosphate (Liljas *et al.*, 2011; Åqvist & Kamerlin, 2015). The structure was adopted from PDB: 4V5L (Voorhees *et al.*, 2010).

silico modeling and molecular dynamics (MD) simulations suggest that it is more likely that the water molecule, possibly through a second water molecule, donates one proton to the γ -phosphate and attacks the phosphate by an in-line mode, which leads to the hydrolysis of the γ -phosphate (Fig. 8; Liljas *et al.*, 2011; Åqvist and Kamerlin, 2015; Mondal and Warshel, 2018). This mechanism is probably universal for the trGTPases. Thus, the phosphate of an adenine in the SRL replaces the GTPase activating protein (GAP) for trGTPases in general.

The role of bL12 in the functions of the trGTPases

The removal of the CTD of bL12 has the same effect as removing the whole protein on the activity of trGTPases (van Aghoven *et al.*, 1975; Koteliansky *et al.*, 1978). Thus, it was concluded that bL12 interacts with the trGTPases through its CTD. Ribosomes with only one bL12 dimer are partially active (Griaznova and Traut, 2000; Mandava *et al.*, 2012). Ribosomes where one CTD of each bL12 dimer was removed also retain activity (Oleinikov *et al.*, 1998). The removal of all bL12 delays the release of inorganic phosphate from EF-G after GTP hydrolysis (Mohr *et al.*, 2002). If the hinge of bL12 is significantly




Fig. 9. Illustration of the interaction sites of the four major trGTPases IF2 (a), EF-Tu (b), EF-G (c) and RF3 (d) on the bL12 CTD as identified by NMR mapping (Helgstrand *et al.*, 2007). The top panel represents the ribbon diagram and the bottom panel represents the surface illustration of bL12-CTD. The interaction sites are highlighted in red.

shortened, the ribosomes behave as if there was no CTD, but essentially doubling the length of the hinge has no negative effect (Gudkov *et al.*, 1991; Bubunencko *et al.*, 1992).

We are faced with the dilemma that bL12 is important for GTP hydrolysis, but apparently not directly involved in the catalytic mechanism. What does it do? Which part of bL12CTD interacts with the factors and where on the factors does the interaction occur?

As mentioned before, bL12 was proposed to be responsible for GTPase activation of EF-G and EF-Tu (Savelsbergh *et al.*, 2000; Mohr *et al.*, 2002; Savelsbergh *et al.*, 2005). However, with the advancement of the field with high-resolution structures and biochemistry, it became evident that bL12 is not a GAP. Rather it could act as a ‘propeller’ like instrument with multiple extended arms for continuous binding and recruitment of the trGTPases, especially EF-Tu to the ribosome (Diaconu *et al.*, 2005). However, so far there is no direct proof to this claim. Only a recent fluorescence-based localization study indicated simultaneous binding of several EF-Tu on each ribosome (Mustafi and Weisshaar, 2018). If true, this would mean that bL12 increases the local concentration of the translation factors, especially EF-Tu in a ternary complex with aminoacylated tRNAs to ensure their constant supply essential for a fast rate of translation.

Mutations of V66, I69, K70, and R73 of the CTD of *E. coli* bL12 lead to an increased K_M for the binding of EF-G but have little effect on GTP hydrolysis. Furthermore, these mutations inhibit the release of inorganic phosphate after GTP hydrolysis by EF-G (Savelsbergh *et al.*, 2005). Similarly, mutations of K65, V66, I69, K70, R73, and K84 have significant effects on ribosome binding of EF-Tu, and as a consequence of that, on GTP hydrolysis (Diaconu *et al.*, 2005). These mutational studies agree well with the NMR experiments suggesting that IF2, EF-Tu, EF-G, and RF3 all interact with helices 4 and 5 of bL12 CTD, in particular with residues V66, A67, V68, K70, G79, L80, E82 (Helgstrand *et al.*, 2007; Fig. 9). This surface corresponds with the conserved surface area of the CTD domain of bL12 (Leijonmarck and Liljas, 1987).

EF-G interacts with the CTD of bL12 through its G' domain as seen by crosslinking, cryo-EM and crystallography (Fig. 10; Datta *et al.*, 2005; Nechifor & Wilson, 2007; Harms *et al.*, 2008; Gao *et al.*, 2009; Tourigny *et al.*, 2013; Brilot *et al.*, 2013; Chen *et al.*, 2013). This is true also for RF3 (Pallesen *et al.*, 2013). For the trGTPases lacking the G'-domain, EF-Tu (Stark *et al.*, 2000), IF2 (Allen *et al.*, 2005; Simonetti *et al.*, 2013; Ge *et al.*, 2018) and BipA (Kumar *et al.*, 2015), bL12 interacts with the G domain.



Fig. 10. The interaction of the G' and G domains of EF-G with bL12CTD (red) on *T. thermophilus* 70S ribosomes (PDB: 4V5F) (Gao *et al.*, 2009).

The main role for the bacterial trGTPase IF2 is to promote efficient ribosomal subunit association. bIF2 forms a preinitiation complex with 30S subunit programmed with mRNA and fMet-tRNA^{fMet}. bL12 is needed for rapid factor-dependent subunit association with a 70S initiation complex (Huang *et al.*, 2010; Mandava *et al.*, 2012). Cryo-EM and crosslinking have shown that bL12 interacts with the G domain of IF2 (Heimark *et al.*, 1976; Allen *et al.*, 2005). Recently, Ge *et al.* (2018) analyzed a range of mutants in bL12 and IF2 to identify the sites involved in subunit association (Fig. 11). Mutations of the conserved K65, K70, R73, and K81 on the $\alpha 4$ - $\alpha 5$ helices of bL12 caused a significant defect in subunit association. Similarly, mutations in D506, D508, E522, and E523 of G4-G5 helix of IF2 were found to be deleterious for fast subunit association. It can be easily noticed that the identified residues of bL12 and IF2 are basic and acidic in nature, respectively. Moreover, mutations to oppositely charged amino acids on both bL12 and IF2 caused largest defects, which could be rescued by using charge-swapped mutants on both the interacting partners. This mutational analysis pinpointed the amino acids on bL12 and IF2, which are the molecular determinants for their interaction. In addition, MD simulations identified the details of the interactions of these residues at the molecular level. It suggested several salt-bridge interactions between the bL12 CTD and the G domain of IF2 (Fig. 11). Based on the analysis, one can conclude that charge complementarity is the basis of




Fig. 11. The interaction of bL12 CTD (red) with IF2 on the 70S ribosome as suggested from the mutational studies combined with MD simulations (Ge *et al.*, 2018).

bL12 and IF2 interaction (Ge *et al.*, 2018). Moreover, it can be inferred that bL12 (CTD) on the 50S and IF2 on 30S-preinitiation complex act like recognition partners; their initial interaction leads to the rapid association of the ribosomal subunits. Based on the similarity of the G-domains of the trGTPases it can be speculated that the other trGTPases too interact with bL12 CTD through a similar mechanism based on electrostatic interaction. The available cryo-EM based pictures of bL12 interaction with RF3 (Pallesen *et al.*, 2013) and EF-G (Gao *et al.*, 2009) support that model.

Carlson *et al.*, (2017) have revisited some of the old experiments after significantly improved purification of the experimental components. First of all they used *E. coli* JE28 ribosomes in which all bL12 proteins carry chromosomally fused C-terminal His-tag (Ederth *et al.*, 2009). With the aid of the His-tag, they ensured that bL12 was completely removed from the ribosomes. They also used an extensive denaturation and renaturation based method for purification of bL12 more extensively. In their study, bL12 did not induce GTPase activity in the isolated EF-G in contrast to what was reported earlier by Savelsbergh *et al.*, 2000. However, the GTPase activity by EF-G on the ribosome requires the presence of bL12. The binding of EF-G in complex with the non-hydrolysable GTP analogue GDPNP to the ribosome was dramatically reduced if bL12 was absent. If the G' domain was removed from EF-G the GTPase activity was much reduced and the presence or absence of bL12 in the ribosome did not make any difference. Likewise, the binding of RF3 and IF2 to ribosomes also required the presence of bL12 (Carlson *et al.*, 2017).

The interaction of P-proteins with trGTPases

In archaeal or eukaryotic translation the P-proteins are essential for the function of the trGTPases. However, all P1/P2 dimers can be removed despite maintaining the GTPase activity. This activity can be maintained by the C-terminus of uL10 alone. However, the lack of some or all of the P1/P2 dimers leads to losses in fidelity and a different subset of mRNAs is translated (Santos & Ballesta, 1994; Remacha *et al.*, 1995; Wawiórka *et al.*, 2017). This may suggest a complex role of these proteins but primarily relates to the decoding process where different complexes of EF1A with GTP and aminoacylated tRNAs are tested against the current codon in the A-site. The rate needed for efficient discrimination of non-cognate and near-cognate tRNAs cannot be met by a single C-terminus.

With regard to the CTD of the P-proteins, there has been a lack of information about its structure and interaction with the




Fig. 12. The C-terminal (C11) peptide of aP1 (red) binds to the archaeal EF2 in a hydrophobic groove between the G (green) and G' (blue) domains (PDB ID: 5H7L) (Tanzawa *et al.*, 2018).

trGTPases. Recently Tanzawa *et al.* (2018) studied the binding of a C-terminal peptide of aP1 to aEF2. The peptide takes the conformation of a helix, which binds to the G4 and G5 elements of the factor similar to the observations in case of bacterial IF2 (Fig. 12) (Ge *et al.*, 2018). However, in contrast to the charge complementarity-based interaction of bL12-CTD and IF2, the 11 amino acid long C-terminal helix of aP1 (aP1C11) binds to aEF2a in a hydrophobic groove.

Ribosome inhibitory proteins high-jack the ribosomal stalk for binding

There is a range of ribosome inhibitory proteins (RIP) that generally modify the part of SRL, which is important for the GTPase activity. In some cases, like ricin, a central nucleotide is depurinated and in other cases, like alpha-sarcin, the large ribosomal RNA is cut. The RIPs are evolutionary related, but the sequence conservation is quite low. Interestingly, without any sequence or structural relationship to the trGTPases, the RIPs like the trGTPases benefit in binding to the ribosome from the presence of the P-proteins. The binding is through hydrophobic interactions of the very C-terminal 6–9 residues of the P-proteins (Shi *et al.*, 2016).

Conclusions

The stalk proteins have very distinctive roles primarily in relation to the trGTPases. However, it remains enigmatic what their actual

role is in relation to the activation of GTP hydrolysis in trGTPases. There are at least two options generally discussed:

- (A) The ribosomal stalk, with multiple arms bearing the C-terminal structures of bL12 or the P-proteins, recruits trGTPases to their binding area on the ribosome by binding and thereby increasing the local concentration of trGTPases. This would, in turn, increase the rate of GTP hydrolysis.
- (B) The ribosomal stalk binds the trGTPases on the ribosome and stabilizes their binding to the factor-binding site at the SRL loop on the ribosome thereby increasing the rate of GTP hydrolysis.

The first mechanism (A) is most widely accepted by the ribosome field at present. However, relevant experimental proof supporting this mechanism is still lacking. If the native-like high rate of GTP hydrolysis could be achieved by adding an increasing concentration of the trGTPases with a bL12-depleted ribosome, then this mechanism would be proven true. That would then justify the need for multiple bL12-CTDs for increasing the local concentration of the trGTPases for high GTPase rates. However, the recent study with super-resolution microscopy infers that the four bL12 proteins are essentially saturated with EF-Tu ternary complexes *in vivo* (Mustafi and Weisshaar, 2018). This observation again speaks in favor of the factor recruitment model.

Experimental evidence for the second mechanism (B) is also scarce. If a high GTPase rate can be achieved by titrating L12-CTDs with a bL12-depleted ribosome, it might support model (B). It would mean that the bL12-CTD interaction is important for stable binding of the trGTPases in the vicinity of the SRL and for attaining the correct conformation for GTPase activation. Carlson *et al.* (2017) found reduced occupancy of EF-G-GDPNP on the ribosome when bL12 is deleted. This result clearly indicates the need of bL12, presumably the CTD, for stable binding of the trGTPases on the ribosome. The case of the initiation factors, bacterial IF2 or archaeal or eukaryotic eIF5B illustrates another aspect of the bL12 CTD (Ge *et al.*, 2018). These factors initially bind to the small subunit and need the stalk proteins for association with the large subunit. These results also clearly support the B alternative. It is also possible that the ribosomal stalk proteins use different or specialized modes of action with respect to different translational GTPases. Its role in the function of other translation factors also remains elusive.

From our discussions of the available literature regarding bL12 it becomes obvious that despite about 50 years of research, the role of the ribosomal stalk remains somewhat enigmatic. On the contrary, the multifaceted research on bL12 and the P proteins has brought us much further into this intriguing story.

Acknowledgements. We are thankful to Dr. Chandra Sekhar Mandava and Dr. Xueliang Ge for helping with the artwork used in this article. SS acknowledges research funding support from Swedish Research Council (2014-4423 and Research Environment Grant 2016-06264), and Knut and Alice Wallenberg Foundation (KAW 2017-0055).

References

- Agrawal RK, Heagle AB, Penczek P, Grassucci RA and Frank J (1999) EF-G-dependent GTP hydrolysis induces translocation accompanied by large conformational changes in the 70S ribosome. *Nature Structural Biology* **6**, 643–647.
- Allen GS, Zavialov A, Gursky R, Ehrenberg M and Frank J (2005) The cryo-EM structure of a translation initiation complex from *Escherichia coli*. *Cell* **121**, 703–712.
- Åqvist J and Kamerlin S (2015) The conformation of a catalytic loop is central to GTPase activity on the ribosome. *Biochemistry* **54**, 546–556.
- Bailey-Serres J, Vangala S, Szick K and Lee CH (1997) Acidic phosphoprotein complex of the 60S ribosomal subunit of maize seedling roots. Components and changes in response to flooding. *Plant Physiology* **114**, 1293–1305.
- Ballesta JP and Remacha M (1996) The large ribosomal subunit stalk as a regulatory element of the eukaryotic translational machinery. *Progress in Nucleic Acid Research and Molecular Biology* **55**, 157–193.
- Ballesta JP, Rodriguez-Gabriel MA, Bou G, Briones E, Zambrano R and Remacha M (1999) Phosphorylation of the yeast ribosomal stalk. Functional effects and enzymes involved in the process. *FEMS Microbiological Reviews* **23**, 537–550.
- Ban N, Nissen P, Hansen J, Moore PB and Steitz TA (2000) The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science* **289**, 905–920.
- Ban N, Beckmann R, Cate JHD, Dinman JD, Dragon F, Ellis SR, Lafontaine DLJ, Lindahl L, Liljas A, Lipton JM, McAleer MA, Moore PB, Noller HF, Ortega J, Panse VG, Ramakrishnan V, Spahn CMT, Steitz TA, Tchorzewski M, Tollervey MD, Warren AJ, Williamson JR, Wilson D, Yonath A and Yusupov M (2014) A new system for naming ribosomal proteins. *Current Opinion in Structural Biology* **24**, 1–5.
- Bargis-Surgey P, Lavergne JP, Gonzalo P, Vard C, Filhol-Cochet O and Reboud JP (1999) Interaction of elongation factor eEF-2 with ribosomal P proteins. *European Journal of Biochemistry* **262**, 606–611.
- Berchtold H, Reshetnikova L, Reiser CO, Schirmer NK, Sprinzl M and Hilgenfeldt R (1993) Crystal structure of active elongation factor Tu reveals major domain rearrangements. *Nature* **365**, 126–132.
- Bernadó P, Modig K, Grell P, Svergun D, Tchorzewski M, Pons M and Akke M (2010) Structure and dynamics of ribosomal protein L12: an ensemble model based on SAXS and NMR relaxation. *Biophysical Journal* **98**, 2374–2382.
- Bocharov EV, Gudkov AT and Arseniev AS (1996) Topology of the secondary structure elements of ribosomal protein L7/L12 from *E. coli* in solution. *FEBS Letters* **379**, 291–294.
- Bocharov EV, Gudkov AT, Budovskaya EV and Arseniev AS (1998) Conformational independence of N- and C-domains in ribosomal protein L7/L12 and in the complex with protein L10. *FEBS Letters* **423**, 347–350.
- Bocharov EV, Sobol AG, Pavlov KV, Korzhnev DM, Jaravine VA, Gudkov AT and Arseniev AS (2004) From structure and dynamics of protein L7/L12 to molecular switching in ribosome. *Journal of Biological Chemistry* **279**, 17697–17706.
- Boublik M, Hellmann W and Roth HE (1976) Localization of ribosomal proteins L7/L12 in the 50S subunit of *Escherichia coli* ribosomes by electron microscopy. *Journal of Molecular Biology* **107**, 479–490.
- Brilot AF, Korostelev AA, Ermolenko DN and Grigorieff N (2013) Structure of the ribosome with elongation factor G trapped in the pretranslocation state. *Proceedings of the National Academy of Sciences USA* **110**, 20994–20999.
- Bubunenko MG, Chuikov SV and Gudkov AT (1992) The length of the interdomain region of the L7/L12 protein is important for its function. *FEBS Letters* **313**, 232–234.
- Bushuev VN, Gudkov AT, Liljas A and Sepetov NF (1989) The flexible region of protein L12 from bacterial ribosomes studied by proton nuclear magnetic resonance. *Journal of Biological Chemistry* **264**, 4498–4505.
- Carlson MA, Bassam GH, Weis AJ, Colby SB, Shelton CD, Wuerth ME, Walter JD and Spiegel Jr PC (2017) Ribosomal protein L7/L12 is required for GTPase translation factors EF-G, RF3, and IF2 to bind in the GTP state to 70S ribosomes. *FEBS Journal*, **284**, 1631–1643.
- Chen Y, Feng S, Kumar V, Ero R and Gao Y-G (2013) Structure of EF-G-ribosome complex in a pretranslocation state. *Nature Structural and Molecular Biology* **20**, 1077–1084.
- Christodoulou J, Larsson G, Fucini P, Connell SR, Pertinhez TA, Hanson CL, Redfield C, Nierhaus KH, Robinson CV, Schleucher J and Dobson CM (2004) Heteronuclear NMR investigations of dynamic regions of intact *Escherichia coli* ribosomes. *Proceedings of the National Academy of Sciences USA* **101**, 10949–10954.
- Cohlberg JA and Nomura M (1976) Reconstitution of *Bacillus stearothermophilus* 50S ribosomal subunits from purified molecular components. *Journal of Biological Chemistry* **251**, 209–221.

- Cowgill CA, Nichols BG, Kenny JW, Butler P, Bradbury EM and Traut RR (1984) Mobile domains in ribosomes revealed by proton nuclear magnetic resonance. *Journal of Biological Chemistry* **259**, 15257–15263.
- Datta PP, Sharma MR, Qi L, Frank J and Agrawal RK (2005) Interaction of the G' domain of elongation factor G and the C-terminal domain of ribosomal protein L7/L12 during translocation as revealed by cryo-EM. *Molecular Cell* **20**, 723–731.
- Dey D, Bochkariov DE, Jokhadze GG and Traut RR (1998) Cross-linking of selected residues in the N- and C-terminal domains of *Escherichia coli* protein L7/L12 to other ribosomal proteins and the effect of elongation factor Tu. *Journal of Biological Chemistry* **273**, 1670–1676.
- Diaconu M, Kothe U, Schlunzen F, Fischer N, Harms JM, Tonevitsky AG, Stark H, Rodnina MV and Wahl MC (2005) Structural basis for the function of the ribosomal L7/L12 stalk in factor binding and GTPase activation. *Cell* **121**, 991–1004.
- Ederth J, Mandava CS, Dasgupta S and Sanyal S (2009) A single-step method for purification of active His-tagged ribosomes from a genetically engineered *Escherichia coli*. *Nucleic Acids Research* **37**, e15.
- Gao H, Zhou Z, Rawat U, Huang C, Bouakaz I, Wang C, Cheng Z, Liu Y, Zavalov A, Gursky R, Sanyal S, Ehrenberg M, Frank J and Song H (2007) RF3 induces ribosomal conformational changes responsible for dissociation of class I release factors. *Cell* **129**, 929–941.
- Gao YG, Selmer M, Dunham CM, Weixlbaumer A, Kelley AC and Ramakrishnan V (2009) The structure of the ribosome with elongation factor G trapped in the posttranslocational state. *Science* **326**, 694–699.
- Ge X, Mandava CS, Lind C, Åqvist J and Sanyal S (2018) Complementary charge-based interaction between the ribosomal-stalk protein L7/L12 and IF2 is the key to rapid subunit association. *Proceedings of the National Academy of Sciences USA* **115**, 4649–4654.
- Gibbs MR and Fredrick K (2018) Roles of elusive translational GTPases come to light and inform on the process of ribosome biogenesis in bacteria. *Molecular Microbiology* **107**, 445–454.
- Gonzalo P and Reboud J-P (2003) The puzzling lateral flexible stalk of the ribosome. *Biology of the Cell* **95**, 179–193.
- Griaznova O and Traut RR (2000) Deletion of C-terminal residues of *Escherichia coli* ribosomal protein L10 causes the loss of binding of one L7/L12 dimer: ribosomes with one L7/L12 dimer are active. *Biochemistry* **39**, 4075–4081.
- Guarinos E, Remacha M and Ballesta JP (2001) Asymmetric interactions between the acidic P1 and P2 proteins in the *Saccharomyces cerevisiae* ribosomal stalk. *Journal of Biological Chemistry* **276**, 32474–32479.
- Gudkov AT (1997) The L7/L12 domain of the ribosome: structural and functional studies. *FEBS Letters* **407**, 253–256.
- Gudkov AT, Gongadze GM, Bushuev VN and Okon MS (1982) Proton nuclear magnetic resonance study of the ribosomal protein L7/L12 *in situ*. *FEBS Letters* **138**, 229–232.
- Gudkov AT, Bubunenkov M and Gryaznova O (1991) Overexpression of L7/L12 protein with mutations in its flexible region. *Biochimie* **73**, 1387–1389.
- Hamel E, Koka M and Nakamoto T (1972) Requirement of an *E. coli* 50S ribosomal protein component for effective interaction of the ribosome with T and G factors with guanosine triphosphate. *Journal of Biological Chemistry* **247**, 805–814.
- Hardy SJS (1975) The stoichiometry of the ribosomal proteins of *Escherichia coli*. *Molecular and General Genetics* **140**, 253–274.
- Harms JM, Wilson DN, Schlunzen F, Connel SL, Stachelhaus T, Zaborowska Z, Spahn CMT and Fucini P (2008) Translation regulation via L11: molecular switches on the ribosome turned on and off by thiostrepton and micrococin. *Cell* **30**, 26–38.
- Heimark RL, Hershey JW and Traut RR (1976) Crosslinking of initiation factor IF2 to proteins L7/L12 in 70S ribosomes of *Escherichia coli*. *Journal of Biological Chemistry* **251**, 7779–7784.
- Helgstrand M, Mandava CS, Mulder FA, Liljas A, Sanyal S and Akke M (2007) The ribosomal stalk binds to translation factors IF2, EF-Tu, EF-G and RF3 via a conserved region of the L12 C-terminal domain. *Journal of Molecular Biology* **365**, 468–479.
- Heller JLE, Kamalampeta R and Wieden H-J (2017) Taking a step back from back-translocation: an integrative view of LepA/EF4's cellular function. *Molecular and Cellular Biology* **37**, e00653–16.
- Huang C, Mandava CS and Sanyal S (2010) The ribosomal stalk plays a key role in IF2-mediated association of the ribosomal subunits. *Journal of Molecular Biology* **399**, 145–153.
- Ilag LL, Videler H, McKay AR, Sobott F, Fucini P, Nierhaus KH and Robinson CV (2005) Heptameric (L12)₆/L10 rather than canonical pentameric complexes are found by tandem MS of intact ribosomes from thermophilic bacteria. *Proceedings of the National Academy of Sciences USA* **102**, 8192–8197.
- Kaltschmidt E and Wittmann HG (1969) Ribosomal proteins: VI. Preparative polyacrylamide gel electrophoresis as applied to the isolation of ribosomal proteins. *Analytical Biochemistry* **36**, 401–412.
- Kaltschmidt E and Wittmann HG (1970) Ribosomal proteins. XII. Number of proteins in small and large ribosomal subunits of *Escherichia coli* as determined by two-dimensional gel electrophoresis. *Proceedings of the National Academy of Sciences USA* **67**, 1276–1281.
- Kischa K, Möller W and Stöffler G (1971) Reconstitution of a GTPase activity by a 50S ribosomal protein from *E. coli*. *Nature New Biology* **233**, 62–63.
- Kotliansky VE, Domogatsky SP and Gudkov AT (1978) Dimer state of protein L7/L12 and EF-G dependent reactions on ribosomes. *European Journal of Biochemistry* **90**, 319–323.
- Kravchenko O, Mitroshin I, Nikonov S, Piendl W and Garber M (2010) Structure of a two-domain N-terminal fragment of ribosomal protein L10 from *Methanococcus jannaschii* reveals a specific piece of the archaeal ribosomal stalk. *Journal of Molecular Biology* **399**, 214–220.
- Kumar V, Chen Y, Ero R, Ahmed T, Tan J, Li Z, Wong AS, Bhushan S and Gao YG (2015) Structure of BipA in GTP form bound to the ratcheted ribosome. *Proceedings of the National Academy of Sciences USA* **112**, 10944–10949.
- Lake JA (1976) Ribosome structure determined by electron microscopy of *Escherichia coli* small subunits, large subunits and monomeric ribosomes. *Journal of Molecular Biology* **105**, 131–139.
- Langen R, Schweins T and Warshel A (1992) On the mechanism of guanine triphosphate hydrolysis in *ras* p21 proteins. *Biochemistry* **31**, 8691–8696.
- Leijonmarck M and Liljas A (1987) Structure of the C-terminal domain of the ribosomal protein L7/L12 from *Escherichia coli* at 1.7 Å. *Journal of Molecular Biology* **195**, 555–579.
- Leijonmarck M, Eriksson S and Liljas A (1980) Crystal structure of a ribosomal component at 2.6 Å resolution. *Nature* **286**, 824–826.
- Liljas A (1982) Structural studies of ribosomes. *Progress in Biophysics and Molecular Biology* **40**, 161–228.
- Liljas A (1991) Comparative biochemistry and biophysics of ribosomal proteins. *International Review of Cytology* **124**, 103–136.
- Liljas A and Ehrenberg M (2013) *Structural Aspects of Protein Synthesis*. 2nd Edn. Singapore: World Scientific, ISBN 978-9814313209.
- Liljas A, Kirsebom LA and Leijonmarck M (1986) Structural studies of the factor binding domain. In *Structure, Function and Genetics of Ribosomes*. B Hardesty and G Kramer (eds), New York: Springer-Verlag, pp. 379–390.
- Liljas A, Ehrenberg M and Åqvist J (2011) Comment on “The mechanism for activation of GTP hydrolysis on the ribosome. *Science* **333**, 37.
- Lin J, Gagnon MG, Buckley D and Steitz TA (2015) Conformational changes of elongation factor G on the ribosome during tRNA translocation. *Cell* **160**, 219–227.
- Maki Y, Hashimoto T, Zhou M, Naganuma T, Ohta J, Nomura T, Robinson CV and Uchiyama T (2007) Three binding sites for stalk protein dimers are generally present in ribosomes from archaeal organism. *Journal of Biological Chemistry* **282**, 32827–32833.
- Mandava CS, Peisker K, Ederth J, Kumar R, Ge X, Szaflarski W and Sanyal S (2012) Bacterial ribosome requires multiple L12 dimers for efficient initiation and elongation of protein synthesis involving IF2 and EF-G. *Nucleic Acids Research* **40**, 2054–2064.
- Moazed D, Robertson JM and Noller HF (1988) Interaction of elongation factors EF-G and EF-Tu with a conserved loop in 23S RNA. *Nature* **334**, 362–364.
- Mohr D, Wintermeyer W and Rodnina MV (2002) GTPase activation of elongation factors Tu and G on the ribosome. *Biochemistry* **41**, 12520–12528.
- Mondal D and Warshel A (2018) EF-Tu and EF-G are activated by allosteric effects. *Proceedings of the National Academy of Sciences USA* **115**, 3386–3391.

- Mulder FAA, Bauakaz L, Lundell A, Liljas A, Akke M and Sanyal SC (2004) Conformation and dynamics of protein L12 in solution and on the ribosome. *Biochemistry* **43**, 5930–5936.
- Mustafi M and Weisshaar JC (2018) Simultaneous binding of multiple EF-Tu copies to translating ribosomes in live *Escherichia coli*. *mBio* **16**, e02143–17.
- Naganuma T, Nomura N, Yao M, Mochizuki M, Uchiyama T and Tanaka I (2010) Structural basis for translation factor recruitment to the eukaryotic/archaeal ribosomes. *Journal of Biological Chemistry* **285**, 4747–4756.
- Nechifor R and Wilson KS (2007) Crosslinking of translation factor EF-G to proteins of the bacterial ribosome before and after translocation. *Journal of Molecular Biology* **368**, 1412–1425.
- Oleinikov AV, Jokhadze GG and Traut RR (1998) A single-headed dimer of *Escherichia coli* ribosomal protein L7/L12 supports protein synthesis. *Proceedings of the National Academy of Sciences USA* **95**, 4215–4218.
- Österberg R, Sjöberg B, Liljas A and Pettersson I (1976) Small angle X-ray scattering and cross-linking study of the proteins from *Escherichia coli* ribosomes. *FEBS letters* **66**, 48–51.
- Österberg R, Sjöberg B, Pettersson I, Liljas A and Kurland CG (1977) Small-angle scattering study of the protein complex of L7/L12 and L10 from *Escherichia coli* ribosomes. *FEBS Letters* **73**, 22–24.
- Pallesen J, Hashem Y, Korkmaz G, Koripella RK, Huang C, Ehrenberg M, Sanyal S and Frank J (2013) Cryo-EM visualization of the ribosome in termination complex with apo-RF3 and RF1. *Elife* **2**, e00411.
- Pettersson I and Kurland CG (1980) Ribosomal protein L7/L12 is required for optimal translation. *Proceedings of the National Academy of Sciences USA* **77**, 4007–4010.
- Pettersson I, Hardy SJS and Liljas A (1976) The ribosomal protein L8 is a complex of L7/L12 and L10. *FEBS Letters* **64**, 135–138.
- Remacha M, Jimenez-Diaz A, Bermejo B, Rodriguez-Gabriel MA, Guarinos E and Ballesta JP (1995) Ribosomal acidic phosphoproteins P1 and P2 are not required for cell viability but regulate the pattern of protein expression in *Saccharomyces cerevisiae*. *Molecular Cellular Biology* **15**, 4754–4762.
- Santos C and Ballesta JPG (1994) Ribosomal protein P0, contrary to phosphoproteins P1 and P2, is required for ribosome activity and *Saccharomyces cerevisiae* viability. *Journal of Biological Chemistry* **269**, 15689–15696.
- Santos C and Ballesta JP (1995) The highly conserved protein P0 carboxyl end is essential for ribosome activity only in the absence of proteins P1 and P2. *Journal of Biological Chemistry* **270**, 20608–20614.
- Sanyal CS and Liljas A (2000) The end of the beginning: structural studies of ribosomal proteins. *Current Opinion in Structural Biology* **10**, 633–636.
- Savelsbergh A, Mohr D, Wilden B, Wintermeyer W and Rodnina MV (2000) Stimulation of the GTPase activity state of translation elongation factor G by protein L7/L12. *Journal of Biological Chemistry* **275**, 890–894.
- Savelsbergh A, Mohr D, Kothe U, Wintermeyer W and Rodnina MV (2005) Control of phosphate release from elongation factor G by ribosomal protein L7/L12. *EMBO Journal* **24**, 4316–4323.
- Schmeing TM, Voorhees RM, Kelley AC, Gao YG, Murphy FV, Weir JR and Ramakrishnan V (2009) The crystal structure of the ribosome bound to EF-Tu and aminoacyl-tRNA. *Science* **326**, 688–694.
- Shi WW, Mak ANS, Wong KB and Shaw PC (2016) Structures and ribosomal interactions of ribosome-inactivating proteins. *Molecules* **21**, 1–13.
- Shimmin LC, Ramirez C, Matheson AT and Dennis PP (1989) Sequence alignment and evolutionary comparison of the L10 equivalent and L12 equivalent ribosomal proteins from archaeobacteria, eubacteria, and eukaryotes. *Journal of Molecular Evolution* **29**, 448–462.
- Simonetti A, Marzi S, Billas IM, Tsai A, Fabbretti A, Myasnikov AG, Roblin P, Vaiana AC, Hazemann I, Eiler D, Steitz TA, Puglisi JD, Gualerzi CO and Klaholz BP (2013) Involvement of protein IF2 N domain in ribosomal subunit joining revealed from architecture and function of the full-length initiation factor. *Proceedings of the National Academy of Sciences USA* **110**, 15656–15661.
- Stark H, Rodnina MV, Wieden HJ, van Heel M and Wintermeyer W (2000) Large-scale movement of elongation factor G and extensive conformational change of the ribosome during translocation. *Cell* **100**, 301–309.
- Stöffler G, Hasenbank R, Bodley JW and Highland JH (1974) Inhibition of protein L7/L12 binding to 50S ribosomal cores by antibodies specific for proteins L6, L10 and L18. *Journal of Molecular Biology* **86**, 171–174.
- Strycharz WA, Nomura M and Lake JA (1978) Ribosomal protein L7/L12 localized at a single region of the large subunit by immune microscopy. *Journal of Molecular Biology* **126**, 123–140.
- Subramanian AR (1975) Copies of proteins L7 and L12 and heterogeneity of the large subunit of *Escherichia coli* ribosomes. *Journal of Molecular Biology* **95**, 1–8.
- Tanzawa T, Kato K, Girodat D, Ose T, Kumakura Y, Wieden HJ, Uchiyama T, Tanaka I and Yao M (2018) The C-terminal helix of ribosomal P stalk recognizes a hydrophobic groove of elongation factor 2 in a novel fashion. *Nucleic Acids Research* **46**, 3232–3244.
- Tchorzewski M (2002) The acidic ribosomal P proteins. *International Journal of Biochemistry and Cellular Biology* **34**, 911–915.
- Tchorzewski M, Boguszewska A, Dukowski P and Grankowski N (2000a) Oligomerization properties of the acidic ribosomal P-proteins from *Saccharomyces cerevisiae*: effect of P1A protein phosphorylation on the formation of the P1A-P2B heterocomplex. *Biochimica et Biophysica Acta* **1499**, 63–73.
- Tchorzewski M, Boldyreff B, Issinger O and Grankowski N (2000b) Analysis of the protein-protein interactions between the human acidic ribosomal P-proteins: evaluation by the two hybrid system. *International Journal of Biochemistry and Cellular Biology* **32**, 737–746.
- Terhorst C, Wittmann-Liebold B and Möller W (1972) 50-S ribosomal proteins. Peptide studies on two acidic proteins, A 1 and A 2, isolated from 50-S ribosomes of *Escherichia coli*. *European Journal of Biochemistry* **25**, 13–19.
- Tourigny DS, Fernández IS, Kelley AC and Ramakrishnan V (2013) Elongation factor G bound to the ribosome in an intermediate of translocation. *Science* **340**, 1235490.
- Traut RR, Dey D, Bochkariov D, Oleinikov AV, Jokhadze GG, Hamman BD and Jameson DM (1995) Location and domain structure of *Escherichia coli* ribosomal protein L7/L12: site specific cysteine crosslinking and attachment of fluorescent probes. *Biochemistry and Cell Biology* **73**, 949–958.
- Tritton TR (1980) Proton NMR observation of the *Escherichia coli* ribosome. *FEBS Letters* **120**, 141–144.
- Uchiyama T, Wabha AJ and Traut RR (1987) Topography and stoichiometry of acidic proteins in large ribosomal subunits from *Artemia salina* as determined by crosslinking. *Proceedings of the National Academy of Sciences USA* **84**, 5580–5584.
- Valle M, Zavialov A, Li W, Stagg SM, Sengupta J, Nielsen RC, Nissen P, Harvey SC, Ehrenberg M and Frank J (2003) Incorporation of aminoacyl-tRNA into the ribosome as seen by cryo-electron microscopy. *Nature Structural Biology* **10**, 899–906.
- van Agthoven A, Maassen JA, Schrier PI and Möller W (1975) Inhibition of EF-G dependent GTPase by an aminoterminal fragment of L7/L12. *Biochemical and Biophysical Research Communication* **64**, 1184–1191.
- Voorhees RM, Schmeing TM, Kelley AC and Ramakrishnan V (2010) The mechanism for activation of GTP hydrolysis on the ribosome. *Science* **330**, 835–838.
- Wahl MC and Möller W (2002) Structure and function of the acid ribosomal stalk proteins. *Current Protein & Peptide Science* **3**, 93–106.
- Wahl MC, Bourenkov GP, Bartunik HD and Huber R (2000) Flexibility, conformational diversity and two dimerization modes in complexes of ribosomal protein L12. *EMBO Journal* **19**, 174–186.
- Wawiórka L, Molestak E, Szahwaj M, Michalec-Wawiórka B, Molon M, Borkiewicz L, Grela P, Boguszewska A and Tchorzewski M (2017) Multiplication of P-stalk proteins contributes to the fidelity of translation. *Molecular Cell Biology* **37**, e00060–17.
- Wool IG, Chan YL, Gluck A and Suzuki K (1991) The primary structure of rat ribosomal proteins P0, P1, and P2 and a proposal for a uniform nomenclature for mammalian and yeast ribosomal proteins. *Biochimie* **73**, 861–870.
- Yusupov MM, Yusupova GZ, Baucom A, Lieberman K, Earnest TN, Cate JH and Noller HF (2001) Crystal structure of the ribosome at 5.5 Å resolution. *Science* **292**, 883–896.
- Zinker S and Warner JR (1976) The ribosomal proteins of *Saccharomyces cerevisiae*. Phosphorylated and exchangeable proteins. *Journal of Biological Chemistry* **251**, 1799–1807.