

## A note on the four-dimensional clasp number of knots

BY PETER FELLER

ETH Zurich, Rämistrasse 101, 8092 Zurich, Switzerland

e-mail: [peter.feller@math.ch](mailto:peter.feller@math.ch)

AND JUNGHWAN PARK

KAIST, 291 Daehak-ro, Daejeon, 34141, South Korea

e-mail: [jungpark0817@kaist.ac.kr](mailto:jungpark0817@kaist.ac.kr)

(Received 25 January 2021; revised 2 June 2021; accepted 14 June 2021)

### Abstract

Among the knots that are the connected sum of two torus knots with cobordism distance 1, we characterise those that have 4-dimensional clasp number at least 2, and we show that their  $n$ -fold connected self-sum has 4-dimensional clasp number at least  $2n$ . Our proof works in the topological category. To contrast this, we build a family of topologically slice knots for which the  $n$ -fold connected self-sum has 4-ball genus  $n$  and 4-dimensional clasp number at least  $2n$ .

2020 Mathematics Subject Classification: 57K10, 57N70 (Primary)

---

### 1. Introduction

Recently the authors determined for which pairs of torus knots  $\{T_{p,q}, T_{p',q'}\}$  the cobordism distance is equal to 1, with one exception [FP19, Theorem 1.2]. This was done by comparing explicit constructions of cobordisms with the lower bound for the cobordism distance using the  $\nu^+$ -invariant [HW16] from the Heegaard Floer knot complex. As an application, the authors determined which pairs of torus knots have Gordian distance 1; see [FP19, Corollary 1.3]. The first result of this paper has two motivations.

Firstly, when determining Gordian distance 1 pairs of torus knots, we relied on  $\nu^+$ , but speculated that the proof could be done using the Tristram–Levine signatures [Tri69, Lev69]; see [FP19, Remark 4.3]. Here, we partially confirm this speculation using signature calculations via the Hirzebruch–Brieskorn formula [Bri66, GG05].

Secondly, recent interest in the 4-dimensional clasp number  $c_4$  and its difference to the 4-ball genus  $g_4$  [KM19, Kro20, JZ20, DS20a, DS20b], made us revisit the lower bounds we had on the Gordian distance of the pairs  $\{T_{p,q}, T_{p',q'}\}$  and consider whether they yield lower bounds on  $c_4(K)$  and  $c_4(\#^n K)$ , where  $K = T_{p,q} \# -T_{p',q'}$ . More precisely, Daemi–Scaduto [DS20a, Theorem 1] used equivariant singular instanton theory to show that, if  $K$  is the knot  $7_4$ , then  $c_4(\#^n K) - g_4(\#^n K) \geq n/5$  (in fact, they showed a stronger statement which we will discuss at the end of this section). Also, Juhász–Zemke [JZ20, Corollary 1.5] used knot Floer homology to show that, if  $K = T_{2,11} \# -T_{4,5}$ , then  $c_4(\#^n K) - g_4(\#^n K) \geq n$ . In this article, we show that it is also possible to establish this behaviour by using Tristram–Levine

signatures, and we provide examples of topologically slice knots with such behaviour, for which the use of smooth tools are a necessity.

We see this paper as shining a spotlight on what can be achieved using a somewhat classical setup of concordance homomorphism, while the new exciting invariants from [KM19, JZ20, DS20a] might be used to discover interesting phenomena beyond.

We write  $-J$  to denote the reverse of the mirror image of a knot  $J$ , and we write  $g_4^{\text{top}}$  and  $c_4^{\text{top}}$  to denote the topological counterparts of  $g_4$  and  $c_4$ , respectively.

**THEOREM 1.1.** *Let  $\{T_{p,q}, T_{p',q'}\}$  be a pair of positive torus knots,  $K = T_{p,q} \# -T_{p',q'}$ , and  $n$  be a positive integer. If  $g_4(K) = 1$  and the Gordian distance between  $T_{p,q}$  and  $T_{p',q'}$  is not 1, then*

$$g_4(\#^n K) = g_4^{\text{top}}(\#^n K) = n \quad \text{and} \quad c_4(\#^n K) \geq c_4^{\text{top}}(\#^n K) \geq 2n.$$

Furthermore, if  $\{T_{p,q}, T_{p',q'}\}$  is either  $\{T_{2,7}, T_{3,4}\}$ ,  $\{T_{2,9}, T_{3,5}\}$ , or  $\{T_{3,7}, T_{4,5}\}$ , then

$$g_4(\#^n K) = g_4^{\text{top}}(\#^n K) = n \quad \text{and} \quad c_4(\#^n K) = c_4^{\text{top}}(\#^n K) = 2n.$$

We comment on the condition  $g_4(K) = 1$ . The point is that  $g_4(K) = 1$  implies that  $\{T_{p,q}, T_{p',q'}\}$  must be one of an infinite family of pairs of torus knots, among which we know which are Gordian distance 1 apart, and, in fact, infinitely many are further apart [FP19]; see Section 3. With this the first part of Theorem 1.1 reduces to showing  $g_4^{\text{top}}(\#^n K) \geq n$  and  $c_4^{\text{top}}(\#^n K) \geq 2n$  for an explicit infinite family of knots. We do this in Section 3 using Tristram–Levine signatures. The ‘furthermore’-part comes down to finding appropriate crossing changes. For the pairs  $\{T_{2,7}, T_{3,4}\}$  and  $\{T_{2,9}, T_{3,5}\}$ ,<sup>1</sup> we believe Theorem 1.1 to be known to experts, but for the pair  $\{T_{3,7}, T_{4,5}\}$  we will exhibit two crossing changes turning one into the other explicitly.

**Question 1.2.** Are there other pairs of torus knots for which  $g_4(K) = 1$  and  $c_4(K) = 2$ ?

We also obtain further equivalent characterizations of Gordian distance 1 pairs of torus knots (compare with [FP19, Corollary 1.3]). Let  $u_s(J)$  denote the slicing number of a knot  $J$ . We have the following chain of inequalities:

$$u_s(J) \geq c_4(J) \geq g_4(J).$$

It is unknown whether there is a knot  $J$  with  $u_s(J) > c_4(J)$ ; see e.g. [OS16, Proposition 6].

**COROLLARY 1.3.** *If  $\{T_{p,q}, T_{p',q'}\}$  is a pair of positive torus knots, then the following statements are equivalent;*

- (i) *the knots  $T_{p,q}$  and  $T_{p',q'}$  have Gordian distance 1;*
- (ii) *the knot  $T_{p,q} \# -T_{p',q'}$  has slicing number 1;*
- (iii) *the knot  $T_{p,q} \# -T_{p',q'}$  has 4-dimensional clasp number 1;*
- (iv) *the pair  $\{T_{p,q}, T_{p',q'}\}$  is one of the following:*
  - (a)  $\{T_{2,2n+1}, T_{2,2n+3}\}$  *for an integer  $n \geq 0$ ;*
  - (b)  $\{T_{3,3n+1}, T_{3,3n+2}\}$  *for an integer  $n \geq 1$ ;*
  - (c)  $\{T_{2,5}, T_{3,4}\}, \{T_{2,7}, T_{3,5}\}$ .

<sup>1</sup>These pairs are familiar in singularity theory as the knots of singularities of the pairs of simple singularities  $\{A_6, E_6\}$  and  $\{A_8, E_8\}$ , which have  $\delta$ -constant deformations to  $A_4$  and  $A_6$ , respectively.

*Question 1.4.* Are there pairs of torus knots for which their Gordian distance is strictly larger than  $c_4(\#^n K)/n$  for some integer  $n \geq 1$ ?

Given that not just  $c_4 - g_4$  but also  $c_4^{\text{top}} - g_4$  can be large even on connected sums of a positive and a negative torus knot (by Theorem 1.1), it seems interesting to find a family of examples of topologically slice knots on which  $c_4 - g_4$  is unbounded. Note that if  $J$  is topologically slice (i.e.  $c_4^{\text{top}}(J) = g_4^{\text{top}}(J) = 0$ ), then the Tristram–Levine signature obstruction for the 4-dimensional clasp number vanishes; hence, one needs to use other invariants. Following in the footsteps of Livingston–Friedl–Zentner [FLZ17] and Juhász–Zemke [JZ20] among others, we use the Ozsváth–Stipsicz–Szabó  $\Upsilon$ -invariant [OSS17] to find such families.

Indeed, let  $D$  denote the positive untwisted Whitehead double of the right-handed trefoil, and let  $J_{p,q}$  denote the  $(p, q)$ -cable of a knot  $J$ , where  $p$  is the longitudinal winding.

**THEOREM 1.5.** *Let  $i \geq 2$  and  $n \geq 1$  be integers. If  $K_i = D_{2,2i+1}\# - T_{2,2i+1}\# - D$ , then  $K_i$  is a topologically slice knot that satisfies*

$$g_4(\#^n K_i) = n \quad \text{and} \quad c_4(\#^n K_i) \geq 2n.$$

The lower bounds for  $c_4$  obtained by using Tristram–Levine signatures and the Ozsváth–Stipsicz–Szabó  $\Upsilon$ -invariant fall into a rather general setup of invariants going back to Livingston [Liv04]; see Section 2. However, the lower bound for  $c_4^{\text{top}}$  cannot be obtained in the exact same way. Regardless, we provide a bound for  $c_4^{\text{top}}$  (see Lemma 2.3) that yields the topological statements of our results above.

By design, the invariants we consider do not (or at least not in an obvious way) allow to show that  $c_4(J) > 2g_4(J)$  or  $c_{4,+}(J) > g_4(J)$ , where  $c_{4,+}$  denotes a version of the 4-dimensional clasp number that only counts positive clasps. In particular, such invariants will not allow to answer the next question.

*Question 1.6.* Are there 4-ball genus 1 knots with arbitrarily large 4-dimensional clasp number? Denoting by  $\mathcal{C}$  the set of smooth concordance classes, is  $\limsup_{[J] \neq 0 \in \mathcal{C}} (c_4(J)/g_4(J)) > 2$ ?

In contrast, the recent work that was part of the inspiration for this paper [KM19, JZ20, DS20a] has the potential to address Question 1.6. In fact, [DS20a, Theorem 1] shows that the gap between  $c_{4,+}$  and  $g_4$  can be made arbitrarily large.

## 2. Lower bounds on $c_4$ and $g_4$ from concordance homomorphism

First, we consider the smooth category. The 4-dimensional clasp number  $c_4(J)$  of a knot  $J$  can be defined as the smallest non-negative integer  $k$  such that  $J$  can be turned into the unknot using a finite sequence of smooth concordances and crossing changes with at most  $k$  crossing changes. Equivalently, one can define the 4-dimensional clasp number as the smallest non-negative integer  $k$  such that there is a smoothly immersed disk with  $k$  transverse double points (also called clasps) in  $B^4$  bounding  $J$  and no other singular points; indeed, any such disk is isotopic to a smoothly immersed disk resulting from stacking a smooth concordance, the trace of  $k$  crossing changes, and a smooth slice disk; see [OS16, Proposition 2.1]. One defines weighted version  $c_{4,+}(J)$  (resp.  $c_{4,-}(J)$ ) as the smallest non-negative integer of

positive-to-negative (resp. negative-to-positive) crossing changes in a sequence as above. In particular, for any knot  $J$ ,

$$c_{4,+}(J) + c_{4,-}(J) \leq c_4(J). \quad (2.1)$$

We recall a setup implicit in [Liv04] and explicit in [Fel14, Lemma 17] and [FLZ17, Lemma 17]. Let  $\mathcal{C}$  denote the smooth knot concordance group.

LEMMA 2.1. *Let  $\nu: \mathcal{C} \rightarrow \mathbb{R}$  be any homomorphism such that:*

- (i)  $\nu(J) \leq g_4(J)$  for every knot  $J$  and;
- (ii) there exists a knot  $J'$  with  $\nu(J') = 1$  such that  $J'$  can be turned into a smoothly slice knot by a positive-to-negative crossing change;

then, for each knot  $J$ , we have

$$|\nu(J)| \leq g_4(J), \quad \nu(J) \leq c_{4,+}(J), \quad \text{and} \quad -\nu(J) \leq c_{4,-}(J). \quad (2.2)$$

For explicit examples of  $\nu$  and their treatment via this approach, we point to [Liv04, Corollary 3] for  $\nu(J) = \tau(J)$ , [Fel14, Lemmas 11 and 17] for  $\nu(J) = -\sigma_\omega(J)/2$  when  $\omega$  is regular, and [Liv17, Theorem 13.1] for  $\nu(J) = -\Upsilon_J(t)/t$ . Here,  $\omega$  is *regular* if  $\omega \in S^1 \setminus \{1\}$  and  $f(\omega) \neq 0$  for all integer coefficient Laurent polynomials  $f$  with  $f(t) = f(t^{-1})$  and  $f(1) = 1$ .

*Remark 2.2.* We note that the above definitions and Lemma 2.1 also work in the topologically locally-flat category by replacing  $\mathcal{C}$  with the topological concordance group and  $g_4$ ,  $c_{4,+}$ ,  $c_{4,-}$ , and  $c_4$  with  $g_4^{\text{top}}$ ,  $c_{4,+}^{\text{top}}$ ,  $c_{4,-}^{\text{top}}$ , and  $c_4^{\text{top}}$ . In particular, for  $\nu(J) = -\sigma_\omega(J)/2$  when  $\omega$  is regular, (2.2) holds in the locally-flat category.

**Warning.** We note the following subtlety. Above we gave two equivalent definitions of  $c_4$  (and implicitly  $c_{4,\pm}$ ) in the smooth category. It is tempting to speculate that the same equivalence of definitions holds in the locally-flat category. However, the authors are not aware of a proof of this, hence the locally-flat version of Lemma 2.1 (see Remark 2.2) is to be read with  $c_4^{\text{top}}$ ,  $c_{4,+}^{\text{top}}$ , and  $c_{4,-}^{\text{top}}$  defined using sequences of concordances and crossing changes. However, from Lemma 2.3 below, we know that the lower bound (2.2) for  $\nu(J) = -\sigma_\omega(J)/2$  with  $\omega$  regular holds also when defining  $c_4^{\text{top}}(J)$ ,  $c_{4,+}^{\text{top}}(J)$ , and  $c_{4,-}^{\text{top}}(J)$  via counting double points in locally-flat immersed disks filling  $J$  with only transversal double points as singularities. Thus, Theorem 1.1 holds for this definition of  $c_4^{\text{top}}$ .

We formulate the following lemma for locally-flat immersed surfaces bounding links since we believe this to be of independent interest; however, for the use in this text it suffices to consider the case of  $L$  being a knot and  $F$  being a disk in the statement (in particular,  $|\eta_\omega(L) - b_0(F) + 1| = b_1(F) = 0$ ). For  $\omega \in S^1$ , we denote by  $\sigma_\omega(L)$  and  $\eta_\omega(L)$  the signature and the nullity, respectively, of  $(1 - \omega)M + (1 - \bar{\omega})M^T$ , where  $M$  is a Seifert matrix for  $L$ .

LEMMA 2.3. *Let  $F$  be an oriented compact surface with boundary, let  $\phi: F \rightarrow B^4$  be a locally-flat proper immersion with  $p$  positive transversal double points and  $n$  negative transversal double points and no other singular points, and let  $L$  be the link  $\phi(\partial F) \subset S^3$ .*

Then, for all regular  $\omega$ , we have

$$\sigma_\omega(L) + |\eta_\omega(L) - b_0(F) + 1| \leq b_1(F) + 2n.$$

We derive Lemma 2.3 from the Tristram–Levine bound:

$$|\sigma_\omega(L)| + |\eta_\omega(L) - b_0(F') + 1| \leq b_1(F') \quad ([\text{Tri69, Lev69, KT76, Pow17, CNT20}] \text{, (2.3)})$$

for every locally-flat embedded proper compact oriented surface  $F'$  with boundary the link  $L \subset S^3$ .

*Proof of Lemma 2.3.* Without loss of generality we assume  $F$  has no closed components (otherwise, consider the restriction of  $\phi$  to the union of the non-closed components of  $F$ ).

We claim that there exists a locally-flat proper embedding of  $F' = F \setminus \{p + n \text{ open disks}\}$  into  $B^4$  with boundary a link  $L'$  that is the union of  $L$  and  $p + n$  meridians,  $p$  of which are link positively with  $L$  and  $n$  of which link negatively with  $L$ . To see this claim, take 4-balls  $N_i$  around double points of  $\phi(F)$  such that  $(N_i, N_i \cap \phi(F))$  is homeomorphic to  $(B^4, B^4 \cap \{(x, y) \in B^4 \subset \mathbb{C}^2 \mid xy = 0\})$  preserving orientations, where the disk  $B^4 \cap \{(x, y) \mid x = 0\}$  carries the orientation induced by the complex orientation and  $B^4 \cap \{(x, y) \mid y = 0\}$  carries the induced complex and anti-complex orientation for a positive and negative double point, respectively. Choose  $p + n$  properly embedded pairwise disjoint arcs  $\alpha_i$  in  $\phi(F) \setminus N_1^\circ \cup \dots \cup N_{p+n}^\circ$  such that  $\alpha_i$  has one endpoint on  $L$  and the other on  $N_i$ . Letting  $N$  be the union of the  $N_i$  and regular neighborhoods of the  $\alpha_i$ , we observe that the pair  $(B^4 \setminus N^\circ, \phi(F) \setminus N^\circ)$  is homeomorphic to  $(B_4, F')$  preserving orientations, where  $F'$  is a locally-flat properly embedded surface with boundary a link  $L'$  as claimed. In particular,  $b_0(F') = b_0(F)$  and  $b_1(F') = b_1(F) + p + n$ .

We note that  $L'$  arises as a  $p + n$  fold connected sum of  $L$  with  $n$  positive Hopf links  $H^+$  and  $p$  negative Hopf links  $H^-$ . Hence, by additivity of signature and nullity using that  $\sigma_\omega(H^\pm) = \mp 1$  and  $\eta_\omega(H^\pm) = 0$ , we find

$$\sigma_\omega(L') = \sigma_\omega(L) - n + p \quad \text{and} \quad \eta_\omega(L') = \eta_\omega(L), \tag{2.4}$$

where  $\eta_\omega$  and  $\sigma_\omega$  denote the nullity and signature, respectively, for every  $\omega \in S^1 \setminus \{1\}$ .

We conclude the proof by calculating that, for every regular  $\omega$ , we have

$$\begin{aligned} \sigma_\omega(L) + |\eta_\omega(L) - b_0(F) + 1| &\stackrel{(2.4)}{=} \sigma_\omega(L') - p + n + |\eta_\omega(L') - b_0(F) + 1| \\ &\stackrel{b_0(F')=b_0(F)}{=} \sigma_\omega(L') + |\eta_\omega(L') - b_0(F') + 1| - p + n \\ &\stackrel{(2.3)}{\leq} b_1(F') - p + n \\ &\stackrel{b_1(F')=b_1(F)+p+n}{=} b_1(F) + 2n. \end{aligned}$$

### 3. Theorem 1.1 via signature calculations and crossing changes

We start by making the family of pairs of torus knots from Theorem 1.1 explicit. Recall that we write  $K := T_{p,q} \# -T_{p',q'}$ . Pairs of torus knots  $\{T_{p,q}, T_{p',q'}\}$  with Gordian distance two or more and  $g_4(K) = 1$  are among the following:

- (I)  $\{T_{3n+1,9n+6}, T_{3n+2,9n+3}\}$  for  $n \geq 1$ ;
- (II)  $\{T_{2n+1,4n+6}, T_{2n+3,4n+2}\}$  for  $n \geq 1$ ; or

- (III)  $\{T_{2,11}, T_{3,7}\}, \{T_{2,13}, T_{3,8}\}, \{T_{2,7}, T_{3,4}\}, \{T_{2,9}, T_{3,5}\}, \{T_{2,11}, T_{4,5}\}, \{T_{3,7}, T_{4,5}\},$   
 $\{T_{3,10}, T_{4,7}\}, \{T_{4,9}, T_{5,7}\}, \{T_{3,14}, T_{5,8}\}.$

In fact, all these pairs have Gordian distance two or more, and, except for  $\{T_{3,14}, T_{5,8}\}$ , all these pairs are known to satisfy  $g_4(K) = 1$ . See [FP19, Theorem 1.2 and Corollary 1.3].

The aim of this section is to prove Theorem 1.1. The following lower bound will be used.

COROLLARY 3.1. *If  $J$  is a knot in  $S^3$ , then*

$$c_4^{\text{top}}(J) \geq \max_{\omega \text{ regular}} \frac{\sigma_\omega(J)}{2} + \max_{\omega \text{ regular}} -\frac{\sigma_\omega(J)}{2}.$$

We note that Corollary 3.1 holds for both definitions of  $c_4^{\text{top}}(J)$  discussed in Section 2.

*Proof.* We first note that  $c_4^{\text{top}}(J) \geq c_{4,+}^{\text{top}}(J) + c_{4,-}^{\text{top}}(J)$ . (This works for either of the two definitions of  $c_4^{\text{top}}, c_{4,+}^{\text{top}}$ , and  $c_{4,-}^{\text{top}}$ .) We show

$$c_{4,+}^{\text{top}}(J) \geq \max_{\omega \text{ regular}} -\frac{\sigma_\omega(J)}{2} \quad \text{and} \quad c_{4,-}^{\text{top}}(J) \geq \max_{\omega \text{ regular}} \frac{\sigma_\omega(J)}{2}. \tag{3.1}$$

In fact, it suffices to establish one of the inequalities since  $c_{4,+}^{\text{top}}(J) = c_{4,-}^{\text{top}}(-J)$  and  $\sigma_\omega(J) = -\sigma_\omega(-J)$  for all  $\omega \in S^1$ .

In case  $c_{4,-}^{\text{top}}(J)$  is defined as the minimal number of negative crossing changes among all finite sequences of concordances and crossing changes turning  $J$  into the unknot, then (3.1) follows from (2.2) for  $\nu = -\sigma_\omega/2$  (which applies by Remark 2.2).

In case  $c_{4,+}^{\text{top}}(J)$  is defined as the minimal number of negative clasps in a immersed disk in  $B^4$  with boundary  $J$ , then (3.1) follows from Lemma 2.3. To see this, let  $D$  be a locally-flat immersed disk that realizes  $c_{4,-}^{\text{top}}(J)$ , that is,  $D$  has only transversal double points as singularities and the number of negative transversal double points of  $D$  is  $c_{4,-}^{\text{top}}(J)$ , and let  $\omega \in S^1$  be regular. By Lemma 2.3, we have

$$\sigma_\omega(J) = \sigma_\omega(J) + |\eta_\omega(J) - b_0(D) + 1| \leq b_1(D) + 2c_{4,-}^{\text{top}}(J) = 2c_{4,+}^{\text{top}}(J).$$

The upper bound will be achieved by explicit constructions.

### 3.1. Proof of Theorem 1.1

In this subsection, we provide the proof of Theorem 1.1 except the necessary signature calculations. The latter are provided in the next subsection.

*Proof of Theorem 1.1.* Let  $K = T_{p,q} \# -T_{p',q'}$  be such that  $g_4(K) = 1$  and there does not exist a crossing change turning  $T_{p,q}$  into  $T_{p',q'}$ ; in particular,  $\{T_{p,q}, T_{p',q'}\}$  must be among (I), (II), or (III). We establish in Examples 3.4, 3.5, and 3.6 below that

$$\max_{\omega \text{ regular}} \sigma_\omega(K) = 2 \quad \text{and} \quad \max_{\omega \text{ regular}} -\sigma_\omega(K) = 2. \tag{3.2}$$

Hence, by Corollary 3.1 and the additivity of the Tristram–Levine signatures, we have

$$c_4^{\text{top}}(\#^n K) \geq 2n.$$




Fig. 1.  $T_{4,5}$  (left, as the closure of a 4-braid) can be turned into  $T_{3,7}$  (right, as the closure of a 3-braid) by two crossing changes (modification in indicated 3-balls (red)). Also indicated ( $=$ ) are braid isotopies (gray for the isotopy between the first and second braid, blue for the one between the second and third braid). Only the braids, rather than their closures, are drawn.

Moreover, for all integers  $n \geq 1$ ,

$$g_4(\#^n K) = g_4^{\text{top}}(\#^n K) = n,$$

since all these quantities are at most  $n$  by  $g_4(K) = 1$ , and the lower bound comes from the Tristram–Levine signatures bound (2.3).

It remains to discuss the ‘furthermore’-paragraph. It suffices to show that for the pairs in question,  $T_{p,q}$  can be turned into  $T_{p',q'}$  by two crossing changes (necessarily of opposite sign by (2.2)). Since then,  $\#^n K$  can be turned into a slice knot with  $2n$  crossing changes,  $n$  of each sign, for all integers  $n \geq 1$ ; hence,

$$2n \geq c_4(\#^n K) \geq c_4^{\text{top}}(\#^n K).$$

For the pairs  $\{T_{2,7}, T_{3,4}\}$  and  $\{T_{2,9}, T_{3,5}\}$  such crossing changes are available in the literature (see e.g. [Fel14]). We end the proof with an explicit sequence of two crossing changes and isotopies turning  $T_{4,5}$  into  $T_{3,7}$ ; see Figure 1.

### 3.2. Tristram–Levine signatures calculation

Denoting by  $\sigma_t(J) = \sigma_{e^{2\pi it}}(J)$  the Tristram–Levine signature for  $\omega = e^{2\pi it} \in S^1$  of a knot  $J$  [Tri69, Lev69], one has for every fixed knot an integer-valued piecewise linear function, which is constant in neighbourhoods  $t$  for which  $\omega = e^{2\pi it}$  is regular. We define  $t_0 \in (0, 1)$  to be a *jump point* of the signature function of  $J$ , if the right limit  $\lim_{t \rightarrow t^+} \sigma_t(J)$  differs from the left limit  $\lim_{t \rightarrow t^-} \sigma_t(J)$ . And say the *jump* at  $t_0 \in (0, 1)$  is  $\lim_{t \rightarrow t^+} \sigma_t(J) - \lim_{t \rightarrow t^-} \sigma_t(J)$ . For a more detailed discussion, and a complete description of what functions arise as signature functions, we refer to [Liv18].

For the proof of Theorem 1.1, it remains to check (3.2) for the pairs from the families (I), (II) and (III). This is done by using the signature formula going back to Hirzebruch and Brieskorn [Bri66] [GG05, Proposition 5.1], which we recall below in the proof of Lemma 3.2. We focus on (I) and (II) since (III) consists of a small finite list of examples for which (3.2) can be checked by hand or computer (see Example 3.6).



LEMMA 3.2. *Let  $0 < p < q$  be coprime integers. The signature function  $\sigma_t(T_{p,q})$  is monotonically decreasing on  $[0, p + q/pq)$ . More precisely, the jump points on  $[0, p + q/pq)$  occur at  $\{\ell/pq \mid p \nmid \ell \text{ and } q \nmid \ell\}$  and each jump is  $-2$ . Furthermore, the jump at  $p + q/pq$  is 2. In other words, for integers  $\ell$  with  $0 \leq \ell < p + q$ , we have*

$$\sigma_t(T_{p,q}) = \begin{cases} -2 \left( t - \left\lfloor \frac{\ell}{q} \right\rfloor - \left\lfloor \frac{\ell}{p} \right\rfloor \right) & \text{for } t \in \left( \frac{\ell}{pq}, \frac{\ell+1}{pq} \right), \\ -2 \left( p + q - 4 - \left\lfloor \frac{q}{p} \right\rfloor \right) & \text{for } t \in \left( \frac{p+q}{pq}, \frac{p+q+1}{pq} \right). \end{cases}$$

The content of Lemma 3.2 might be well known to experts. In particular, the first author previously discussed the monotonicity of the signature function of the torus knot  $T_{p,q}$  on  $[0, p + q/pq)$  with Charles Livingston. For completeness, we provide a proof.

*Proof of Lemma 3.2.* We recall the torus knot signature version of the Hirzebruch–Brieskorn signature formula [Bri66] as explicitly stated in [Lit79, Proposition 1]; compare also [GG05, Proposition 5.1]. For coprime integers  $p, q > 0$  and

$$S := \left\{ \frac{k}{p} + \frac{j}{q} \mid k, j \in \mathbb{Z}, 0 < k < p, \text{ and } 0 < j < q \right\} \subset (0, 2),$$

we have

$$\sigma_{e^{2\pi i t}}(T_{p,q}) = \sigma_t(T_{p,q}) = \#\{S \setminus [t, 1 + t]\} - \#\{S \cap (t, 1 + t)\} \quad \text{for all } t \in [0, 1].$$

We start by using this formula to prove the following four observations:

- (i)  $\min S = p + q/pq$ ;
- (ii)  $1 + p + q/pq \notin S$ ;
- (iii)  $1 + \ell/pq \in S$  for all  $0 < \ell < p + q$  with  $p \nmid \ell$  and  $q \nmid \ell$ ;
- (iv)  $\ell/pq, 1 + \ell/pq \notin S$  for all  $0 < \ell < pq$  with  $p \mid \ell$  or  $q \mid \ell$ .

Note that (i) follows immediately from the definition of  $S$ . Towards establishing (ii)-(iv), we note the following claim.

CLAIM. *For  $0 < \ell < pq$ , exactly one of  $\ell/pq$  and  $1 + \ell/pq$  is in  $S$  if  $p \nmid \ell$  and  $q \nmid \ell$ , while neither is the case if  $p \mid \ell$  or  $q \mid \ell$ .*

To see this claim, for  $0 < \ell < pq$ , note that  $\ell \equiv kq + jp \pmod{pq}$  for some  $0 \leq k < p$  and  $0 \leq j < q$ . Moreover,  $p \nmid \ell$  and  $q \nmid \ell$  if and only if  $k \neq 0$  and  $j \neq 0$ , which implies (iv). Since  $0 \leq kq + jp < 2pq$ , we see that  $\ell$  is either  $kq + jp$  or  $kq + jp - pq$ . If  $\ell = kq + jp$  with  $0 < k < p$  and  $0 < j < q$ , then  $\ell/pq \in S$ . For the sake of contradiction, assume  $1 + \ell/pq \in S$ , then  $pq + \ell = k'q + j'p$  for some  $0 < k' < p$  and  $0 < j' < q$ . This implies that  $(k' - k)q + (j' - j)p = pq$  and  $p \mid (k' - k)$  and  $q \mid (j' - j)$ . In particular, we have that  $k = k'$  and  $j = j'$ , which lead us to a contradiction. A similar argument shows that if  $\ell = kq + jp - pq$  with  $0 < k < p$  and  $0 < j < q$ , then  $1 + \ell/pq \in S$  and  $\ell/pq \notin S$ .

With the claim, observation (ii) follows from (i) since  $(p + q)/pq \in S$ . With the claim observation (iii) also follows in a similar way as (ii) since for  $0 < \ell < p + q$ , we have that  $\ell/pq \notin S$ . Finally, observation (vi) is contained in the claim. We have thus established observations (i)-(iv).

It remains to show that (i)-(iv) imply Lemma 3.2. First we note that (iii) and (iv) imply that the jump points on  $[0, p + q/pq)$  occur at  $\{\ell/pq \mid p \nmid \ell \text{ and } q \nmid \ell\}$  and each jump is  $-2$ ;



in particular, the signature function  $\sigma_t(T_{p,q})$  is monotonically decreasing on  $[0, p + q/pq]$ . Secondly, (i) and (ii) show that the jump at  $p + q/pq$  is 2. That last sentence of Lemma 3.2 is a formulaic reformulation of the two former statements.

**COROLLARY 3.3.** *Let  $0 < p < q$  and  $0 < p' < q'$  be pairs of coprime integers such that  $pq = p'q'$  and  $p < p'$ , then*

$$\sigma_t(T_{p,q}\# - T_{p',q'}) = \begin{cases} 0 & \text{for } t \in \left[0, \frac{1}{q}\right), \\ 2 & \text{for } t \in \left(\frac{1}{q}, \frac{1}{q} + \frac{1}{pq}\right), \\ 2 \left( \left\lfloor \frac{p'+q'}{q} \right\rfloor + \left\lfloor \frac{p'+q'}{p} \right\rfloor - 4 - \left\lfloor \frac{q'}{p'} \right\rfloor \right) & \text{for } t \in \left(\frac{p'+q'}{qp}, \frac{p'+q'+1}{qp}\right). \end{cases}$$

**Example 3.4.** [Family (I)] Fix an integer  $n \geq 1$  and set  $p = 3n + 1, q = 9n + 6, p' = 3n + 2, q' = 9n + 3$ .

If  $t \in \left(\frac{1}{9n+6}, \frac{1}{9n+6} + \frac{1}{3(3n+1)(3n+2)}\right)$ , then  $\sigma_t(T_{p,q}\# - T_{p',q'}) = 2$ .

If  $t \in \left(\frac{12n+5}{3(3n+1)(3n+2)}, \frac{12n+6}{3(3n+1)(3n+2)}\right)$ , then

$$\begin{aligned} \sigma_t(T_{p,q}\# - T_{p',q'}) &= 2 \left( \left\lfloor \frac{12n+5}{9n+6} \right\rfloor + \left\lfloor \frac{12n+5}{3n+1} \right\rfloor - 4 - \left\lfloor \frac{9n+3}{3n+2} \right\rfloor \right) \\ &= 2(1 + 4 - 4 - 2) = -2. \end{aligned}$$

**Example 3.5.** [Family (II)] Fix an integer  $n \geq 1$  and set  $p = 2n + 1, q = 4n + 6, p' = 2n + 3, q' = 4n + 2$ .

If  $t \in \left(\frac{1}{4n+6}, \frac{1}{4n+6} + \frac{1}{2(2n+1)(2n+3)}\right)$ , then  $\sigma_t(T_{p,q}\# - T_{p',q'}) = 2$ .

If  $t \in \left(\frac{6n+5}{2(2n+1)(2n+3)}, \frac{6n+6}{2(2n+1)(2n+3)}\right)$ , then

$$\begin{aligned} \sigma_t(T_{p,q}\# - T_{p',q'}) &= 2 \left( \left\lfloor \frac{6n+5}{4n+6} \right\rfloor + \left\lfloor \frac{6n+5}{2n+1} \right\rfloor - 4 - \left\lfloor \frac{4n+2}{2n+3} \right\rfloor \right) \\ &= 2(1 + 3 - 4 - 1) = -2. \end{aligned}$$

**Example 3.6.** [Family (III)] For all  $K = T_{p,q}\# - T_{p',q'}$  from family (III), we provide some  $\omega = e^{2\pi it}$  for which the maximum 2 and the minimum  $-2$  of the Tristram–Levine signatures are realized. Slightly more concretely, we note that  $pq > p'q'$  in all these examples, which immediately yields that  $\min_{\omega \text{ regular}} \sigma_{\omega}(K) \leq -2$ . The reader might argue for the maximum being 2 using Lemma 3.2. However, this seems artificial given the availability of the signature functions, which in particular yield the following:

$$\begin{aligned} \{T_{2,11}, T_{3,7}\}: \sigma_t(K) &= 2 \quad \text{for } t \in \left(\frac{2}{21}, \frac{3}{22}\right) \text{ and } \sigma_t(K) = -2 \quad \text{for } t \in \left(\frac{1}{22}, \frac{1}{21}\right), \\ \{T_{2,13}, T_{3,8}\}: \sigma_t(K) &= 2 \quad \text{for } t \in \left(\frac{2}{26}, \frac{3}{24}\right) \text{ and } \sigma_t(K) = -2 \quad \text{for } t \in \left(\frac{1}{26}, \frac{1}{24}\right), \\ \{T_{2,7}, T_{3,4}\}: \sigma_t(K) &= 2 \quad \text{for } t \in \left(\frac{2}{12}, \frac{3}{14}\right) \text{ and } \sigma_t(K) = -2 \quad \text{for } t \in \left(\frac{1}{14}, \frac{1}{12}\right), \end{aligned}$$

- $\{T_{2,9}, T_{3,5}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{2}{15}, \frac{3}{18})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{18}, \frac{1}{15})$ ,
- $\{T_{2,11}, T_{4,5}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{2}{20}, \frac{3}{22})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{22}, \frac{1}{20})$ ,
- $\{T_{3,7}, T_{4,5}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{3}{20}, \frac{4}{21})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{21}, \frac{1}{20})$ ,
- $\{T_{3,10}, T_{4,7}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{3}{28}, \frac{4}{30})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{30}, \frac{1}{28})$ ,
- $\{T_{4,9}, T_{5,7}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{4}{35}, \frac{5}{36})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{36}, \frac{1}{35})$ ,
- $\{T_{3,14}, T_{5,8}\}$ :  $\sigma_t(K) = 2$  for  $t \in (\frac{3}{40}, \frac{4}{42})$  and  $\sigma_t(K) = -2$  for  $t \in (\frac{1}{42}, \frac{1}{40})$ .

4. Theorem 1.5 via Heegaard Floer invariants and cabling

The aim of this section is to prove Theorem 1.5. We start with a proposition that summarizes the properties of the  $\Upsilon$ -invariant [OSS17] that we use in the proof.

PROPOSITION 4.1. *For any knot  $J$ , the  $\Upsilon$ -invariant  $\Upsilon_J$  is a piecewise linear function  $\Upsilon_J : [0, 2] \rightarrow \mathbb{R}$  with the following properties:*

- (i) [OSS17, Corollary 1.12]  $\Upsilon$  is a concordance invariant and, for all knots  $J$  and  $J'$ ,

$$\Upsilon_{J\#J'}(t) = \Upsilon_J(t) + \Upsilon_{J'}(t);$$

- (ii) [OSS17, Theorem 1.11] For  $0 < t \leq 1$ ,  $|\Upsilon_J(t)/t| \leq g_4(J)$ ;
- (iii) [OSS17, Theorem 1.14] For a positive integer  $i$ ,

$$\Upsilon_{T_{2,2i+1}}(t) = -i \cdot t \quad \text{for } t \in [0, 1].$$

Proposition 4.1 and Lemma 2.1 imply the following lower bound for the 4-dimensional clasp number. Compare with [Liv17, Theorem 13.1] for Lemma 4.2 and its proof via Lemma 2.1, and compare with [JZ20, Proposition 2.1], where Lemma 4.2 is derived from the stronger bound using  $v^+$  given in [HW16, BCG17]. More on  $v^+$  below.

LEMMA 4.2. *If  $J$  is a knot in  $S^3$ , then*

$$c_4(J) \geq \max_{t \in (0,1]} \Upsilon_J(t)/t + \max_{t \in (0,1]} -\Upsilon_J(t)/t.$$

In [HW16], Hom and Wu define a non-negative integer valued smooth concordance invariant  $v^+$  for knots in  $S^3$ . Following [KP18] (see also [Hom17]), we say two knots  $J$  and  $J'$  are  $v^+$ -equivalent if

$$v^+(J\# - J') = v^+(J'\# - J) = 0,$$

and it is straight forward to verify that this forms a equivalence relation on the set of concordance classes of knots. By [OSS17, Proposition 4.7], we have that  $v^+$ -equivalent knots have the same  $\Upsilon$ -invariant. Recall that  $D$  denotes the positive untwisted Whitehead double of the right-handed trefoil. The key fact that we will use is that  $D$  and  $T_{2,3}$  are  $v^+$ -equivalent [HKL16, Proposition 6.1]. Furthermore, we have the following proposition. Recall that  $J_{p,q}$  denotes the  $(p, q)$ -cable of a knot  $J$ , where  $p$  is the longitudinal winding.

PROPOSITION 4.3. *If  $K_i = D_{2,2i+1}\# - T_{2,2i+1}\# - D$ , then  $K_i$  and  $(T_{2,3})_{2,2i+1}\# - T_{2,2i+3}$  are  $v^+$ -equivalent.*

*Proof.* As mentioned above  $D$  and  $T_{2,3}$  are  $v^+$ -equivalent [HKL16, Proposition 6·1]. Furthermore, for positive integer  $n$ , we have that  $\#^n T_{2,3}$  and  $T_{2,2n+1}$  are also  $v^+$ -equivalent [HKL16, Theorem B·1]. Moreover, [KP18, Theorem B] implies that  $(T_{2,3})_{2,2i+1}$  and  $D_{2,2i+1}$  are  $v^+$ -equivalent. Hence the proof is complete by noting that  $v^+$ -equivalence forms an equivalence relation on the set of concordance classes of knots and the fact that  $K_1 \# K_2$  and  $J_1 \# J_2$  are  $v^+$ -equivalent if  $K_i$  and  $J_i$  are  $v^+$ -equivalent for  $i = 0, 1$  (see e.g. [KKP19, Prop. 3.12]).

Finally, we compute the  $\Upsilon$ -invariant of  $D_{2,2i+1} \# -T_{2,2i+1} \# -D$ . By Proposition 4·3, we only need to compute the  $\Upsilon$ -invariant of  $(T_{2,3})_{2,2i+1} \# -T_{2,2i+3}$ . Note that  $(T_{2,3})_{2,2i+1}$  and  $T_{2,2i+3}$  are both  $L$ -space knots and for each  $L$ -space knot  $J$  there is a formal semigroup  $S_J$ , a subset of  $\mathbb{Z}_{\geq 0}$ , associated to  $J$  [Wan18]. For positive integers  $a_1, a_2, \dots, a_\ell$ , let

$$\langle a_1, a_2, \dots, a_\ell \rangle := \{c_1 a_1 + c_2 a_2 + \dots + c_\ell a_\ell \mid c_i \in \mathbb{Z}_{\geq 0} \text{ for } i = 1, 2, \dots, n\}.$$

For instance, the formal semigroup associated to a positive torus knot  $T_{p,q}$  is  $\langle p, q \rangle$ . More generally, the formal semigroups of iterated cables of torus knots that are  $L$ -space knots can be computed. We only state the simplest case.

LEMMA 4·4 ([Wan18, Proposition 2·7]). *If  $(T_{p,q})_{r,s}$  is an  $L$ -space knot, then the formal semigroup  $S_{(T_{p,q})_{r,s}}$  for  $(T_{p,q})_{r,s}$  is  $\langle pr, qr, s \rangle$ .*

The  $\Upsilon$ -invariant of an  $L$ -space knot can be computed in terms of its formal semigroup (see also [BL16, Propotion 4.4] for algebraic knots).

LEMMA 4·5 ([Wan18, Proposition 3·2]). *If  $J$  is an  $L$ -space knot with  $g_4(J) = g$  and  $S_J$  is the formal semigroup associated to  $J$ , then*

$$\Upsilon_J(t) = \max_{m \in \{0, \dots, 2g\}} \{-2\#(S_J \cap [0, m]) + (m - g) \cdot t\}.$$

We need one last computational lemma before we prove Theorem 1·5.

LEMMA 4·6. *If  $K_i = D_{2,2i+1} \# -T_{2,2i+1} \# -D$  where  $i > 1$ , then*

$$\Upsilon_{K_i}(t) = \begin{cases} -t & \text{for } t \in [0, 1/2], \\ -2 + 3t & \text{for } t \in [1/2, 1]. \end{cases}$$

*Proof.* By Proposition 4·3,  $K_i$  and  $(T_{2,3})_{2,2i+1} \# -T_{2,2i+3}$  are  $v^+$ -equivalent and hence have the same  $\Upsilon$ -invariant. Moreover, by Proposition 4·1 (i) and (iii), it suffices to compute the  $\Upsilon$ -invariant of  $(T_{2,3})_{2,2i+1}$ . For  $i > 1$ , note that  $(T_{2,3})_{2,2i+1}$  is an  $L$ -space knot since  $2i + 1 \geq 2(2g_4(T_{2,3}) - 1) = 2$  [Hed09, Theorem 1·10] and note that  $g_4((T_{2,3})_{2,2i+1}) = i + 2$ . Hence, by Lemma 4·4 and Lemma 4·5, we obtain

$$\Upsilon_{(T_{2,3})_{2,2i+1}}(t) = \max_{m \in \{0, \dots, 2i+4\}} \{-2\#(\langle 4, 6, 2i + 1 \rangle \cap [0, m]) + (m - i - 2) \cdot t\}.$$

We claim that for  $t \in [0, 1]$ ,

$$\max_{m \in \{1, \dots, 2i+4\}} \{-2\#(\langle 4, 6, 2i + 1 \rangle \cap [0, m]) + (m - i - 2) \cdot t\} = -2 + (2 - i) \cdot t. \tag{4·1}$$

Indeed, if  $1 \leq m \leq 3$  and  $t \in [0, 1]$ , then

$$\begin{aligned} -2\#((4, 6, 2i + 1) \cap [0, m)) + (m - i - 2) \cdot t &= -2 + (m - i - 2) \cdot t \\ &\leq -2 + (2 - i) \cdot t. \end{aligned}$$

If instead  $m = 4$  and  $t \in [0, 1]$ , then

$$-2\#((4, 6, 2i + 1) \cap [0, m)) + (m - i - 2) \cdot t = -2 + (2 - i) \cdot t.$$

Finally, if  $5 \leq m \leq 2i + 4$  and  $t \in [0, 1]$ , since

$$\langle 4, 6 \rangle = \{4, 6, 8, 10, 12, \dots\} \subset \langle 4, 6, 2i + 1 \rangle$$

we have

$$\begin{aligned} -2\#((4, 6, 2i + 1) \cap [0, m)) + (m - i - 2) \cdot t &\leq -2(\lceil m/2 \rceil - 1) + (m - i - 2) \cdot t \\ &\leq -2\lceil m/2 \rceil + 2 + (2\lceil m/2 \rceil - i - 2) \cdot t \\ &\leq -2 + (2 - i) \cdot t, \end{aligned}$$

where the last inequality follows from the fact that if  $5 \leq m \leq 2i + 4$  and  $t \in [0, 1]$ , then

$$(-2\lceil m/2 \rceil + 4) \leq (-2\lceil m/2 \rceil + 4) \cdot t.$$

Hence we have verified equation (4.1). Finally, note that this implies that

$$\begin{aligned} \Upsilon_{(T_{2,3})_{2,2i+1}}(t) &= \max_{m \in \{0,4\}} \{-2\#((4, 6, 2i + 1) \cap [0, m)) + (m - i - 2) \cdot t\} \\ &= \begin{cases} -(i + 2) \cdot t & \text{for } t \in [0, 1/2], \\ -2 + (2 - i) \cdot t & \text{for } t \in [1/2, 1]. \end{cases} \end{aligned}$$

Combining the above with Proposition 4.1 (i) and (iii) yields the desired formula.

*Proof of Theorem 1.5.* Let  $K_i = D_{2,2i+1} \# -T_{2,2i+1} \# -D$ , where  $i > 1$ . The knot  $D$  is a topologically slice knot [Fre82], which implies that  $K_i$  is topologically concordant to  $T_{2,2i+1} \# -T_{2,2i+1}$  which is slice. Hence  $K_i$  is topologically slice.

We claim that  $g_4(\#^n K_i) = n$ . First, we show that  $g_4(K_i) = 1$  by following the same argument as in [HW16, Lemma 3.3]. We consider a genus  $i + 2$  Seifert surface  $\Sigma$  for  $D_{2,2i+1}$  obtained by taking two parallel copies of the genus one Seifert surface for  $D$  and connecting them with  $i$  half-twisted bands. Note that there is a genus  $i + 1$  Seifert surface  $\Sigma'$  for the knot  $D \# T_{2,2i+1}$  embedded in  $\Sigma$ . The Seifert surface  $\Sigma'$  is obtained by taking the connected sum with the slightly pushed in Seifert surface for  $D$  and the surface which is obtained by pushing in the half-twisted bands. Now, we consider a Seifert surface  $\tilde{\Sigma}$  for  $K_i$  obtained by taking the boundary connected sum of  $\Sigma$  with the Seifert surface for  $-T_{2,2i+1} \# -D$  with genus  $i + 1$ . Hence the genus of  $\tilde{\Sigma}$  is  $2i + 3$ . Moreover, taking the boundary connected sum of  $\Sigma'$  with a slightly pushed in Seifert surface for  $-T_{2,2i+1} \# -D$  yields a Seifert surface  $\tilde{\Sigma}'$  for  $J = T_{2,2i+1} \# D \# -T_{2,2i+1} \# -D$  embedded in  $\tilde{\Sigma}$  with genus  $2i + 2$ . Performing surgery along  $J$  on  $\tilde{\Sigma}$  in  $B^4$  yields a genus 1 surface for  $K_i$ . Hence we conclude that  $g_4(K_i) \leq 1$ . Moreover, by Lemma 4.6 and Proposition 4.1 (i) and (ii), we have  $g_4(\#^n K_i) = n$  for any positive integer  $n$ .

We conclude the proof by applying Lemma 4.2 to

$$\Upsilon_{K_i}(t)/t = -1 \quad \text{for } t \in (0, 1/2] \quad \text{and} \quad \Upsilon_{K_i}(t)/t = 1 \quad \text{for } t = 1 \quad (\text{Lemma 4.6}).$$

*Acknowledgements.* We thank Jennifer Hom, Allison Miller, Patrick Orson and Mark Powell for helpful discussions. Also, we are grateful to our anonymous referee for thoughtful suggestions. PF gratefully acknowledges support by the SNSF Grant 181199.

## REFERENCES

- [BCG17] J. BODNÁR, D. CELORIA, and M. GOLLA. A note on cobordisms of algebraic knots. *Algebr. Geom. Topol.*, **17**(4):2543–2564, 2017.
- [BL16] M. BORODZIK and C. LIVINGSTON. Semigroups,  $d$ -invariants and deformations of cuspidal singular points of plane curves. *J. Lond. Math. Soc. (2)*, **93**(2):439–463, 2016.
- [Bri66] E. BRIESKORN. Beispiele zur Differentialtopologie von Singularitäten. *Invent. Math.*, **2** (1966), 1–14.
- [CNT20] A. CONWAY, M. NAGEL, and E. TOFFOLI. Multivariable signatures, genus bounds, and 0.5-solvable cobordisms. *Michigan Math. J.*, **69**(2) (2020), 381–427.
- [DS20a] A. DAEMI and C. SCADUTO. Chern-simons functional, singular instantons, and the four-dimensional clasp number. arXiv:2007.13160, (2020).
- [DS20b] A. DAEMI and C. SCADUTO. Equivariant singular instanton homology, virtual talk in nearly carbon neutral geometry and topology conference. Available at <http://ncngt.org/2020/05/16/floer-theory-and-low-dimensional-topology-gauge-theory/>, (June 2020).
- [Fel14] P. FELLER. Gordian adjacency for torus knots. *Algebr. Geom. Topol.*, **14**(2) (2014), 769–793.
- [FLZ17] S. FRIEDL, C. LIVINGSTON, and R. ZENTNER. Knot concordances and alternating knots. *Michigan Math. J.*, **66**(2) (2017), 421–432.
- [FP19] P. FELLER and J. PARK. Genus one cobordisms between torus knots. arXiv:1910.01672 to appear: *Int. Math. Res. Not. IMRN*, (2019).
- [Fre82] M. H. FREEDMAN. The topology of four-dimensional manifolds. *J. Differential Geom.*, **17**(3) (1982), 357–453.
- [GG05] J.-M. GAMBAUDO and É. GHYS. Braids and signatures. *Bull. Soc. Math. France*, **133**(4) (2005), 541–579.
- [Hed09] M. HEDDEN. On knot Floer homology and cabling. II. *Int. Math. Res. Not. IMRN*, (12) (2009), 2248–2274.
- [HKL16] M. HEDDEN, S.-G. KIM, and C. LIVINGSTON. Topologically slice knots of smooth concordance order two. *J. Differential Geom.*, **102**(3) (2016), 353–393.
- [Hom17] J. HOM. A survey on Heegaard Floer homology and concordance. *J. Knot Theory Ramifications*, **26**(2) (2017), 1740015, 24.
- [HW16] J. HOM and Z. WU. Four-ball genus bounds and a refinement of the Ozváth-Szabó tau invariant. *J. Symplectic Geom.*, **14**(1) (2016), 305–323.
- [JZ20] A. JUHÁSZ and I. ZEMKE. Concordance invariants with applications to the 4-dimensional clasp number. arXiv:2007.07106, (2020).
- [KKP19] M. H. KIM, D. KRATOVICH, and J. PARK. Links with non-trivial Alexander polynomial which are topologically concordant to the Hopf link. *Trans. Amer. Math. Soc.*, **371**(8) (2019), 5379–5400.
- [KM19] P. B. KRONHEIMER and T. S. MROWKA. Instantons and some concordance invariants of knots. arXiv:1910.11129, (2019).
- [KP18] M. H. KIM and K. PARK. An infinite-rank summand of knots with trivial Alexander polynomial. *J. Symplectic Geom.*, **16**(6) (2018), 1749–1771.
- [Kro20] P. B. KRONHEIMER. Genus versus double-points for immersed surfaces, virtual talk in the regensburg low-dimensional geometry and topology seminar. Available at <https://www-app.uni-regensburg.de/Fakultaeten/MAT/sfb-higher-invariants/index.php/Regensb-urg-low-dimensional-geometry-and-topology-seminar>, (May 2020).
- [KT76] L. H. KAUFFMAN and L. R. TAYLOR. Signature of links. *Trans. Amer. Math. Soc.*, **216** (1976), 351–365.
- [Lev69] J. LEVINE. Knot cobordism groups in codimension two. *Comment. Math. Helv.*, **44** 1969, 229–244.

- [Lit79] R. A. LITHERLAND. Signatures of iterated torus knots. In *Topology of low-dimensional manifolds* (Proc. Second Sussex Conf., Chelwood Gate, 1977), *Lecture Notes in Math.* vol. 722 (Springer, Berlin, 1979), pages 71–84.
- [Liv04] C. LIVINGSTON. Computations of the Ozsváth–Szabó knot concordance invariant. *Geom. Topol.*, **8** (2004), 735–742.
- [Liv17] C. LIVINGSTON. Notes on the knot concordance invariant  $\epsilon$ . *Algebr. Geom. Topol.*, **17**(1) (2017), 111–130.
- [Liv18] C. LIVINGSTON. Signature functions of knots. *Proc. Amer. Math. Soc.*, **146**(10) (2018), 4513–4520.
- [OS16] B. OWENS and S. STRLE. Immersed disks, slicing numbers and concordance unknotting numbers. *Comm. Anal. Geom.*, **24**(5) (2016), 1107–1138.
- [OSS17] P. OZSVÁTH, A. I. STIPSICZ, and Z. SZABÓ. Concordance homomorphisms from knot Floer homology. *Adv. Math.*, **315** (2017), 366–426.
- [Pow17] M. POWELL. The four-genus of a link, Levine–Tristram signatures and satellites. *J. Knot Theory Ramifications*, **26**(2) (2017), 1740008, 28.
- [Tri69] A. G. TRISTRAM. Some cobordism invariants for links. *Proc. Camb. Philo. Soc.*, **66** (1969), 251–264.
- [Wan18] S. WANG. Semigroups of  $L$ -space knots and nonalgebraic iterated torus knots. *Math. Res. Lett.*, **25**(1) (2018), 335–346.