

# Reference genes validation for qPCR normalization in *Deschampsia antarctica* during abiotic stresses

HYOUNGSEOK LEE, JI HYUN KIM, MIRA PARK, IL-CHAN KIM, JOUNG HAN YIM and HONG KUM LEE

*Polar BioCenter, Korea Polar Research Institute (KOPRI), KORDI, Incheon 406-840, Korea*  
*soulaid@kopri.re.kr*

**Abstract:** Quantitative real time PCR is the most sensitive and widely used method for the analysis of gene expression. The choice of one or several reference genes is very important for a normalization process, which should not fluctuate under stress conditions, to reduce error rate and bias during experimental procedure. In the present study, the expression stability of nine reference genes (two actins, two tubulins, two elongation factor 1 $\alpha$ , two ubiquitins, and cyclophilin) during abiotic stresses such as cold, salt, and PEG treatments, was evaluated on *Deschampsia antarctica* plants using geNorm software. Results from various experimental conditions indicated that cyclophilin and elongation factor 1 $\alpha$  were the most stable genes in the leaf and the root, respectively. The expression of the other reference genes varied under stress. The relative quantification of the TACR7 gene varied according to the kind and the number of reference genes used, suggesting the importance of considering the implications of a combination of reference genes under different stress conditions and in different tissues.

Received 21 September 2009, accepted 20 May 2010, first published online 30 July 2010

**Key words:** adaptation, Antarctic, cold, PEG, salt, TACR7

## Introduction

The Antarctic has the most extreme environment for terrestrial plants with 86% of the earth's ice (Alberdi *et al.* 2002). In the maritime Antarctic the climate is less severe than on the continent. Summer air temperatures typically vary over 25°C, with daily means between -1.9°C and 2.3°C whilst summer precipitation is 42 mm in November to 68 mm in March (Rakusa-Suszczewski 1993) for King George Island. *Deschampsia antarctica* Desv., the Antarctic hairgrass, is the only native grass species growing in the maritime Antarctic (Edwards & Smith 1988). Since *D. antarctica* in King George Island usually grows along the seashore, it is confronted by frequent flooding with fresh (from a melting glacier) and salty water (during storms). Due to its habitat and environmental factors in the Antarctic, *D. antarctica* is usually exposed to cold, salt, and desiccant winds, and it might be an ideal model to study adaptation and response of plants to various abiotic stresses.

Transcriptome studies are widely used to understand the molecular mechanism for plant stress responses and to discover novel stress-responsive genes. To accomplish these purposes, sensitive, precise, and reproducible methods are essential to quantify specific mRNA sequences. Among several techniques, qPCR (quantitative real time PCR) is the most sensitive method for the detection of low abundance mRNAs (Gachon *et al.* 2004). It has been widely used for different applications in plant biology, such as expression profiling in developmental processes and under stress conditions (Song *et al.* 2009), detecting GMOs in foods (Marmioli *et al.* 2008), determining copy number and zygosity in transgenic plants (Bubner &

Baldwin 2004), and plant pathogen genotyping (Abd-Elsalam 2003).

The most important element for qPCR analysis is the selection of proper reference genes to avoid bias. Ideally, the conditions of the experiment should not influence the expression of the reference gene (Schmittgen & Zakrajsek 2000). However, many previous studies have shown that reference genes widely used for the quantification of mRNA expression, could vary with the experimental conditions (Thellin *et al.* 1999, Stürzenbaum & Kille 2001, Radonic *et al.* 2004). According to Thellin *et al.* (1999) and Vandesompele *et al.* (2002), at least two or three reference genes should be used as internal standards because the use of a single gene for normalization could lead to relatively large errors. Currently, some reference genes have been well described for the normalization of expression signals. The most common genes are actin, tubulin, and elongation factor 1 $\alpha$  (Gutierrez *et al.* 2008). Ribosomal RNA, cyclophilin, and ubiquitin may also be used (Thomas *et al.* 2003, Nakashima *et al.* 2008, Gutjahr *et al.* 2008).

Recently, a growing number of studies have focussed on the importance of reference genes and the need for their validation for each particular experimental system. Although most studies deal with human or animal tissues, a few have concerned model plants or crop species, for example, wheat (Paolacci *et al.* 2009), barley (Burton *et al.* 2004), rice (Kim *et al.* 2003), poplar (Brunner *et al.* 2004), potato (Nicot *et al.* 2005), grape (Reid *et al.* 2006), peach (Tong *et al.* 2009), Arabidopsis (Czechowski *et al.* 2005), and tobacco (Volkov *et al.* 2003). To our knowledge, no reports have so far described the suitability of reference

**Table 1.** Primer sequences for internal genes used in this study.

Gene symbol	Accession no.	Similarity	Subject	Highest BLASTX similarity	Sequence of primer pair	Amplicon size (bp)
ACT1	HM208294	100	AAN59956.1, actin ( <i>Hordeum vulgare</i> )		F: GCTCAAACAATCCAAACCCAAACCA R: CTGCATGTGCAAGTGCCAAAGTACA	107
ACT2	HM208295	99	BAD23897.1, actin ( <i>Triticum aestivum</i> )		F: TCTGTGGACGTGGACCAATGTTA R: AGACTGAGTCGCTGAAACCAACT	132
aTUB	HM208296	100	ABD92939.1, alpha tubulin-5D ( <i>Triticum aestivum</i> )		F: GACCATCAGAGTACACGAAACCGAT R: CAGGTGCTTAATTGGGTGGCATT	139
bTUB	HM208297	100	AAD10489.1, beta-tubulin 3 ( <i>Triticum aestivum</i> )		F: TGTCCGATTTGGTGGTCTTGTGGT R: ATCAGATTCCGACGCTCAAAGCA	143
CYP	HM208298	96	ACF49505.1, cyclophilin B-D ( <i>Aegilops tauschii</i> )		F: GCGCATCACCTCCAAATTTCCCA R: ATAGAACTACCGTGCCTGGTTGT	165
EF1a1	HM208299	98	CAA80666.1, elongation factor-1 alpha ( <i>Hordeum vulgare</i> subsp. <i>vulgare</i> )		F: TTTGTCCACTGCTACACTCGTGGT R: TCGAAGCTGACCGACACATAACCAA	98
EF1a2	HM208300	100	ADB28897.1, elongation factor 1-alpha-like protein ( <i>Lolium perenne</i> )		F: ACTTTAACCGAGCATCGAGGACA R: TACACACTCAGTGGTGGCGGTTTA	170
UBQ1	HM208301	99	ACA24140.1, ubiquitin ( <i>Lolium temulentum</i> )		F: CGAGAACTGAAITGAGCTTGGT R: TCGGTGCGCATCATCTGAGTAACA	128
UBQ2	HM208302	100	AAA62699.1, ubiquitin ( <i>Hordeum vulgare</i> subsp. <i>vulgare</i> )		F: CGTCCATCGTATTGAAAGTAGCA R: TGATAATATCCAGCACGACACGA	102
TACR7	FL687117	69	AAC37511.2, pTACR7 ( <i>Triticum aestivum</i> )		F: TTTAGACTCGAGTAGCACCCGC R: ACAGACGCTCAATAAAGGATCACCGG	93

genes for qPCR studies of differential expression of genes from plants in extreme environmental conditions, even though they could be the best targets for studying stress responsive gene expression. In the present work, the expression variability of nine genes is reported in order to assess their value as reference genes in gene expression studies on abiotic stresses from cold, salt, and PEG treatment.

## Materials and methods

### Plant materials and stress treatments

*Deschampsia antarctica* Desv. (Poaceae) plants were collected close to the Korean King Sejong Antarctic Station (62°14'29"S, 58°44'18"W), on the Barton Peninsula of King George Island, during two Antarctic summers (January 2006 and 2007). Plants were placed in plastic containers and transported to Korea, where they were cultivated in a growing chamber in clay sand with 16h/8h of day/night cycles at 15°C. Cold treatment of plants was performed at 4°C with regard to the control at 15°C. For salt and PEG treatments, plants were treated with 250 mM NaCl and 40% PEG6000 respectively, whereas the control was treated with water. Plant leaves and roots were sampled at indicated time points to determine the results of stress conditions.

### Total RNA extraction


Total RNA extraction was performed for three biological replicates by the RNeasy plant mini kit (Qiagen, Leusden, The Netherlands) according to the manufacturer's instructions. The quantity and quality of total RNA was determined by spectroscopic measurements at 230, 260 and 280 nm using ND-1000 spectrophotometer (NanoDrop Technologies, USA), and the integrity was checked by electrophoresis in 1% agarose gel.

### Primer design

Specific primers were designed for reference genes with the PrimerQuest<sup>SM</sup> software in IDT homepage (<http://eu.idtdna.com/scitools/applications/primerquest/default.aspx>, accessed March 2009). Each primer sequence was chosen because it had no similarity with any gene sequence in *Arabidopsis* and rice databases available in NCBI with 170 bp maximum length, optimal  $T_m$  at 60°C, GC% between 40% and 60% (Table 1).

### Reverse transcription and qPCR


Two micrograms of each RNA sample was reverse transcribed to cDNA with MMLV reverse transcriptase (Enzymomics, Korea) using random hexamers after treatment with RNase-free DNase (Promega, USA). Oligo primer molecules and dNTPs were eliminated by using PCR purification kit according to the manufacturer's instructions (Intron biotech, Korea). The cDNA



**Fig. 1.** RNA transcription levels of reference genes tested, presented as mean of C<sub>q</sub> value in the different *D. antarctica* samples. **a.–c.** Leaves exposed to cold, salt, and PEG treatment for indicated time, and their corresponding controls. **d.–f.** Roots exposed to cold, salt, and PEG treatment for indicated time, and their corresponding controls. C<sub>q</sub> values are mean of three replicates.

concentration was checked using ND-1000 spectrophotometer (NanoDrop Technologies, USA). PCR was performed using SYBR Green I technology on Rotor-Gene 6000 Real-time PCR cyclor (Corbett, Australia). A master mix for each PCR run was prepared with SYBR Premix Ex Taq II (TaKaRa, Japan). Two ng of cDNA were added for 10 ul reaction. 300 nM each for specific sense and anti-sense primers were used. PCR cycling with an annealing temperature of 55°C were performed in four replicates from the same RNA preparation and the mean value was considered. The real-time PCR efficiency was determined

for each gene and each stress with the slope of a linear regression model (Pfaffl 2001). For each gene, PCR efficiency was determined by measuring the C<sub>q</sub> to a specific threshold (Walker 2002). For this process, each of the cDNA samples were mixed and then used as PCR templates in a range of 20, 10, 5, 2, 1, 0.5, 0.2 and 0.1 ng. The standard curve, C<sub>q</sub> value and PCR efficiency for each gene were acquired automatically using Rotor-Gene 6000 Series Software ver. 1.7.40. All PCRs displayed efficiencies between 1.9 and 2.1. PCR product sizes were checked on a 3% agarose gel. All corresponded to the




**Fig. 2.** Average expression stability values of control genes by geNorm analysis. M values for each reference gene in the leaf and the root under **a.** & **e.** cold, **b.** & **f.** salt, and **c.** & **g.** PEG treatment, respectively. **d.** & **h.** Determination of the optimal number of control genes for normalization by geNorm analysis in the leaf and the root.

expected size. Melting curves showed a single amplified product. In order to verify the amplification products, their sequences were determined using amplicon specific primers, and analysed with BLAST 2 sequences software (<http://www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html>, accessed June 2009). All amplified sequences had 100% identities with sequences in database.

### Gene expression analysis

Expression levels were determined as the number of cycles needed for the amplification to reach a threshold fixed in the exponential phase of PCR reaction ( $C_q$ ). The  $C_q$  were transformed into quantity using PCR efficiency according to Vandesompele *et al.* (2002). Average expression stability



**Fig. 3.** Relative quantification of *D. antarctica* TACR7 gene expression using various combinations of reference genes as the internal control. **a. & d.** During cold stress, **b. & e.** During salt stress, **c. & f.** During PEG treatment from leaves and roots, respectively.

values and the optimal number of control genes for normalization were calculated by using geNorm software (<http://allserv.ugent.be/jvdesomp/genorm/index.html>, accessed June 2009). The TACR7 expression levels were normalized using the geometric mean of selected reference gene quantities and SD values were calculated with the mean values of three biological replicates in Microsoft Excel following the guidelines described in the geNorm manual ([http://medgen.ugent.be/~jvdesomp/genorm/geNorm\\_manual.pdf](http://medgen.ugent.be/~jvdesomp/genorm/geNorm_manual.pdf), accessed June 2009).

**Results**

*Variations of reference gene expression*

In this study, the expression levels of nine genes were assessed in six types of *D. antarctica* samples representing different tissues under various abiotic stresses. These genes, selected because of their common use as references in qPCR analysis, were ACT1, ACT2, aTUB, bTUB, CYP, EF1a1, EF1a2, UBQ1, and UBQ2 (Table I). Each primer sequence was designed from 3' UTR with no homology to

any sequence in NCBI database to avoid non-specific amplification.

To evaluate the stability of expression of the nine reference genes, RNA transcription levels for all samples were measured for each stress (Fig. 1). One of elongation factor 1 $\alpha$ , EF1a2, showed the highest expression level (low  $C_q$  value) and aTUB showed the lowest expression in all conditions tested. Values are represented as mean of  $C_q$  from 12 experiments comprised of three biological replications and four technical replications. Variation of experiments was assessed as SD for each reference gene. As a result of calculation with  $C_q$  in all the experiments for leaves and roots under various stresses, CYP showed the lowest SD value of 0.47, while aTUB showed the highest value of 0.73, and others were between 0.48 and 0.58. In order to validate and select the best reference genes, two parameters of M (average expression stability) and V (pairwise variation) were used to quantify the reference gene stability. A low M value is indicative of a more stable expression, hence, increasing the suitability of a particular gene as a control gene. Based on a cut-off value of 0.15 for the pairwise variation below which the inclusion of an additional control gene is not required, we found that the most stable reference genes differed depending on the stress condition and the tissue of *D. antarctica* (Fig. 2).

In the cold stress in leaves, the most stable genes were CYP and EF1a1 (Fig. 2a). The M value obtained for these two genes was 0.26. Since the pairwise variation was higher than 0.15 (0.191) (Fig. 2d), two additional genes should be added to get V value below the cut-off. The two most stable genes in leaves for the salt stress were also CYP and EF1a1 with M value of 0.414 (Fig. 2b). For the PEG treatment in leaves, ACT1 and EF1a1 were the two most stable genes with M value of 0.22 (Fig. 2c). In salt conditions, a third gene should be added to make pairwise variation below 0.15, and in PEG treatment, two genes would be sufficient (Fig. 2d). EF1a1 showed the most stable expression in all three conditions. The least stable genes in leaves during cold, salt, and PEG treatments were ACT2, aTUB, and UBQ1 (Fig. 2a–c).

In the root tissues, the stability of reference genes varied depending on the stress. For the salt stress, the most stable genes were ACT1 and CYP with the M value of 0.36 (Fig. 2f). Since the pairwise variation was higher than 0.15 (0.193) (Fig. 2h), two more genes are needed to make V value below the cut-off. The two most stable genes for cold stress were ACT1 and EF1a2 (Fig. 2e), with aTUB and bTUB for PEG treatment (Fig. 2g). The M values were 0.346 and 0.321 for cold and PEG treatment, respectively (Fig. 2h). These two conditions had pairwise variations below the cut-off (0.134 and 0.112), thus only two genes would be sufficient as the reference.

#### *TACR7* gene expression

The expression level for TACR7 as the gene of interest was quantified. Its expression was greatly increased by the

cold stress. The gene expression level was the highest in leaves after 6 hr of treatment and had increased over 100-fold compared with the control (Fig. 3a). After 12 hr of treatment, the level decreased but was still 50-fold higher compared with the control. Though the levels were much lower than those from leaves were, the gene expression in roots increased approximately 12-fold compared with the control (Fig. 3d). For the other treatments of salt and PEG, the level of TACR7 expression remained constant or showed a slight fluctuation independent of the length of treatment (Fig. 3b, c, e, & f).

CYP and ACT1 were the most stable reference genes for most stresses in this study. To investigate the difference between single and multiple genes used as reference, the expression level of the TACR7 gene was quantified with CYP or ACT1 singly, and with multiple reference genes. As a result, differences in quantification were detected according to the combination of reference genes used (Fig. 3). Compared with the use of a unique internal control, using multiple reference genes as the internal control did not significantly change the quantification or the pattern of the expression level of TACR7 (Fig. 3). However, when CYP was used as single reference gene, the quantification of the gene expression was slightly overestimated for cold and salt stresses in the leaf tissues (Fig. 3a & b).

#### Discussion

The reliability of qPCR data would be greatly improved by inclusion of a reference gene whose transcription level is invariable in the different experimental conditions. Expression stability analyses showed that the most stable reference gene identified by geNorm software was not the same for the six datasets analysed (Fig. 2). As shown in previous reports, it is not possible to find an ideal reference gene for all purposes (Reid *et al.* 2006, Paolacci *et al.* 2009, Tong *et al.* 2009). Instead, Gutierrez *et al.* (2008) highlighted the importance of pilot studies to select the most suitable reference gene for specific experimental conditions.

When the six datasets originating from the combination of three treatments and two plant tissues were analysed, geNorm identified EF1a1 and ACT1 as the most stable gene in leaves under various abiotic stresses (Fig. 2a & b). Besides ACT1, bTUB and EF1a2 were ranked in the five most stable genes in all three different stresses to leaves. However, in roots, EF1a1 was the only gene ranked in the five most stable genes in all three different stresses (Fig. 2e–g). For the cold and the salt stress for roots, ACT1 and CYP belonged to the three most stable genes (Fig. 2e & f). However, they are not suitable as control genes for analysing the samples that were subjected to PEG treatment because they were the two least stable genes, wherein aTUB and bTUB performed much better (Fig. 2g). Comparing expression stabilities in different tissues for the same stress, CYP and ACT1, showed high stabilities in the

cold, CYP and EF1a1 for the salt stress, and bTUB for the PEG treatment (Fig. 2).

For the TACR gene, relative quantification varied depending on the reference gene used (Fig. 3). When compared with the use of a unique internal control, the use of multiple reference genes as the internal control did not change significantly the quantification or the pattern of the expression level of TACR7. Accordingly, the use of single gene as the internal standard could be a valuable alternative to quantify a gene of interest, keeping in mind that it could overestimate the variations of expression in the cold stress or the salt stress.

TACR7 was originally isolated as a gene designated as *Triticum aestivum* cold regulated 7 from hard red winter wheat where its expression increased in freeze-resistant plants relative to freeze-susceptible plants (Gana *et al.* 1997). Its expression was not induced by ABA or stresses such as salt, dehydration, and heat, so the gene was grouped as low-temperature specific. In addition, the TACR7-homologous gene of *D. antarctica*, has shown a 60-fold increase of transcription level in plants from an Antarctic environment compared to plants grown in 15°C (Lee *et al.* 2008). In this study TACR7 homologue of *D. antarctica* was implicated in cold stress with a maximum of expression at 6 hr after treatment and its expression level was highly maintained 12 hr after treatment in the leaf tissue (Fig. 3a). Expression level in the root increased almost ten times from 1 hr to 12 hr after cold stress (Fig. 3d). The results of cold stress were similar to those shown by Gana *et al.* (1997), where the gene transcripts accumulated during cold exposure in a freeze resistant wheat species. This suggests that the gene product is probably functional in the process of cold adaptation. Furthermore, TACR7 expression remained constant in the leaf under PEG treatment, or showed a small range of fluctuation in the leaf under salt stress, in roots under salt stress and PEG treatment. Thus, this study indicated that TACR7 in *D. antarctica* might be implicated in the cold adaptation and should be designated as low-temperature-specific gene.

Although in some of our experimental conditions a single reference gene was adequate to normalize the qPCR data, the use of multiple reference genes should be considered in future studies to improve the accuracy and reliability of gene expression analysis. The specific primer set of qPCR developed in this study for the stable reference genes for each stress conditions and plant tissue types, would enable us to perform more accurate normalization and quantification of gene expression in *D. antarctica*.

### Acknowledgements

We thank the referees whose comments helped improve this manuscript. This work was supported by a grant PE09050 from the Korea Polar Research Institute.

### References

- ABD-ELSALAM, K.A. 2003. Non-gel based techniques for plant pathogen genotyping. *Acta Microbiologica Polonica*, **52**, 329–341.
- ALBERDI, M., BRAVO, L.A., GUTIÉRREZ, A.H., GIDEKEL, M. & CORCUERA, L.J. 2002. Ecophysiology of Antarctic vascular plants. *Physiologia Plantarum*, **115**, 479–486.
- BRUNNER, A.M., YAKOVLEV, I.A. & STRAUSS, S.H. 2004. Validating internal controls for quantitative plant gene expression studies. *BMC Plant Biology*, **4**, 14.
- BUBNER, B. & BALDWIN, I.T. 2004. Use of real-time PCR for determining copy number and zygosity in transgenic plants. *Plant Cell Reports*, **5**, 263–271.
- BURTON, R.A., SHIRLEY, N.J., KING, B.J., HARVEY, A.J. & FINCHER, G.B. 2004. The *CesA* gene family of barley. Quantitative analysis of transcripts reveals two groups of co-expressed genes. *Plant Physiology*, **134**, 224–236.
- CZECHOWSKI, T., STITT, M., ALTMANN, T., UDVARDI, M.K. & SCHEIBLE, W.-R. 2005. Genome-wide identification and testing of superior reference genes for transcript normalization in arabidopsis. *Plant Physiology*, **139**, 5–17.
- EDWARDS, J.A. & SMITH, R.I.L. 1988. Photosynthesis and respiration of *Colobanthus quitensis* and *Deschampsia antarctica* from the maritime Antarctic. *British Antarctic Survey Bulletin*, No. 81, 43–63.
- GACHON, C., MINGAM, A. & CHARRIER, B. 2004. Real-time PCR: what relevance to plant studies. *Journal of Experimental Botany*, **55**, 1445–1454.
- GANA, J.A., SUTTON, F. & KENEFICK, D.G. 1997. cDNA structure and expression patterns of a low-temperature-specific wheat gene *TACR7*. *Plant Molecular Biology*, **34**, 643–650.
- GUTIERREZ, L., MAURIAT, M., GUÉNIN, S., PELLOUX, J., LEFEBVRE, J.F., LOUVET, R., RUSTERUCCI, C., MORITZ, T., GUERINEAU, F., BELLINI, C. & VAN WUYTSWINKEL, O. 2008. The lack of a systematic validation of reference genes: a serious pitfall undervalued in reverse transcription-polymerase chain reaction (RT-PCR) analysis in plants. *Plant Biotechnology Journal*, **6**, 609–618.
- GUTJAHR, C., BANBA, M., CROSET, V., AN, K., MIYAO, A., AN, G., HIROCHIKA, H., IMAIZUMI-ANRAKU, H. & PASZKOWSKI, U. 2008. Arbuscular mycorrhiza-specific signaling in rice transcends the common symbiosis signaling pathway. *Plant Cell*, **20**, 2989–3005.
- KIM, B.R., NAM, H.Y., KIM, S.U., KIM, S.I. & CHANG, Y.J. 2003. Normalization of reverse transcription quantitative-PCR with housekeeping genes in rice. *Biotechnology Letters*, **25**, 1869–1872.
- LEE, H., CHO, H.H., KIM, I.C., YIM, J.H., LEE, H.K. & LEE, Y.K. 2008. Expressed sequence tag analysis of Antarctic hairgrass *Deschampsia antarctica* from King George Island, Antarctica. *Molecules and Cells*, **25**, 258–264.
- MARMIROLI, N., MAESTRI, E., GULLI, M., MALCEVSCHI, A., PEANO, C., BORDONI, R. & DE BELLIS, G. 2008. Methods for detection of GMOs in food and feed. *Analytical and Bioanalytical Chemistry*, **392**, 369–384.
- NAKASHIMA, A., CHEN, L., PHUONG THAO, N., FUJIWARA, M., WONG, H.L., KUWANO, M., UMEMURA, K., SHIRASU, K., KAWASAKI, U. & SHIMAMOTO, K. 2008. RACK1 functions in rice innate immunity by interacting with the Rac1 immune complex. *Plant Cell*, **20**, 2265–2279.
- NICOT, N., HAUSMAN, J.F., HOFFMANN, L. & EVERS, D. 2005. Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. *Journal of Experimental Botany*, **56**, 2907–2914.
- PAOLACCI, A., TANZARELLA, O., PORCEDDU, E. & CIAFFI, M. 2009. Identification and validation of reference genes for quantitative RT-PCR normalization in wheat. *BMC Molecular Biology*, **10**, 11.
- PEAFFL, M.W. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, **29**, 2002–2007.
- RADONIC, A., THULKE, S., MACKAY, I.M., LANDT, O., SIEGERT, W. & NITSCHKE, A. 2004. Guideline to reference gene selection for quantitative real-time PCR. *Biochemical and Biophysical Research Communications*, **313**, 856–862.
- RAKUSA-SUSZCZEWSKI, S., ed. 1993. *The maritime Antarctic coastal ecosystem of Admiralty Bay*. Warsaw: Department of Antarctic Biology, Polish Academy of Sciences, 216 pp.

- REID, K.E., OLSSON, N., SCHLOSSER, J., PENG, F. & LUND, S.T. 2006. An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development. *BMC Plant Biology*, **14**, 6–27.
- SCHMITTGEN, T. & ZAKRAJSEK, B.A. 2000. Effect of experimental treatment on housekeeping gene expression: validation by real-time, quantitative RT-PCR. *Journal of Biochemical and Biophysical Methods*, **46**, 69–81.
- SONG, Y., WANG, L. & XIONG, L. 2009. Comprehensive expression profiling analysis of *OsLAA* gene family in developmental processes and in response to phytohormone and stress treatments. *Planta*, **229**, 577–591.
- STÜRZENBAUM, S.R. & KILLE, P. 2001. Control genes in quantitative molecular biological techniques: the variability of invariance. *Comparative Biochemistry and Physiology*, **B130**, 281–289.
- THELLIN, O., ZORZI, W., LAKAYE, B., DE BORMAN, B., COUMANS, B., HENNEN, G., GRISAR, T., IGOUT, A. & HEINEN, E. 1999. Housekeeping genes as internal standards: use and limits. *Journal of Biotechnology*, **75**, 291–295.
- THOMAS, C., MEYER, D., WOLFF, M., HIMBER, C., ALIOUA, M. & STEINMETZ, A. 2003. Molecular characterization and spatial expression of the sunflower *ABP1* gene. *Plant Molecular Biology*, **52**, 1025–1036.
- TONG, Z., GAO, Z., WANG, F., ZHOU, J. & ZHANG, Z. 2009. Selection of reliable reference genes for gene expression studies in peach using real-time PCR. *BMC Molecular Biology*, **10**, 71.
- VANDESOMPELE, J., DE PRETER, K., PATTYN, F., POPPE, B., VAN ROY, N., DE PAEPE, A. & SPELEMAN, F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, **3**, research0034.1.
- VOLKOV, R.A., PANCHUK, I.I. & SCHÖFFL, F. 2003. Heat-stress-dependency and developmental modulation of gene expression: the potential of house-keeping genes as internal standards in mRNA expression profiling using real-time RT-PCR. *Journal of Experimental Botany*, **54**, 2343–2349.
- WALKER, N.J. 2002. A technique whose time has come. *Science*, **296**, 557–559.