

Zero-full law for well approximable sets in missing digit sets

BY BING LI

School of Mathematics, South China University of Technology, Guangzhou, China, 510641.

e-mail: scbingli@scut.edu.cn

RUOFAN LI

Department of Mathematics, Jinan University, Guangzhou, China, 510632.

e-mail: liruofan@jnu.edu.cn

AND YUFENG WU

School of Mathematics and Statistics, Central South University, Changsha, China, 410083.

e-mail: yufengwu.wu@gmail.com

(Received 16 January 2023; revised 22 January 2024; accepted 27 November 2024)

Abstract

Let $b \geq 3$ be an integer and $C(b, D)$ be the set of real numbers in $[0, 1]$ whose base b expansion only consists of digits in a set $D \subseteq \{0, \dots, b-1\}$. We study how close can numbers in $C(b, D)$ be approximated by rational numbers with denominators being powers of some integer t and obtain a zero-full law for its Hausdorff measure in several circumstances. When b and t are multiplicatively dependent, our results correct an error of Levesley, Salp and Velani (*Math. Ann.* **338** (2007), 97–118) and generalise their theorem. When b and t are multiplicatively independent but have the same prime divisors, we obtain a partial result on the Hausdorff measure and bounds for the Hausdorff dimension, which are close to the multiplicatively dependent case. Based on these results, several conjectures are proposed.

2020 Mathematics Subject Classification: 11J83 (Primary); 11J82, 11K55 (Secondary)

1. Introduction

In an influential paper [18], Mahler asked “How close can irrational elements of Cantor’s set be approximated by rational numbers?” This question has inspired a wide range of research, such as [5, 10, 17, 21] and references therein. In [16], Levesley, Salp and Velani showed that there exist numbers which are not Liouville numbers in the middle-third Cantor set C that can be very well approximated by rational numbers whose denominators are powers of 3. More precisely, let $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function and define

$$W_3(\psi) = \left\{ x \in [0, 1] : \left| x - \frac{p}{3^n} \right| < \psi(n) \text{ for i.m. } (p, n) \in \mathbb{N}^2 \right\},$$

where i.m. is short for “infinitely many”. They proved that the f -Hausdorff measure \mathcal{H}^f (we refer to Section 2 for terminologies) of $W_3(\psi) \cap C$ satisfies a zero-full law.

THEOREM 1.1 ([16, theorem 1]). *Let f be a dimension function such that $f(r)r^{-\log 2/\log 3}$ is monotonic and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then*

$$\mathcal{H}^f(W_3(\psi) \cap C) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} f(\psi(n)) \times 3^{n \log 2/\log 3} < \infty, \\ \mathcal{H}^f(C), & \text{if } \sum_{n=1}^{\infty} f(\psi(n)) \times 3^{n \log 2/\log 3} = \infty. \end{cases}$$

In the same paper, the authors claimed their method can also yield a generalisation of Theorem 1.1. For any integer $b \geq 3$ and set $D \subseteq \{0, \dots, b - 1\}$ with cardinality between 2 and $b - 1$, the *missing digit set* $C(b, D)$ is defined as the set of real numbers in $[0, 1]$ whose base b expansions only consist of digits in D .

Claim 1.2 ([16, theorem 4]). *Let $C(b, D)$ be a missing digit set with Hausdorff dimension γ , f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then*

$$\mathcal{H}^f(W_b(\psi) \cap C(b, D)) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} f(\psi(n))b^{n\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)), & \text{if } \sum_{n=1}^{\infty} f(\psi(n))b^{n\gamma} = \infty. \end{cases}$$

Unfortunately, this claim is not always valid. When the set D does not contain 0 and $b - 1$, all rational numbers of the form pb^{-n} are not in the missing digit set $C(b, D)$, so it is possible to find ψ such that $\sum_{n=1}^{\infty} f(\psi(n)) \times b^{n\gamma}$ diverges while $W_b(\psi) \cap C(b, D) = \emptyset$; see Example 3.1 for a concrete example. The correction of Claim 1.2 is as follows.

THEOREM 1.3. *Let $C(b, D)$ be a missing digit set with Hausdorff dimension γ , $m = \min\{\min D, b - 1 - \max D\}$, f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic, and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then*

$$\begin{aligned} &\mathcal{H}^f(W_b(\psi) \cap C(b, D)) \\ &= \begin{cases} 0, & \text{if } \sum_{n \geq 1: \psi(n) > \frac{m}{(b-1)b^n}} f\left(\psi(n) - \frac{m}{(b-1)b^n}\right) b^{n\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)), & \text{if } \sum_{n \geq 1: \psi(n) > \frac{m}{(b-1)b^n}} f\left(\psi(n) - \frac{m}{(b-1)b^n}\right) b^{n\gamma} = \infty. \end{cases} \end{aligned}$$

It is crucial in Theorem 1.3 to approximate numbers in $C(b, D)$ by rational numbers with denominators b^n , as this matches the structure of $C(b, D)$. One naturally wonders if similar zero-full laws hold when denominators are powers of other numbers. Let $t \geq 2$ be an integer, we consider the Hausdorff measure of $W_t(\psi) \cap C(b, D)$, where

$$W_t(\psi) = \left\{ x \in [0, 1] : \left| x - \frac{p}{t^n} \right| < \psi(n) \text{ for i.m. } (p, n) \in \mathbb{N}^2 \right\}.$$

For $t = 2$, Velani proposed the following conjecture.

CONJECTURE 1.4 (Velani’s conjecture; see [2]). *Suppose $\psi : \mathbb{N} \rightarrow (0, \infty)$ is monotonic, then*

$$\mathcal{H}^{\log 2 / \log 3}(W_2(\psi) \cap C) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} \psi(n) \times 2^n < \infty, \\ 1, & \text{if } \sum_{n=1}^{\infty} \psi(n) \times 2^n = \infty. \end{cases}$$

Some progress towards proving Conjecture 1.4 has been made by various authors; see [1–3]. The main difficulty is that we know very little about how the dyadic rationals are distributed around the middle-third Cantor set, this is similar to the Furstenberg’s conjecture on times two and times three [11].

We are going to consider the case that b and t are multiplicatively dependent, that is, $\log t / \log b \in \mathbb{Q}$. Under this assumption, we prove a zero-full law.

THEOREM 1.5. *Let $C(b, D)$ be a missing digit set with Hausdorff dimension γ such that D contains at least one of 0 and $b - 1$, $t \geq 2$ be an integer which is multiplicatively dependent with b , f be a dimension function such that $r^{-\gamma} f(r)$ is monotonic, and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then*

$$\mathcal{H}^f(W_t(\psi) \cap C(b, D)) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} f(\psi(n))t^{n\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)), & \text{if } \sum_{n=1}^{\infty} f(\psi(n))t^{n\gamma} = \infty. \end{cases}$$

Unlike Theorem 1.3, here we need the assumption that D contains at least one of 0 and $b - 1$ to obtain a complete zero-full law. If this condition is dropped, we are still able to deduce a result for $\mathcal{H}^f(W_t(\psi) \cap C(b, D))$, but the two series for the divergence and convergence parts may be different. Indeed, our method is applicable to more general approximable sets in the case that b and t have the same prime divisors, which is weaker than that p and t are multiplicatively dependent.

Suppose $A = (a_n)_{n \geq 1}$ is a sequence of positive integers and define

$$W_{t,A}(\psi) = \left\{ x \in [0, 1] : \left| x - \frac{p}{t^{a_n}} \right| < \psi(n) \text{ for i.m. } (p, n) \in \mathbb{N}^2 \right\}.$$

Let

$$I(A) = \{i \in \mathbb{N} : a_n = i \text{ for some } n\}.$$

That is, $I(A)$ is the set of elements of the sequence A . For $i \in I(A)$, let

$$\psi_A(i) = \max \{ \psi(n) : a_n = i \}.$$

If the prime divisors of b and t are the same, denote

$$\alpha_1(b, t) = \min \left\{ \frac{v_q(t)}{v_q(b)} : q \text{ is a prime divisor of } b \right\},$$

$$\alpha_2(b, t) = \max \left\{ \frac{v_q(t)}{v_q(b)} : q \text{ is a prime divisor of } b \right\},$$

where $v_q(b)$ means the greatest integer such that $q^{v_q(b)}$ divides b . For the sake of clarity, we will simply write $I(A)$, $\alpha_1(b, t)$ and $\alpha_2(b, t)$ as I , α_1 and α_2 respectively when there

is no confusion. We use the notations $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ to mean the floor and ceiling functions respectively.

THEOREM 1.6. *Suppose $C(b, D)$ is a missing digit set with Hausdorff dimension γ , m is the greatest integer such that*

$$D \subseteq \{m, m + 1, \dots, b - 1 - m\},$$

$t \geq 2$ is an integer that has the same prime divisors as b , and $A = (a_n)_{n \geq 1}$ is an unbounded non-decreasing sequence of positive integers. Let f be a dimension function such that $r^{-\gamma} f(r)$ is monotonic, and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then

$$\mathcal{H}^f(W_{t,A}(\psi) \cap C(b, D)) = \begin{cases} 0 & \text{if } \sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lceil i\alpha_2 \rceil}}} f\left(\psi_A(i) - \frac{m}{(b-1)b^{\lceil i\alpha_2 \rceil}}\right) b^{i\alpha_2\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)) & \text{if } \sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lceil i\alpha_1 \rceil}}} f\left(\psi_A(i) - \frac{m}{(b-1)b^{\lceil i\alpha_1 \rceil}}\right) b^{i\alpha_1\gamma} = \infty. \end{cases}$$

Remarks.

- (i) When b and t are multiplicatively dependent, we have

$$\alpha_1 = \alpha_2 = \frac{\log t}{\log b}$$

and thus Theorem 1.5 follows from Theorem 1.6 by taking $A = (n)_{n \geq 1}$ and using $m = 0$.

- (ii) If $b = t$ and $A = (n)_{n \geq 1}$, then $\alpha_1 = \alpha_2 = 1$ and hence $\lceil n\alpha_2 \rceil = \lfloor n\alpha_1 \rfloor = n$. Therefore Theorem 1.6 also implies Theorem 1.3.
- (iii) In general, the two series in Theorem 1.6 are different, so our formula is inconclusive in the case that the first series diverges while the second series converges.

The rest of this paper is structured as follows. Section 2 includes terminologies and tools needed. The main reason that Claim 1.2 is wrong and what modification is needed are discussed in Section 3. We prove a variation of Theorem 1.6 in Section 4 and then use it to obtain Theorem 1.6 in Section 5. Finally, we discuss Hausdorff dimension and the large intersection property of $W_t(\psi) \cap C(b, D)$ and propose a conjecture for multiplicatively independent case in Section 6.

2. Preliminaries

2.1. Hausdorff measure and dimension

A function $f : (0, \infty) \rightarrow (0, \infty)$ is called a *dimension function* if it is continuous, non-decreasing and $\lim_{r \rightarrow 0} f(r) = 0$. For a set $S \subseteq \mathbb{R}^k$, we say a countable collection of balls $\{B_i\}$ in \mathbb{R}^k is a ρ -cover of S if $S \subseteq \cup_i B_i$ and their radii are not larger than ρ . The *Hausdorff f -measure* \mathcal{H}^f of S is

$$\mathcal{H}^f(S) = \lim_{\rho \rightarrow 0} \mathcal{H}_\rho^f(S),$$

where

$$\mathcal{H}_\rho^f(S) = \inf \left\{ \sum_i f(r(B_i)) : \{B_i\} \text{ is a } \rho\text{-cover of } S \right\},$$

and $r(B_i)$ means the radius of ball B_i .

When $f(r) = r^s$ for some $s \geq 0$, we write \mathcal{H}^f as \mathcal{H}^s . The Hausdorff dimension of a set S is

$$\dim_H S = \inf\{s : \mathcal{H}^s(S) = 0\}.$$

It is known that a missing digit set $C(b, D)$ has Hausdorff dimension $\log \#D / \log b$, where $\#D$ denotes the cardinality of D . This number will be used frequently and we denote it by γ . More properties of Hausdorff measure and dimension can be found in [8].

2.2. Mass transference principle

For two positive numbers x and y , we write $x \ll y$ if there exists a constant $K > 0$ such that $x \leq Ky$. The relation $x \gg y$ is defined similarly and we write $x \asymp y$ if $x \ll y$ and $x \gg y$.

Let X be a compact subset of \mathbb{R}^k and μ be a Borel measure on X . We say μ is δ -Ahlfors regular if there exists constant $r_0 > 0$ such that for any ball $B(x, r) \subseteq X$ with $x \in X$ and radius $r \leq r_0$, we have

$$\mu(B(x, r)) \asymp r^\delta.$$

When X is a missing digit set $C(b, D)$ with dimension γ , the measure $\mathcal{H}^\gamma|_{C(b,D)}$ is γ -Ahlfors regular; see for example [19]. This allows us to use the mass transference principle, a widely-used tool in computing Hausdorff dimension of limsup sets.

THEOREM 2.1 (Mass transference principle, [4]). *Let X be a compact subset of \mathbb{R}^k equipped with a δ -Ahlfors regular measure μ . Let $(B_n)_{n \geq 1}$ be a sequence of balls in X with $r(B_n) \rightarrow 0$ as $n \rightarrow \infty$. Suppose f is a dimension function such that $r^{-\delta}f(r)$ is monotonic. For a ball $B = B(x, r)$, denote $B^f = B(x, f(r)^{1/\delta})$. If for any ball B in X , we have*

$$\mathcal{H}^\delta(B \cap \limsup_{n \rightarrow \infty} B_n^f) = \mathcal{H}^\delta(B),$$

then

$$\mathcal{H}^f(B \cap \limsup_{n \rightarrow \infty} B_n) = \mathcal{H}^f(B)$$

for any ball B in X .

2.3. Measure theoretic lemmas

In this subsection we state several lemmas on measures. The first one is about when a subset has the same measure as the whole set.

LEMMA 2.2 ([16, lemma 1]). *Let X be a compact set in \mathbb{R}^k and μ be a finite measure on X such that all open sets are measurable and $\mu(B(x, 2r)) \ll \mu(B(x, r))$ for all balls $B(x, r)$ with center in X . Suppose E is a Borel subset of X and there exist positive constants r_0, c_0 such that for any ball B with radius $r(B) < r_0$ and center in X , we have $\mu(E \cap B) \geq c_0\mu(B)$. Then*

$$\mu(E) = \mu(X).$$

The second lemma is a generalisation of the divergence part of the Borellonger -Cantelli lemma.

LEMMA 2.3 ([22, lemma 5]; see also [16, lemma 2]). *Let X be a compact set in \mathbb{R}^k and let μ be a finite measure on X . Also, let E_n be a sequence of μ -measurable sets such that $\sum_{n=1}^\infty \mu(E_n) = \infty$. Then*

$$\mu(\limsup_{n \rightarrow \infty} E_n) \geq \limsup_{Q \rightarrow \infty} \frac{\left(\sum_{0 < s \leq Q} \mu(E_s)\right)^2}{\sum_{0 < s, t \leq Q} \mu(E_s \cap E_t)}.$$

In Theorem 1.6, an infinite countable set I is used, and we have the following variation of Lemma 2.3: if $\sum_{i \in I} \mu(E_i) = \infty$, then

$$\mu(\limsup_{i \rightarrow \infty, i \in I} E_i) \geq \limsup_{Q \rightarrow \infty} \frac{\left(\sum_{0 < s \leq Q, s \in I} \mu(E_s)\right)^2}{\sum_{0 < s, t \leq Q, s, t \in I} \mu(E_s \cap E_t)}. \tag{2.1}$$

3. Intersection of balls and the missing digit set

The set $W_b(\psi)$ is the limsup of balls of the form $B(pb^{-n}, \psi(n))$, so we investigate what is the intersection of those balls and $C(b, D)$. We start with an example illustrating why Claim 1.2 is false.

Example 3.1. Let $b = 5, D = \{1, 2\}, \gamma = \log 2 / \log 5$ be the dimension of the missing digit set $C(5, \{1, 2\})$ and

$$\psi(n) = \sum_{i=n+1}^\infty \frac{1}{5^i} = \frac{1}{4 \times 5^n}.$$

Since

$$\sum_{n=1}^\infty \left(\frac{1}{4 \times 5^n}\right)^\gamma \times 5^{n\gamma} = \sum_{n=1}^\infty \frac{1}{4^\gamma} = \infty,$$

Claim 1.2 says

$$\mathcal{H}^\gamma(W_5(\psi) \cap C(5, \{1, 2\})) = \mathcal{H}^\gamma(C(5, \{1, 2\})) > 0.$$

Let $x \in [0, 1]$ be a number in $B(p5^{-n}, \psi(n))$ for some $(p, n) \in \mathbb{N}^2$, then

$$\frac{p-1}{5^n} + \sum_{i=n+1}^\infty \frac{3}{5^i} < x < \frac{p}{5^n} + \sum_{i=n+1}^\infty \frac{1}{5^i},$$

hence the base 5 expansion of x must contain a digit 0 or 4, and thus $x \notin C(5, \{1, 2\})$. Therefore in this case

$$W_5(\psi) \cap C(5, \{1, 2\}) = \emptyset$$

cannot have positive measure.

When the set D does not contain 0 and $b - 1$, we have $pb^{-n} \notin C(b, D)$ for any $(p, n) \in \mathbb{N}^2$, so the intersections of balls $B(pb^{-n}, \psi(n))$ and $C(b, D)$ are all empty unless $\psi(n)$ is not too small; see the proof of Lemma 3.2 for details. To better understand how those intersections are, we introduce several notations. For any $n \geq 1$, let $C_n(b, D)$ be the n th level of $C(b, D)$, which consists of $b^{n\gamma}$ intervals of length b^{-n} . More precisely,

$$C_n(b, D) = \left\{ \sum_{i=1}^{\infty} \frac{x_i}{b^i} \in [0, 1] : x_i \in D \text{ for } i = 1, \dots, n \right\}.$$

Denote the set of all left endpoints of the intervals in $C_n(b, D)$ by L_n and the set of all right endpoints of the intervals in $C_n(b, D)$ by R_n . Note that a point can be both a left and right endpoint. For instance, in Example 3.1, we have $L_1 = \{1/5, 2/5\}$ and $R_1 = \{2/5, 3/5\}$, hence $2/5$ is in both L_1 and R_1 . The two quantities below are used to measure how many digits of $\{0, \dots, b - 1\}$ are missing in D from left and right respectively.

$$m_l = \min\{D\} \quad \text{and} \quad m_r = b - 1 - \max\{D\}.$$

Recall that $m = \min\{m_l, m_r\}$.

LEMMA 3.2 Suppose $\psi(n) < b^{-n}/2$. Denote

$$d_{l,n} = \frac{m_l}{(b - 1)b^n} \quad \text{and} \quad d_{r,n} = \frac{m_r}{(b - 1)b^n}.$$

We make the convention that an open ball with non-positive radius is regarded as an empty set.

(i) If $pb^{-n} \in L_n \setminus R_n$, then

$$B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) = B\left(\frac{p}{b^n} + d_{l,n}, \psi(n) - d_{l,n}\right) \cap C(b, D).$$

(ii) If $pb^{-n} \in R_n \setminus L_n$, then

$$B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) = B\left(\frac{p}{b^n} - d_{r,n}, \psi(n) - d_{r,n}\right) \cap C(b, D).$$

(iii) If $pb^{-n} \in L_n \cup R_n$, then

$$\begin{aligned} & B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) \\ &= \left(B\left(\frac{p}{b^n} + d_{l,n}, \psi(n) - d_{l,n}\right) \cup B\left(\frac{p}{b^n} - d_{r,n}, \psi(n) - d_{r,n}\right) \right) \\ & \quad \cap C(b, D). \end{aligned}$$

(iv) If $pb^{-n} \notin L_n \cup R_n$, then

$$B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) = \emptyset.$$

Proof.

(i) If $m_l = 0$, then $d_{l,n} = 0$ and the equality holds trivially. So we assume $m_l > 0$ without loss of generality. Let $x \in B(pb^{-n}, \psi(n))$. If $x < pb^{-n}$, then $\psi(n) < b^{-n}/2$ implies that

$x \notin C_n(b, D) \supseteq C(b, D)$. If $pb^{-n} \leq x < pb^{-n} + d_{l,n}$, then

$$\frac{p}{b^n} \leq x < \frac{p}{b^n} + \frac{m_l}{(b-1)b^n} = \frac{p}{b^n} + \sum_{i=n+1}^{\infty} \frac{m_l}{b^i},$$

so the base b expansion of x contains a digit between 0 and $m_l - 1$, hence $x \notin C(b, D)$ by the definition of m_l . Then $x \in C(b, D)$ only happens when $x \geq pb^{-n} + d_{l,n}$, which implies that $\psi(n) > d_{l,n}$. Also note that if $x \geq pb^{-n} + d_{l,n}$, then

$$0 \leq x - \left(\frac{p}{b^n} + d_{l,n}\right) = x - \frac{p}{b^n} - d_{l,n} < \psi(n) - d_{l,n},$$

so $x \in B(pb^{-n} + d_{l,n}, \psi(n) - d_{l,n})$. Therefore if $\psi(n) > d_{l,n}$, then

$$B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) \subseteq B\left(\frac{p}{b^n} + d_{l,n}, \psi(n) - d_{l,n}\right) \cap C(b, D)$$

and these two sets are equal since

$$B\left(\frac{p}{b^n} + d_{l,n}, \psi(n) - d_{l,n}\right) \subseteq B\left(\frac{p}{b^n}, \psi(n)\right)$$

is trivial.

- (ii) Again, we may assume $m_r > 0$ without loss of generality. Let $x \in B(pb^{-n}, \psi(n))$. If $x > pb^{-n}$, then $x \notin C_n(b, D)$ since $\psi(n) < b^{-n}/2$. If $pb^{-n} - d_{r,n} \leq x < pb^{-n}$, then

$$\frac{p-1}{b^n} + \sum_{i=n+1}^{\infty} \frac{b-1-m_r}{b^i} \leq x < \frac{p}{b^n},$$

so the base b expansion of x contains a digit between $b - m_l$ and $b - 1$, hence $x \notin C(b, D)$ by the definition of m_r . Then a similar argument as in the previous case shows that

$$B\left(\frac{p}{b^n}, \psi(n)\right) \cap C(b, D) = B\left(\frac{p}{b^n} - d_{r,n}, \psi(n) - d_{r,n}\right) \cap C(b, D)$$

if $\psi(n) > d_{r,n}$ and the intersection on the left is empty when $\psi(n) \leq d_{r,n}$.

- (iii) This part is treated by combining the arguments in previous two cases, hence we skip the details.
- (iv) If $pb^{-n} \notin R_n \cup L_n$, then $\psi(n) < b^{-n}/2$ implies that

$$B(pb^{-n}, \psi(n)) \cap C_n(b, D) = \emptyset,$$

and thus $B(pb^{-n}, \psi(n)) \cap C(b, D) = \emptyset$.

By the definition of m_l and m_r , the new ball centers $pb^{-n} + d_{l,n}$ and $pb^{-n} - d_{l,n}$ are points in the missing digit set $C(b, D)$, hence if $\mu = \mathcal{H}^\gamma|_{C(b, D)}$, then

$$\mu\left(B\left(\frac{p}{b^n} + d_{l,n}, \psi(n) - d_{l,n}\right)\right) \asymp (\psi(n) - d_{l,n})^\gamma, \tag{3.1}$$

$$\mu\left(B\left(\frac{p}{b^n} - d_{r,n}, \psi(n) - d_{r,n}\right)\right) \asymp (\psi(n) - d_{r,n})^\gamma. \tag{3.2}$$

4. A special case

In this section we prove a special case of Theorem 1.6 where $b = t$. It will be used later to deduce Theorem 1.6.

LEMMA 4.1. *Suppose $C(b, D)$ is a missing digit set with Hausdorff dimension γ , $m = \min\{\min D, b - 1 - \max D\}$, and $A = (a_n)_{n \geq 1}$ is an unbounded non-decreasing sequence of positive integers. Let f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic, and $\psi : \mathbb{N} \rightarrow (0, \infty)$ be a function. Then*

$$\mathcal{H}^f(W_{b,A}(\psi) \cap C(b, D)) = \begin{cases} 0, & \text{if } \sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)), & \text{if } \sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} = \infty. \end{cases}$$

Remark. We can assume $\psi_A(i) < b^{-i}/2$ for all i without loss of generality. Indeed, suppose there exists an infinite set I_0 such that $\psi_A(i) \geq b^{-i}/2$ for all $i \in I_0$, thus $W_{b,A}(\psi) = [0, 1]$. Since the cardinality of D is at least 2, we have $m < (b - 1)/2$. Then

$$\begin{aligned} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} &\geq f\left(\frac{1}{2b^i} - \frac{m}{(b-1)b^i}\right) b^{i\gamma} \\ &= f\left(\frac{b-1-2m}{2(b-1)b^i}\right) b^{i\gamma} \\ &\geq f\left(\frac{1}{2(b-1)b^i}\right) b^{i\gamma} \\ &\geq b^{-\gamma} f\left(\frac{1}{2b^{i+1}}\right) b^{(i+1)\gamma} \end{aligned} \tag{4.1}$$

for all $i \in I_0$. For any $\rho > 0$ and any integer i_0 big enough, we have

$$\mathcal{H}_\rho^f(C(b, D)) \ll \sum_{i \geq i_0: i \in I_0} f\left(\frac{1}{2b^i}\right) b^{i\gamma}.$$

If $\mathcal{H}^f(C(b, D)) = 0$, then Theorem 4.1 is trivial. Otherwise,

$$\sum_{i \in I_0} f\left(\frac{1}{2b^i}\right) b^{i\gamma} = \infty,$$

which implies

$$\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} = \infty$$

by (4.1) and the monotonicity of $r^{-\gamma}f(r)$. So in this case, Theorem 4.1 is also valid.

The proof of Theorem 4.1 naturally splits into two parts: the convergence part and the divergence part. We start with the convergence part, which involves finding covers of $W_{b,A}(\psi) \cap C(b, D)$ of arbitrarily small measure.

LEMMA 4.2. *If*

$$\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} < \infty,$$

then

$$\mathcal{H}^f(W_{b,A}(\psi) \cap C(b, D)) = 0.$$

Proof. For $i \in I$, let

$$S_i = \bigcup_{0 \leq p \leq b^i} B\left(\frac{p}{b^i}, \psi_A(i)\right) \cap C(b, D).$$

For each $0 \leq p \leq b^i$, we have

$$B\left(\frac{p}{b^i}, \psi_A(i)\right) = \bigcup_{n: a_n=i} B\left(\frac{p}{b^{a_n}}, \psi(n)\right),$$

since all balls on the right-hand side have the same center and $\psi_A(i)$ is the maximum of their radii. So

$$S_i = \bigcup_{a_n=i} \bigcup_{0 \leq p \leq b^i} B\left(\frac{p}{b^{a_n}}, \psi(n)\right) \cap C(b, D).$$

Therefore

$$\begin{aligned} W_{b,A}(\psi) \cap C(b, D) &= \limsup_{n \rightarrow \infty} \bigcup_{0 \leq p \leq b^{a_n}} B\left(\frac{p}{b^{a_n}}, \psi(n)\right) \cap C(b, D) \\ &= \limsup_{i \rightarrow \infty} S_i. \end{aligned}$$

Lemma 3.2 implies that S_i is a subset of

$$\bigcup_{\frac{p}{b^i} \in L_i \cup R_i} B\left(\frac{p}{b^i} + d_{l,i}, \psi_A(i) - d_{l,i}\right) \cup B\left(\frac{p}{b^i} - d_{r,i}, \psi_A(i) - d_{r,i}\right),$$

where

$$d_{l,i} = \frac{m_l}{(b-1)b^i} \quad \text{and} \quad d_{r,i} = \frac{m_r}{(b-1)b^i}.$$

Recall that $m = \min\{m_l, m_r\}$, so

$$f(\psi_A(i) - d_{l,i}) + f(\psi_A(i) - d_{r,i}) \leq 2f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right).$$

Then for any $\rho > 0$ and any integer i_0 big enough, we have

$$\begin{aligned} & \mathcal{H}_\rho^f(W_{b,A}(\psi) \cap C(b, D)) \\ & \ll \sum_{i \geq i_0: i \in I} \mathcal{H}_\rho^f(S_i) \\ & \ll \sum_{i \geq i_0: i \in I, \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) \times \#(L_i \cup R_i) \\ & \ll \sum_{i \geq i_0: i \in I, \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) \times b^{i\gamma}. \end{aligned}$$

Let $i_0 \rightarrow \infty$ and then $\rho \rightarrow 0$, we deduce $\mathcal{H}^f(W_{b,A}(\psi) \cap C(b, D)) = 0$.

Now we turn to the more difficult divergence part. We are going to use Lemma 3.2 to rewrite balls $B(pb^{-n}, \psi(n))$ as balls with center in $C(b, D)$. Note that the new balls could have different radii depending on whether pb^{-n} is a left or right endpoint of $C_n(b, D)$, and we will deal with these two cases separately. For any $i \geq 1$, let

$$\begin{aligned} L_i^* &= \left\{ \frac{p}{b^i} \in L_i : \frac{p}{b^i} + d_{l,i} \neq \frac{q}{b^j} + d_{l,j} \text{ for any } q \text{ and } j < i \right\}, \\ LS_i^* &= \bigcup_{\frac{p}{b^i} \in L_i^*} B\left(\frac{p}{b^i} + d_{l,i}, \psi_A(i) - d_{l,i}\right), \end{aligned}$$

and $LW_{b,A}^*(\psi) = \limsup_{i \rightarrow \infty} LS_i^*$. Replacing L_i by R_i , the set $RW_{b,A}^*(\psi)$ is defined in a similar way, and Lemma 3.2 implies that

$$LW_{b,A}^*(\psi) \cup RW_{b,A}^*(\psi) \subseteq W_{b,A}(\psi). \tag{4.2}$$

Let $\mu = \mathcal{H}^\gamma|_{C(b,D)}$, B be an arbitrary ball with center in $C(b, D)$ and

$$LS_i^*(B) = B \cap LS_i^*.$$

Recall that μ is γ -Ahlfors regular, so there exists a constant $r_0 > 0$ such that for any ball $B(x, r_1)$ with $x \in C(b, D)$ and $r_1 < r_0$, we have

$$\mu(B(x, r_1)) \asymp r_1^\gamma. \tag{4.3}$$

Suppose the radius of B satisfies that $r(B) < r_0/2$, so (4.3) implies that $\mu(2B) \asymp \mu(B)$. For ease of notation, we write

$$B_i^*(\psi_A) := B\left(\frac{p}{b^i} + d_{l,i}, \psi_A(i) - d_{l,i}\right) \text{ and } B_i^* := B\left(\frac{p}{b^i} + d_{l,i}, \frac{1}{2b^i}\right)$$

when the value of p is unimportant. Then

$$\begin{aligned} & \#\{B_i^*(\psi_A) \subseteq B : B_i^*(\psi_A) \cap C(b, D) \neq \emptyset\} \\ & \asymp \#\{B_i^* \subseteq B : B_i^* \cap C(b, D) \neq \emptyset\} \\ & \asymp \frac{\mu(B)}{\mu(B_i^*)} \text{ since } B_i^* \text{ are disjoint,} \\ & \asymp \mu(B)b^{i\gamma}. \end{aligned}$$

Therefore

$$\mu(LS_i^*(B)) \asymp \mu(B)b^{i\gamma} \mu(B_i^*(\psi_A)) \asymp \mu(B)b^{i\gamma} (\psi_A(i) - d_{l,i})^\gamma. \tag{4.4}$$

Next we show that the $\mu(LS_i^*(B))$ satisfies a quasi-independence relation.

LEMMA 4.3. *Suppose $\psi_A(i) \leq b^{-i}/2$ for all i . Let t_0 be a sufficiently large integer satisfying $b^{-t_0} < r(B)$. Then there exists a constant $K > 0$ such that for any $i > j > t_0$,*

$$\mu(LS_i^*(B) \cap LS_j^*(B)) \leq \frac{K}{\mu(B)} \mu(LS_i^*(B))\mu(LS_j^*(B)).$$

Proof. We first consider the case $\psi_A(j) - d_{l,j} \leq b^{-i}/2$. Let $p_1b^{-i} \in L_i^*$ and $p_2b^{-j} \in L_j^*$. By the definition of L_i^* , the two ball centers $p_1b^{-i} + d_{l,i}$ and $p_2b^{-j} + d_{l,j}$ are distinct. Recall that

$$d_{l,i} = \frac{m_l}{(b-1)b^i},$$

so the distance between the two centers is

$$\begin{aligned} \left| \frac{p_1}{b^i} + d_{l,i} - \frac{p_2}{b^j} - d_{l,j} \right| &= \left| \frac{p_1}{b^i} - \frac{m_l}{(b-1)b^i} - \frac{p_2}{b^j} + \frac{m_l}{(b-1)b^j} \right| \\ &= \left| \frac{p_1 - p_2b^{i-j}}{b^i} - \frac{m_l(b^{i-j} - 1)}{(b-1)b^i} \right| \\ &\geq \frac{1}{b^i} \end{aligned}$$

since $b-1$ divides $b^{i-j} - 1$. Note that both radii $\psi_A(i) - d_{l,i}$ and $\psi_A(j) - d_{l,j}$ are not greater than $b^{-i}/2$, hence

$$B\left(\frac{p_1}{b^i} + d_{l,i}, \psi_A(i) - d_{l,i}\right) \cap B\left(\frac{p_2}{b^j} + d_{l,j}, \psi_A(j) - d_{l,j}\right) = \emptyset.$$

Therefore

$$LS_i^*(B) \cap LS_j^*(B) = \emptyset$$

and thus

$$\mu(LS_i^*(B) \cap LS_j^*(B)) = 0 \leq \frac{K}{\mu(B)} \mu(LS_i^*(B))\mu(LS_j^*(B))$$

for any constant $K > 0$.

Now assume $\psi_A(j) - d_{l,j} > b^{-i}/2$. We have

$$\begin{aligned} &\mu(LS_i^*(B) \cap LS_j^*(B)) \\ &= \mu\left(LS_i^*(B) \cap B \cap \bigcup_{bp^{-j} \in L_j^*} B\left(\frac{p}{b^j}, \psi_A(j)\right)\right) \\ &\leq \mathcal{N}(j) \mu\left(LS_i^*(B) \cap B_j^*(\psi_A)\right), \end{aligned} \tag{4.5}$$

where

$$\mathcal{N}(j) = \# \left\{ B_j^*(\psi_A) : B_j^*(\psi_A) \cap B \cap C(b, D) \neq \emptyset \right\}.$$

Let $2B$ denotes the ball with same center as B but twice the radius. Since $2\psi_A(j) \leq b^{-j} \leq b^{-t_0} < r(B)$, we have

$$\begin{aligned} \mathcal{N}(j) &\leq \# \left\{ B_j^*(\psi_A) \subseteq 2B : B_j^*(\psi_A) \cap C(b, D) \neq \emptyset \right\} \\ &\leq \# \left\{ B_j^* \subseteq 2B : B_j^* \cap C(b, D) \neq \emptyset \right\} \\ &\leq \frac{\mu(2B)}{\mu(B_j^*)} \quad \text{because } B_j^* \text{ are disjoint,} \\ &\ll \mu(B)b^{j\gamma} \quad (\text{by } \gamma\text{-Ahlfors regularity}). \end{aligned} \tag{4.6}$$

Similarly, for any fixed j ,

$$\begin{aligned} &\# \left\{ B_i^*(\psi_A) : B_i^*(\psi_A) \cap B_j^*(\psi_A) \cap C(b, D) \neq \emptyset \right\} \\ &\leq \# \left\{ B_i^* : B_i^* \cap B_j^*(\psi_A) \cap C(b, D) \neq \emptyset \right\} \\ &\leq 2 + \frac{\mu(B_j^*(\psi_A))}{\mu(B_i^*)} \\ &\ll 2 + (\psi_A(j) - d_{l_j})^\gamma b^{i\gamma}, \end{aligned}$$

where the number 2 is for the possible existence of those B_i^* which intersect with but are not contained in $B_j^*(\psi_A)$. Then

$$\begin{aligned} &\mu \left(LS_i^*(B) \cap B_j^*(\psi_A) \right) \\ &\ll \mu(B_i^*(\psi_A))(2 + (\psi_A(j) - d_{l_j})^\gamma b^{i\gamma}) \\ &\ll (\psi_A(i) - d_{l_i})^\gamma + (\psi_A(i) - d_{l_i})^\gamma (\psi_A(j) - d_{l_j})^\gamma b^{i\gamma} \\ &\ll (\psi_A(i) - d_{l_i})^\gamma (\psi_A(j) - d_{l_j})^\gamma b^{i\gamma} \end{aligned} \tag{4.7}$$

since $(\psi_A(j) - d_{l_j})b^j > 1/2$.

Now (4.4), (4.5), (4.6) and (4.7) give that

$$\begin{aligned} \mu(LS_i^*(B) \cap LS_j^*(B)) &\ll \mu(B)b^{j\gamma} (\psi_A(i) - d_{l_i})^\gamma (\psi_A(j) - d_{l_j})^\gamma b^{i\gamma} \\ &\ll \frac{\mu(LS_i^*(B))\mu(LS_j^*(B))}{\mu(B)}. \end{aligned}$$

PROPOSITION 4.4. Let $\mu = \mathcal{H}^\gamma|_{C(b,D)}$. If $\psi_A(i) \leq b^{-i}/2$ for all $i \in I$ and

$$\sum_{i \in I: \psi_A(i) > d_{l_i}} (\psi_A(i) - d_{l_i})^\gamma b^{i\gamma} = \infty,$$

then

$$\mu(LW_{b,A}^*(\psi)) = \mu(C(b, D)).$$

Proof. Let t_0 be the number as in Lemma 4.3. For all $i > t_0$, (2.1), (4.4) and Lemma 4.3 imply that

$$\mu(\limsup_{i \rightarrow \infty, i \in I} \mathcal{S}_i^*(B)) \geq \frac{\mu(B)}{C}.$$

Applying Lemma 2.2 and noting that

$$\limsup_{i \rightarrow \infty, i \in I} LS_i^*(B) = B \cap \limsup_{i \rightarrow \infty, i \in I} LS_i^* = B \cap LW_{b,A}^*(\psi),$$

we have $\mu(LW_{b,A}^*(\psi)) = \mu(C(b, D))$.

Now we extends to f -Hausdorff measure by applying the mass transference principle (Theorem 2.1).

LEMMA 4.5. *Let f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic. If $\psi_A(i) < b^{-i}/2$ for all $i \in I$ and*

$$\sum_{i \in I: \psi_A(i) > d_{l,i}} f(\psi_A(i) - d_{l,i}) b^{i\gamma} = \infty,$$

then

$$\mathcal{H}^f(LW_{b,A}^*(\psi) \cap C(b, D)) = \mathcal{H}^f(C(b, D)).$$

Proof. Define a function $\theta : \mathbb{N} \rightarrow (0, \infty)$ by

$$\theta(n) = \begin{cases} f(\psi(n) - d_{l,a_n})^{1/\gamma} + d_{l,a_n}, & \text{if } \psi(n) > d_{l,a_n}, \\ d_{l,a_n}/2, & \text{otherwise.} \end{cases}$$

Let

$$\theta_A(i) = \max \{ \theta(n) : a_n = i \},$$

then

$$\sum_{i \in I: \theta_A(i) > d_{l,i}} (\theta_A(i) - d_{l,i})^\gamma \times b^{i\gamma} = \sum_{i \in I: \psi_A(i) > d_{l,i}} f(\psi_A(i) - d_{l,i}) \times b^{i\gamma} = \infty.$$

Now Proposition 4.4 says that

$$\mu(LW_{b,A}^*(\theta)) = \mu(C(b, D)),$$

which is equivalent to

$$\mathcal{H}^\gamma(LW_{b,A}^*(\theta) \cap C(b, D)) = \mathcal{H}^\gamma(C(b, D)).$$

Hence Theorem 2.1 implies that

$$\mathcal{H}^f(LW_{b,A}^*(\psi) \cap C(b, D)) = \mathcal{H}^f(C(b, D)).$$

A similar result for $RW_{b,A}^*(\psi)$ can be proved by the same method.

LEMMA 4.6. *Let f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic. If $\psi_A(i) < b^{-i}/2$ for all i and*

$$\sum_{i \in I: \psi_A(i) > d_{r,i}} f(\psi_A(i) - d_{r,i}) b^{i\gamma} = \infty,$$

then

$$\mathcal{H}^f(RW_{b,A}^*(\psi)) = \mathcal{H}^f(C(b, D)).$$

Note that $m = \min\{m_l, m_r\}$, so

$$\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^i}} f\left(\psi_A(i) - \frac{m}{(b-1)b^i}\right) b^{i\gamma} = \infty,$$

implies either

$$\sum_{i \in I: \psi_A(i) > d_{l,i}} f(\psi_A(i) - d_{l,i}) b^{i\gamma} = \infty,$$

or

$$\sum_{i \in I: \psi_A(i) > d_{r,i}} f(\psi_A(i) - d_{r,i}) b^{i\gamma} = \infty.$$

Therefore the divergence part of Theorem 4.1 is a consequence of Lemma 4.5 and Lemma 4.6 since $LW_{b,A}^*(\psi) \cup RW_{b,A}^*(\psi) \subseteq W_{b,A}(\psi)$.

5. Proof of Theorem 1.6

For any prime divisor q of b , we have

$$v_q(b^{\lfloor \alpha_1 a_n \rfloor}) = \lfloor \alpha_1 a_n \rfloor v_q(b) \leq \alpha_1 a_n v_q(b) \leq a_n v_q(t) = v_q(t^{a_n}).$$

Hence $b^{\lfloor \alpha_1 a_n \rfloor} \mid t^{a_n}$ and thus

$$\bigcup_{0 \leq p \leq b^{\lfloor \alpha_1 a_n \rfloor}} B\left(\frac{p}{b^{\lfloor \alpha_1 a_n \rfloor}}, \psi(n)\right) \subseteq \bigcup_{0 \leq p \leq t^{a_n}} B\left(\frac{p}{t^{a_n}}, \psi(n)\right).$$

Therefore

$$W_{b, (\lfloor \alpha_1 a_n \rfloor)_{n \geq 1}}(\psi) \subseteq W_{t, (a_n)_{n \geq 1}}(\psi). \tag{5.1}$$

Let $J_1 = I(\{\lfloor i\alpha_1 \rfloor\}_{i \in I})$ be the set of integers appearing in the sequence $\{\lfloor i\alpha_1 \rfloor\}_{i \in I}$. Note that

$$\begin{aligned} \psi_{(\lfloor a_n \alpha_1 \rfloor)_{n \geq 1}}(j) &= \max \{ \psi(n) : \lfloor a_n \alpha_1 \rfloor = j \} \\ &= \max \{ \psi_A(i) : \lfloor i\alpha_1 \rfloor = j \}. \end{aligned}$$

Then

$$\begin{aligned} &\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lfloor i\alpha_1 \rfloor}}} f\left(\psi_A(i) - \frac{m}{(b-1)b^{\lfloor i\alpha_1 \rfloor}}\right) b^{i\alpha_1\gamma} \\ &= \sum_{j \in J_1} \sum_{\substack{i: \lfloor i\alpha_1 \rfloor = j \\ \psi_A(i) > \frac{m}{(b-1)b^{\lfloor i\alpha_1 \rfloor}}} f\left(\psi_A(i) - \frac{m}{(b-1)b^{\lfloor i\alpha_1 \rfloor}}\right) b^{i\alpha_1\gamma} \end{aligned}$$

$$\begin{aligned} &\leq \sum_{\substack{j \in J_1 \\ \psi_{(\lfloor a_n \alpha_1 \rfloor)_{n \geq 1}}(j) > \frac{m}{(b-1)b^j}}} \sum_{i: \lfloor i \alpha_1 \rfloor = j} f \left(\psi_{(\lfloor a_n \alpha_1 \rfloor)_{n \geq 1}}(j) - \frac{m}{(b-1)b^{\lfloor i \alpha_1 \rfloor}} \right) b^{i \alpha_1 \gamma} \\ &\ll \sum_{j \in J_1: \psi_{(\lfloor a_n \alpha_1 \rfloor)_{n \geq 1}}(j) > \frac{m}{(b-1)b^j}} f \left(\psi_{(\lfloor a_n \alpha_1 \rfloor)_{n \geq 1}}(j) - \frac{m}{(b-1)b^j} \right) b^{j \gamma}. \end{aligned}$$

Therefore Theorem 4.1 and (5.1) imply that

$$\mathcal{H}^f(W_t(\psi) \cap C(b, D)) = \mathcal{H}^f(C(b, D))$$

if

$$\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lfloor i \alpha_1 \rfloor}}} f \left(\psi_A(i) - \frac{m}{(b-1)b^{\lfloor i \alpha_1 \rfloor}} \right) b^{i \alpha_1 \gamma} = \infty.$$

The other half of the theorem is proved similarly using $\lceil \alpha_2 a_n \rceil$. For any prime divisor q of b , we have

$$v_q(b^{\lceil \alpha_2 a_n \rceil}) = \lceil \alpha_2 a_n \rceil v_q(b) \geq \alpha_2 a_n v_q(b) \geq a_n v_q(t) = v_q(t^{a_n}),$$

which implies $t^{a_n} \mid b^{\lceil \alpha_2 a_n \rceil}$. Therefore

$$\bigcup_{0 \leq p \leq t^{a_n}} B \left(\frac{p}{t^{a_n}}, \psi(a_n) \right) \subseteq \bigcup_{0 \leq p \leq b^{\lceil \alpha_2 a_n \rceil}} B \left(\frac{p}{b^{\lceil \alpha_2 a_n \rceil}}, \psi(a_n) \right)$$

and hence

$$W_{t, (a_n)_{n \geq 1}}(\psi) \subseteq W_{b, (\lceil \alpha_2 a_n \rceil)_{n \geq 1}}(\psi). \tag{5.2}$$

Let $J_2 = I(\{\lceil i \alpha_2 \rceil\}_{i \in I})$ be the set of integers appearing in $\{\lceil i \alpha_2 \rceil\}_{i \in I}$. Then

$$\begin{aligned} &\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lceil i \alpha_2 \rceil}}} f \left(\psi_A(i) - \frac{m}{(b-1)b^{\lceil i \alpha_2 \rceil}} \right) b^{i \alpha_2 \gamma} \\ &= \sum_{j \in J_2} \sum_{\substack{i: \lceil i \alpha_2 \rceil = j \\ \psi_A(i) > \frac{m}{(b-1)b^{\lceil i \alpha_2 \rceil}}} f \left(\psi_A(i) - \frac{m}{(b-1)b^{\lceil i \alpha_2 \rceil}} \right) b^{i \alpha_2 \gamma} \\ &\geq \sum_{j \in J_2: \psi_{(\lceil a_n \alpha_2 \rceil)_{n \geq 1}}(j) > \frac{m}{(b-1)b^j}} f \left(\psi_{(\lceil a_n \alpha_2 \rceil)_{n \geq 1}}(j) - \frac{m}{(b-1)b^{\lceil i \alpha_2 \rceil}} \right) b^{i \alpha_2 \gamma} \\ &\gg \sum_{j \in J_2: \psi_{(\lceil a_n \alpha_2 \rceil)_{n \geq 1}}(j) > \frac{m}{(b-1)b^j}} f \left(\psi_{(\lceil a_n \alpha_2 \rceil)_{n \geq 1}}(j) - \frac{m}{(b-1)b^j} \right) b^{j \gamma} \end{aligned}$$

Now Theorem 4.1 and (5.2) imply that

$$\mathcal{H}^f(W_t(\psi) \cap C(b, D)) = 0$$

if

$$\sum_{i \in I: \psi_A(i) > \frac{m}{(b-1)b^{\lceil i\alpha_2 \rceil}}} f\left(\psi_A(i) - \frac{m}{(b-1)b^{\lceil i\alpha_2 \rceil}}\right) b^{i\alpha_2\gamma} < \infty.$$

6. Further discussion

6.1. Hausdorff dimension

When b and t are multiplicatively dependent, we have computed the Hausdorff f -measure of $W_t(\psi) \cap C(b, D)$ for any dimension function such that $r^{-\gamma}f(r)$ is monotonic (Theorem 1.5). In particular, if the set D contains at least one of 0 and $b - 1$, take $f(r) = r^s$ for any $s \geq 0$ and we have

$$\begin{aligned} \mathcal{H}^s(W_t(\psi) \cap C(b, D)) &= \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} \psi(n)^s \times t^{n\gamma} < \infty, \\ \mathcal{H}^s(C(b, D)), & \text{if } \sum_{n=1}^{\infty} \psi(n)^s \times t^{n\gamma} = \infty. \end{cases} \end{aligned}$$

Note that

$$\sum_{n=1}^{\infty} \psi(n)^s \times t^{n\gamma} = \sum_{n=1}^{\infty} t^{n\left(\gamma + s \frac{\log \psi(n)}{n \log t}\right)}.$$

Since $t \geq 2$, the above series converges if

$$s > \limsup_{n \rightarrow \infty} \frac{-\gamma n \log t}{\log \psi(n)}$$

and diverges if

$$s < \limsup_{n \rightarrow \infty} \frac{-\gamma n \log t}{\log \psi(n)}.$$

Let

$$\lambda_\psi = \liminf_{n \rightarrow \infty} \frac{-\log \psi(n)}{n \log t}.$$

For the rest of this paper, we assume $\lambda_\psi \geq 1$ as otherwise the situation is trivial. By computations above, we deduce the Hausdorff dimension of $W_t(\psi) \cap C(b, D)$.

COROLLARY 6.1. *Suppose b and t are multiplicatively dependent, D contains at least one of 0 and $b - 1$ and $\lambda_\psi \geq 1$. Then*

$$\dim_{\text{H}} W_t(\psi) \cap C(b, D) = \frac{\gamma}{\lambda_\psi}.$$

6.2. *Multiplicatively independent case*

We first consider the situation that b and t have the same prime divisors but not necessarily multiplicatively dependent. Using Theorem 1-6, we deduce that

$$\mathcal{H}^s(W_t(\psi) \cap C(b, D)) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} \psi(n)^s \times b^{n\alpha_2\gamma} < \infty, \\ \mathcal{H}^s(C(b, D)), & \text{if } \sum_{n=1}^{\infty} \psi(n)^s \times b^{n\alpha_1\gamma} = \infty \end{cases}$$

if D contains at least one of 0 and $b - 1$. Then a similar computation as in the previous section yields

COROLLARY 6.2. *Suppose b and t have the same prime divisors and D contains at least one of 0 and $b - 1$. If $\lambda_\psi \geq 1$, then*

$$\frac{\alpha_1 \log b}{\log t} \frac{\gamma}{\lambda_\psi} \leq \dim_H W_t(\psi) \cap C(b, D) \leq \frac{\alpha_2 \log b}{\log t} \frac{\gamma}{\lambda_\psi}.$$

Based on Corollary 6.1 and note that

$$\frac{\alpha_1 \log b}{\log t} \leq 1 \leq \frac{\alpha_2 \log b}{\log t},$$

we propose the following conjecture.

CONJECTURE 6.3. *Suppose b and t have the same prime divisors, D contains at least one of 0 and $b - 1$ and $\lambda_\psi \geq 1$, then*

$$\dim_H W_t(\psi) \cap C(b, D) = \frac{\gamma}{\lambda_\psi}.$$

In [2], the authors gave a heuristic of Conjecture 1.4. Here we modify their argument to formulate a conjecture for $\mathcal{H}^f(W_t(\psi) \cap C(b, D))$ when the sets of prime divisors of b and t are different and D contains at least one of 0 and $b - 1$. For any big integer n , choose another integer m such that $b^{-m} \asymp \psi(n)$. Divide $[0, 1]$ into b^m intervals of length b^{-m} , among them there are approximately $b^{m\gamma}$ intervals that intersect with $C(b, D)$. If the sets of prime divisors of b and t are different, the distribution of points pt^{-n} should be random with respect to those length b^{-m} intervals. So in the union

$$\bigcup_{0 \leq p \leq t^n} B\left(\frac{p}{t^n}, \psi(n)\right),$$

there are about $t^n b^{m(\gamma-1)}$ balls that intersect with $C(b, D)$. Therefore

$$\begin{aligned} \mathcal{H}^f(W_t(\psi) \cap C(b, D)) &\ll \sum_{n=n_0}^{\infty} f(\psi(n)) t^n b^{m(\gamma-1)} \\ &\ll \sum_{n=n_0}^{\infty} f(\psi(n)) t^n \psi(n)^{1-\gamma} \end{aligned}$$

for any $n_0 > 0$. Based on above heuristic argument, we propose the following conjecture.

CONJECTURE 6.4. Let $C(b, D)$ be a missing digit set with Hausdorff dimension γ , the set D contains at least one of 0 and $b - 1$, $t \geq 2$ be an integer satisfying the sets of prime divisors of b and t are different, f be a dimension function such that $r^{-\gamma}f(r)$ is monotonic, and $\psi: \mathbb{N} \rightarrow (0, \infty)$ be a function. Then

$$\mathcal{H}^f(W_t(\psi) \cap C(b, D)) = \begin{cases} 0, & \text{if } \sum_{n=1}^{\infty} f(\psi(n))t^n \psi(n)^{1-\gamma} < \infty, \\ \mathcal{H}^f(C(b, D)), & \text{if } \sum_{n=1}^{\infty} f(\psi(n))t^n \psi(n)^{1-\gamma} = \infty. \end{cases}$$

Assuming Conjecture 6.4, the Hausdorff dimension of $W_t(\psi) \cap C(b, D)$ can be computed in the same way as we did in Section 6.1, so the following conjecture holds if Conjecture 6.4 is valid.

CONJECTURE 6.5. Let $C(b, D)$ be a missing digit set with Hausdorff dimension γ , with D containing at least one of 0 and $b - 1$, $t \geq 2$ be an integer satisfying the sets of prime divisors of b and t are different. If $\lambda_\psi \geq 1$, then

$$\dim_{\text{H}}(W_t(\psi) \cap C(b, D)) = \max \left\{ \frac{1}{\lambda_\psi} + \gamma - 1, 0 \right\}. \tag{6.1}$$

When $b = 3$ and t is not a power of 3, formula (6.1) was already conjectured by Bugeaud and Durand [6, conjecture 1.2]. Note that this formula is very different with the multiplicatively dependent case Corollary 6.1 and the same prime divisors case Corollary 6.2. It is known that $\dim_{\text{H}}(W_t(\psi)) = 1/\lambda_\psi$ (see [12]), so Corollary 6.1 says that the Hausdorff dimension of the intersection $W_t(\psi) \cap C(b, D)$ is the product of two dimensions when b and t are multiplicatively dependent, while Conjecture 6.5 predicts that the Hausdorff dimension of the intersection is the sum of two dimensions minus one when the set of prime divisors of b and t are different. For two unrelated fractals, the Hausdorff dimension of their intersection is likely equal to the sum of dimensions minus the dimension of the ambient space (see [8, chapter 8]), so Conjecture 6.5 is consistent with the intuition that $W_t(\psi)$ and $C(b, D)$ are “unrelated” when the sets of prime divisors of b and t are different. This kind of formula also appears in [13, 23] as well as in the study of other limsup sets induced by recurrence of orbits in dynamical systems [7, 15]. Note that if b and t are multiplicatively independent but have the same prime divisors, then Corollary 6.2 implies that

$$\dim_{\text{H}} W_t(\psi) \cap C(b, D) \quad \text{and} \quad \max \left\{ \frac{1}{\lambda_\psi} + \gamma - 1, 0 \right\}$$

are not necessarily equal since the right -hand side becomes 0 when λ_ψ is large while the lower bound in Corollary 6.2 is always positive.

6.3. Large intersection property

We are also able to show the large intersection property of $W_t(\psi) \cap C(b, D)$ when $\psi(n) = t^{-\theta n}$ for some $\theta > 1$. The large intersection property was introduced by Falconer [9] and has many applications, we refer to [14] and references therein. Let (X, \mathcal{B}, μ, d) be a compact metric space such that μ is γ -Ahlfors regular. When $X = \mathbb{R}^d$, the set $\mathcal{G}^s(X)$ is defined to be the class of all G_δ sets F in X such that

$$\dim_{\text{H}} \bigcap_{n=1}^{\infty} g_n(F) \geq s$$

holds for all sequences of similarity transformations $\{g_n\}_{n \geq 1}$. Definition of $\mathcal{G}^s(X)$ for general metric spaces is more complicated and can be found in [20]. For any Borel set U , define

$$I_s(\mu, U) = \int_U \int_U |x - y|^{-s} d\mu(x) d\mu(y).$$

Next we state two results that enable us to deduce the large intersection property of $W_t(\psi) \cap C(b, D)$.

THEOREM 6.6 ([14 theorem 1·1]). *Suppose $\{B_n\}_{n \geq 1}$ is a sequence of balls in X whose radii decrease to 0 as $n \rightarrow \infty$. For each $n \geq 1$, let E_n be an open subset of B_n and define*

$$\lambda = \sup \left\{ s \geq 0 : \sup_{n \geq 1} \frac{\mu(B_n) I_s(\mu, E_n)}{\mu(E_n)^2} < \infty \right\}.$$

Then $\mu(\limsup_{n \rightarrow \infty} B_n) = \mu(X)$ implies $\limsup_{n \rightarrow \infty} E_n \in \mathcal{G}^\lambda(X)$.

LEMMA 6.7 ([14, lemma 3·1]). *Suppose $0 \leq s < \gamma$ and U is a Borel set with diameter $\text{diam}(U) > 0$. Then*

$$I_s(\mu, U) \ll \text{diam}(U)^{\gamma-s} \mu(U).$$

In our setting, we take $X = C(b, D)$, $\mu = \mathcal{H}^\gamma|_{C(b,D)}$ and obtain the following result.

COROLLARY 6.8. *Suppose the set D contains at least one of 0 and $b - 1$, $\psi(n) = t^{-\theta n}$ for some $\theta > 1$, $s < \gamma/\theta$ and $\log t / \log b = \alpha \in \mathbb{Q}$, then*

$$W_t(\psi) \in \mathcal{G}^s(C(b, D)).$$

Proof. Since $\theta > 1$, we have $\theta \alpha n \geq [\alpha n]$ for n big enough. Hence

$$\psi(n) = \frac{1}{t^{\theta n}} = \frac{1}{b^{\theta \alpha n}} \leq \frac{1}{b^{[\alpha n]}}$$

for n big enough. Then for all n big enough and any $0 \leq p \leq t^n$, we have

$$B\left(\frac{p}{t^n}, \psi(n)\right) \subseteq B\left(\frac{q}{b^{[\alpha n]}}, \frac{1}{b^{[\alpha n]}}\right)$$

for some $0 \leq q \leq b^{[\alpha n]}$, since $t^n \mid b^{[\alpha n]}$.

If $pt^{-n} = qb^{-[\alpha n]} \notin C(b, D)$, then both balls above are disjoint with $C(b, D)$. Otherwise, the fact that μ is γ -Ahlfors regular and D contains at least one of 0 or $b - 1$ imply that for n big enough, we have

$$\mu\left(B\left(\frac{q}{b^{[\alpha n]}}, \frac{1}{b^{[\alpha n]}}\right)\right) \ll b^{-\gamma[\alpha n]} \ll t^{-\gamma n}, \tag{6.2}$$

$$\mu\left(B\left(\frac{p}{t^n}, \psi(n)\right)\right) \asymp \psi(n)^\gamma. \tag{6.3}$$

Then

$$\begin{aligned} & \mu \left(B \left(\frac{q}{b^{\lceil \alpha n \rceil}}, \frac{1}{b^{\lceil \alpha n \rceil}} \right) \right) I_s \left(\mu, B \left(\frac{p}{t^n}, \psi(n) \right) \right) \\ & \ll t^{-\gamma n} \psi(n)^{\gamma-s} \mu \left(B \left(\frac{p}{t^n}, \psi(n) \right) \right) \quad \text{by (6.2) and Lemma (6.7),} \\ & \ll t^{-\gamma n} \psi(n)^{-s} \mu \left(B \left(\frac{p}{t^n}, \psi(n) \right) \right)^2 \quad \text{by (6.3),} \\ & \ll t^{n(-\gamma+\theta s)} \mu \left(B \left(\frac{p}{t^n}, \psi(n) \right) \right)^2. \end{aligned}$$

For any $s < \gamma/\theta$, we have $\lim_{n \rightarrow \infty} t^{n(-\gamma+\theta s)} = 0$. Note that

$$\limsup_{n \rightarrow \infty, 0 \leq q \leq b^{\lceil \alpha n \rceil}} B \left(\frac{q}{b^{\lceil \alpha n \rceil}}, \frac{1}{b^{\lceil \alpha n \rceil}} \right) \cap C(b, D) = C(b, D).$$

Therefore Theorem 6.6 implies that

$$W_t(\psi) \in \mathcal{G}^s(C(b, D)).$$

Acknowledgements. The authors thank Bo Wang for helpful discussions, in particular an observation that led to Corollary 6.2. We also thank the anonymous referee for several suggestions that improved the expression of this paper. B. Li was supported by NSFC 12271176 and Guangdong Natural Science Foundation 2024A1515010946. R. Li (corresponding author) was supported by NSFC 12401006 and Guangdong Basic and Applied Basic Research Foundation 2023A1515110272. Y. F. Wu was supported by NSFC 12301110 and Natural Science Foundation of Hunan Province 2023JJ40700.

REFERENCES

- [1] D. ALLEN, S. BAKER, S. CHOW and H. YU. A note on dyadic approximation in Cantor's set. *Indag. Math.* **34** (2023), 190–197.
- [2] D. ALLEN, S. CHOW and H. YU. Dyadic approximation in the middle-third Cantor set. *Selecta Math. (N.S.)*, **29** (11) (2023).
- [3] S. BAKER. Approximating elements of the middle third Cantor set with dyadic rationals. *Israel J. Math.* (2024).
- [4] V. BERESNEVICH and S. VELANI. A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures. *Ann. of Math. (2)*. (3) **164** (2006), 971–992.
- [5] Y. BUGEAUD. Diophantine approximation and Cantor sets. *Math. Ann.* **341** (2008), 677–684.
- [6] Y. BUGEAUD and A. DURAND. Metric Diophantine approximation on the middle-third Cantor set. *J. Eur. Math. Soc.* (6) **18** (2016), 1233–1272.
- [7] Y. CHANG, M. WU and W. WU. Quantitative recurrence properties and homogeneous self-similar sets. *Proc. Amer. Math. Soc.* (4) **147**(2019), 1453–1465.
- [8] K. FALCONER. *Fractal Geometry: Mathematical Foundations and Applications* (Wiley, 1990).
- [9] K. FALCONER. Sets with large intersection properties. *J. London Math. Soc.* (2). **49** (1994), 267–280.
- [10] L. FISHMAN and D. SIMMONS. Intrinsic approximation for fractals defined by rational iterated function systems: Mahler's research suggestion. *Proc. London Math. Soc.* (1) **109** (2014), 189–212.
- [11] H. FURSTENBERG. Intersections of Cantor sets and transversality of semigroups. *Problems in Analysis (Sympos. Salomon Bochner, Princeton Univ., Princeton, N.J., 1969)*, (1970) 41–59.
- [12] R. HILL and S. VELANI. The ergodic theory of shrinking targets. *Invent. Math.* (1) **119** (1995), 175–198.
- [13] Z. HU, B. LI and Y. XIAO. On the intersection of dynamical covering sets with fractals. *Math. Z.* (1) **301** (2022), 485–513.

- [14] Z. HU, B. LI and L. YANG. Large intersection property for limsup sets in metric space. *Potential Anal.* **60** (2024), 341–363. <https://doi.org/10.1007/s11118-022-10052-7>.
- [15] M. HUSSAIN, B. LI, D. SIMMONS and B. WANG. Dynamical Borel-Cantelli lemma for recurrence theory. *Ergodic Theory Dynam. Systems* (6) **42** (2022), 1994–2008.
- [16] J. LEVESLEY, C. SALP and S. VELANI. On a problem of K. Mahler: Diophantine approximation and Cantor sets. *Math. Ann.* **338** (2007), 97–118.
- [17] B. LI, R. LI and Y. WU. Rational numbers in $\times b$ -invariant sets. *Proc. Amer. Math. Soc.* **151** (2023), 1877–1887.
- [18] K. MAHLER. Some suggestions for further research. *Bull. Aust. Math. Soc.* **29** (1984), 101–108.
- [19] P. MATTILA. *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability* (Cambridge University Press, 1995).
- [20] F. NEGREIRA and E. SEQUEIRA. Sets with large intersection properties in metric spaces. *J. Math. Anal. Appl.* (1) **511** (2022), 126064.
- [21] J. SCHLEISCHITZ. On intrinsic and extrinsic rational approximation to Cantor sets. *Ergodic Theory Dynam. Systems* (41) **6** (2021), 1560–1589.
- [22] V.G. SPRINDŽUK. *Metric Theory of Diophantine Approximation* (translated by R. A. Silverman) (Wiley, 1979).
- [23] B. WANG, J. WU and J. XU. Dynamical covering problems on the triadic Cantor set. *C. R. Math. Acad. Sci. Paris* (7) **355**(2017), 738–743.