

## Guest Editorial

# The interface of autism and (borderline) personality disorder

Orestis Zavlis and Peter Tyrer

**Summary**

Prominent clinical perspectives posit that the interface of autism and (borderline) personality disorder manifests as either a misdiagnosis of the former as the latter or a comorbidity of both. In this editorial, we integrate these disparate viewpoints by arguing that personality difficulties are inherent to the autistic spectrum.

**Keywords**

Autism; personality disorder; misdiagnosis; comorbidity; diagnostic medicine.

**Copyright and usage**

© The Author(s), 2024. Published by Cambridge University Press on behalf of Royal College of Psychiatrists.

The interface of autism spectrum disorder (ASD) and personality disorder, particularly borderline personality disorder, has been a topic of recent debate.<sup>1</sup> The two conditions were shown to overlap in various domains, including the interpersonal (relational instabilities and detachment), intrapersonal (instability in self-perception), anankastic (obsessional rigidity) and affective (mood instability).<sup>1–3</sup> How these overlaps ought to be interpreted, though, remains unclear. Some researchers have suggested that putative comorbidities between ASD and personality disorder should be read as cautionary tales of misdiagnoses, since the two conditions are fundamentally different.<sup>1</sup> Yet others have argued that the two conditions are sufficiently similar to warrant the possibility of comorbidity.<sup>2</sup> In this editorial, we closely examine these perspectives and present an alternative that can unify them. Namely, the view that personality psychopathology is an inherent feature of the autistic spectrum, but not vice versa.

**A note about terminology and focus**

Before elaborating this alternative viewpoint, we make some notes on the terminology and focus of our editorial. First, with ASD we refer to high-functioning autism (that is, autism without intellectual disability – formerly known as Asperger's syndrome), since this is the kind of autism that is frequently misdiagnosed as (borderline) personality disorder, particularly in females.<sup>4</sup> Second, by female/male, we mean people who were assigned female/male sex at birth (although it is worth noting that most researchers do not distinguish between gender and sex and no research on transgender individuals exists in this line of inquiry, at the time of writing). Finally, although most research in this area has focused on 'borderline' personality disorder, in this editorial we focus on its dimensional counterpart for at least two reasons. First, borderline personality disorder is so heterogeneous that it falls on the border of virtually all other psychiatric conditions, not just ASD.<sup>5</sup> Second, accruing evidence suggests that borderline personality disorder can be subsumed under a general personality disorder spectrum (much like Asperger's can be subsumed under the autistic spectrum).<sup>5</sup> Thus, focusing on the broader personality disorder spectrum, as opposed to borderline personality or any other putative personality 'type', presents a fruitful conceptual shift. With these notes being made, we move on to examine the two alternative viewpoints on the interface of ASD and general personality pathology.

**Misdiagnosis thesis**

According to the first viewpoint, cases of high-functioning ASD can sometimes be misdiagnosed as (borderline) personality disorder,

given the inherent overlap of the two conditions (particularly in females).<sup>1,4</sup> Since being first expressed in the early 2000s, this argument has been supported by diagnostic asymmetries in the two conditions. In particular, the relative underdiagnosis of ASD (and overdiagnosis of 'borderline' personality disorder) in females versus males has prompted several researchers to argue that 'female autism' could be conflated with (borderline) personality disorder.<sup>1,4</sup> Indeed, compared with male autism, female autism is less well-defined, more camouflaged and has a higher likelihood of being misdiagnosed (with borderline personality disorder being among the main misdiagnoses).<sup>1,4</sup>

The misdiagnosis thesis clearly bears merit and is supported not only by diagnostic asymmetries but also by notable phenomenological overlaps across the two conditions (e.g. in relational and emotional instabilities).<sup>1</sup> Nevertheless, extreme variations of this viewpoint may perhaps better be avoided since they risk promoting the notion that ASD and personality disorder can never co-occur, something that cannot be justified by recent clinical findings.

**Comorbidity thesis**

Indeed, in light of recent findings revealing genuine comorbidities of ASD and personality disorder,<sup>4</sup> researchers have argued that personality disorder and autism can often present together.<sup>2</sup> Unlike those endorsing the misdiagnosis perspective, proponents of this viewpoint emphasise points of convergence, rather than divergence, of the two phenotypes and attempt to shed light on their clinical presentation. Two notable findings can be reported from this literature. First, epidemiologically, although the comorbidity rate of the two disorders matches their population estimates (1–5%), the prevalence of personality disorder traits (such as instability of self, emotional reactivity and impulsivity) in those with ASD is higher than the prevalence of ASD features (such as alexithymic or systemising tendencies) in those with (borderline) personality disorder.<sup>1,2</sup> Second, clinically speaking, in comorbid cases, the neurodevelopmental symptoms of ASD are considered primary and the personality features secondary (since they could stem from autistic neurodivergence).<sup>1,4</sup>

In this sense, the personality disorder phenotype can be thought to be closer to the ASD one (than vice versa). In light of this observation, the comorbidity and misdiagnosis theses can be fruitfully united.

**The autistic and personality disorder spectra**

Given that the comorbidity and misdiagnosis accounts are not mutually exclusive (and both may be endorsed in conjunction by

most clinicians), they could be conjoined by embracing recent advances in the field of personality disorder.

First, contemporary classification systems (including both ICD-11, with its new personality disorder classification, and DSM-5, with its alternative model of personality dysfunction) diagnose personality disorder in dimensional, not categorical, terms.<sup>5</sup> ICD-11, in particular, highlights that personality disorder falls on a spectrum of severity that ranges from personality difficulty (in the subsyndromal case) to mild, moderate and severe personality disorder.<sup>5</sup>

Second, maladaptive personality traits (including the ICD-11 ones and their DSM-5 counterparts) are omnipresent in various psychopathologies, particularly neurodevelopmental ones.<sup>3</sup>

Finally, although somewhat controversial, the diagnosis of personality disorder can now be made at any age once a pattern of personality pathology has been demonstrated for at least 2 years.<sup>5</sup> This is in keeping with the new lifespan approach to diagnosis that is being followed in both ICD-11 and DSM-5. The option of making an assessment of personality pathology across severity and age groups allows many relational psychopathologies (particularly neurodevelopmental ones such as ASD or attention-deficit hyperactivity disorder) to be characterised by levels of (at least) personality difficulty, which if recognised during early development can be addressed before major personality disorder becomes manifest.<sup>2,3</sup>

Given these points, we argue that some level of personality disorder is intrinsic to ASD, but not vice versa. In other words, certain personality problems (that is, maladaptive ways of relating to oneself and others) are inherent to the autistic phenotype; but autistic traits are not inherent to those with personality disorder. This conclusion is supported by (a) epidemiological findings indicating high rates (48–68%) of personality difficulties in those with ASD (but not vice versa)<sup>1,2</sup> and (b) developmental findings showing that personality difficulties are a common clinical sequela of neurodivergence.<sup>3</sup>

This conclusion unifies the abovementioned perspectives by highlighting that even though neurodiversity could give rise to maladaptive personality traits (potentially birthing a comorbid personality disorder),<sup>3</sup> (sub)threshold personality pathology could also dominate the clinical picture (yielding a misdiagnosis of ASD as personality disorder).<sup>1</sup>

## Clinical and research implications

Based on the evidence synthesis above, several research and clinical implications may follow. First and foremost, ASD may be considered as a differential diagnosis of personality disorder, particularly for females, for whom ASD is currently under-recognised.<sup>4</sup> Although this point may seem intuitive clinically, it is worth reiterating because personality problems are often a consequence of other issues, including complex trauma, social deprivation and, of course, neurodiversity.<sup>5</sup> Examining these other issues (and often formulating them as primary) is crucial for an accurate personality diagnosis (particularly when the medical history of a patient is unknown).<sup>5</sup>

This point leads to the second implication of the reviewed evidence above: a more scientifically accurate and clinically helpful way of conceiving the interface of ASD and personality disorder is in dimensional, not categorical terms. That is, researchers should focus on identifying the personality traits, not personality ‘disorders’, that are most likely to ensue in people with ASD. For instance, affective volatility, relational instability, behavioural rigidity and detached relationships are found in many psychiatric conditions, including ASD.<sup>3,5</sup> Examining how and why these, as well as related, traits proliferate in some, but not all, individuals with ASD could help clarify the intimate link between neurodiversity and personality psychopathology.<sup>2,3</sup>

Finally, in light of this intimate link, the treatment of ASD could be extended to interventions that have been constrained to

personality disorder. This includes environment-based interventions (which focus on altering patients’ environments rather than their life-long illnesses)<sup>6</sup> but also dialectical behavioural interventions (which focus on teaching coping skills and acceptance practices). Although thought to be inapplicable to people with autism, such interventions are increasingly found (in modified forms) to be useful.<sup>1</sup>

## Conclusions

In sum, conceptualising autism and personality disorder as dimensional spectra appears to unify disparate viewpoints on their interface. Doing so further reveals that personality difficulties are somewhat inherent to the autistic spectrum (but not vice versa) and can yield either a misdiagnosis (when they dominate the clinical picture) or a comorbidity (when they reach diagnostic status that is acknowledged alongside neurodiversity). In light of these points, a fruitful way forward can be to examine the interface of ASD and personality disorder dimensionally, identifying the personality difficulties that individuals with ASD typically face, as well as the conditions under which they emerge. Treating those ASD difficulties using therapies that have been constrained to personality disorder appears a promising clinical avenue.

**Orestis Zavlis** , Department of Psychiatry, University of Oxford, Oxford, UK; and Department of Psychology and Language Sciences, University College London, London, UK; **Peter Tyrer** , Centre of Psychiatry, Imperial College London, London, UK

**Correspondence:** Orestis Zavlis. Email: orestis.research@gmail.com

First received 2 Oct 2023, final revision 14 Mar 2024, accepted 16 Mar 2024

## Data availability

Data availability is not applicable to this article as no new data were created or analysed in this study.

## Author contributions

O.Z.: writing – original draft; O.Z. and P.T.: writing – review and editing; P.T.: supervision.

## Funding

This study received no specific grant from any funding agency, commercial or not-for-profit sectors.

## Declaration of interest

P.T. is the chair of a registered charity, NIDUS-UK, that promotes the development of nidotherapy.

## References

- 1 May T, Pilkington PD, Younan R, Williams K. Overlap of autism spectrum disorder and borderline personality disorder: a systematic review and meta-analysis. *Autism Res* 2021; **14**: 2688–710.
- 2 Lugnegård T, Hallerback MU, Gillberg C. Personality disorders and autism spectrum disorders: what are the connections? *Compr Psychiatry* 2012; **53**: 333–40.
- 3 Anckarsäter H, Stahlberg O, Larson T, Hakansson C, Jutblad SB, Niklasson L, et al. The impact of ADHD and autism spectrum disorders on temperament, character, and personality development. *Am J Psychiatry* 2006; **163**: 1239–44.
- 4 Allely CS, Woodhouse E, Mukherjee RA. Autism spectrum disorder and personality disorders: how do clinicians carry out a differential diagnosis? *Autism* 2023; **27**: 1847–50.
- 5 Tyrer PJ, Mulder R. *Personality Disorder: From Evidence to Understanding*. Cambridge University Press, 2022.
- 6 Tyrer P, Tarabi S, Bassett P, Liedtka N, Hall R, Nagar J, et al. Nidotherapy compared with enhanced care programme approach training for adults with aggressive challenging behaviour and intellectual disability (NIDABID): cluster-randomised controlled trial. *J Intellect Disabil Res* 2017; **61**: 521–31.