

Hormonal nature of seed responses to fluctuating temperatures in *Cynara cardunculus* (L.)

H. Roberto Huarte^{1,2*} and Roberto L. Benech-Arnold³

¹Faculty of Agricultural Sciences, Argentine Catholic University, 183 Ramón Freire, 1426, CABA, Argentina; ²Faculty of Agricultural Sciences, National University of Lomas de Zamora, Kilometer 2 Route 4, 1836 Llavallol, Buenos Aires Province, Argentina; ³IFEVA/Cátedra de Cultivos Industriales, CONICET/Faculty of Agronomy, University of Buenos Aires, 4453 San Martín Avenue, 1417DSE, CABA, Argentina

(Received 20 April 2009; accepted after revision 12 November 2009)

Abstract

Cynara cardunculus (L.) seeds require incubation at fluctuating temperatures to terminate dormancy. In this study, we analysed the physiological mechanisms underlying such a requirement, focusing on the role of abscisic acid (ABA) and gibberellin (GA). As a conceptual framework, we considered the possibility that fluctuating temperatures and light trigger a similar set of hormonal processes after stimulus perception. To test this possibility, we (1) carried out hydrotime analysis of germination in seeds exposed to fluctuating temperatures (25/15°C) and constant temperature (20°C) with or without gibberellin (GA₃) or red light; (2) determined the responses of seeds incubated at fluctuating or constant temperature to ABA, GA₃, fluridone, an inhibitor of ABA biosynthesis, and paclobutrazol, an inhibitor of GA biosynthesis; and (3) determined the ABA content of seeds incubated at fluctuating or constant temperature. Incubation at 25/15°C or 20°C in the presence of GA₃ reduced the mean base water potential [$\psi_b(50)$] of the population to a similar extent, compared to that observed with seeds incubated at 20°C without GA₃. Irradiation with red light also reduced $\psi_b(50)$ to a lesser extent than incubation in the presence of GA₃. At all concentrations tested, exogenously applied GA₃ did not promote germination of seeds incubated at 25/15°C. However, paclobutrazol inhibited germination, suggesting that fluctuating temperatures terminate dormancy through *de novo* GA biosynthesis. Fluctuating temperatures enhanced seed germination in the presence of ABA, but ABA content did not differ between seeds incubated at fluctuating and constant temperatures. This study provides clear evidence for the involvement of

hormonal regulation in dormancy termination by fluctuating temperatures.

Keywords: abscisic acid, *Cynara cardunculus*, fluctuating temperatures, fluridone, gibberellins, paclobutrazol

Introduction

Temperature fluctuation is an absolute requirement for germination of many species (Probert, 1992). The ecological significance of this feature is associated with the strategies of seeds to detect canopy gaps, depth of burial of seeds in the soil or their positions under water (Roberts and Totterdell, 1981; Thompson and Grime, 1983; Pons and Schroder, 1985). Near to the soil surface, diurnal temperatures fluctuate less below a vegetation cover, or under water, than on bare soils (Balisky and Burton, 1993). While seed responses to fluctuating temperatures are widespread among many species, physiological and biochemical mechanisms underlying such responses are still largely unknown. Benech-Arnold *et al.* (1995) reported that fluctuating temperatures stimulate germination of immature dormant sorghum caryopses by reducing embryo sensitivity to abscisic acid (ABA). More recently, Huarte and Benech-Arnold (2005) and Huarte (2006) performed a water relation analysis of seed germination at fluctuating and constant temperatures using Gummerson's hydrotime model (Gummerson, 1986). In such an assessment, the authors found that incubation at fluctuating temperatures reduced seed mean base water potential $\psi_b(50)$. This finding implies that fluctuating temperatures promote germination through an enhancement of embryo potential to overcome a physical restraint for germination.

Germination is promoted by gibberellins (GAs) and inhibited by ABA. GA *de novo* synthesis appears to be

*Correspondence

Fax: + 54-11-4-552-2721-2711

Email: robertohuarte@uca.edu.ar

an important requirement for dormancy release (Jacobsen and Olszewski, 1993). Inhibitory effects of GA biosynthesis inhibitors such as paclobutrazol (PCB) and uniconazole on seed germination have been observed (Nambara *et al.*, 1991). Kucera *et al.* (2005) defined ABA as a positive regulator of dormancy induction. ABA plays a crucial role both in the acquisition and maintenance of seed dormancy (Le Page-Degivry *et al.*, 1996; Grappin *et al.*, 2000). The role of ABA in dormancy induction was clearly demonstrated by the reduced dormancy in ABA-deficient or -insensitive mutants (Li and Foley, 1997). Dormancy maintenance in dormant ecotypes of *Arabidopsis* (Ali-Rachedi *et al.*, 2004) and other species requires *de novo* ABA synthesis upon imbibition (Le Page-Degivry *et al.*, 1997; Grappin *et al.*, 2000). ABA is synthesized from a C₄₀ carotenoid precursor; hence, chemical inhibitors of carotenoid biosynthesis, such as fluridone, inhibit ABA accumulation (Zeevaart, 1988). ABA effects are related to its endogenous level and seed sensitivity to its action (Corbineau *et al.*, 2002). From these and many other results, it was concluded that the maintenance of a dormancy state depends on high ABA–GA ratios and/or high ABA sensitivity. Dormancy release is associated with increased GA biosynthesis and ABA degradation, both of which contribute to low ABA–GA ratios, and a high GA sensitivity (Finch-Savage and Leubner-Metzger, 2006). This balance is also modulated by environmental factors. Toyomasu *et al.* (1993, 1998) and Yamaguchi *et al.* (1998) have found that the promotion of germination by Pfr, the active form of phytochrome, after seed exposure to red light, is mediated by an increase in GA biosynthesis, due to the up-regulation of genes encoding GA₃ β-hydroxylase, a key enzyme in the production of active GAs. This increases embryo growth potential and decreases a physical restraint for germination by the covering tissues of the seed. Pfr enhancement of embryo capacity to overcome a physical restraint for germination is also through a reduction in embryo sensitivity to ABA (Sánchez and Mella, 2004) and an increase in embryo sensitivity to GA (Yang *et al.*, 1995; Arana *et al.*, 2006). Since fluctuating temperatures promote germination through an enhancement of embryo capacity to overcome a physical restraint, it is possible that this effect is also mediated by hormonal regulation. We hypothesized that seed responses to fluctuating temperatures in terms of dormancy termination are through the elicitation of a set of physiological processes that are similar to those underlying seed responses to red light.

In this study, we aimed to determine: (1) ABA content in and sensitivity of *Cynara cardunculus* seeds during incubation at fluctuating or constant temperatures; (2) seed responses to exogenous GA at constant or fluctuating temperatures; and (3) whether a

fluctuating temperature requirement can be cancelled by red light or GA.

Materials and methods

Plant materials

C. cardunculus (L.) mature achenes (hereafter termed as seeds) were hand collected during January 2006 from an infested roadside at Alejandro Petion, Buenos Aires Province, Argentina (34°59'S, 58°40'W). After the initial cleaning, seeds were kept for 3 months in paper bags at room temperature (20 ± 2°C) before use. Three independent experiments were conducted during 2006.

Germination tests

Germination was expressed as cumulative percentage of total seeds. Each value was mean ± SE of three replicates of 25 seeds each. Germination was scored during 14 consecutive days. Seeds with visible radicle protrusion were considered to have germinated and were removed. Germination tests were performed in germination chambers at 20°C (constant temperature) or at fluctuating temperatures (25°C, 12 h/15°C, 12 h). Seeds were incubated in the dark in 9-cm Petri dishes covered with plastic film to prevent evaporation.

Hydrotime analysis

To determine the effects of fluctuating temperatures (25/15°C) on hydrotime parameters, seeds were placed on dishes containing water or different concentrations of polyethylene glycol solutions (PEG8000, Anedra, Buenos Aires, Argentina), which were prepared according to the Michel (1983) equation. The resulting osmotic potential was measured with a vapour-pressure osmometer (VPO, Model 5100 C, Wescor Inc., Logan, Utah, USA) calibrated with sodium chloride solutions. PEG solutions were replaced after 24 h and every 6 d thereafter. Hydrotime parameters were also determined in seeds incubated at a constant temperature (20°C). Under these conditions, the parameters were also calculated for red-light-treated, GA₃-treated (throughout incubation) and dark-treated seeds. Irradiated seeds were pre-incubated in distilled water in darkness at 20°C for 24 h. After pre-incubation, seeds were irradiated for 2 h with a red light provided by Philips 40/15 40 W fluorescent lamps (Philips, Eindhoven, The Netherlands) to obtain a calculated proportion of Pfr of 0.87 (Casal *et al.*, 1991). After light exposure, seeds were transferred to Petri dishes containing distilled water or the different

concentrations of PEG solutions. All practices including germination counting were carried out under a green safety light. To determine the effects of GA₃ on hydrotime parameters, a mixed solution of GA₃ plus PEG was prepared by dissolving the PEG required according to the Michel equation in 100 μM GA₃. GA₃ plus PEG solution was replaced after 24 h and every 6 d thereafter.

Germination time courses were analysed according to the hydrotime model using Solver Tool of 2003 Microsoft Excel[®]. This module allows maximizing the fit between simulated values and experimentally recorded values. The optimization criterion used to obtain the best fit was minimum root-mean-square error (RMSE) between simulated and experimentally obtained data.

ABA, GA₃, fluridone and paclobutrazol treatments

Sensitivity of seeds incubated at either fluctuating or constant temperatures to (+)-*cis*, *trans*-ABA (Sigma Chemical Company, St. Louis, Missouri, USA) was evaluated through incubation in 6 ml of ABA solutions at different concentrations (0, 1, 50 and 100 μM). Sensitivity of seeds to GA₃ at fluctuating or constant temperature was determined through incubation in 6 ml of GA₃ solutions at different concentrations (0, 1, 25, 50 and 100 μM). Seeds were also incubated in the presence of 6 ml of 50 μM fluridone {1-methyl-3-phenyl-5-[3-trifluoromethyl-(phenyl)]-(4-(1H)-pyridinone)} (Phytotechnology Laboratories, Shawnee Mission, Kansas, USA) and 68 μM paclobutrazol (PCB) [2RS, 3RS-1-(4-chlorofenil)-4,4-dimetil-2-(1H,2,4-Triazol-1-il)pentan-3ol] (CRESTAR, Syngenta Crop Protection AG, Birsfelden, Switzerland). Fluridone solutions were prepared by dissolving the compound in 0.1% acetone until complete dissolution and then diluting it with water. Control experiments showed no acetone effect on germination. Seed germination in a mixed solution of PCB (68 μM) plus GA₃ (50 μM) was also tested for seeds incubated at 25/15°C.

In ABA experiments, the treatments were factorial combinations of four ABA doses and two thermal conditions (20°C vs. 25/15°C). In GA experiments the combinations were five GA doses and two thermal conditions (20°C vs. 25/15°C). The germination data were subjected to analysis of variance (Statistix 8.0, Analytical Software, Tallahassee, Florida, USA). Tukey's test at 5% level of probability was used for comparison between means.

ABA quantification

To quantify ABA content in seeds incubated at fluctuating (25/15°C) or constant (20°C) temperatures, three replicates of approximately 100 mg seeds were

collected with 12-h intervals for the first 4 d of incubation. Ungerminated seeds incubated at 20°C were also sampled at 24-h intervals from the fourth day of incubation onwards. The samples were immediately frozen in liquid nitrogen. Seeds were then lyophilized, powdered, weighed and stored at -18°C until assayed for ABA content with a radioimmunoassay using the monoclonal antibody MAC 252, as described by Steinbach *et al.* (1995). The results presented are the means of three measurements ± SE.

Results

Hydrotime analysis of germination of seeds incubated at fluctuating temperatures and seeds incubated at constant temperatures with red light or GA treatment

Fluctuating temperatures (25/15°C) promoted the final germination percentage of seeds compared to that of seeds incubated at a constant temperature (20°C) (Fig. 1a and d). Seeds incubated at reduced osmotic potentials displayed a decrease in total germination, except for GA₃-treated seeds incubated at -0.3 MPa, (Fig. 1c). Germination data observed for each treatment was analysed using the hydrotime model in order to quantify changes in water relation parameters. The resulting parameters were used to predict seed germination time courses according to: $\text{probit}(g) = [\psi - (\theta_H/t_g) - \psi_b(50)]/\sigma_{\psi_b(50)}$, converting probit values back to germination percentages. Incubation at fluctuating temperatures or at constant temperature (20°C) in the presence of GA₃ reduced the $\psi_b(50)$ of the population to a similar extent, compared to that observed for seeds incubated at 20°C without the hormone (Fig. 1h and g). Irradiation with red light prior to incubation at 20°C reduced $\psi_b(50)$ to a lesser extent than incubation at 25/15°C or in the presence of GA₃ (Fig. 1e and f). The reduction in $\psi_b(50)$ by fluctuating temperatures and GA₃ was accompanied by a reduction of their θ_H compared to the 20°C treatment. On the other hand, there were no major changes in σ_{ψ_b} between the dark control and the rest of the treatments.

Role of ABA in seed responses to constant or fluctuating temperatures

The two main effects, ABA doses and the thermal treatment on germination, were significant ($P < 0.001$). On the other hand, the interaction between ABA doses and thermal treatments was not significant ($P = 0.96$). Seeds incubated at fluctuating temperatures exhibited higher germination compared to that of seeds incubated at 20°C at the same ABA




Figure 1. Cumulative germination time courses of *Cynara cardunculus* seeds incubated at (a) 20°C under water potentials of 0, -0.3 and -0.6 MPa; (b) at 20°C under water potentials of 0, -0.3 and -0.6 MPa after red light exposure; (c) at 20°C in the presence of 100 μM GA₃; and (d) at 25/15°C under water potentials of 0, -0.3 and 0.6 MPa. The symbols represent the experimental data, and the lines are the time courses predicted by the hydrotime model. (e) to (h) Normal distributions showing the relative frequencies of Ψ_b (g) values corresponding to panels (a) to (d), respectively. The median, standard deviation (δ_{Ψ_b}), and hydrotime constant (θ_H) values are shown.

concentrations (Fig. 2). Total germination in water (ABA 0 μM) was $96.4 \pm 1.7\%$ (mean \pm SE) at 25/15°C and $36.8 \pm 8.3\%$ at 20°C, respectively. The final germination percentage at 25/15°C dropped from $96.4 \pm 1.7\%$ in water to $56.6 \pm 8.3\%$ in 100 μM ABA. In contrast, when incubation was performed at 20°C, the final germination percentages in water and 100 μM ABA were $36.8 \pm 8.3\%$ and $1.6 \pm 1.6\%$, respectively (Fig. 2). Fluridone (50 μM) promoted germination of seeds incubated at 20°C ($85.6 \pm 3.8\%$) (Fig. 2), suggesting that ABA biosynthesis upon seed imbibition might be attributable to the inhibition of germination at 20°C. To examine this possibility, we measured ABA content in seeds incubated at constant and fluctuating temperatures throughout imbibition. ABA content in dry seeds was $19.71 \pm 1.16 \text{ pg mg}^{-1}$ seeds (0 h of imbibition) (Fig. 3). ABA contents in seeds incubated at 25/15°C and 20°C were similar until 60 h after imbibition (Fig. 3). ABA content decreased in seeds incubated at 25/15°C prior to radicle emergence (Fig. 3; timing of radicle emergence indicated by an arrow). While ABA contents in seeds incubated at 20°C were $27.7 \pm 3.4 \text{ pg mg}^{-1}$ seeds at 72 h and $26.3 \pm 4.19 \text{ pg mg}^{-1}$ seeds at 84 h, ABA contents in seeds incubated at 25/15°C were $20.3 \pm 1.41 \text{ pg mg}^{-1}$ seeds at 72 h and $19.37 \pm 3.12 \text{ pg mg}^{-1}$ seeds at 84 h. A steady ABA content was observed in seeds incubated at 20°C after 72 h.

Role of GA in seed responses to constant and fluctuating temperatures

The two main effects, thermal treatment (20°C and 25/15°C) and seed responses to GA doses, and their interaction, were statistically significant ($P < 0.001$). Germination did not differ between 25/15°C and 20°C when seeds were incubated in the presence of GA




Figure 2. Final germination percentages of *Cynara cardunculus* seeds incubated at different ABA concentrations. Open bars show germination under fluctuating temperatures (25/15°C) and filled bars show germination scored at a constant temperature (20°C). Error bars indicate SEs. The germination experiments were performed for 14 d.




Figure 3. ABA content (pg mg^{-1} dry seeds) in *Cynara cardunculus* seeds at different time points of incubation. Open symbols represent ABA content of seeds incubated at 25/15°C, closed symbols represent ABA content of seeds incubated at 20°C. Data are means of triplicates \pm SEs.

(1–100 μM). Germination of these seeds in the presence of GA was similar to that observed in seeds incubated in water at 25/15°C (Fig. 4). On the other hand, a reduced total germination was scored both in water at 20°C ($31.6 \pm 9.2\%$) (mean \pm SE) and PCB (68 μM) ($21.3 \pm 3.3\%$ and $16 \pm 4\%$ for 20°C and 25/15°C, respectively) (Fig. 4). Incubation in the mixture solution of PCB (68 μM) plus GA₃ (50 μM) at 25/15°C restored germination ($88.8 \pm 8\%$) to a similar extent to that observed in water ($90.8 \pm 3.3\%$) (Fig. 4).




Figure 4. Final germination percentages of *Cynara cardunculus* seeds incubated at fluctuating temperatures (25/15°C) (open bars) or at a constant temperature (20°C) (filled bars) incubated in different solutions (see legend beneath horizontal axis). Vertical bars indicate the SEs. Germination experiments were performed for 14 d. Different letters at the top of each bar indicate significant differences according Tukey's test ($\alpha = 0.05$).

Discussion

Fluctuating temperatures and light are environmental factors that terminate seed dormancy in many species (Batlla *et al.*, 2005). A large body of information is available for light signalling, including its perception and signal transduction. In contrast, little is known about physiological and biochemical mechanisms involved in the responses of seeds to fluctuating temperatures. Our previous work on the enhancement of germination of *C. cardunculus* by fluctuating temperatures clearly demonstrated that none of the single (constant) temperatures included in the fluctuating temperatures but fluctuation *per se* released *C. cardunculus* seeds from dormancy (Huarte and Benech-Arnold, 2005). The hydrotime model indicated that the promotion of germination by fluctuating temperatures was accompanied by a reduction in ψ_b , which provides an important physiological implication that fluctuating temperatures stimulate dormancy termination through an enhancement of embryo capacity to overcome either physical or osmotic restraint. Embryo growth potential is positively or negatively affected by GA or ABA, respectively (Karssen *et al.*, 1989; Sánchez and de Miguel, 1997; da Silva *et al.*, 2004). Hormone action could be explained by its content in seed tissues and cells and/or by tissue or cell sensitivity to hormones. In the present study, a clear reduction of ABA sensitivity at fluctuating temperatures was demonstrated (Fig. 2). This is in full agreement with the work of Benech-Arnold *et al.* (1995), Romagosa *et al.* (2001) and Corbineau *et al.* (2002). These authors also found a reduction in ABA sensitivity and its possible contribution to dormancy release. In contrast, only slight differences in the ABA content were found during seed incubation, which do not seem to be instrumental to explaining the different behaviour of *C. cardunculus* seeds at fluctuating and constant temperatures. So, ABA synthesis could be important to maintain seed dormancy at a constant temperature. Nevertheless, evidence to support the involvement of ABA in thermodormancy provided by previous reports showed more dramatic changes in ABA content than that reported in this paper (Yoshioka *et al.*, 1998; Ali-Rachedi *et al.*, 2004; Benech-Arnold *et al.*, 2006). Hence, our results suggest that ABA deactivation is not the major mechanism in termination of *C. cardunculus* dormancy. This is somewhat contradictory to the enhancement of germination observed with fluridone-treated seeds incubated at 20°C (Fig. 2), because fluridone is known to inhibit ABA biosynthesis by blocking carotenoid biosynthesis. However, at least one report points out that fluridone enhances germination of dormant seeds without modifying ABA content (Benech-Arnold *et al.*, 1999). The application of exogenous GA₃ to seeds incubated

at fluctuating temperatures did not increase their germination percentage. On the other hand, GA₃ treatment enhanced germination of seeds incubated at a constant temperature (Fig. 4). That is, GA₃ replaced the requirement of fluctuating temperatures for seeds to germinate. This suggests that GA biosynthesis is involved in the promotion of germination at fluctuating temperatures. The importance of GA biosynthesis to promote germination under 25/15°C is also supported by the strong inhibition of germination by PCB, which was then completely reversed by co-incubation with GA₃. Moreover, the results of the hydrotime analysis, where GA₃-treated seeds at 20°C exhibited hydrotime parameters very similar to those obtained for seeds incubated at 25/15°C, strongly support the proposition that GAs are involved in the process. Taking these results together, we propose that fluctuating temperatures terminate dormancy of *C. cardunculus* seeds mainly by the promotion of GA biosynthesis and a reduction in ABA sensitivity.

Acknowledgements

We are grateful to Silvina Enciso for her guidance to H.R.H. in ABA determination, Sebastian Staltari and María Susana Pereyra for their help during the course of the experiments and Eduardo Fernandez for his help with the statistical analyses.

References

- Ali-Rachedi, S., Bouinot, D., Wagner, M.H., Bonnet, M., Sotta, B., Grappin, P. and Jullien, M. (2004) Changes in endogenous abscisic acid levels during dormancy release and maintenance of mature seeds: studies with the Cape Verde Ecotype, the dormant model of *Arabidopsis thaliana*. *Planta* **219**, 479–488.
- Arana, M.V., de Miguel, L.C. and Sánchez, R.A. (2006) A phytochrome-dependent embryonic factor modulates gibberellin responses in the embryo and micropylar endosperm of *Datura ferox* seeds. *Planta* **223**, 847–857.
- Balisky, A.C. and Burton, P.J. (1993) Distinction of soil thermal regimes under various experimental vegetation covers. *Canadian Journal of Plant Sciences* **73**, 411–420.
- Batlla, D., Kruk, B.C. and Benech-Arnold, R.L. (2005) Modeling changes in dormancy in weed soil banks: implications for the prediction of weed emergence. pp. 245–270 in Benech-Arnold, R.L.; Sánchez, R.A. (Eds) *Handbook of seed physiology applications to agriculture*. New York, Haworth Press.
- Benech-Arnold, R.L., Steinbach, H.S., Kristof, G. and Sánchez, R.A. (1995) Fluctuating temperatures have different effects on embryonic sensitivity to ABA in Sorghum varieties with contrasting pre-harvest susceptibility. *Journal of Experimental Botany* **46**, 711–717.
- Benech-Arnold, R.L., Enciso, S. and Sánchez, R.A. (1999) Fluridone stimulus of dormant sorghum seeds germination at low temperatures is not accompanied by changes in ABA content. pp. 76–80 in Weipert, D. (Ed.) *Pre-harvest*

- sprouting in cereals (Part II). Detmold, Association of Cereal Research.
- Benech-Arnold, R.L., Gualano, N., Leymarie, J., Come, D. and Corbineau, F.** (2006) Hypoxia interferes with ABA metabolism and increases ABA sensitivity in embryos of dormant barley grains. *Journal of Experimental Botany* **57**, 1423–1430.
- Casal, J.J., Sánchez, R.A., Di Benedetto, A.H. and Miguel, L.C.** (1991) Light promotion of seed germination in *Datura ferox* is mediated by a highly stable pool of phytochrome. *Photochemistry and Photobiology* **53**, 249–254.
- Corbineau, F., Bianco, J., Garello, G. and Come, D.** (2002) Breakage of *Pseudotsuga menziesii* seed dormancy by cold treatment as related to changes in seed ABA sensitivity and ABA levels. *Physiologia Plantarum* **114**, 313–319.
- da Silva, E.A., Toorop, P.E., van Aelst, A.C. and Hilhorst, H.W.M.** (2004) Abscisic acid controls embryo growth potential and endosperm cap weakening during coffee (*Coffea arabica* cv. Rubi) seed germination. *Planta* **220**, 251–264.
- Finch-Savage, W.E. and Leubner-Metzger, G.** (2006) Seed dormancy and the control of germination. *New Phytologist* **171**, 501–523.
- Grappin, P., Bouinot, D., Sotta, B., Miginiac, E. and Jullien, M.** (2000) Control of seed dormancy in *Nicotiana plumbaginifolia* post imbibition abscisic acid synthesis imposes dormancy maintenance. *Planta* **210**, 279–285.
- Gummerson, R.J.** (1986) The effect of constant temperatures and osmotic potential on the germination of sugar beet. *Journal of Experimental Botany* **41**, 1431–1439.
- Huarte, H.R.** (2006) Hydrotime analysis of the effect of fluctuating temperatures on seed germination in several non-cultivated species. *Seed Science and Technology* **34**, 555–569.
- Huarte, H.R. and Benech-Arnold, R.L.** (2005) Incubation under fluctuating temperatures reduces mean base water potential for seed germination in several non-cultivated species. *Seed Science Research* **15**, 89–97.
- Jacobsen, S.E. and Olszewski, N.E.** (1993) Mutations at the SPINDLY locus of *Arabidopsis* alter gibberellins signal transduction. *Plant Cell* **5**, 887–895.
- Karszen, C.M., Zagorski, S., Kepczynski, J. and Groot, S.P.C.** (1989) Key role for endogenous gibberellins in the control of seed germination. *Annals of Botany* **63**, 71–80.
- Kucera, B., Cohn, M.A. and Leubner-Metzger, G.** (2005) Plant hormone interactions during seed dormancy release and germination. *Seed Science Research* **15**, 281–307.
- Le Page-Degivry, M.T., Bianco, J., Barthe, P. and Garello, G.** (1996) Change in hormone sensitivity in relation to the onset and breaking of sunflower dormancy. pp. 221–231 in Lang, G.A. (Ed.) *Plant dormancy physiology, biochemistry and molecular biology*. Wallingford, UK, CAB International.
- Le Page-Degivry, M.T., Bianco, J. and Barthe, P.** (1997) Changes in abscisic acid biosynthesis and catabolism during dormancy breaking in *Fagus sylvatica* embryo. *Journal of Plant Growth Regulation* **16**, 57–61.
- Li, B.L. and Foley, M.E.** (1997) Genetic and molecular control of seed dormancy. *Trends in Plant Science* **2**, 384–389.
- Michel, B.E.** (1983) Evaluation of water potential of solutions of polyethylene glycol 8000 both in the absence and presence of other solutes. *Plant Physiology* **72**, 66–70.
- Nambara, E., Akazawa, T. and McCourt, P.** (1991) Effects of the gibberellin biosynthetic inhibitor uniconazole on mutants of *Arabidopsis*. *Plant Physiology* **97**, 736–738.
- Pons, T.L. and Schroder, H.F.J.M.** (1985) Significance of temperature fluctuation and oxygen concentration for the germination of the rice field weeds *Fimbristylis littoralis* and *Scirpus juncooides*. *Oecologia* **69**, 315–319.
- Probert, R.J.** (1992) The role of temperature in germination ecophysiology. pp. 285–325 in Fenner, M. (Ed.) *Seeds. The ecology of regeneration in plant communities*. Wallingford, UK, CAB International.
- Roberts, E.H. and Totterdell, S.** (1981) Seed dormancy in *Rumex* species in response to environmental factors. *Plant, Cell and Environment* **4**, 97–106.
- Romagosa, I., Prada, D., Moralejo, M.A., Sopena, A., Muñoz, P., Casas, A.M., Swanston, J.S. and Molinacano, J.L.** (2001) Dormancy, ABA content and sensitivity of a barley mutant to ABA application during seed development and after ripening. *Journal of Experimental Botany* **52**, 1499–1506.
- Sánchez, R.A. and de Miguel, L.** (1997) Phytochrome promotion of mannan-degrading enzyme activities in the micropylar endosperm of *Datura ferox* seeds requires the presence of the embryo and gibberellin synthesis. *Seed Science Research* **7**, 27–34.
- Sánchez, R.A. and Mella, R.A.** (2004) The exit from dormancy and the induction of germination. Physiological and molecular aspects. pp. 221–235 in Benech-Arnold, R.L.; Sánchez, R.A. (Eds) *Handbook of seed physiology applications to agriculture*. New York, Haworth Press.
- Steinbach, H.S., Benech-Arnold, R.L., Kristof, G., Sánchez, R.A. and Marcucci Pultri, S.** (1995) Physiological basis of pre-harvest sprouting resistance in *Sorghum bicolor* (L.) Moench. ABA level and sensitivity in developing embryos of sprouting resistance and sprouting susceptible varieties. *Journal of Experimental Botany* **45**, 701–709.
- Thompson, K. and Grime, J.** (1983) A comparative study of germination responses to diurnally fluctuating temperatures. *Journal of Applied Ecology* **20**, 141–156.
- Toyomasu, T., Tsuji, H., Yamane, H., Nakayama, M., Yamaguchi, I., Murofushi, N., Takahashi, N. and Inoue, Y.** (1993) Light effects on endogenous levels of gibberellins in photoblastic lettuce seeds. *Journal of Plant Growth Regulation* **12**, 85–90.
- Toyomasu, T., Kawaide, H., Mitsushashi, W., Inoue, Y. and Kamiya, Y.** (1998) Phytochrome regulates gibberellin biosynthesis during germination of photoblastic lettuce seeds. *Plant Physiology* **118**, 1517–1523.
- Yamaguchi, S., Smith, M., Brown, R., Kamiya, Y. and Sun, T.** (1998) Phytochrome regulation and differential expression of gibberellin 3 β -hydroxylase genes in germinating *Arabidopsis* seeds. *Plant Cell* **10**, 2115–2126.
- Yang, Y., Nagatani, A., Zhao, Y., Kendrick, R. and Kamiya, Y.** (1995) Effects of gibberellins on seed germination of phytochrome deficient mutants of *Arabidopsis thaliana*. *Plant Cell and Physiology* **36**, 1205–1211.
- Yoshioka, T., Endo, T. and Satoh, S.** (1998) Restoration of seed germination at supraoptimal temperatures by fluridone, an inhibitor of abscisic acid biosynthesis. *Plant Cell and Physiology* **39**, 307–312.
- Zeevaart, J.A.D.** (1988) Metabolism and physiology of abscisic acid. *Annual Review of Plant Physiology and Plant Molecular Biology* **39**, 439–473.