

# Collection, characterization, conservation and utilization of *Cordia sinensis* Lam.: An underexploited multipurpose fruit species of hot arid regions

Rahul Dev<sup>1\*</sup>, M. Sureshkumar<sup>1</sup>, Sushil Kumar<sup>1</sup>, K. Venkatesan<sup>3</sup>, Traloki Singh<sup>1</sup>, Arvind Singh Tetarwal<sup>1</sup>, Anil Patidar<sup>4</sup>, Devi Dayal<sup>1</sup> and P. R. Meghwal<sup>2</sup>

<sup>1</sup>ICAR-Central Arid Zone Research Institute, Regional Research Station, Kukma-Bhuj 370 105, Gujarat, India, <sup>2</sup>ICAR-Central Arid Zone Research Institute, Jodhpur 342 003, Rajasthan, India, <sup>3</sup>ICAR-Central Island Agricultural Research Institute, Port Blair 744 105, Andaman and Nicobar Islands, India and <sup>4</sup>ICAR-Central Arid Zone Research Institute, Regional Research Station, Jaisalmer 345 001, Rajasthan, India

Received 18 January 2020; Accepted 23 October 2020 – First published online 2 December 2020

## Abstract

*Cordia sinensis*, locally known as ‘Goondi’ in India, is an underexploited multipurpose fruit species found in hot arid regions that is well adapted to drought, salt and hot conditions. The present study was undertaken to collect fruit samples from different locations in the Kachchh region of Gujarat, India, and to determine their field establishment for characterization, conservation and utilization. The maximum distribution of the species was observed in Bhuj (45%) and Mandvi (25%). Field boundaries (35%) and scrub forests (30%) had greater frequencies, whereas backyards had rarer frequencies (10%). The species most commonly occurred on levelled topography (60%) with a soil pH in the range of 8–8.5 (63%). Morphological data of three-year-old plants in the field gene bank showed a maximum coefficient of variation in the number of leaves per plant (66.6), followed by the number of branches per plant (45.62) and collar diameter (27.69). Wide variations were recorded in plant height (121.67–212 cm), spread (118–223 cm<sup>2</sup>) and the number of branches per plant (6–24.33). Specific accessions were identified for fodder (CBCG-12, CBCG-13 and CBCG-16), early flowering and fruiting (CBCG-12, CBCG-13 and CBCG-14), easier propagation by seeds (CBCG-12 and CBCG-13) and salt tolerance (CBCG-15 and CBCG-16). Preliminary findings and information provided about this species’ utilization and other aspects might be useful for future research on its domestication, sole plantation and conservation aspects, improving the exploitation of this species by present and future generations.

**Keywords:** arid Kachchh, conservation, *Cordia*, fodder, germplasm

## Introduction

The Indian Thar Desert occupies 89.6% (31.7 million ha) of the hot arid zone, mainly covering the states of western Rajasthan, Gujarat, Punjab and Haryana. Approximately

81% of this area is located in the western Rajasthan and Gujarat states of India (Sharma and Tewari, 2005). The nature of vegetation in the Thar Desert is mostly dominated by mixed shrubs and annual and perennial grasses with sparse/limited tree density (Kumar and Parveen, 2004; Sharma and Tewari, 2005). The region is endowed with multipurpose and highly useful plant species, which are all well adapted to arid agroecological habitats. Different

\*Corresponding author. E-mail: rahul2iari@gmail.com

parts of the plant species, such as the ripe fruits of *Cordia sinensis*, *Ziziphus nummularia*, *Grewia tenax* and *Capparis decidua*; the flower buds of *Calligonum polygoides* and *C. decidua*; the unripe fruits of *C. decidua* and *Leptadenia pyrotechnica* and the seeds of *Haloxylon salicornicum* are harvested from the wild for food purposes in arid regions of India (Singh et al., 2008). People are transitioning to be more health conscious and are trying to avoid chemical/synthetic foods, hence preferring natural therapeutic nutrition through natural products (Kumar et al., 2008).

The demand for underutilized fruit crops such as Bael is increasing daily in many parts of the world. Specifically, fruit pulp sharbat has become a popular drink in India due to its nutritional and health benefits (Kumar et al., 2008; Dhakar et al., 2019). An underutilized fruit species locally called September yellow (*Stauntonia obovatifolia* Hayata subsp. *urophylla*) has become a new potential edible fruit crop in China due to its precious nutritional value and pleasant taste (Zou et al., 2020). Its fruit is rich in total sugar (40.26%), crude protein (1.28%), vitamin C (1.55 mg/100 g) and reducing sugar (Wang et al., 2015) and is traditionally utilized by local villagers for medicinal purposes, such as anti-inflammatory and pain-killing properties. Ronoh et al. (2019) reported that African nightshade (*Solanum scabrum* Mill.) played a significant role in improving food and nutrition security in sub-Saharan Africa. In India, Singh and Meghwal (2020) emphasized the importance of underutilized *Ziziphus* species in the rural livelihood security of the local people of Rajasthan. *Z. mauritiana* is a valuable food, fodder, fuel and fencing source, thereby improving the socio-economic life of the local people. The leaves of naturally distributed *Z. nummularia*, a wild relative of *Z. mauritiana*, are excellent sources of fodder for sheep, goats, camels and cattle. *Z. rotundifolia* is a source of quality timber and rootstock promoting the large-scale multiplication of *Z. mauritiana*. The fruits of Blue honeysuckle (*Lonicera caerulea* L.) are sweet to sour or bitter in taste and are traditionally utilized for food purposes in Russia, China, Japan, and recently in North America (Chaovanalikit et al., 2004).

The genus *Cordia* (family *Boraginaceae*), with >300 species, has widely been grown in dry and hot regions of the world (Bouby et al., 2011). *Cordia myxa* and *C. sinensis* have grown naturally in arid regions of Rajasthan and Gujarat states (Meghwal et al., 2014). *C. sinensis* is one of the most important underutilized multipurpose fruit species with high potential as a useful source of food, fodder, traditional medicine for animals and humans, fibre and wood. Its leaves are rich in dry matter (41.2%) and crude protein (13.8%), with high amounts of energy and minerals such as potassium, phosphorus, calcium, zinc and iron, whereas its fruits are rich in dietary fibre (11.6%), energy (18.9 kcal/100 g dry wt.) and crude protein (12.6%)


(Kuria et al., 2005). It is a multi-stemmed shrub or small tree, growing up to 12 m high. Leaves are opposite, ovate to obovate, and light green with pale hairs on both surfaces. Flowers are born in terminal and axillary cymes, white or cream, and are sweetly scented (Warfa, 1990; Orwa et al., 2009). The fruit is small, drupe and bright red or yellow-orange. One fruit contains 1–4 hard and rough seeds.

The robust tap root system and other adaptive features, such as waxy leaves, hairiness, sunken and covered stomata in leaves, water binding mechanism and tolerance to salinity, drought and heat, make this species hardier and essential in arid regions (Meghwal et al., 2014). Although *C. sinensis* has become a favourable species for domestication and utilization in arid regions due to its multiple uses and drought hardiness characteristics, it has received limited research attention regarding its improvement and utilization. Thus, the present paper presents research that was an attempt to gather information on its geographical distribution, morphological diversity, multiplication and *ex situ* conservation and provides a preliminary evaluation regarding its use as potential food and fodder to stimulate interest and promote the sustainable utilization of the species.

## Material and methods

### Field survey, germplasm and passport data collection

Field surveys were conducted in five administrative talukas (is an administrative division within a district) of the Kachchh region of Gujarat during two consecutive years from June to December in 2015 and 2016 at the fruiting stage, and *C. sinensis* germplasm was collected (Fig. 1 and online Supplementary Table S1). The five administrative talukas and the regions surveyed are as follows: Bhuj (Tapkeshwari hill, Gangeshwar hill, Kotda Chakar region, etc.), Nakhatrana (Dhinodhar hill region, Chhari Dand region, etc.), Rapar (Pragpar, Adesar, Chitrod, etc.), Mandvi (Vijay Vilas Palace, Dahisara, Meghpar, Kera, etc.) and Bhachau (Amardi, Bhudarmora, etc.). Propagules such as fruits and hardwood cuttings were collected through a selective sampling strategy. Hardwood cuttings were kept in moist sand in a small plastic bag to avoid moisture loss. In total, 20 diverse germplasms were collected during the exploration trip, and the collector's number was allotted to each germplasm. First-hand field data on passport information (site information, global positioning system data, habitat, topography, etc.), ethnobotanical/ethnoveterinary information (traditional medicinal uses) and other important economic characteristics of species (fruit/fodder value) were collected using a semi-structured questionnaire, household survey and individual interviews of the village



**Fig. 1.** The collection sites of *C. sinensis* in the Kachchh region (1, Kotda Ugamana/Bhuj; 2, Bhujodi/Bhuj; 3, Madhapar/Bhuj; 4, Pragpar/Rapar; 5, Pragpar/Rapar; 6, Tapkeshwari/Bhuj; 7, Tapkeshwari/Bhuj; 8, Tapkeshwari/Bhuj; 9, Madhapar/Bhuj; 10, Madhapar/Bhuj; 11, Saiyedpar/Bhuj; 12, Dahisar/Mandvi; 13, Koday/Mandvi; 14, Kodaypul/Mandvi; 15, VRTI/Mandvi; 16, Vijay Vilas Palace/Mandvi; 17, Kharoi/Bhachau; 18, Ustiya/Abdasa; 19, Kothara/Abdasa; 20, Ranara Mota/Nakhatrana).

elders. The soil samples were collected at two depths (0–15 and 16–30 cm) from the sampling sites along with plant propagules.

### **Habitat and associated vegetation study**

The climate of the Kachchh region of Gujarat is hot to semi-arid, with a maximum temperature of 39–45°C during May–June and a minimum temperature of 1–8°C during December–January (Mangalassery *et al.*, 2017). The average rainfall in Kachchh is 385 mm (average of 1998–2016), and most rainy days occur from July to September with high evapotranspiration (1500–2000 mm per year). Rainfall is scant, erratic and irregular with a high coefficient of variation (CV) (71.25%). The collection sites represented different habitats, including grasslands, rocky land, barren land, piedmont plain, lower hills and valleys, scrub forest and agricultural field boundaries in the Kachchh region. Data on the local habitat, associated vegetation and soils of each collection site were recorded and analysed to determine the influence of local factors on the status of the plants.

### **Analysis of soil samples**

The collected soil samples were analysed for essential soil characteristics and micronutrient status (soil pH, EC, OC, N, P, K, Na<sup>+</sup>, Ca<sup>+</sup> and Mg<sup>2+</sup>). Soil pH and EC (dS/m) were determined by taking the supernatant solution with a 1:2 soil/water ratio (w/v) using a laboratory pH meter and conductivity meter (Jackson, 1973). An atomic absorption spectrophotometer (AA500, PG instrument, UK) was used to quantify the Na<sup>+</sup>, K<sup>+</sup> and Mg<sup>2+</sup> concentrations in the soil samples, whereas ethylenediaminetetraacetic acid (EDTA) complexometry was used to determine the concentration of Ca<sup>+</sup>. The organic carbon (%) and available phosphorus (%) of soil samples were determined by Walkley and Black's rapid titration method (Walkley and Black, 1934) and Olsen's method (Olsen *et al.*, 1954), respectively.

### **Standardization of propagation and nursery study**

Collected fruits and hardwood cuttings of *C. sinensis* were brought to the horticulture nursery of Regional Research Station, ICAR-Central Arid Zone Research Institute,

Kukma-Bhuj, Gujarat (India), where the hardwood cuttings (1.0–1.5 cm thickness and 15–20 cm length) were washed under running water and treated with 0.2% solution of Bavistin to remove the surface impurities and then immediately planted in polythene bags. For the propagation of seeds, freshly collected ripe fruits were rubbed with sand to clear away any gummy mucilage and then dried in the shade. Dried seeds were soaked in water overnight before sowing and were disinfected using Bavistin solution (2%) for disinfection. Seeds and hardwood cuttings were sown/planted in black poly bags (15 × 10 cm height) filled with clay, sand and FYM (2:4:1) and kept under a green shade net (50% shade) where regular intercultural operations followed. Data including days to sprouting, sprouting (%), seed germination (%), days to germination and survival (%) were recorded in the nursery stage.

### **Transplanting and preliminary field evaluation**

The 16 best germplasms were selected among one-year-old plants on the basis of the growth rate. Transplantation (pit size – 0.6 × 0.6 × 0.6 m) occurred in the field during August–September 2016 with five replications at a spacing of 6 × 6 m under a randomized complete block design. For morphological characterization, data on 15 growth parameters, including plant height (cm), plant spread (cm<sup>2</sup>), number of branches, leaves per plant, collar diameter (mm), leaf area (cm<sup>2</sup>), leaf length (cm), leaf width (cm), petiole length (mm), petiole width (mm) and fresh and dry weights (g) of the ten leaves were recorded. Furthermore, data on dry matter (%), moisture content (%) and crude protein (%) were recorded using leaf samples collected from three-year-old plants.

### **Statistical analysis**

The collected data were subjected to statistical analysis including basic descriptive statistics and Pearson's correlation analysis with the help of the software package 'PAST3' (Hammer *et al.*, 2001).

## **Results and discussion**


### **Collection, distribution and diversity**

The distribution pattern of the collected germplasm of *C. sinensis* based on habitat, location (taluka), soil status (pH) and topography are presented in Fig. 2. Field surveys revealed the occurrence of *C. sinensis* at the range of 22°84' 624–23°56'288N° latitude to 68°95'730–70°73'100E° longitude and 0–221 m altitude with maximum distribution in Bhuj (Kotda Ugamana, Bhujodi, Madhapar, Tapekshwari hill region, Saiyedpar), followed by Mandvi (Dahisar,

Koday, Kodaypul, nearby VRTI and Vijay Vilas Palace), Rapar (Pragpar), Abdasa (Kothada), Nakhatrana (Ranara Mota) and Bhachau talukas. The effects of local habitat, soil status and site topography on the species existence of *C. sinensis* were indicated by their distribution pattern. The sites with the highest collection values belonged to taluka Bhuj (9), followed by Mandvi (5), Abdasa (2), Rapar (2), Bhachau (1) and Nakhatrana (1) of Kachchh (Gujarat). Based on habitat and topography, maximum collections came from cultivated field boundaries (8), followed by five in each of scrub forests and roadsides (online Supplementary Fig.S1(a)–(d)). Approximately 45% of germplasm collections belonged to taluka Bhuj, followed by Mandvi (25%), Rapar (10%) and Abdasa (10%), with a minimum of 5% each in Nakhatrana and Bhachau. Field surveys revealed the presence of a good amount of natural density (%) in habitats such as field boundaries (35), rainfed scrub forests (30), roadsides (25) and backyards (10). Maximum collections occurred in areas with level topography (60%) with a soil pH in the range of 8–8.5 (63%), whereas only 6% of the collections came from sites with soil pH > 8.5.

*C. sinensis* is widely distributed in arid and semiarid regions of America, Asia and Africa, mainly in Egypt, Ethiopia, India, Israel, Jordan, Kenya, Madagascar, Mozambique, Namibia, Pakistan, Senegal, Somalia, South Africa, Sri Lanka, Sudan, Tanzania, Yemen and Zimbabwe (Orwa *et al.*, 2009). Due to its suitable adaptive mechanism and long tap root system, the species can survive in rainfed, arid and semiarid habitats (Meghwal *et al.*, 2014). In India, this species is commonly found in the arid and semiarid parts of Gujarat, Rajasthan (Jaisalmer, Jodhpur, Pratapgarh), Maharashtra (Nasik, Pune, Thane), Karnataka (Bangalore, Dharwar, Chikmaglour, North Karnataka), Tamil Nadu (Coimbatore, Cuddalore, Dharamapuri, Salem, Tiruchhirapalli, Tirunelveli, Vellore) and Kerala (Idakki) (Renu *et al.*, 2016). Its distribution in different parts of Gujarat was reported in earlier studies as follows: Panchmahal district (Qureshimatva *et al.*, 2016), Ambaji forest of Banaskantha district (Patel, 2015), Bhandaria Forest Area of Bhavnagar district (Chavda and Mehta, 2015), Mehsana district (Barot *et al.*, 2017), Thol Lake Wildlife Sanctuary of Mehsana district (Vyas and Patel, 2015), Rampara forest in Saurashtra region (Panchal and Pandey, 2004), Surat district (Gadhia *et al.*, 1999) and in terrestrial areas of Ghogha, Jalapor, Jambusar, Olpad, Talja, Umbagaon, Gandeve, Jafrabad, Khambhat, Kodinar, Mahuva, Una talukas situated around the Gulf of Khambhat (Gujarat Ecology Commission, 2011) as well as in Bhachau and Mandvi talukas of the Gulf of Kachchh (Gujarat Ecology Commission, 2014).

During the field survey, wide genetic diversity was observed in natural populations of *C. sinensis* with respect to plant height, flowering (time and colour), fruiting (time



**Fig. 2.** Distribution pattern of collected germplasm of *C. sinensis* on the basis of habitat, collection region, soil pH and topography.

and colour), and growth habits of plants from the inland region of Rapar, Bhuj, Bhachau to the coastal region of Mandvi and Abdasa taluka. In addition to *C. sinensis*, fair diversity was observed in other *Cordia* species, such as *C. myxa* L, *C. monoica* Roxb., *C. perrottetii* Wt. and *C. sebestena* L. The vigorous growth of plants with generous bearing capacity noted in the coastal region might be attributed to the presence of high atmospheric humidity at the collection sites. Early flowering and fruiting (February to April and May to August, respectively) were noted in the coastal talukas (Mandvi and Abdasa) compared to the other inland sites of Bhuj, Rapar and Bhachau talukas. A considerable amount of variation was observed in the fruit size (small to large), colour of ripe fruits (light orange to dark orange) and fruit-bearing potential (light to a profuse bearing) (online Supplementary Fig. S1(e)–(j)).

A tree census on non-forest areas of Gujarat by the Gujarat forest department reported the majority of the *C. sinensis* population was in the South Gujarat region, followed by the Kachchh and Saurashtra (315,000) regions (Khanna and Singh, 2010). In addition to *C. sinensis*, the Kachchh region of Gujarat had rich diversity of other underutilized plant species, such as *G. tenax*, *G. villosa*, *G. flavescens* and *C. myxa*. The rich diversity of *Cordia* species, including *C. sinensis*, was observed in the Kotda Chakar (Bhuj), Dahisara and Koday (Mandvi) regions of Kachchh (Dev *et al.*, 2016). Earlier studies have reported the vast genetic diversity of *C. myxa* L. in natural populations regarding its morphological traits (Pareek and Sharma, 1993; Saini *et al.*, 2002; Malik *et al.*, 2010). A morphological and RAPD marker study on 22 *Cordia* accessions of the Indian arid zone revealed a considerable amount of genetic diversity (Sivalingam *et al.*, 2012).

Wide variability has also been reported for 14 accessions of *C. myxa* collected from arid regions of the Rajasthan and Gujarat states in terms of fruit characteristics such as bunch weight (14.1–137.4 g), fruit weight (1.96–10.5 g), fruit numbers per bunch (4.62–14.8) and pulp: stone ratio (1.1–6.96) (Meghwal et al., 2014).

### Primary soil status, habitat and associated vegetation

The soil analysis of the surveyed sites (Bhuj, Mandvi, Nakhatarana, Rapar, Bhachau and Abdasa) represented sodic soil pH (7.1–9.2) with deficient organic carbon (0.01–0.48%). The available cations Na<sup>+</sup>, Ca<sup>+</sup> and K<sup>+</sup> (ppm) ranged from 65 to 437.14, 13.33 to 123.33 and 40 to 470.40, respectively (online Supplementary Table S2). High population densities were observed for *C. sinensis* and *C. myxa* at soil pH values of 8.5 and 8.2, respectively. *C. sinensis* can be grown on almost all types of soils but prefers clay, clay sand or alluvial soil, red loam and rocky soils (Warfa, 1990). Accessions collected from Mandvi taluka (CBCG-15 and CBCG-16) were identified as salt-tolerant accessions that require further study for their utilization in the future. A higher leaf Na<sup>+</sup> ion content was reported in *C. sinensis* (230–920 ppm), followed by *C. myxa* (200–830 ppm) collected from the Kachchh region (Dev et al., 2019). The domestication process in this species is still not yet completed compared to *C. myxa*. This was evidenced by the distribution pattern and soil status of species observed during the field survey, as a high amount of the overall population (63%) belonged to habitats such as scrub forest, field boundaries and roadsides with soil pH values in the range of 8.0–8.5, whereas less of the overall lower population (6%) was found on sites near human settlements or as backyard plantations (10%) with soil pH > 8.5.

Field surveys revealed the underutilized shrub and tree species including Ingorio (*Balanites aegyptica* L), Nano Akado (*Calotropis procera*), Kair (*C. decidua* (Forsk.) Edgew.), Avar (*Cassia auriculata* L.), Tankaro (*Clerodendron plomidis* L.), Guggal (*Commiphora wightii* (Arn.) Bhandari), Danda-thor (*Euphorbia caducifolia* Hains.), Gangeti (*G. tenax* (Forsk.) Fiori), Luska (*Grewia villosa* Willd.), Kundher (*Premna resinosa* Schau), Mithi Jar (*Salvadora oleoides* Dence), Khari Jar (*S. persica* L.) and Bordi (*Z. nummularia* (Burm.f.) Wight & Arn.) as close associates of *C. sinensis*. Its natural preference for moist habitat has been evidenced by its wide occurrence near the coastal area of the Gulf of Khambhat and Gulf of Kachchh of Gujarat (Gujarat Ecology Commission, 2011; 2014). However, its preference for dry riverine land, open bushland (Orwa et al., 2009), riverbanks, seasonally flooded areas and in-land depressions, moist grounds or

sometimes gardens (Warfa, 1990) has also been reported in association with *Balanities*, *Combretum*, *Ficus*, *Grewia*, *Salix*, *Salvadora*, *Tamarindus*, *Thespesia* and *Ziziphus* species (Maundu et al., 1999; Eltayeib and Ishag, 2015).

### Propagation

Data on seed germination and sprouting in cuttings of *C. sinensis* were observed on the 12th and 17th days of sowing and planting, respectively (online Supplementary Table S3). The results showed low seed germination (28–42%) and sprouting in cuttings (17–32%). Two accessions (CBCG-12 and CBCG-13) showed early seed germination (8 d) in the nursery. Although sprouting was noted in hardwood cuttings, plants were dried subsequently due to rooting failure. Similar observations were recorded by Meghwal and Vashistha (1998) in the propagation of *C. decidua* through hardwood cuttings. Hence, only 16 seed-propagated accessions could be used for field establishment and further study. Meghwal (2007) reported the seed as the standard propagation method of *Cordia* species, including *C. sinensis*, who also noted higher seed germination (50–60%) for *C. sinensis* than for *C. myxa* (only 20–30%).

### Preliminary field evaluation

#### Range of variation

The morphological characterization of 16 *C. sinensis* genotypes using 15 growth parameters is presented in Table 1. The mean plant survival after two and a half years of planting was 96% in the field. The morphological characterization revealed the presence of a considerable amount of variation expressed as the CV (%). The maximum CV (%) was observed for the number of leaves per plant (66.6), followed by the number of branches per plant (45.62), collar diameter (27.69), percentage of dry matter (25.34) and minimum fresh weight of 10 leaves (6.01). The range and mean values of essential characteristics such as plant height (121.67–212.0 and 166.83 cm), plant spread (118–223 and 170.5 cm<sup>2</sup>), number of branches per plant (6–24.33 and 15.16), number of leaves per plant (322.33–2991.33 and 1656.5) and collar diameter (17.53–45.99 and 27.0 mm). Accession 'CBCG-11' collected from Saiyedpar village of the Bhuj taluka showed maximum plant growth in terms of plant height (212 cm), canopy spread (223 cm<sup>2</sup>) and collar diameter (45.99 mm). The maximum number of branches (24.33) and leaves per plant (299.33) were recorded for the accession 'CBCG-16'. Three accessions, CBCG-13, CBCG-16 and CBCG-8, showed higher values in leaf parameters such as leaf area, leaf length, leaf width, fresh weight of ten leaves and lateral veins per

**Table 1.** Results of the characterization of *C. sinensis* germplasm in the field

Parameters	Range		Mean	SD	CV (%)	SE ( $\pm$ )
	Min	Max				
Plant height (cm)	121.67	212.00	166.83	40.99	24.47	10.25
Plant spread (cm <sup>2</sup> )	118.00	223.00	170.50	31.36	17.59	7.84
No. of branch/plant	6.00	24.33	15.16	6.06	45.62	1.52
No. of leaves/plant	322.33	2991.33	1656.50	846.91	66.60	211.73
Collar diameter (mm)	17.53	45.99	27.00	7.48	27.69	1.87
Leaf area (cm <sup>2</sup> )	11.14	20.82	14.60	3.03	20.73	0.76
Leaf length (cm)	5.21	6.63	5.78	0.40	6.95	0.10
Leaf width (cm)	2.27	2.87	2.62	0.16	6.01	0.04
Petiole length (cm)	0.59	0.88	0.75	0.08	10.94	0.02
Petiole width (mm)	0.97	1.25	1.10	0.08	7.53	0.02
Leaf fresh wt. (10 no's)	1.85	3.15	2.43	0.34	14.14	0.09
No of lateral vein/leaf	9.20	15.00	11.85	1.35	11.41	0.34
Dry matter (%)	18.49	49.16	31.19	7.90	25.34	1.98
Moisture content (%)	50.84	81.51	68.82	7.90	11.48	1.98
Crude protein (%)	3.86	7.22	5.43	0.97	17.82	0.24

leaf. In some accessions of *C. sinensis* (CBCG-12, CBCG-13 and CBCG-14), flowering was noted late in the third year during the third week of February and fruiting was noted in March–April 2019. Meghwal *et al.* (2014) also reported significant variations in flowering pattern, fruiting and fruit yield among 14 on-farm conserved *C. myxa* genotypes collected from Rajasthan and Gujarat.

### Variation in leaf qualitative traits

Five leaf qualitative traits of collected germplasm are presented in Table 2 and online Supplementary Fig. S2. There was considerable variation in leaf apex (acute, 8; obtuse, 6; retuse, 2), leaf base (cuneate, 9; obtuse, 5; oblique, 2) and leaf shape (lanceolate, 1; elliptic, 3; ovate, 4; oblanceolate, 5; oblong, 2; obovate, 1), whereas no variation was observed on the 'entire' leaf margin or 'simple' leaf type.

### Reproductive biology

Although flowers of *C. sinensis* are hermaphroditic in nature, functional male flowers are also present with rare female flowers (Warfa, 1990). Flowering and fruiting occur twice per year, during September–October and February–March. Its flowers are entomophilous mainly by honeybees, houseflies and yellow wasps due to their sweet fragrance. Plants of this species are deciduous, as they remain in physiological dormancy for almost two months during winter (December–January) when leaf shedding

occurs. The fruits are harvested upon changing colour from green to orange.

### Traditional utilization

Different aspects of *C. sinensis* utilization, such as direct consumption of fruits, leaves as fodder and wood as timber, have been reported worldwide. Bright red or orange coloured, fully ripened fruits are edible and sold in local markets of Kenya (Maundu *et al.*, 1999) and some parts of arid Rajasthan (Rathore, 2009). The chemical composition of fruits (online Supplementary Table S4) indicates its good nutritive (Kuria *et al.*, 2005) and fodder (Murray *et al.*, 2001) value. The sweet and sticky pulps were consumed fresh and sometimes utilized in porridge as a sugar substitute. Children or adults in rural areas have consumed fruits during the fruiting season or during food shortages. In Kenya, fruits are sun-dried and stored in a wooden container for future uses (Maundu *et al.*, 1999). Fresh fruit pulps are squeezed in water, mixed with tamarind (*Tamarindus indica*) juice and kept for fermentation to brew local beer (Orwa *et al.*, 2009).

Generally, leaves of *C. sinensis* are used as browse materials by goats, sheep, cattle and camels during dry and wet seasons in arid and semiarid areas (Kuria *et al.*, 2005; Orwa *et al.*, 2009; Deng *et al.*, 2017b). Abdullah *et al.* (2017) classified arid species based on their palatability as highly palatable (*Prosopis cineraria* and *Acacia nilotica*), moderately palatable (*C. polygonoides*, *Suaeda fruticosa*, *Salsola baryosma*, *Haloxylon*

**Table 2.** Variation in the leaf qualitative traits of collected *C. sinensis* germplasm

Germplasm	Leave apex	Leaf base	Leaf margin	Leaf shape	Leaf type
CBCG-1	Acute	Cuneate	Entire	Lanceolate	Simple
CBCG-2	Obtuse	Obtuse	Entire	Elliptic	Simple
CBCG-3	Obtuse	Cuneate	Entire	Elliptic	Simple
CBCG-4	Acute	Oblique	Entire	Ovate	Simple
CBCG-5	Obtuse	Obtuse	Entire	Ovate	Simple
CBCG-6	Obtuse	Cuneate	Entire	Ovate	Simple
CBCG-7	Obtuse	Cuneate	Entire	Oblanceolate	Simple
CBCG-8	Acute	Cuneate	Entire	Oblanceolate	Simple
CBCG-9	Acute	Oblique	Entire	Oblanceolate	Simple
CBCG-10	Acute	Cuneate	Entire	Oblanceolate	Simple
CBCG-11	Obtuse	Obtuse	Entire	Ovate	Simple
CBCG-12	Acute	Cuneate	Entire	Oblanceolate	Simple
CBCG-13	Acute	Obtuse	Entire	Oblong	Simple
CBCG-14	Acute	Cuneate	Entire	Obovate	Simple
CBCG-15	Retuse	Cuneate	Entire	Elliptic	Simple
CBCG-16	Retuse	Obtuse	Entire	Oblong	Simple

*recurvum*, *C. decidua*, *C. procera* and *Tamarix aphylla*) and less palatable (*H. salicornicum*) in the Cholistan Desert of Pakistan. Analysis of the nutrient composition of *C. sinensis* showed high year-round moisture and crude protein (13.8–15.93%) with high energy and mineral (potassium, phosphorus, calcium, zinc and iron) contents (Kuria et al., 2005; Deng et al., 2017a; Derero and Kitaw, 2018).

Traditionally, *C. sinensis* was used to treat different ailments and for wound-healing purposes (online Supplementary Table S5). It has the potential to cure gastric disorders, malaria, rheumatism, bladder diseases, gastric ulcers and infectious diseases (Orwa et al., 2009). The use of different plant parts, such as root and bark decoction, has been reported to treat malaria, stomach disorders and chest pains; bark has been used to treat gastric and respiratory disturbances and mixtures of roots and bark have been used to treat conjunctivitis in cattle (Orwa et al., 2009; Al-Musayeb et al., 2011). Leaves and ripe fruits were used alone or in combination with other medicinal plants to treat fever and dry cough, respectively (Adnan et al., 2014). Among the ten bioactive compounds isolated from *C. sinensis*, two compounds [kaempferide-3-O-D-glucopyranoside (62.4%) and kaempferol-3-O-D-glucopyranoside (59.6%)] were found to have potential as diclofenac (57.6%), which also showed significant anti-inflammatory activity (Thirupathi et al., 2008; Nyambe, 2014).

The hardwood of *C. sinensis* is locally used for making furniture, agricultural implements such as tool handles, walking sticks, clubs, wooden spoons, stirrers and stools as well as in the construction of local huts. Fibre from the inner bark is useful for making ropes, caulking boats, etc.

An edible gum from the bark may be used to adulterate gum Arabic (*Acacia senegal*). Its poles are sold for construction purposes in the local markets of Pokot and Turkana town of Kenya (Maundu et al., 1999).

*Cordia sinensis* is a preferred plant species in arid regions due to its multiple uses as food, fodder, medicine, gum and timber. Despite its numerous uses, its utilization and domestication process is still minimal. Poor domestication of this species was evidenced from its high population frequency between soil pH 8.0–8.5 on level topography and in habitats such as scrub forests, field boundaries, roadsides and its low population frequency on sites near a human settlement or as backyard plantation in the present study. The extensive genetic diversity noted on natural populations of species in Bhuj and Mandvi taluka of Kachchh could be useful for future studies. Based on the preliminary evaluation, accessions selected for different purposes were as follows: fast growth and fodder (CBCG-11 and CBCG-16); early flowering and fruiting (CBCG-12, CBCG-13 and CBCG-14); most natural propagation by seeds (CBCG-12 and CBCG-13) and salt tolerance (CBCG-15 and CBCG-16). Future research works should be focused on its exploration, sole plantation, standardization of propagation techniques and characterization and conservation aspects, which still exist at the infancy stage; thus, information about this valuable species could enhance its utilization by present and future generations.

#### Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1479262120000453>



## Acknowledgements

The author is thankful to the Director of ICAR-CAZRI for providing the necessary facility for conducting research.

## Funding

Not applicable.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## References

- Abdullah M, Rafay M, Khan T, Ahmad H, Tahir U, Rasheed F, Ruby T and Khalil S (2017) Nutritive potential and palatability preference of browse foliage by livestock in arid rangelands of Cholistan desert (Pakistan). *Journal of Animal and Plant Sciences* 27: 1656–1664.
- Adnan M, Ullah I, Tariq A, Murad W, Azizullah A, Khan AL and Ali N (2014) Ethnomedicine use in the war affected region of northwest Pakistan. *Journal of Ethnobiology and Ethnomedicine* 10: 16.
- Al-Musayeb N, Perveen S, Fatima I, Nasir M and Hussain A (2011) Antioxidant, anti-glycation and anti-inflammatory activities of phenolic constituents from *Cordia sinensis*. *Molecules* 16: 10214–10226.
- Barot CJ, Patel VA and Dabgar YB (2017) Floristic study of some wetlands and its corridor of mehsana district, North Gujarat. *International Journal for Research Trends and Innovation*. 2: 171–180. <http://www.ijrti.org/papers/IJRTI1707027.pdf> accessed on 19 March 2019.
- Bouby L, Bouchette A and Figueiral I (2011) *Sebasten* fruits (*Cordia myxa* L.) in Gallia Narbonensis (Southern France): a trade item from the Eastern Mediterranean. *Vegetation History and Archaeobotany* 20: 397–404.
- Chaovanalikit A, Thompson MM and Wrolstad RE (2004) Characterization and quantification of anthocyanins and polyphenolics in blue honeysuckle (*Lonicera caerulea* L.) *Journal of Agricultural and Food Chemistry* 52: 848–852.
- Chavda NH and Mehta SK (2015) Study of species diversity of trees and shrubs in Bhandaria Forest Area, District-Bhavnagar, Gujarat. *International Journal of Pure & Applied Bioscience* 3: 356–361.
- Deng MT, Ondiek JO and Onjoro PA (2017a) Chemical composition and *in Vitro* gas production of lesser known South Sudan browse species. *Livestock Research for Rural Development* 29: 81. Retrieved July 4, 2019, from <http://www.lrrd.org/lrrd29/4/mame29081.html>.
- Deng MT, Ondiek JO and Onjoro PA (2017b) Intake, relative palatability index and preference class of selected Bor, South Sudan browse species fed to crossbred growing goats. *Livestock Research for Rural Development*. 29: 171. Retrieved July 5, 2019, from <http://www.lrrd.org/lrrd29/9/mame29171.html>.
- Derero A and Kitaw G (2018) Nutritive values of seven high priority indigenous fodder tree species in pastoral and agro-pastoral areas in Eastern Ethiopia. *Agriculture & Food Security* 7: 68. <https://doi.org/10.1186/s40066-018-0216-y>.
- Dev R, Kumar MS, Kumar S and Dayal D (2016) Survey, collection and utilization of grewia and cordia species for human and animal nutrition in Arid Kachchh, Gujarat. In IAC Abstracts Book. 1st International Agrobiodiversity Congress, November 6–9, 2016, New Delhi, India, 344 p.
- Dev R, Sureshkumar M, Singh T and Dayal D (2019) Distribution, adaptation and utilization of local underutilized grewia and cordia species in the salt-affected Kachchh Region for better livelihood and fodder security. In: Sharma PC, Yadav RK, Bundela DS, Anil Kumar, Sanwal SK, Meena RL, Rakesh Banyal, Jogendra Singh, Raj Kumar, Fagodiya RK, Arijit Barman, Anshuman Singh, Ashim Datta, Priyanka Chandra and Awtar Singh. Golden Jubilee International Salinity Conference on Resilient Agriculture in Saline Environments under Changing Climate: Challenges & Opportunities. Indian Society of Soil Salinity and Water Quality, Karnal, Haryana. 187pp.
- Dhakar MK, Das B, Nath V, Sarkar PK and Singh AK (2019) Genotypic diversity for fruit characteristics in bael [*Aegle marmelos* (L.) Corr.] based on principal component analysis. *Genetic Resources and Crop Evolution* 66: 951–964. <https://doi.org/10.1007/s10722-019-00757-x>
- Eltayeib AA and Ishag W (2015) Phytochemical screening, antimicrobial, antioxidant and cytotoxicity activities of bark's crude extracts of *Cordia sinensis*. *Advancement in Medicinal Plant Research* 3: 39–45.
- Gadhia PK, Gadhia M and Parabia MH (1999) Dietary, health and biotic baseline survey around Kakrapar atomic power station. Report published by South Gujarat University, Surat in December, 1999. p. 136.
- GEC (2014). Ecological profile for coastal Talukas of Gulf of Kachchh. Gujarat Ecology Commission, Gandhinagar, 120p.
- GEC (Gujarat Ecology Commission) (2011) Ecological profile of coastal talukas around gulf of Khambhat. Gujarat Ecology Commission, Gandhinagar. 32p. [www.gec.gov.in](http://www.gec.gov.in).
- Hammer O, Harper DAT and Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 9. [http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm).
- Jackson ML (1973) *Soil Chemical Analysis*. New Delhi: Prentice Hall of India Pvt. Ltd., pp. 38–56.
- Khanna P and Singh HS (2010) Tree wealth of the non-forest areas of Gujarat. A report tree census in non-forest areas-2009. Forest department, Gujarat state. p. 47.
- Kumar S and Parveen F (2004) Traditional ethno-medicinal plants in the Indian Thar: their status in nature and possibilities of their cultivation and trading. Final report of DST Project 216.
- Kumar V, Pal V, Sapra P and Umrao V (2008) Variability evaluation in Beal (*Aegle marmelos* Correa) accessions. *Annals of Horticulture* 1: 23–26.
- Kuria SG, Wanyoike MM, Gachuii CK and Wahome RG (2005) Nutritive value of important range forage species for camels in Marsabit district, Kenya. *Tropical and Subtropical Agroecosystems* 5: 15–24.
- Malik SK, Chaudhury R, Dhariwal OP and Bhandari DC (2010) *Genetic Resources of Tropical Underutilized Fruits in India*. New Delhi: NBPGR, p. 168.
- Mangalassery S, Dayal D, Arvind K, Bhatt K, Nakar DR, Kumar A, Singh J and Misra A (2017) Pattern of salt accumulation and its impact on salinity tolerance in two halophyte grasses in extreme saline desert in India. *Indian Journal of Experimental Biology* 55: 542–548.
- Maundu PM, Ngugi GW and Kabuye CHS (1999) Traditional food plants of Kenya. Kenya Resources Centre for Indigenous Knowledge. National Museums of Kenya, 270 p.
- Meghwal PR (2007) Propagation studies in lehsua (*Cordia myxa*). *The Indian Journal of Agricultural Sciences* 77: 765–767.

- Meghwal PR and Vashishtha BB (1998) Effect of time of planting and auxin on rooting of cuttings of ker (*Capparis decidua* L. Forsk). *Annals of Arid Zone* 37: 401–404.
- Meghwal PR, Singh A, Kumar P and Morwal BR (2014) Diversity, distribution and horticultural potential of *Cordia myxa* L.: a promising underutilized fruit species of arid and semi-arid regions of India. *Genetic Resources and Crop Evolution* 61: 1633–1643.
- Murray SS, Schoeninger MJ, Bunn HT, Pickering TR and Marlett JA (2001) Nutritional composition of some wild plant foods and honey used by hadza foragers of Tanzania. *Journal of food composition and analysis* 14: 13.
- Nyambe MM (2014) Phytochemical and antibacterial analysis of indigenous chewing sticks, *Diospyros lycioides* and *Euclea divinorum*. M.Sc. Thesis, The University of Namibia, Namibia.
- Olsen SR, Cole CV, Watanabe FS and Dean LA (1954) Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939.
- Orwa C, Mutua A, Kindt R, Jamnadass R and Anthony S (2009) Agroforestry Database: a tree reference and selection guide version 4.0. <http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>.
- Panchal NS and Pandey AN (2004) Analysis of vegetation of Rampara forest in Saurashtra region of Gujarat state of India. *Tropical Ecology* 45: 223–231. [http://www.tropecol.com/pdf/open/PDF\\_45\\_2/45204.pdf](http://www.tropecol.com/pdf/open/PDF_45_2/45204.pdf). accessed on 20.03.2019.
- Pareek OP and Sharma S (1993) Underutilized fruits. *Indian Horticulture* 38: 47–56.
- Patel RS (2015) An ethnobotanical study of medicinal plants used in sacred groves of Ambaji forest, Gujarat, India. *International Journal of Advanced Technology in Engineering and Science* 3: 2348–2350. [http://ijates.com/images/short\\_pdf/1425811703\\_474.pdf](http://ijates.com/images/short_pdf/1425811703_474.pdf). accessed on 19.03.19.
- Qureshmatva UM, Maurya RR, Gamit SB and Solanki HA (2016) Checklist of flowering plants in Panchmahal District, Gujarat, India. *Forest Research* 5: 176.
- Rathore M (2009) Nutrient content of important fruit trees from arid zone of Rajasthan. *Journal of Horticulture and Forestry* 1: 103–108.
- Renu G, Thilakar SJ and Narasimhan D (2016) *Cordia gharaf* Ascherson. [online] India Biodiversity Portal, Species Page: {*Cordia gharaf* Ascherson} Available at: <http://indiabiodiversity.org/biodiv/species/show/31221>. accessed on 21 August 2016.
- Ronoh R, Linde M, Winkelmann T, Abukutsa-Onyango M, Dinssa FF and Debener T (2019) Morphological characterization, genetic diversity and population structure of African nightshades (section *Solanum* L.). *Genetic Resources and Crop Evolution* 66: 105–120. <https://doi.org/10.1007/s10722-018-0700-z>
- Saini RS, Kaushik RA and Singh S (2002) Research note on the evaluation of lasora (*Cordia myxa* Roxb.) germplasm for vegetative growth characters under semi-arid conditions. *Haryana Journal of Horticultural Sciences* 31: 62–63.
- Sharma AK and Tewari JC (2005) Arid zone forestry with special reference to Indian hot arid zone, in forests and forest plants, from Encyclopedia of life support systems (EOLSS). Developed under the auspices of the UNESCO, Eolss Publishers, Oxford, UK. <http://www.eolss.net>.
- Singh A and Meghwal PR (2020) Socio-economic and horticultural potential of *Ziziphus* species in arid regions of Rajasthan India. *Genetic Resources and Crop Evolution* 67: 1301–1313. <https://doi.org/10.1007/s10722-020-00891-x>
- Singh JP, Rathore VS and Beniwal RK (2008). Diversity and conservation of shrubs of hot arid region of Rajasthan, India. In: Singh NP and Chhonkar PK (eds), *Biodiversity for Sustainable Development*. Samakriti, New Delhi, pp. 51–71.
- Sivalingam PN, Singh D and Chauhan S (2012) Morphological and molecular diversity of an underutilized fruit crop – *Cordia myxa* L. germplasm from the arid region of Rajasthan, India. *Genetic Resources and Crop Evolution* 59: 305.
- Thirupathi K, Kumar SS, Raju VS, Ravikumar B, Krishna DR and Mohan GK (2008) A review of medicinal plants of the genus *Cordia*: their chemistry and pharmacological uses. *Journal of Natural Remedies* 8: 1–10.
- Vyas DN and Patel AH (2015) Tree species diversity and their phytosociological evaluation in ‘Thol Lake Wildlife Sanctuary, North Gujarat. *International Journal of Scientific Research* 4: 713–716.
- Walkley A and Black IA (1934) An examination of Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science* 37: 29–37.
- Wang Y, Xing W, He X, Duan W, Huang W and Gong C (2015) Differential comparison among different geographical provenances of *Stauntonia urophylla*. *Journal of Central South University of Forestry & Technology* 35: 55–60.
- Warfa AM (1990) Taxonomy and distribution of *Cordia sinensis* and *C. nevillii* (Boraginaceae), a widespread species pair in Africa and Asia. *Nordic Journal of Botany* 9: 649–656.
- Zou S, Yao X, Zhong C, Gao P, Wang Z and Huang H (2020) Phenotypic characterization of *Stauntonia obovatifoliola* Hayata subsp. *urophylla* germplasm: a potential new fruit crop. *Genetic Resources and Crop Evolution* 67: 1037–1050. <https://doi.org/10.1007/s10722-020-00885-9>