

Two new foliicolous species of *Enterographa* (Roccellaceae) from Kenya

Kumelachew YESHITELA, Eberhard FISCHER,
Dorothee KILLMANN and Emmanuël SÉRUSIAUX

Abstract: *Enterographa fellhaneroides* and *E. meklitiae* are described as new to science from Kakamega Forest in Western Kenya. The latter overgrows the thallus of a foliicolous non-filamentous *Coenogonium* and the former most probably has the same ecology. Both species appear to be closely related to *E. brezhonega* and *E. epiphyllum*.

Key words: *Coenogonium*, *Enterographa fellhaneroides*, *Enterographa meklitiae*, lichens, *Plectocarpon*, *Sclerophyton*

Introduction

The lichen genus *Enterographa* Fée (Roccellaceae) has been expertly monographed by Sparrius (2004) and is characterized by the following characters: crustose growth habit, photobiont *Trentepohlia* or *Phycopeltis*, rounded to elongate or punctiform ascospores, most usually immersed in the thallus, exciple poorly developed, hamothecium of branched and anastomosing paraphysoids, ellipsoid to cylindrical-clavate asci of the *Opegrapha*-type, ascospores usually fusiform with thin septa and perispore, without enlarged terminal cells, conidiomata punctiform, and bacilliform or filiform microconidia. As circumscribed, the genus comprises 35 species, while the related genus *Sclerophyton* Eschw. includes species with macrocephalic ascospores with relatively thick septa and ascospores included in pseudostromata, with well-known and widespread species such as *Enterographa elegans* (Eschw.) Tuck, and *E.*

extenuatum (Nyl.) Redinger being transferred to *Sclerophyton*. Since the monograph by Sparrius (2004), 8 further new species have been described in the genus *Enterographa* and one species transferred into it (Ertz *et al.* 2005; Sparrius *et al.* 2006; Aptroot *et al.* 2007; Cáceres 2007; Sparrius & Aptroot 2007; Sparrius & Björk 2008; Jagadeesh Ram *et al.* 2008).

In spite of several detailed studies (Grube 1998; Myllys *et al.* 1999; Tehler 1990; Tehler *et al.* 1997) generic delimitation within the Roccellaceae is unsatisfactory and several genera are most probably polyphyletic or paraphyletic (D. Ertz, pers. comm.). A detailed phylogenetic study of two genes by Tehler & Irestedt (2007) could even demonstrate parallel evolution of the fruticose growth form in the supposedly well-circumscribed genera *Roccella* and *Roccellina*. The recent inclusion of *Chiodecton epiphyllum* Sérus., a species growing over a foliicolous species of *Coenogonium* in Papua New Guinea and the Neotropics, in *Enterographa* as a “better solution” than its earlier assignment to *Plectocarpon* Fée, shows that more data should be gathered to reach a well-supported delimitation of the genus (Ertz *et al.* 2005).

Santesson (1952) recognized three foliicolous species of *Enterographa*. Since then

K. Yeshitela, E. Fischer and D. Killmann: Institute for Integrated Natural Sciences, Department of Biology, University of Koblenz-Landau, Universitätsstraße 1, D-56070 Koblenz, Germany.

E. Sérusiaux (corresponding author): Plant Taxonomy & Conservation Biology Unit, University of Liège, Sart Tilman B22, B-4000 Liège, Belgium. Email: e.serusiaux@ulg.ac.be

Vězda (1975), Sérusiaux (1984), Lücking (1991), Sérusiaux (in Aptroot *et al.* 1995), Lücking & Matzer (1996), Matzer (1996), Lücking *et al.* (1998), Herrera-Campos & Lücking (2002), Lücking *et al.* (2003), Lücking & Henssen (in Sparrius 2004) and Sparrius & Björk (2008) described further new species. A total of 15 foliicolous species of *Enterographa* are currently known, including 2 lichenicolous fungi on foliicolous lichens (Sparrius 2004; Ertz *et al.* 2005); most of them distributed in the Neotropics and eastern Palaeotropics. Only two species (*E. multiseptata* and *E. vezdai*) have so far been recorded from the African continent.

Here we describe two further foliicolous species belonging to *Enterographa*, discovered by the first author in Kakamega Forest (W Kenya) in August 2006. One was clearly overgrowing the thallus of a species of *Coenogonium* and the second most probably had the same ecology.

Materials and Methods

Morphological characters were studied on dry specimens under a dissecting microscope. Anatomical characters were measured using light and interference contrast microscopy on hand-cut sections and squash preparations mounted in water. Water solution of KOH (10%) and lactophenol-cotton blue were used for detailed observation of asci and hamathecial elements. Amyloidy of the tholus of asci and hymenium were tested with Lugol's solution.

The study site

Kakamega Forest is located in Shinyalu Division of the Kakamega District in the Western Province of Kenya. It is situated about 40 km north-west of Lake Victoria between 0°10' and 0°21' N latitude and 34°47' and 34°58' E longitude and is made up of a main forest block covering 8500 ha surrounded by five forest fragments of various sizes (130–1400 ha; Farwig *et al.* 2006). The vegetation consists of a mosaic of primary rainforest, secondary forests at different stages of development, swamp and riverine forest, selectively logged forest, plantations of indigenous and exotic tree species, natural glades, and clearings resulting from anthropogenic disturbance (Tattersfield *et al.* 2001; Althof 2005). Kakamega Forest is the easternmost patch of the equatorial forests distributed over the Congo basin and the only tropical rainforest with a mixture of Guineo-Congolian and afro-montane species in Kenya (Althof 2005).

The Species

Enterographa fellhaneroides Yeshitela, Eb. Fisch., Killmann & Sérus. sp. nov.

Differt ab omnibus speciebus foliicolis generis *Enterographa* ascomata apothecioideis vel breve lirelliformibus, disco fusco cum margine albido et ascosporibus 6-septatis fusiformibus apicibusque rotundatis. *Enterographa brezhonega* differt ab *E. fellhaneroides* ascomata longe lirelliformibus saepe ramificatis et disco atrofusco. *Enterographa seawardii* differt sporibus 7–11-septatis anguste fusiformibus et thallo cum Pd+ luteo.

Typus: Kenya, Western Province, Kakamega Forest, 00°21'.276" N and 034°51'.519" E, 1609 m, on living leaves of *Dracaena fragrans*, 14 August 2006, Yeshitela 349 (LG—holotypus).

(Figs 1 & 2)

Thallus absent, ascomata overgrowing the thallus of a non-filamentous *Coenogonium* which is continuous, smooth, greyish green, usually with a whitish prothallus, photobiont *Trentepohlia* with cells angular-rounded, 12–20 × 5–8 µm.

Ascomata sessile, ± aggregated, rounded to shortly lirelliform, constricted at base, 0.2–0.8 mm diam., rarely up to 0.6 mm long, 50–60 µm high; disc exposed, pale to dark brown; margin 40–50 µm thick, slightly raised, whitish. *Excipulum* 10–15 µm wide, formed by densely interwoven hyphae filled with large crystals, hyaline or pale straw in section. *Hypothecium* 120–155 µm high including a 100–130 µm socle (formed by densely intricate hyphae filled with large, angular and hyaline oxalate crystals), hyaline, K–. *Hymenium* 50–75 µm high, light brown in the upper part (c. 25 µm), otherwise hyaline, I+ blue, rapidly turning red, KI+ blue. *Epithecium* 5–8 µm high, orange-brown; K–. *Hamathecium* of branched and anastomosing paraphysoids, c. 1.5 µm thick, not swollen apically. *Asci* of the *Opegrapha*-type, ellipsoid to clavate, bitunicate, 30–35 × 7.5–10 µm, with apical KI+ blue ring. *Ascospores* 4–6 per ascus, fusiform, with rounded ends, 6-septate, not constricted at the septa, the median cell usually slightly enlarged, hyaline, 15–18 × 2.5–3 µm, perispore up to 0.5 µm thick.

Conidiomata not found.




FIG. 1. *Enterographa fellhaneroides* (holotype). A–C, habit of apothecia; D, section through the apothecium. Scales: A–C = 0.2 mm; D = 50 μ m.




FIG. 2. *Enterographa fellhaneroides*, ascospores. Scale = 3 μ m.

Chemistry. No compounds detected by TLC.

Etymology. When we first observed the apothecia of this species, we considered it to be a poorly developed specimen of *Fellhanera* Vězda until the anatomical features pointed to *Enterographa*. Hence, the specific epithet is chosen to indicate this resemblance.

Notes. *Enterographa fellhaneroides* is easily distinguished from other foliicolous species assigned to the genus by its apothecioid or short lirelliform ascomata, with a pale brown disc and a whitish margin, and 6-septate, fusiform ascospores with rounded ends. Its closest related species appears to be the recently described *E. brezhonega* (Sparrius & Aptroot 2007), overgrowing the thallus of epiphytic *Porina rosei* in W France, and reported from the New Forest (S England) on the same host (Hitch 2008). This species is distinguished by its lirelliform, often branched ascomata with a dark brown to black disc. Otherwise both species are very similar, especially the ascospores.

Morphologically, *Enterographa fellhaneroides* looks similar to *E. seawardii* Lücking & Henssen, a recently described foliicolous species (in Sparrius 2004) and only known from the Seychelles, because of similar prominent, open, angular-rounded to shortly lirellate ascomata with pale orange discs. However, *Enterographa seawardii* is easily distinguished by its 7–11-septate, narrowly fusi-

form ascospores, 30–40 \times 3–4 μ m and its Pd+ yellow (probably psoromic acid) thallus.

Ecology and distribution. *Enterographa fellhaneroides* is so far known only from Kakamega Forest in Western Kenya in near primary and middle-aged secondary forest fragments. It is a lichenicolous species overgrowing thalli assigned to a *Coenogonium* sp. (poorly developed but typical apothecia seen). The species is a component of the foliicolous lichen flora of the understorey.

Selected specimens examined. **Kenya:** Western Province: Kakamega Forest, Kisere fragment 00°22'.966" N, 034°53'.751" E, 1594 m, on living leaves of *Chrysophyllum albidum*, 2006, *Yeshitela* 373 (KOBL); Kisere fragment, 00°23'.151" N, 034°53'.595" E, 1580 m, on living leaves of *Teclea nobilis*, *Dracaena fragrans* and *Cassipourea ruwenzorensis*, 2006, *Yeshitela* 538, 539 & 537 (KOBL); Isecheno fragment, 00°14'.522" N, 034°51'.959" E, 1580 m, on living leaves of *Cassipourea ruwenzorensis*, 2006, *Yeshitela* 484 & 493 (KOBL).

***Enterographa meklitiae* Yeshitela, Eb. Fisch., Killmann & Sérus. sp. nov.**

Differt ab omnibus speciebus foliicolis generis *Enterographa* ascomata numerosis punctiformibus in pseudostromata immersis. *Enterographa epiphylla* differt ab *E. meklitiae* pseudostromata convexibus punctiformibus ad vix lirellatis vel irregularibus, ascomata nigricantibus et ascosporibus majoribus 6-septatis.

Typus: Kenya, Western Province, Kisere fragment of Kakamega Forest, 00°23'.151" N and 034°53'.595" E, 1612 m, on living leaves of *Dracaena fragrans*, 24 August 2006, *Yeshitela* 539 (LG—holotypus).

(Figs 3 & 4)

Thallus assumed to be absent, ascomata most probably overgrowing the thallus of a non-filamentous *Coenogonium* which is continuous, smooth, greyish green, photobiont *Trentepohlia* with cells angular-rounded, 12–20 \times 3–7 μ m.

Ascomata punctiform, 0.05–0.1 mm diam., arranged in groups of 8–12(–20) in pseudostromata, visible as punctiform (30–50 μ m diam.) brownish spots. *Pseudostromata* rounded or irregular in outline, imarginate, convex to \pm plane, 0.2–0.5 mm diam., up to 75 μ m high, surface greyish green to whitish, not constricted at base, in section encrusted with large, angular and




FIG. 3. *Enterographa meklitiae* (holotype). A–C, habit of apothecia; D, section through the apothecium. Scales: A & B = 0.2 mm; C = 100 μ m; D = 50 μ m.




FIG. 4. *Enterograppha meklitiae*, ascospores. Scale = 3 μ m.

hyaline Ca oxalate crystals, K⁻. *Excipulum* very thin, c. 5 μ m wide, pale straw. *Hypothecium* 10–15 μ m high, pale straw, K⁻, I+ blue rapidly turning red, KI+ blue. *Hymenium* hyaline, I+ blue rapidly turning red, KI+ blue, 50–60 μ m high. *Hamathecium* of branched and anastomosing paraphysoids, c. 1.5 μ m thick, not swollen apically. *Epithecium* indistinct. *Asci* of the *Opegrapha*-type, ellipsoid to clavate, bitunicate, 37–50 \times 7–10 μ m, with apical KI+ blue ring. *Ascospores* 4 per ascus, 4–5-septate, ellipsoid to fusiform, with rounded ends, the median cell sometimes slightly enlarged, (13–)15–17 \times 2.5–3 μ m, not constricted at the septa, perispore < 5 μ m thick.

Conidiomata not observed. *Conidia* sometimes produced from tissue inside ascomata (probably the excipulum), simple, c. 2.5 \times 1 μ m, bacilliform, hyaline.

Chemistry. No compounds detected by TLC.

Etymology. This species is dedicated to Meklit, the daughter of the first author.

Notes. *Enterograppha meklitiae* can easily be distinguished from other foliicolous species of *Enterograppha* by its numerous, punctiform ascomata immersed in pseudostromata. At first glance, it looks like the conidiomata of the Neotropical *Phyllobathelium leguminosae* (Cavalc. & A. A. Silva) Lücking & Sérus. (see fig. 7E in Lücking *et al.* 1998). It is close to *Enterograppha epiphylla* (Sérus.) Ertz *et al.*, which overgrows the foliicolous lichen *Coenogonium flavicans*, and which can be dis-

tinguished by its convex pseudostromata, punctiform to slightly lirellate, or irregular, and blackish ascomata, and its larger ascospores (17–22 \times 3–4.5 μ m) with consistently 6 septa.

Ecology and distribution. *Enterograppha meklitiae* is so far recorded only from Kakamega Forest in Western Kenya. The type locality is a near-primary rain forest dominated by *Strychnos usambarensis*, *Uvariopsis congensis*, *Pouteria altissima* (= *Aninngeria altissima*), *Cassipourea ruwensorensis*, *Dracaena fragrans*, *Funtumia africana*, *Diospyros abyssinica*, and *Heinsenia dieverilleoides*. There is no strong evidence that it is a lichenicolous species overgrowing the thallus of a non-filamentous *Coenogonium*, but it is assumed to have such an ecology. Indeed, its ‘thallus’ is very similar to that of *Enterograppha fellhaneroides* (in one collection, both species grow over the same thallus), and all genuinely lichenized foliicolous species of *Enterograppha* have *Phycopeltis* as photobiont (Sparrius 2004).

Selected specimens examined. **Kenya**: Western Province: Kakamega Forest, Ischcheno, 00°14'.522" N, 034°51'.959" E, 1580 m, on living leaves of *Cassipourea ruwensorensis*, 2006, Yeshitela 484 (KOBL).

Discussion

As with most other genera in the *Roccellaceae*, the delimitation of the genus *Enterograppha* requires further studies, hopefully based on molecular sequences from several genes. Nevertheless, the four species described or discussed in this paper (*Enterograppha epiphylla*, *E. brezhonega*, *E. fellhaneroides* and *E. meklitiae*) characterized by their lichenicolous habit on thalli with *Trentepohlia* as photobiont and 4(–6) fusiform ascospores with rounded ends per ascus, an even number of septa and the median cell slightly enlarged, seem to be closely related. It is premature to suggest the taxonomic recognition of that group but equally, its inclusion in *Enterograppha*, *Plectocarpon* or *Sclerophyton* is not fully satisfactory.

We wish to thank very warmly Dr. Damien Ertz from the National Botanical Garden of Belgium for important

comments and suggestions; we further thank both anonymous referees for their interesting suggestions. The study was funded by the German Ministry of Education and Research (BMBF) within the BIOTA East project E04 (LC0625C1). The first author was funded by the German Catholic Academic Exchange Service (KAAD). Thanks are also due to the Kenya Wildlife Service (KWS) in Nairobi for a research permit in Kakamega Forest. We would also like to thank the staff of the East African Herbarium (EA) at the National Museums of Kenya (NMK), Nairobi and the Kenyan Wildlife Service field office in Kakamega for logistic support.

REFERENCES

- Althof, A. (2005) *Human impact on flora and vegetation of Kakamega Forest, Kenya*. Ph.D. Thesis, University of Koblenz-Landau.
- Aptroot, A., Diederich, P., Sérusiaux, E. & Sipman, H. J. M. (1995) Lichens and lichenicolous fungi of Laing Island (Papua New Guinea). *Bibliotheca Lichenologica* **57**: 19–48.
- Aptroot, A., Saipunkaew, W., Sipman, H. J. M., Sparrius, L. B. & Wolseley, P. A. (2007) New lichens from Thailand, mainly microlichens from Chiang Mai. *Fungal Diversity* **24**: 75–134.
- Cáceres, Marcela Eugenia da Silva (2007) Corticolous crustose and microfoliose lichens of northeastern Brazil. *Libri Botanici* **22**: 1–168.
- Ertz, D., Christnach, C., Wedin, M. & Diederich, P. (2005) A world monograph of the genus *Plectocarpon* (Roccellaceae, Arthoniales). *Bibliotheca Lichenologica* **91**: 1–155.
- Farwig, N., Böhning-Gaese, K. & Bleher, B. (2006) Enhanced seed dispersal of *Prunus africana* in fragmented and disturbed forests. *Oecologia* **147**: 238–252.
- Grube, M. (1998) Classification and phylogeny in the Arthoniales (lichenized Ascomycetes). *Bryologist* **101**: 377–391.
- Herrera-Campos, M. A. & Lücking, R. (2002) The foliicolous lichen flora of Mexico. I. New species from Los Tuxtlas Tropical Biology Station, Veracruz. *Lichenologist* **34**: 211–222.
- Hitch, C. (2008) New, rare or interesting lichens. *British Lichen Society Bulletin* **102**: 24–38.
- Jagadeesh Ram, T. A. M., Sinha, G. P., Singh, K. P. (2008) New species and new records of *Enterographa* (Roccellaceae) from India. *Lichenologist* **40**: 415–418.
- Lücking, R. (1991) Neue Arten foliikoler Flechten aus Costa Rica, Zentralamerika. *Nova Hedwigia* **52**: 267–304.
- Lücking, R. & Matzer, M. (1996) Ergänzungen und Verbesserungen zur Kenntnis der foliikolen Flechtenflora Costa Ricas. Die Familie Opegraphaceae. *Nova Hedwigia* **63**: 109–144.
- Lücking, R., Sérusiaux, E., Maia, L. C. & Pereira, C. G. (1998) A revision of the names of foliicolous lichenized fungi published by Batista and co-workers between 1960 and 1975. *Lichenologist* **30**: 121–191.
- Lücking, R., Wirth, V., Ferraro, L. I. & Cáceres, M. E. S. (2003) Foliicolous lichens from Valdivian temperate rain forest of Chile and Argentina: evidence of an austral element, with the description of seven new taxa. *Global Ecology & Biogeography* **12**: 21–36.
- Matzer, M. (1996) Lichenicolous ascomycetes with fis-situnicate asci on foliicolous lichens. *Mycological Papers* **111**: 1–202.
- Myllys, L., Lohtander, K., Källersjö, M. & Tehler, A. (1999) Applicability of ITS data in Roccellaceae (Arthoniales, Euascomycetes) phylogeny. *Lichenologist* **31**: 461–476.
- Santesson, R. (1952) Foliicolous lichens I. A revision of the taxonomy of the obligately foliicolous, lichenized fungi. *Symbolae Botanicae Upsalienses* **12**(1): 1–590.
- Sérusiaux, E. (1984) New species or interesting records of foliicolous lichens. *Mycotaxon* **20**: 283–306.
- Sparrius, L. B. (2004) A monograph of *Enterographa* and *Sclerophyton*. *Bibliotheca Lichenologica* **89**: 1–141.
- Sparrius, L. B. & Aptroot, A. (2007) A new lichenicolous *Enterographa* species from Brittany (France). *Lichenologist* **39**: 315–317.
- Sparrius, L. B. & Björk, C. R. (2008) *Enterographa oregonensis* (Roccellaceae), a new foliicolous species from the northwest coast of North America. *Bryologist* **111**: 487–489.
- Sparrius, L. B., Saipunkaew, W., Wolseley, P. A. & Aptroot, A. (2006) New species of *Bactrospora*, *Enterographa*, *Graphidastra* and *Lecanographa* from northern Thailand and Vietnam. *Lichenologist* **38**: 27–36.
- Tattersfield, P., Seddon, M. B. & Lange, C. N. (2001) Land-snail faunas in indigenous rainforest and commercial forestry plantations in Kakamega forest, western Kenya. *Biodiversity and Conservation* **10**: 1809–1829.
- Tehler, A. (1990) A new approach to the phylogeny of euascomycetes with a cladistic outline of Arthoniales focusing on Roccellaceae. *Canadian Journal of Botany* **68**: 2458–2492.
- Tehler, A., Lohtander, K., Myllys, L. & Sundin, R. (1997) On the identity of the genera *Hubbsia* and *Reinkella* (Roccellaceae). In *Lichen Studies Dedicated to Rolf Santesson* (L. Tibell, & I. Hedberg, eds): 255–265. *Symbolae Botanicae Upsalienses*, Acta Universitatis Upsaliensis, Uppsala.
- Tehler, A. & Irestedt, M. (2007) Parallel evolution of lichen growth forms in the family Roccellaceae (Arthoniales, Ascomycota). *Cladistics* **23**: 1–23.
- Vězda, A. (1975) Foliicole Flechten aus Tanzania. *Folia Geobotanica Phytotaxonomica, Praha* **10**: 383–432.

Accepted for publication 29 September 2008

