

NUMBER SYSTEMS WITH SIMPLICITY HIERARCHIES: A GENERALIZATION OF CONWAY'S THEORY OF SURREAL NUMBERS II

PHILIP EHRLICH AND ELLIOT KAPLAN

Abstract. In [16], the algebraico-tree-theoretic simplicity hierarchical structure of J. H. Conway's ordered field \mathbf{No} of surreal numbers was brought to the fore and employed to provide necessary and sufficient conditions for an ordered field to be isomorphic to an initial subfield of \mathbf{No} , i.e., a subfield of \mathbf{No} that is an initial subtree of \mathbf{No} . In this sequel to [16], analogous results for ordered abelian groups and ordered domains are established which in turn are employed to characterize the convex subgroups and convex subdomains of initial subfields of \mathbf{No} that are themselves initial. It is further shown that an initial subdomain of \mathbf{No} is discrete if and only if it is a subdomain of \mathbf{No} 's canonical integer part \mathbf{Oz} of omnific integers. Finally, making use of class models the results of [16] are extended by showing that the theories of nontrivial divisible ordered abelian groups and real-closed ordered fields are the sole theories of nontrivial densely ordered abelian groups and ordered fields all of whose models are isomorphic to initial subgroups and initial subfields of \mathbf{No} .

§1. Introduction. J. H. Conway [10] introduced a real-closed field of *surreal numbers* embracing the reals and the ordinals as well as a great many less familiar numbers including $-\omega$, $\omega/2$, $1/\omega$, $\sqrt{\omega}$, and e^ω , to name only a few. This particular real-closed field, which Conway calls \mathbf{No} , is so remarkably inclusive that, subject to the proviso that numbers—construed here as members of ordered fields—be individually definable in terms of sets of von Neumann-Bernays-Gödel set theory with global choice (NBG) [24], it may be said to contain “All Numbers Great and Small.” In this respect, \mathbf{No} bears much the same relation to ordered fields that the ordered field \mathbb{R} of real numbers bears to Archimedean ordered fields [14, 16, 20].

In addition to its inclusive structure as an ordered field, \mathbf{No} has a rich *simplicity hierarchical* (or *s-hierarchical*) *structure* [15, 16], that depends upon its structure as a *lexicographically ordered full binary tree* and arises from the fact that the sums and products of any two members of the tree are the simplest possible elements of the tree consistent with \mathbf{No} 's structure as an ordered group and an ordered field, respectively, it being understood that x is *simpler than* y just in case x is a predecessor of y in the tree.

Among the striking s-hierarchical features of \mathbf{No} that emerged from [16] is that much as the surreal numbers emerge from the empty set of surreal numbers by means of a transfinite recursion that provides an unfolding of the entire spectrum

Received January 11, 2017.

2010 *Mathematics Subject Classification.* Primary 06A05, 03C64, Secondary 12J15, 06F20, 06F25.

Key words and phrases. surreal numbers, ordered abelian groups, ordered domains, ordered fields.

© 2018, Association for Symbolic Logic
0022-4812/18/8302-0011
DOI:10.1017/jsl.2017.9

of numbers great and small (modulo the aforementioned provisos), the recursive process of defining \mathbf{No} 's arithmetic in turn provides an unfolding of the entire spectrum of ordered abelian groups (ordered fields) in such a way that an isomorphic copy of every such system either emerges as an initial subtree of \mathbf{No} or is contained in a theoretically distinguished instance of such a system that does. In particular, it was shown that every divisible ordered abelian group (real-closed ordered field) is isomorphic to an *initial subgroup (initial subfield)* of \mathbf{No} .

The divisible ordered abelian groups and real-closed ordered fields, however, do not exhaust the ordered abelian groups and ordered fields that are isomorphic to initial subgroups and subfields of \mathbf{No} . For example, every 2-divisible Archimedean ordered abelian group has an initial isomorphic copy in \mathbf{No} , as does every Archimedean ordered field [16, Theorem 8], and these groups and fields of course are not in general divisible or real-closed. In the case of ordered fields, more generally, in [16, Theorem 18] it was shown that:

An ordered field is isomorphic to an initial subfield of \mathbf{No} if and only if it is isomorphic to a truncation closed, cross sectional subfield of a power series field $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ where Γ is isomorphic to an initial subgroup of \mathbf{No} .

The present paper is a sequel to [16]. Following some preliminary material, in Sections 5 and 6 we generalize for ordered abelian groups and ordered domains the just-said result for ordered fields, and in Section 7 we employ these generalizations to characterize the convex subgroups and convex subdomains of initial subfields of \mathbf{No} that are themselves initial. We further show that an initial subdomain of \mathbf{No} is discrete if and only if it is a subdomain of \mathbf{No} 's canonical integer part \mathbf{Oz} of *omnific integers*. And in Section 8, making use of class models, we extend results of [16] by showing that the theories of nontrivial divisible ordered abelian groups and real-closed ordered fields are the *sole* theories of nontrivial densely ordered abelian groups and ordered fields *all* of whose models are isomorphic to initial subgroups and initial subfields of \mathbf{No} . Finally, in Section 9 we state a pair of open questions regarding s-hierarchical ordered algebraic systems that supplement an open question raised in Section 8 about the dispensability of the just-said reliance on class models.

Throughout the paper, the underlying set theory is assumed to be NBG and as such by *class* we mean set or proper class, the latter of which, in virtue of the axioms of global choice and foundation, always has the "cardinality" of the universe of sets. For additional information on formalizing the theory of surreal numbers in NBG, we refer the reader to [13].

The authors gratefully acknowledge the referee's helpful comments which led to a number of improvements in the exposition.

§2. Preliminaries I: Lexicographically ordered binary trees and surreal numbers.

A *tree* $\langle A, <_s \rangle$ is a partially ordered class such that for each $x \in A$, the class $\{y \in A : y <_s x\}$ of *predecessors* of x , written ' $pr_A(x)$ ', is a set well ordered by $<_s$. A maximal subclass of A well ordered by $<_s$ is called a *branch* of the tree. Two elements x and y of A are said to be *incomparable* if $x \neq y$, $x \not<_s y$, and $y \not<_s x$. An *initial subtree* of $\langle A, <_s \rangle$ is a subclass A' of A with the induced order such that for each $x \in A'$, $pr_{A'}(x) = pr_A(x)$. The *tree-rank* of $x \in A$, written ' $\rho_A(x)$ ', is the

ordinal corresponding to the well-ordered set $\langle pr_A(x), <_s \rangle$; the α th level of A is $\{x \in A : \rho_A(x) = \alpha\}$; and a root of A is a member of the zeroth level. If $x, y \in A$, then y is said to be an *immediate successor* of x if $x <_s y$ and $\rho_A(y) = \rho_A(x) + 1$; and if $(x_\alpha)_{\alpha < \beta}$ is a chain in A (i.e., a subclass of A totally ordered by $<_s$), then y is said to be an *immediate successor of the chain* if $x_\alpha <_s y$ for all $\alpha < \beta$ and $\rho_A(y)$ is the least ordinal greater than the tree-ranks of the members of the chain. The *length* of a chain $(x_\alpha)_{\alpha < \beta}$ in A is the ordinal β .

A tree $\langle A, <_s \rangle$ is said to be *binary* if each member of A has at most two immediate successors and every chain in A of limit length has at most one immediate successor. If every member of A has two immediate successors and every chain in A of limit length (including the empty chain) has an immediate successor, then the binary tree is said to be *full*. Since a full binary tree has a level for each ordinal, the universe of a full binary tree is a proper class.

Following [16, Definition 1], a binary tree $\langle A, <_s \rangle$ together with a total ordering $<$ defined on A will be said to be *lexicographically ordered* if for all $x, y \in A$, x is incomparable with y if and only if x and y have a common predecessor lying between them (i.e., there is a $z \in A$ such that $z <_s x, z <_s y$ and either $x < z < y$ or $y < z < x$). The appellation “lexicographically ordered” is motivated by the fact that: $\langle A, <, <_s \rangle$ is a lexicographically ordered binary tree if and only if $\langle A, <, <_s \rangle$ is isomorphic to an initial ordered subtree of the *lexicographically ordered canonical full binary tree* $\langle B, <_{lex(B)}, <_B \rangle$, where B is the class of all sequences of $-$ s and $+$ s indexed over some ordinal, $x <_B y$ signifies that x is a proper initial subsequence of y , and $(x_\alpha)_{\alpha < \mu} <_{lex(B)} (y_\alpha)_{\alpha < \sigma}$ if and only if $x_\beta = y_\beta$ for all $\beta < \text{some } \delta$, but $x_\delta < y_\delta$, it being understood that $- < \text{undefined} < +$ [16, Theorem 1].

NOTATION CONVENTIONS 2.1. *Let $\langle A, <, <_s \rangle$ be a lexicographically ordered binary tree. If (L, R) is a pair of subclasses of A for which every member of L precedes every member of R , then we will write ‘ $L < R$ ’. Also, if x and y are members of A , then ‘ $x <_s y$ ’ will be read “ x is simpler than y ”; and if there is an $x \in I = \{y \in A : L < \{y\} < R\}$ such that $x <_s y$ for all $y \in I - \{x\}$, then we will denote this simplest member of A lying between the members of L and the members of R by ‘ $\{L \mid R\}$ ’. Finally, by ‘ $L_{s(x)}$ ’ we mean $\{a \in A : a <_s x \text{ and } a < x\}$ and by ‘ $R_{s(x)}$ ’ we mean $\{a \in A : a <_s x \text{ and } x < a\}$.*

The following three propositions collect together a number of properties of, or results about, lexicographically ordered binary trees that will be appealed to in subsequent portions of the paper.

PROPOSITION 2.2 ([16, Theorem 2]). *Let $\langle A, <, <_s \rangle$ be a lexicographically ordered binary tree. (i) For all $x \in A, x = \{L_{s(x)} \mid R_{s(x)}\}$; (ii) for all $x, y \in A, x <_s y$ if and only if $L_{s(x)} < \{y\} < R_{s(x)}$ and $y \neq x$; (iii) for all $x \in A$ and all $L, R \subseteq A, x = \{L \mid R\}$ if and only if L is cofinal with $L_{s(x)}$ and R is coinital with $R_{s(x)}$ if and only if $L < \{x\} < R$ and $\{y \in A : L < \{y\} < R\} \subseteq \{y \in A : L_{s(x)} < \{y\} < R_{s(x)}\}$.*

Let $\langle \mathbf{No}, <, <_s \rangle$ be the *lexicographically ordered binary tree of surreal numbers* constructed in any of the manners found in the literature [11, 15–17, 20, 25], including simply letting $\langle \mathbf{No}, <, <_s \rangle = \langle B, <_{lex(B)}, <_B \rangle$.¹ Central to the development of

¹Conway’s cut construction [10] and the related construction based on Cuesta Dutari cuts introduced by the first author in [12] (and adopted in [3] and [4]), do not include \mathbf{No} ’s lexicographically ordered

the s-hierarchical theory of surreal numbers is the following result where a lexicographically ordered binary tree $\langle A, <, <_s \rangle$ is said to be *complete* [16, Definition 6], if whenever L and R are subsets of A for which $L < R$, there is an $x \in A$ such that $x = \{L \mid R\}$.

PROPOSITION 2.3 ([16, Theorem 4], [18]). *A lexicographically ordered binary tree is complete if and only if it is full if and only if it is isomorphic to $\langle \mathbf{No}, <, <_s \rangle$.*

An immediate consequence of Proposition 2.3 is

PROPOSITION 2.4. *Let $\langle A, <, <_s \rangle$ be a lexicographically ordered binary tree. $\langle A, <_s \rangle$ is full if and only if $\langle A, < \rangle$ is an $\eta_{\mathbf{On}}$ -ordering (i.e., whenever L and R are subsets of A for which $L < R$, there is an $x \in A$ such that $L < x < R$).*

§3. Preliminaries II: Conway names. Let \mathbb{D} be the set of all surreal numbers having finite tree-rank, and

$$\mathbb{R} = \mathbb{D} \cup \{ \{L \mid R\} : (L, R) \text{ is a Dedekind gap in } \mathbb{D} \}.$$

The following result regarding the structure of \mathbb{R} is essentially due to Conway [10, pages 12, 23–25].

PROPOSITION 3.1. *\mathbb{R} (with $+$, $-$, \cdot and $<$ defined à la \mathbf{No}) is isomorphic to the ordered field of real numbers defined in any of the more familiar ways, \mathbb{D} being \mathbf{No} 's ring of dyadic rationals (i.e., rationals of the form $m/2^n$ where m and n are integers); $n = \{0, \dots, n - 1 \mid \emptyset\}$ for each positive integer n , $-n = \{\emptyset \mid -(n - 1), \dots, 0\}$ for each positive integer n , $0 = \{\emptyset \mid \emptyset\}$, and the remainder of the dyadics are the arithmetic means of their left and right predecessors of greatest tree-rank; e.g., $1/2 = \{0 \mid 1\}$. The systems of natural numbers and integers so defined are henceforth denoted \mathbb{N} and \mathbb{Z} , respectively.*

\mathbf{No} 's canonical class \mathbf{On} of ordinals consists of the members of the “rightmost” branch of $\langle \mathbf{No}, <, <_s \rangle$, i.e., the unique branch of $\langle \mathbf{No}, <, <_s \rangle$ whose members satisfy the condition: $x < y$ if and only if $x <_s y$. In those formulations where surreal numbers are pairs (L, R) of sets of previously defined surreal numbers [3, 10, 20], the ordinals are the surreal numbers of the form (L, \emptyset) ; and in the formulation [21] where surreal numbers are sign-expansions (see Section 2), the ordinals are the sequences (including the empty sequence) consisting solely of $+$'s.

A striking s-hierarchical feature of \mathbf{No} is that every surreal number can be assigned a canonical “proper name” (or normal form) that is a reflection of its characteristic s-hierarchical properties. These *Conway names*, as we call them, are expressed as formal sums of the form $\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha$ where β is an ordinal, $(y_\alpha)_{\alpha < \beta}$ is a strictly decreasing sequence of surreals, and $(r_\alpha)_{\alpha < \beta}$ is a sequence of nonzero real numbers, the Conway name of an ordinal being just its Cantor normal form [10, pages 31–33], [16, Sections 3.1 and 5].

binary tree structure. However, as was noted in [15, page 257], they admit relational extensions to the ordered tree structure vis-à-vis the definition: for all $x = (L, R), y \in \mathbf{No}$, $x <_s y$ if and only if $L < \{y\} < R$ and $y \neq x$.

The identification $\langle \mathbf{No}, <, <_s \rangle = \langle B, <_{lex(B)}, <_B \rangle$, which is employed in [1, 6, 11, 16, 23], is simply a relational extension of the familiar (non tree-theoretic) construction of \mathbf{No} based on *sign-expansions*—the members of B —introduced by Conway [10, page 65], and made popular by Gonshor [21].

The surreal numbers having Conway names of the form ω^y are called *leaders* since they denote the simplest positive members of the various Archimedean classes of **No**. More formally, they may be inductively defined by formula

$$\omega^y = \left\{ 0, n\omega^{y^L} \mid \frac{1}{2^n}\omega^{y^R} \right\}, \tag{1}$$

where n ranges over the positive integers, and y^L and y^R range over the elements of $L_{s(y)}$ and $R_{s(y)}$, respectively.

There are a number of significant relations between surreal numbers that are reflected in terms of relations between their respective Conway names. The following collection of such results, which are known from the literature, will be appealed to in the subsequent discussion.

PROPOSITION 3.2 ([16, Theorems 11 and 15]). (i) For all $x, y \in \mathbf{No}$, $\omega^x <_s \omega^y$ if and only if $x <_s y$; (ii) $\sum_{\alpha < \mu} \omega^{y^\alpha} \cdot r_\alpha <_s \sum_{\alpha < \beta} \omega^{y^\alpha} \cdot r_\alpha$ whenever $\mu <_s \beta$;

$$\text{(iii) } \sum_{\alpha < \beta} \omega^{y^\alpha} \cdot r_\alpha = \left\{ \sum_{\alpha < \mu} \omega^{y^\alpha} \cdot r_\alpha + \omega^{y^\mu} \cdot \left(r_\mu - \frac{1}{2^n} \right) \mid \sum_{\alpha < \mu} \omega^{y^\alpha} \cdot r_\alpha + \omega^{y^\mu} \cdot \left(r_\mu + \frac{1}{2^n} \right) \right\},$$

if β is a limit ordinal (where n and μ range over all positive integers and all ordinals less than β , respectively).

We shall also appeal to the following compilation of results regarding Conway names which, while new to the literature, essentially consists of corollaries of results from the first author’s analysis of the surreal number tree [19] or improvements (based on that analysis) of a result of Gonshor [21, Lemma 5.8(a)].

LEMMA 3.3. Suppose $a, b, r \in \mathbb{R}$ and $x, y \in \mathbf{No}$. Then: (i) $\omega^y \cdot a <_s \omega^y \cdot b$ whenever $a <_s b$. (ii) If either $r \in \mathbb{R} - \mathbb{D}$, or $r \in \mathbb{D} - \mathbb{Z}$ and $L_{s(y)} = \emptyset$, then

$$\omega^y \cdot r = \{ \omega^y \cdot r^L \mid \omega^y \cdot r^R \};$$

moreover, in virtue of (i), $\omega^y \cdot r^L <_s \omega^y \cdot r$ and $\omega^y \cdot r^R <_s \omega^y \cdot r$ for all $r^L \in L_{s(r)}$ and $r^R \in R_{s(r)}$. (iii) If $r \in \mathbb{D} - \mathbb{Z}$ and $L_{s(y)} \neq \emptyset$, then

$$\omega^y \cdot r = \{ \omega^y \cdot r^L + \omega^{y^L} \cdot n \mid \omega^y \cdot r^R - \omega^{y^L} \cdot n \},$$

where n ranges over \mathbb{N} ; moreover, $\omega^y \cdot r^L + \omega^{y^L} \cdot n <_s \omega^y \cdot r$ and $\omega^y \cdot r^R - \omega^{y^L} \cdot n <_s \omega^y \cdot r$, for all $y^L \in L_{s(y)}$, $r^L \in L_{s(r)}$ and $r^R \in R_{s(r)}$. (iv) For all $n \in \mathbb{N}$, $\frac{1}{2^n}\omega^y <_s \omega^x$ if $y \in R_{s(x)}$.

PROOF. (i) follows immediately from [19, Theorems 3.13 and 3.16], by considering a number $c \in \mathbb{R} - \mathbb{D}$ such that $b = c$ or $b <_s c$. Necessarily, $\omega^y \cdot c$ is the immediate successor of a chain of limit length having a cofinal subchain of the form $(\omega^y \cdot c_n)_{n < \omega}$ where c is the immediate successor of $(c_n)_{n < \omega}$. As $a \in (c_n)_{n < \omega}$ and either $b \in (c_n)_{n < \omega}$ or $b = c$, it is the case that $\omega^y \cdot a \in (\omega^y \cdot c_n)_{n < \omega}$ and either $\omega^y \cdot b \in (\omega^y \cdot c_n)_{n < \omega}$ or $\omega^y \cdot b = \omega^y \cdot c$. In either case, $\omega^y \cdot a <_s \omega^y \cdot b$. For (ii), if $r \in \mathbb{R} - \mathbb{D}$, the result can be proved from (i) by simply forming the specified cut in the just-said cofinal subchain; and if $L_{s(y)} = \emptyset$ and $r \in \mathbb{D} - \mathbb{Z}$, the result follows from (i) and the second part

of Gonshor’s Lemma 5.8(a) from [21]. (iii) follows from the first part of Gonshor’s Lemma 5.8(a) from [21] and repeated applications of [19, Theorem 3.18(ii)]. For (iv), suppose without loss of generality that x is the immediate left successor of y ; for if it is not, the immediate left successor of y is in $R_{s(x)}$ and if we show that $\frac{1}{2^n}\omega^y$ is simpler than the immediate left successor of y for each n , then it follows that $\frac{1}{2^n}\omega^y$ is simpler than x as well. Consider the chain $pr_{\mathbf{No}}(\omega^x)$ of predecessors of ω^x . Clearly, this is a chain of limit length, as ω^x is the immediate successor of another surreal number if and only if $x = 0$, in which case $R_{s(x)} = \emptyset$. By [19, Theorems 3.13 and 3.16], we may conclude that $pr_{\mathbf{No}}(\omega^x)$ contains a cofinal chain of the form $(\omega^y \cdot a_n)_{n < \omega}$ where $a_n = \frac{1}{2^n}$ for each n . As this chain is contained in $pr_{\mathbf{No}}(\omega^x)$, $\frac{1}{2^n}\omega^y <_s \omega^x$ for each n . \dashv

Let $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ be the ordered group (ordered domain; ordered field) of power series (defined á la Hahn [22]) consisting of all formal power series of the form $\sum_{\alpha < \beta} r_\alpha t^{y_\alpha}$ where $(y_\alpha)_{\alpha < \beta \in \mathbf{On}}$ is a possibly empty descending sequence of elements of an ordered class (ordered commutative monoid; ordered abelian group) Γ and $r_\alpha \in \mathbb{R} - \{0\}$ for each $\alpha < \beta$. $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ is a set (often simply written $\mathbb{R}((t^\Gamma))$) if Γ is a set, and a proper class otherwise. An element $x \in \mathbb{R}((t^\Gamma))_{\mathbf{On}}$ is said to be a *proper truncation* of $\sum_{\alpha < \beta} r_\alpha t^{y_\alpha} \in \mathbb{R}((t^\Gamma))_{\mathbf{On}}$ if $x = \sum_{\alpha < \sigma} r_\alpha t^{y_\alpha}$ for some $\sigma < \beta$. A subgroup (subdomain; subfield) A of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ is said to be *truncation closed* if every proper truncation of every member of A is itself a member of A . A subgroup (subdomain; subfield) A of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ is said to be *cross sectional* if $\{t^y : y \in \Gamma\} \subseteq A$. For a truncation closed, cross sectional subgroup A of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$, the set $\mathbb{R}_y = \{r \in \mathbb{R} : rt^y \in A\}$ is an Archimedean ordered group, which we will call the *y-coefficient group* of A . If A is also a domain, then $\mathbb{R}_x \cdot \mathbb{R}_y \subseteq \mathbb{R}_{x+y}$ for all $x, y \in \Gamma$, and if A is an ordered field, then \mathbb{R}_0 is a subfield of \mathbb{R} .

PROPOSITION 3.4 ([16, 20]). *There is a canonical isomorphism of ordered groups from \mathbf{No} onto $\mathbb{R}((t^{\mathbf{No}}))_{\mathbf{On}}$ that sends each surreal number $\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha$ to $\sum_{\alpha < \beta} r_\alpha t^{y_\alpha}$. The isomorphism is in fact an isomorphism of ordered domains and, hence, of ordered fields.*

Following [16, Definition 14], a class B of surreal numbers is said to be *approximation complete* if $\sum_{\alpha < \sigma} \omega^{y_\alpha} \cdot r_\alpha \in B$ whenever $\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha \in B$ and $\sigma < \beta$. In virtue of the canonical isomorphism referred to in Proposition 3.4, approximation completeness is the analog for surreal numbers of truncation closure for formal power series.

§4. Preliminaries III: s-hierarchical ordered structures. Following [16, Definition 2], $\langle A, +, <, <_s, 0 \rangle$ is said to be an *s-hierarchical ordered group* if (i) $\langle A, +, <, 0 \rangle$ is an ordered abelian group; (ii) $\langle A, <, <_s \rangle$ is a lexicographically ordered binary tree; and (iii) for all $x, y \in A$

$$x + y = \{x^L + y, x + y^L \mid x^R + y, x + y^R\}.$$

$\langle A, +, \cdot, <, <_s, 0, 1 \rangle$ will be said to be an *s-hierarchical ordered domain* if (i) $\langle A, +, \cdot, <, 0, 1 \rangle$ is an ordered domain; (ii) $\langle A, +, <, <_s, 0 \rangle$ is an s-hierarchical ordered group; and (iii) for all $x, y \in A$

$$xy = \{x^L y + xy^L - x^L y^L, x^R y + xy^R - x^R y^R \mid x^L y + xy^R - x^L y^R, x^R y + xy^L - x^R y^L\}.$$

Moreover, $\langle A, +, \cdot, <, <_s, 0 \rangle$ will be said to be an *s-hierarchical ordered K-module* if (i) *K* is an s-hierarchical ordered domain, (ii) *A* is an s-hierarchical ordered group, and (iii) *A* is an ordered *K*-module in which for all $x \in K$ and all $y \in A$

$$xy = \{x^L y + xy^L - x^L y^L, x^R y + xy^R - x^R y^R \mid x^L y + xy^R - x^L y^R, x^R y + xy^L - x^R y^L\}.$$

s-hierarchical ordered domains and modules are generalizations of the s-hierarchical ordered fields and vector spaces introduced in [16]. In virtue of Conway’s field operations, $\langle \mathbf{No}, +, \cdot, <, <_s, 0, 1 \rangle$ is an s-hierarchical ordered domain. In fact, it is (up to isomorphism) the unique universal and unique maximal s-hierarchical ordered domain (in the sense of [16, page 1239]). Moreover, if *K* is an s-hierarchical ordered subdomain of **No**, then **No** is an s-hierarchical ordered *K*-module. Furthermore, extending the argument for s-hierarchical ordered subfields and subspaces of **No** from [16, page 1236], it is evident that

PROPOSITION 4.1. *Every initial subdomain (resp. initial submodule) of an s-hierarchical domain (resp. s-hierarchical module) is itself s-hierarchical.*

Extending the notation employed in [16], if *A* is an ordered module and $B \subseteq A$, then by $(B)_A$ we mean *the ordered submodule of A generated by B*.

The next preparatory result is a modest generalization of [16, Theorem 6].

LEMMA 4.2. *Let M' be an s-hierarchical ordered K-module and M be an initial submodule of M'. If (L, R) is a partition of M and $b = \{L \mid R\}^{M'}$, then $(M \cup \{b\})_{M'}$ is an initial submodule of M'.*

PROOF. Except for replacing the references to “*K*-vector spaces” and “subspaces” with references to “*K*-modules” and “submodules”, the proof is the same as the proof of [16, Theorem 6]. ⊖

§5. Initial subgroups and submodules of No. To fully characterize the initial subgroups of **No**, we must first characterize the initial subgroups of \mathbb{R} .

LEMMA 5.1. *An ordered group G is an initial subgroup of \mathbb{R} if and only if either $G = \{0\}$, $\mathbb{D} \subseteq G \subseteq \mathbb{R}$ or $G = \{\frac{z}{2^m} : z \in \mathbb{Z}\}$ for some $m \in \mathbb{N}$.*

PROOF. It follows from the definition of \mathbb{R} and Proposition 3.1 that the subgroups of \mathbb{R} specified in the statement of the lemma are initial. Now suppose *G* is a nontrivial initial subgroup of \mathbb{R} that does not contain \mathbb{D} . Then there is a greatest $m \in \mathbb{N}$ such that $\frac{z}{2^m} \in G$ for some odd $z \in \mathbb{Z}$. Using closure under subtraction and the fact that we must always have 1 and, therefore \mathbb{Z} in any nontrivial initial subgroup of **No**, it follows that $\frac{1}{2^m} \in G$ and, by closure under addition and subtraction, $\{\frac{z}{2^m} : z \in \mathbb{Z}\} \subseteq G$. But since *m* is the greatest member of \mathbb{N} for which $\frac{z}{2^m} \in G$ for some odd $z \in \mathbb{Z}$, the inclusion is not proper, thereby proving the lemma. ⊖

Our final preparatory result follows immediately from the definitions.

PROPOSITION 5.2. *If G is a cross sectional, truncation closed subgroup of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ and*

$$Z = \left\{ \sum_{\mu < \nu} r_\mu t^{y_\mu} \in G : \nu \text{ is an infinite limit ordinal and } r_0 = 1 \right\},$$

then $\{rt^y : y \in \Gamma, r \in \mathbb{R}_y\} \cup Z$ constitutes a class of generators for G considered as a \mathbb{Z} -module.

As is noted above, \mathbf{No} is isomorphic to $\mathbb{R}((t^{\mathbf{No}}))_{\mathbf{On}}$. We now prove more generally

THEOREM 5.3. *A subgroup of \mathbf{No} is initial if and only if it is isomorphic (via the canonical isomorphism) to a truncation closed, cross sectional subgroup G of a power series group $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$, where (i) Γ is an initial ordered subclass of \mathbf{No} , (ii) every y -coefficient group \mathbb{R}_y of G is an initial subgroup of \mathbb{R} , and (iii) $\mathbb{D} \subseteq \mathbb{R}_y$ for all $x, y \in \Gamma$ where $y \in R_{s(x)}$.*

PROOF. Let A be an initial subgroup of \mathbf{No} , $Lead(A)$ be the class of leaders in A , and $\Gamma = \{y : \omega^y \in Lead(A)\}$. Since A is initial, it follows from parts (i) and (ii) of Proposition 3.2 that A is approximation complete (see Section 3) and Γ is initial. Moreover, since A is a group, for each $y \in \Gamma$ the set $\mathbb{R}_y = \{r \in \mathbb{R} : \omega^y \cdot r \in A\}$ is a subgroup of \mathbb{R} . Furthermore, since A is initial, it follows from Lemma 3.3(i) that \mathbb{R}_y is itself initial. Now suppose $x, y \in \Gamma$ where $y \in R_{s(x)}$. In virtue of Lemma 3.3(iv), $\frac{1}{2^n} \omega^y <_s \omega^x$ for each $n \in \mathbb{N}$. Therefore, $\frac{1}{2^n} \in \mathbb{R}_y$ for each $n \in \mathbb{N}$; and thus, since groups are closed under addition and subtraction, $\mathbb{D} \subseteq \mathbb{R}_y$. Accordingly, by appealing to the restriction to A of the canonical isomorphism specified in Proposition 3.4, it is evident that $G = \{\sum_{\alpha < \beta} r_\alpha t^{y_\alpha} \in \mathbb{R}((t^\Gamma))_{\mathbf{On}} : \sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha \in A\}$ is a truncation closed, cross sectional subgroup of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ having the requisite properties listed in the statement of the theorem, thereby establishing the “only if” portion of the theorem.

Aspects of the “if” portion of the proof borrow from the first author’s proof of [16, Theorem 18]. However, whereas the latter proof concerns an ordered subfield F of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$, which is treated as an ordered vector space over the Archimedean ordered field $\{r \in \mathbb{R} : rt^0 \in F\}$, here we are concerned with an ordered subgroup G of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ that we may only assume to be an ordered \mathbb{Z} -module, which complicates the argument. To keep the argument largely self-contained, however, we repeat with modifications portions of the earlier proof.

Let G be a subgroup of $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$ satisfying the conditions specified in the statement of the theorem and let A be the isomorphic copy of G in \mathbf{No} that is the image of the restriction to G of the inverse of the canonical mapping referred to in Proposition 3.4. That is, let $A = \{\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha \in \mathbf{No} : \sum_{\alpha < \beta} r_\alpha t^{y_\alpha} \in G\}$. To show that A is an initial subgroup of \mathbf{No} , it suffices to show that $\langle A, <_s \upharpoonright A \rangle$ (where $<_s \upharpoonright A$ is the restriction of $<_s$ to A) is an initial subtree of $\langle \mathbf{No}, <_s \rangle$. We do this by induction on Γ .

Let $a_0, \dots, a_\alpha, \dots (\alpha < \beta)$ be a well-ordering of Γ such that $\rho_{\mathbf{No}}(a_\mu) \leq \rho_{\mathbf{No}}(a_\nu)$ whenever $\mu < \nu < \beta$. We consider A as an ordered \mathbb{Z} -module. Let A_α be the submodule of A containing 0 as well as all of the elements in A with exponents only from $\Gamma_\alpha = \{a_\delta : \delta \leq \alpha\}$. We see that, considered as an ordered \mathbb{Z} -module, $A = \bigcup_{\alpha < \beta} A_\alpha$. Notice also that since $a_0 = 0$, $A_0 = \mathbb{R}_0$, which, by condition (ii), is an initial subgroup of \mathbf{No} . Therefore, $\langle A_0, <_s \upharpoonright A_0 \rangle$ is an initial subtree of \mathbf{No} . To complete the proof that A is an initial subtree of \mathbf{No} , it remains to show that for all $0 < \alpha < \beta$, $\langle A_\alpha, <_s \upharpoonright A_\alpha \rangle$ is an initial subtree of \mathbf{No} , if $\langle \bigcup_{\mu < \alpha} A_\mu, <_s \upharpoonright \bigcup_{\mu < \alpha} A_\mu \rangle$ is an initial subtree of \mathbf{No} .

Accordingly, let $0 < \alpha < \beta$ and suppose $\langle \bigcup_{\mu < \alpha} A_\mu, <_s \upharpoonright \bigcup_{\mu < \alpha} A_\mu \rangle$ is an initial subtree of \mathbf{No} . Also, let $Z_\alpha =$:

$$\left\{ \sum_{\mu < \nu} \omega^{y_\mu} \cdot r_\mu \in A_\alpha - \bigcup_{\mu < \alpha} A_\mu : \nu \text{ is an infinite limit ordinal and } r_0 = 1 \right\},$$

and let $b_0, \dots, b_\sigma, \dots$ ($\sigma < \tau$) be a well-ordering of $\{\omega^{a_\alpha} \cdot r_\alpha : r_\alpha \in \mathbb{R}_{a_\alpha} - \{0\}\} \cup Z_\alpha$ such that for all $\gamma < \delta < \tau$, either:

1. $b_\gamma, b_\delta \in \{\omega^{a_\alpha} \cdot r_\alpha : r_\alpha \in \mathbb{R}_{a_\alpha} - \{0\}\}$ and $\rho_{\mathbf{No}}(b_\gamma) \leq \rho_{\mathbf{No}}(b_\delta)$;
2. $b_\gamma \in \{\omega^{a_\alpha} \cdot r_\alpha : r_\alpha \in \mathbb{R}_{a_\alpha} - \{0\}\}$ and $b_\delta \in Z_\alpha$; or
3. $b_\gamma, b_\delta \in Z_\alpha$ and the initial sequence of ordinals over which the exponents in b_γ are indexed is contained in the initial sequence of ordinals over which the exponents in b_δ are indexed.

By appealing to Proposition 5.2 (and recalling that $(X)_A$ denotes the ordered submodule of A generated by X), it is easy to see that $\{\omega^{a_\alpha} \cdot r_\alpha : r_\alpha \in \mathbb{R}_{a_\alpha} - \{0\}\} \cup Z_\alpha \cup \bigcup_{\mu < \alpha} A_\mu$ is a class of generators for A_α , and hence that, $A_\alpha = \bigcup_{\sigma < \tau} B_\sigma$, where $B_0 = \left(\{b_0\} \cup \bigcup_{\mu < \alpha} A_\mu\right)_A$ and $B_\sigma = \left(\{b_\sigma\} \cup \bigcup_{\delta < \sigma} B_\delta\right)_A$ for $0 < \sigma < \tau$. Thus to show that A_α is an initial subtree of \mathbf{No} , it suffices to show that B_σ is an initial subtree of \mathbf{No} for each $\sigma < \tau$. Moreover, since $B_\sigma = \bigcup_{\delta < \sigma} B_\delta$, whenever $b_\sigma \in \bigcup_{\delta < \sigma} B_\delta$, henceforth we need only consider those $b_\sigma \notin \bigcup_{\delta < \sigma} B_\delta$.

We proceed by transfinite induction. First, note that $b_0 = \omega^{a_\alpha}$. Moreover, since Γ is assumed to be initial, both $L_{s(a_\alpha)}$ and $R_{s(a_\alpha)} \subseteq \{a_\delta : \delta < \alpha\}$. Let a^L and a^R be typical elements of $L_{s(a_\alpha)}$ and $R_{s(a_\alpha)}$, respectively. It follows from Equation (1) (see Section 3) that

$$b_0 = \omega^{a_\alpha} = \left\{ 0, n\omega^{a^L} \mid \frac{1}{2^n}\omega^{a^R} \right\}$$

where n ranges over the positive integers. As every element of $\bigcup_{\mu < \alpha} A_\mu - \{0\}$ is Archimedean equivalent to a unique member of $\{\omega^{a_\delta} : \delta < \alpha\} \subseteq \bigcup_{\mu < \alpha} A_\mu - \{0\}$, there is a unique partition (L_α, R_α) of $\bigcup_{\mu < \alpha} A_\mu$ where $L_\alpha < R_\alpha$, $\{0, n\omega^{a^L}\}$ is cofinal with L_α and $\{\frac{1}{2^n}\omega^{a^R}\}$ is coinital with R_α . In virtue of the well ordering, each $n\omega^{a^L}$ is in $\bigcup_{\mu < \alpha} A_\mu$ as is each ω^{a^R} . Plainly then, $\{n\omega^{a^L}\} \subseteq L_\alpha$. Moreover, in virtue of the well ordering and the fact that condition (iii) in the statement of the theorem requires $\mathbb{D} \subseteq \mathbb{R}_{a^R}$ for each a^R , it follows that $\{\frac{1}{2^n}\omega^{a^R}\} \subseteq R_\alpha$. But then by Proposition 2.2, $b_0 = \{L_\alpha \mid R_\alpha\}$; and so, by Lemma 4.2, B_0 is an initial subtree of \mathbf{No} .

Now let $\sigma > 0$ and suppose B_δ is an initial subtree of \mathbf{No} for all $\delta < \sigma$. As was mentioned above, we may assume $b_\sigma \notin \bigcup_{\delta < \sigma} B_\delta$. Since $\{\omega^{a_\alpha} \cdot r_\alpha : r_\alpha \in \mathbb{Z}\} \subseteq B_0$, there are three cases that remain to consider.

CASE 1: $b_\sigma = \omega^{a_\alpha} \cdot r_\alpha$ where either $r_\alpha \in \mathbb{D} - \mathbb{Z}$ and $L_{s(a_\alpha)} = \emptyset$ or $r_\alpha \in \mathbb{R} \setminus \mathbb{D}$. By the relevant portion of Lemma 3.3(ii), we have

$$b_\sigma = \{\omega^{a_\alpha} \cdot r_\alpha^L \mid \omega^{a_\alpha} \cdot r_\alpha^R\},$$

where $\omega^{a_\alpha} \cdot r_\alpha^L, \omega^{a_\alpha} \cdot r_\alpha^R <_s b_\sigma$; and this together with the nature of our well ordering implies $\omega^{a_\alpha} \cdot r_\alpha^L, \omega^{a_\alpha} \cdot r_\alpha^R \in \bigcup_{\delta < \sigma} B_\delta$. In virtue of Proposition 2.2(iii), we have that there is a partition (L'_σ, R'_σ) of $\bigcup_{\delta < \sigma} B_\delta$ such that $b_\sigma = \{L'_\sigma \mid R'_\sigma\}$; and so, by Lemma 4.2, B_σ is an initial subtree of \mathbf{No} .

CASE 2: $b_\sigma = \omega^{a_\alpha} \cdot r_\alpha$ where $r_\alpha \in \mathbb{D} - \mathbb{Z}$ and $L_{s(a_\alpha)} \neq \emptyset$. By Lemma 3.3(iii), we have

$$b_\sigma = \{ \omega^{a_\alpha} \cdot r_\alpha^L + \omega^{a_\alpha^L} \cdot n \mid \omega^{a_\alpha} \cdot r_\alpha^R - \omega^{a_\alpha^L} \cdot n \},$$

where $\omega^y \cdot r^L + \omega^{y^L} \cdot n <_s \omega^y \cdot r$ and $\omega^y \cdot r^R - \omega^{y^L} \cdot n <_s \omega^y \cdot r$, for all $y^L \in L_{s(y)}$, $r^L \in L_{s(r)}$, $r^R \in R_{s(r)}$ and $n \in \mathbb{N}$; and, again, this together with the nature of our well ordering implies that all those options are contained in $\bigcup_{\delta < \sigma} B_\delta$. Again, by Proposition 2.2 (iii) and Lemma 4.2, B_σ is an initial subtree of **No**.

CASE 3: $b_\sigma \in Z_\alpha$. In this case, b_σ has a Conway name of the form $\sum_{\alpha < \pi} \omega^{y_\alpha} \cdot r_\alpha$, where π is an infinite limit ordinal and $r_0 = 1$. Moreover, by part (ii) of Proposition 3.2, $b_\sigma = \{L \mid R\}$ where

$$L = \left\{ \sum_{\alpha < \mu} \omega^{y_\alpha} \cdot r_\alpha + \omega^{y_\mu} \cdot \left(r_\mu - \frac{1}{2^n} \right) \right\}_{0 < n < \omega, \mu < \pi}$$

and

$$R = \left\{ \sum_{\alpha < \mu} \omega^{y_\alpha} \cdot r_\alpha + \omega^{y_\mu} \cdot \left(r_\mu + \frac{1}{2^n} \right) \right\}_{0 < n < \omega, \mu < \pi}.$$

But since, by construction, $L \cup R \subseteq \bigcup_{\delta < \sigma} B_\delta$ and $b_\sigma \notin \bigcup_{\delta < \sigma} B_\delta$, there is a partition (L'_σ, R'_σ) of $\bigcup_{\delta < \sigma} B_\delta$ such that

$$b_\sigma = \{L'_\sigma \mid R'_\sigma\}.$$

Once again by Lemma 4.2, we see that B_σ is an initial subtree of **No**.

In showing that B_σ is an initial subtree of **No** for each $\sigma < \tau$, we have shown that A_α and, hence, $\langle A, <_s \upharpoonright A \rangle$ are initial subtrees of **No**; thereby proving the theorem. \dashv

Theorem 5.3 has the virtue that it relates the initial subgroups of **No** to the important concept of a Hahn group. However, as the following immediate consequence of Theorem 5.3 makes clear, it is possible to provide an alternative characterization of the initial subgroups of **No** based solely on structures internal to **No**.

COROLLARY 5.4. *A subgroup G of **No** is initial if and only if (i) G is approximation complete; (ii) $\omega^y \in G$ whenever $\omega^y \cdot r \in G$ for some $r \in \mathbb{R} - \{0\}$; (iii) $\Gamma = \{y : \omega^y \in G\}$ is initial; (iv) for each $\omega^y \in G$, $\mathbb{R}_y = \{r \in \mathbb{R} : \omega^y \cdot r \in G\}$ is initial; (v) $\mathbb{D} \subseteq \mathbb{R}_y$ for all $x, y \in \Gamma$ where $y \in R_{s(x)}$.*

REMARK 5.5. Using the axiom of choice or global choice (if the class is a proper class), it is a routine matter to prove that every ordered class is isomorphic to an initial ordered subclass of **No**. This might seem to suggest that in the statement of Theorem 5.3 one may omit the assumption that Γ is an initial ordered subclass of **No**. However, the presence of condition (iii) precludes the omission of that statement.

5.1. Densely and discretely ordered initial subgroups of No. A nontrivial ordered group $\langle G, <, +, 0 \rangle$ is said to be *discrete or discretely ordered* if it contains a least positive member, and it is said to be *dense or densely ordered* if for all $a, b \in G$ where $a < b$ there is a $c \in G$ such that $a < c < b$. A nontrivial ordered group $\langle G, <, +, 0 \rangle$ is dense if and only if it is not discrete.

The following proposition provides a simple means of distinguishing between nontrivial initial subgroups of **No** that are discrete and those that are dense.

PROPOSITION 5.6. *A nontrivial initial subgroup G of \mathbf{No} is discrete if and only if there is a member of G of the form $\frac{1}{2^n}\omega^{-\alpha}$ (where $n \in \mathbb{N}$ and $\alpha \in On$) having no left immediate successor in G .*

PROOF. Note that elements of \mathbf{No} of the form $\frac{1}{2^n}\omega^{-\alpha}$ are precisely those having 0 as their sole left predecessor. We must show that for any $g \in G$, g is the least positive element of G if and only if $L_{s(g)} = \{0\}$ and g has no left immediate successor.

First, suppose there is a least positive $g \in G$ for which $L_{s(g)} \neq \{0\}$. If $0 \notin L_{s(g)}$, then g is not positive, a contradiction. If there is an $a \in L_{s(g)}$ where $a \neq 0$, then a is a positive element less than g , another contradiction.

Next, notice that any least positive $g \in G$ must have no immediate left successor, since, if g has a left immediate successor, this successor must be a positive element less than g .

To show the other direction, suppose G is initial. Further suppose g has no left immediate successor, $L_{s(g)} = \{0\}$ and there is an $a \in G$ where $0 < a < g$. Since G is lexicographically ordered, it follows that for any $x, y \in G$ where $x < y$, x is incomparable with y if and only if x and y have a common predecessor z such that $x < z < y$ (see Section 2). Clearly, a must be incomparable with g , as the only surreal less than g and comparable with g is zero. Therefore, they must have a common predecessor z such that $a < z < g$, but by the assumption that $L_{s(g)} = \{0\}$, z must equal 0 and a must be negative, which is impossible. \dashv

§6. Initial subdomains of \mathbf{No} . We now turn to the characterization of initial subdomains of \mathbf{No} beginning with the initial subdomains of \mathbb{R} .

LEMMA 6.1. *The initial subdomains of \mathbb{R} are \mathbb{Z} and the subdomains of \mathbb{R} containing \mathbb{D} . Every initial subdomain of \mathbf{No} is an extension of an initial subdomain of \mathbb{R} .*

PROOF. Let K be a subdomain of \mathbb{R} . If $K = \mathbb{Z}$, then by Proposition 3.1 K is initial. Now suppose $K \neq \mathbb{Z}$. If K is initial, then $\mathbb{Z} \subset K$ and, by Proposition 3.1, K must contain some element of the form $z + \frac{1}{2}$ where $z \in \mathbb{Z}$. By subtracting z , we see that $\frac{1}{2} \in K$. But since the domain \mathbb{D} is generated by $\frac{1}{2}$, $\mathbb{D} \subseteq K$. Finally, if $\mathbb{D} \subseteq K$, then K is initial since every predecessor of a member of $\mathbb{R} - \mathbb{D}$ is a member of \mathbb{D} , and \mathbb{D} is initial. The second part of the lemma is trivial. \dashv

THEOREM 6.2. *A subdomain of \mathbf{No} is initial if and only if it is isomorphic (via the canonical isomorphism) to a truncation closed, cross sectional subdomain K of a power series domain $\mathbb{R}((t^\Gamma))_{On}$, where Γ is an initial submonoid of \mathbf{No} , every y -coefficient group \mathbb{R}_y of K is an initial ordered subgroup of \mathbb{R} , and $\mathbb{D} \subseteq \mathbb{R}_y$ for any $x, y \in \Gamma$ where $y \in R_{s(x)}$.*

PROOF. First note that the above conditions are precisely the conditions for initial groups, with the exception of the stipulation that the initial subclass Γ of \mathbf{No} must be a monoid. The latter condition is necessary since: (i) $0 \in \Gamma$ insofar as Γ initial; and (ii) given that the domain K is cross sectional, it follows that for all $x, y \in \Gamma$, t^x and t^y are in K , so $t^x \cdot t^y = t^{x+y} \in K$ and, hence, $x + y \in \Gamma$. Of course, the rest of the conditions are necessary for integral domains as well, as they are necessary for abelian groups. In order to show that the conditions are also sufficient, we may treat K as a \mathbb{Z} -module and repeat the second part of the proof of Theorem 5.3. \dashv

6.1. Densely ordered initial subdomains of \mathbf{No} . An ordered domain is said to be *dense* if its ordered additive group is dense.

COROLLARY 6.3. *A densely ordered subdomain of \mathbf{No} is initial if and only if it is isomorphic (via the canonical isomorphism) to a truncation closed, cross sectional subdomain K of a power series domain $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$, where Γ is an initial submonoid of \mathbf{No} and \mathbb{D} is a subdomain of K .*

PROOF. Let K be a dense initial subdomain of \mathbf{No} . In light of Theorem 6.2, to prove the corollary it suffices to show that \mathbb{D} is a subdomain of K if and only if (i) every y -coefficient group \mathbb{R}_y of K is an initial ordered subset of \mathbb{R} , and (ii) $\mathbb{D} \subseteq \mathbb{R}_y$ for any $x, y \in \Gamma$ where $y \in R_{s(x)}$. Suppose \mathbb{D} is a subdomain of K . Then $\mathbb{D} \subseteq \mathbb{R}_0$. Moreover, since K is cross sectional, $\mathbb{R}_0 \subseteq \mathbb{R}_y$ for all $y \in \Gamma$, since $t^y \in K$ and $r \cdot t^0 \in K$ for all $r \in \mathbb{R}_0$, which implies $r \cdot t^y \in K$ for all $r \in \mathbb{R}_0$. This implies (ii) is satisfied, which along with Lemma 6.1 implies (i) is satisfied as well. Now suppose (i) and (ii) are the case. If \mathbb{D} is not a subdomain of K , then by (i) and Lemma 6.1, $\mathbb{R}_0 = \mathbb{Z}$. Moreover, since K is both dense and initial, the simplest member of \mathbf{No} lying between 0 and 1, namely $\frac{1}{2}$, is in K . But this implies \mathbb{D} is a subdomain of K , contrary to assumption. \dashv

The following result, which is a special case of Corollary 6.3, is the aforementioned result (see Section 1) categorizing the initial subfields of \mathbf{No} established in [16]. Since the special case is about ordered fields, the ordered monoid Γ must be an ordered abelian group and the reference to \mathbb{D} may be deleted since every ordered field is an extension of an isomorphic copy of \mathbb{D} , the latter of which is initial in \mathbf{No} .

COROLLARY 6.4. *A subfield of \mathbf{No} is initial if and only if it is isomorphic (via the canonical isomorphism) to a truncation closed, cross sectional subfield of a power series field $\mathbb{R}((t^\Gamma))_{\mathbf{On}}$, where Γ is an initial subgroup of \mathbf{No} .*

6.2. Discretely ordered initial subdomains of \mathbf{No} . An ordered domain is said to be *discrete* if its additive group is discrete. Accordingly, an ordered domain is discrete if and only if it is not dense. The least positive member of an ordered domain is its multiplicative identity 1.

A discrete subdomain A of an ordered field B is said to be an *integer part* if every member of B is at most a distance 1 from a member of A . Conway introduced a canonical integer part \mathbf{Oz} of \mathbf{No} consisting of the surreal numbers of the form $x = \{x - 1 \mid x + 1\}$. These *omnific integers*, as Conway calls them, are precisely the surreal numbers having Conway names of the form $\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha$ where $y_\alpha \geq 0$ for all $\alpha < \beta$, and r_α is an integer if $y_\alpha = 0$.

As we will now see, \mathbf{Oz} is in fact (up to isomorphism) the unique discrete s-hierarchical ordered domain containing an initial isomorphic copy of every discrete s-hierarchical ordered domain.

THEOREM 6.5. *An initial subdomain of \mathbf{No} is discrete if and only if it is a subdomain of \mathbf{Oz} .*

PROOF. Suppose K is an initial subdomain of \mathbf{Oz} . Since \mathbf{Oz} is discrete [19, p. 3: Note 2], K must be discrete as well. For the converse, suppose there is a discrete initial subdomain K of \mathbf{No} containing some element $a \notin \mathbf{Oz}$. Let $\sum_{\alpha < \beta} \omega^{y_\alpha} \cdot r_\alpha$ be the Conway name of a . Also let $b = \sum_{\alpha < \beta} \omega^{y'_\alpha} \cdot r'_\alpha$ where $r'_\alpha = r_\alpha$ if $y_\alpha > 0$, $r'_\alpha = 0$ if $y_\alpha < 0$ and r'_α is largest integer less than r_α if $y_\alpha = 0$. Note that $b \in \mathbf{Oz}$, so $b = \{b - 1 \mid b + 1\}$. Moreover, since $b - 1 < a < b + 1$, $b <_s a$; and so, as K is initial and $a \in K$, $b \in K$. As K is a domain, $a - b \in K$; but $0 < a - b < 1$, which contradicts the assumption that K is discrete. \dashv

§7. Initial subgroups and subdomains that are convex. Among the important subgroups and subdomains of ordered groups and fields are those that are convex. Using Theorems 5.3 and 6.2, we now identify the convex subgroups of initial subgroups of **No** that are themselves initial as well as the convex subdomains of initial subfields of **No** that are likewise initial. As we shall see, unlike the convex subgroups and subdomains of ordinary ordered groups and ordered fields, the initial convex subgroups and subdomains of initial subgroups and subfields of **No** are always well ordered by inclusion.

Let A be a nontrivial initial subgroup of **No** and $\mathbf{On}(A) =: \mathbf{On} \cap A$ be its subtree of ordinals (see Section 3). Following [16, Definition 19], A is said to be α -Archimedean if α is the height of $\mathbf{On}(A)$ considered as a subtree of A . By [16, Theorem 24], A is α -Archimedean if and only if α is the least ordinal such that for each $x \in A$ there is an ordinal $\beta < \alpha$ such that $-\beta < x < \beta$.

Ordinals of the form ω^ϕ are said to be *additively indecomposable* since they are precisely the ordinals $\lambda > 0$ such that $\mu + \nu < \lambda$ for all ordinals $\mu, \nu < \lambda$, and ordinals of the form ω^ω are said to be *multiplicatively indecomposable* since they are precisely the ordinals $\lambda > 2$ such that $\mu\nu < \lambda$ for all ordinals $\mu, \nu < \lambda$, where the just-said sums and products of ordinals are the familiar Cantorian operations. Every nontrivial initial subgroup (resp. initial subdomain) is ω^ϕ -Archimedean for some nonzero ordinal (resp. nonzero additively indecomposable ordinal) ϕ ; moreover, if A is an ordered field, then A is Archimedean if and only if A is ω -Archimedean [16, Theorem 24].

Let A be an ω^ϕ -Archimedean initial subgroup of **No** and for each nonzero ordinal $\tau \leq \phi$, let

$$A[\omega^\tau] =: \{x \in A : -\alpha < x < \alpha \text{ for some } \alpha < \omega^\tau\}.$$

PROPOSITION 7.1. *Let A be an ω^ϕ -Archimedean initial subgroup of **No** and τ be a nonzero ordinal $\leq \phi \leq \mathbf{On}$.*

- (i) $A[\omega^\tau] = A$ if and only if $\tau = \phi$.
- (ii) $A[\omega^\tau] = \{x \in A : -n\alpha < x < n\alpha \text{ for some } n \in \mathbb{N} \text{ and some } \alpha < \omega^\tau\}$.
- (iii) *The class of ordinals $< \omega^\tau$ is a cofinal subclass of $A[\omega^\tau]$.*
- (iv) *For each leader $\omega^y \in A$ there is a unique leader $\omega^\alpha \in A$ where α is an ordinal $< \phi$ such that $\omega^y \leq \omega^\alpha$ and $\omega^\alpha \leq_s \omega^y$. If $\omega^y \neq \omega^\alpha$, then ω^α is the least ordinal $> \omega^y$. Moreover, if $\omega^x <_s \omega^y$, then $\omega^x \leq \omega^\alpha$.*

PROOF. (i) is trivial, (ii) follows from the additively indecomposable nature of ω^τ , and (iii) follows from the definition of $A[\omega^\tau]$ and the initial nature of A . For (iv), note that for each $z > 0$ in **No**, there is a unique ordinal α such that $z \leq \alpha$ and $\alpha \leq_s z$; moreover, if $y \neq \alpha$, then α is the least ordinal $> y$. But by [16, Theorem 11] and [19, Proposition 3.6(i)] for all $z, y \in \mathbf{No}$, $z <_s y$ if and only if $\omega^z <_s \omega^y$, which implies the first two parts of (iv). Finally, suppose $\omega^z <_s \omega^y$. If z is an ordinal, then plainly $\omega^z \leq \omega^\alpha$; and if z is not an ordinal, then its sign-expansion begins with ω^α pluses followed by a minus, which implies it is $< \omega^\alpha$, thereby completing the proof. \dashv

THEOREM 7.2. *Let A be an ω^ϕ -Archimedean initial subgroup of **No**. Then K is a nontrivial initial convex subgroup of A if and only if $K = A[\omega^\tau]$ for some additively indecomposable infinite ordinal $\omega^\tau \leq \omega^\phi$.*

PROOF. First suppose K is a nontrivial initial convex subgroup of A . Since K is a nontrivial initial subgroup of A , K is ω^τ -Archimedean for some nonzero ordinal $\tau \leq \phi$. Moreover, since K is a convex subgroup of A , $A[\omega^\tau] \subseteq K$. Furthermore, if $A[\omega^\tau] \neq K$, there is an $x \in K$ such that $A[\omega^\tau] < \{x\} < \omega^\tau$. But then $\omega^\tau <_s x$, which implies K is not initial, since $\omega^\tau \notin K$; and so $K = A[\omega^\tau]$.

Now suppose $K = A[\omega^\tau]$ for some additively indecomposable infinite ordinal $\omega^\tau \leq \omega^\phi$. If $\tau = \phi$, then $K = A$, which is a trivial initial convex subgroup of A . Now suppose $1 \leq \tau < \phi$. In virtue of the definition of $A[\omega^\tau]$, K is a convex subclass of A . Moreover, by Proposition 7.1(ii) and the convex nature of K , every element of A that is Archimedean equivalent to some $x \in K$ is likewise in K . Plainly then, $x + y \in K$ whenever $x, y \in K$, since $x + y$ is in the Archimedean class containing ω^z , where z is the maximal member in the supports of x and the supports of y , which shows K is a group. To establish K is initial, first note that since A is cross sectional and closed under truncation, it follows from the fact that K is a convex subgroup of A (in which every element of A that is Archimedean equivalent to some $x \in K$ is likewise in K) that K is also cross sectional and closed under truncation. In addition, since K is a convex subgroup of A and A is initial, the y -coefficient groups of K satisfy conditions (ii) and (iii) of Theorem 5.3. Therefore, in virtue of Theorem 5.3, to complete the proof it remains to show $\{y \in \mathbf{No} : \omega^y \in K\}$ is an initial subtree of \mathbf{No} . For this purpose, suppose $\omega^y \in K$ and further suppose $\omega^x <_s \omega^y$. Since $x <_s y$ if and only if $\omega^x <_s \omega^y$ for all $x, y \in \mathbf{No}$, it suffices to show $\omega^x \in K$. To this end, note that by Proposition 7.1(iii), there is an ordinal $\beta \in K$ such that $\omega^y < \beta < \omega^\tau$. Moreover, by Proposition 7.1(iv), there is an ordinal ω^α such that either $\omega^\alpha = \omega^y$ or ω^α is the least ordinal $> \omega^y$ and $\omega^x \leq \omega^\alpha$. Plainly then, $\omega^x \leq \omega^\alpha \leq \beta$. And so, since K is a convex subgroup of A , $\omega^x \in K$. \dashv

THEOREM 7.3. *Let A be an ω^ϕ -Archimedean initial subfield of \mathbf{No} . Then K is an initial convex subdomain of A if and only if $K = A[\omega^\tau]$ for some multiplicatively indecomposable ordinal $\omega^\tau \leq \omega^\phi$.*

PROOF. Since A is a field, it follows that for each $\omega^y \in K$, $\{r \in \mathbb{R} : \omega^y \cdot r \in K\} = \{r \in \mathbb{R} : \omega^0 \cdot r \in K\}$ and so, for each leader $\omega^y \in K$, \mathbb{D} is a subdomain of $\{r \in \mathbb{R} : \omega^y \cdot r \in K\}$. Accordingly, in virtue of Theorems 7.2 and 6.2, to complete the proof it suffices to show that K (which is convex) is a subdomain of A if and only if $K = A[\omega^\tau]$ for some multiplicatively indecomposable ordinal $\omega^\tau \leq \omega^\phi$. If ω^τ is not multiplicatively indecomposable, there are ordinals $\alpha, \beta \in K$ where $\alpha\beta > \omega^\tau$ and, hence, $\alpha\beta \notin K$, which shows K is not a domain. Now suppose ω^τ is multiplicatively indecomposable. If $\tau = 1$, K consists of the finitely bounded members of A , in which case K is obviously a subdomain of A . Moreover, if $1 < \tau < \phi$, then $\{\omega^\beta : \beta < \tau\}$ is a cofinal subclass of K without a greatest member. Accordingly, if $x, y \in K$, $|x| \leq \text{some } \omega^\beta \in K$ and $|y| \leq \text{some } \omega^\gamma \in K$, where β, γ are ordinals $< \tau$, and so $|xy| \leq \omega^{\beta+\gamma} < \omega^\tau$. But then $\omega^{\beta+\gamma}$ and, hence, $|xy|$ are in K , which suffices to show K is a domain. \dashv

REMARK 7.4. Since every real-closed ordered field is isomorphic to an initial subfield of \mathbf{No} , it is natural to inquire if this is true for real-closed ordered domains more generally, the latter of which coincide with the convex subdomains of real-closed ordered fields [8]. However, since the convex subdomains of real-closed ordered fields are not in general well-ordered by inclusion, Theorem 7.3 implies this is not the case.

§8. Optimality results and an open question. As we mentioned above, in [16] it was shown that every divisible ordered abelian group (real-closed ordered field) is isomorphic to an initial subgroup (initial subfield) of **No**. The following result shows that in an important sense these results are optimal.

Let T^D and T^{DIV} be the theories of nontrivial densely ordered abelian groups and nontrivial divisible ordered abelian groups in the language $\{\leq, +, 0\}$ of ordered additive groups, and let T^{OF} and T^{RCF} be the theories of ordered fields and real-closed ordered fields in the language $\{\leq, +, \cdot, 0, 1\}$ of ordered fields.

THEOREM 8.1. (i) *If T is a theory in $\{\leq, +, 0\}$ containing T^D , then every model of T is isomorphic to an initial subgroup of **No** if and only if $T = T^{DIV}$.* (ii) *If T is a theory in $\{\leq, +, \cdot, 0, 1\}$ containing T^{OF} , then every model of T is isomorphic to an initial subfield of **No** if and only if $T = T^{RCF}$.*

PROOF. In light of the above-mentioned results on divisible ordered abelian groups and real-closed fields from [16], it remains to consider the cases where $T \neq T^{DIV}$ and $T \neq T^{RCF}$. Let $T^D \subseteq T \neq T^{DIV}$ where T is a theory in $\{\leq, +, 0\}$ and let M be a countable model of T . There is an elementary chain M_α ($\alpha < \mathbf{On}$) of models of T such that for each α , M_α is an $\omega_{\alpha+1}$ -saturated elementary extension of M of power 2^{ω_α} [9, Lemma 5.1.4]. Since each M_α is an $\eta_{\alpha+1}$ -ordering ([9, Page 369]; also see [5]), the union M' of the chain is a model of T that is an $\eta_{\mathbf{On}}$ -ordering (see Proposition 2.4). However, since **No** is a lexicographically ordered full binary tree and no lexicographically ordered binary tree contains a proper initial ordered subtree that is isomorphic to itself [16, Lemmas 1 and 2], it follows from Propositions 2.3 and 2.4 that an initial ordered subtree of **No** is an $\eta_{\mathbf{On}}$ -ordering if and only if it is **No** itself. But since **No** is divisible and M' is not, there is no isomorphism of M' onto an initial subgroup of **No**. Except for trivial modifications, the same argument applies to (ii), which suffices to prove the theorem. \dashv

The above proof of the “only if” portion of Theorem 8.1 makes critical use of class models. This naturally suggests:

QUESTION 8.2. *Can the “only if” portion of Theorem 8.1 be established in NBG appealing solely to models whose universes are sets?*

An ordered field K is said to be *n-real-closed* [7, page 327] if every polynomial of degree at most n admitting a root in a real closure of K admits a root in K . Boughattas [7] has shown that, for each positive integer n , there is a model of the theory of n -real-closed fields whose universe is a set that does not have an integer part. Therefore, since every initial subfield of **No** has a canonical integer part [16, Theorem 25], it follows that for each positive integer n , there is a model of the theory of n -real-closed fields whose universe is a set that is not isomorphic to an initial subfield of **No**. This, however, does not provide a positive answer to the field portion of Question 1 since there are theories of ordered fields that are not equivalent to the theory of n -real-closed fields for any n . One can also find theories of ordered abelian groups having models whose universes are sets that are not isomorphic to initial subgroups of **No**. For example, if we say that a nontrivial ordered abelian group is *n-divisible* if every element is divisible by n , then for each prime $p > 2$ the p -divisible ordered abelian group generated by 1 is not isomorphic to an initial subgroup of **No**. However, we are not aware of a proof that applies to every theory

of ordered abelian groups lacking full divisibility, which leaves the group portion of Question 8.2 open as well.

§9. Further open questions. Unlike discrete initial subdomains of **No**, discrete initial subgroups of **No** need not be subgroups of **Oz**. A case in point is the subgroup of **No** consisting of all elements of the form $d + (a/\omega)$, where $d \in \mathbb{D}$ and $a \in \mathbb{Z}$. On the other hand, this discrete initial subgroup of **No** is isomorphic to an initial subgroup of **Oz**, as is evident from the mapping $f(d + (a/\omega)) = \omega.d + a$ for all $d \in \mathbb{D}$ and all $a \in \mathbb{Z}$. This motivates

QUESTION 9.1. *Is every discrete initial subgroup of **No** isomorphic to an initial subgroup of **Oz**?*

QUESTION 9.2. *What is a set of conditions that are individually necessary and collectively sufficient for an arbitrary ordered commutative monoid to be isomorphic to an initial submonoid of **No**?*

REFERENCES

- [1] M. ASCHENBRENNER, L. VAN DEN DRIES, and J. VAN DER HOEVEN, *The surreal numbers as a universal H-field*. *Journal of the European Mathematical Society*, forthcoming.
- [2] N. ALLING, *Foundations of Analysis Over Surreal Number Fields*, North-Holland, Amsterdam, 1987.
- [3] N. ALLING and P. EHRLICH, Sections 4.02 and 4.03 of [2].
- [4] ———, *An Alternative Construction of Conway's Surreal Numbers*, *Comptes rendus mathématiques – Mathematical Reports, Academy of Science, Canada*, vol. 8 (1986), pp. 241–246.
- [5] N. ALLING and S. KUHLMANN, *On η_α -groups and fields*. *Order*, vol. 11 (1994), pp. 85–92.
- [6] A. BERARDUCCI and V. MANTOVA, *Surreal numbers, derivations and transseries*. *Journal of the European Mathematical Society*, forthcoming.
- [7] S. BOUGHATTAS, *Résultats optimaux sur l'existence d'une partie entière dans les corps ordonnés*, this JOURNAL, vol. 58 (1993), pp. 326–333.
- [8] G. CHERLIN and M. DICKMANN, *Real Closed Rings II. Model Theory*. *Annals of Pure and Applied Logic*, vol. 25 (1983), pp. 213–231.
- [9] C. C. CHANG and H. J. KEISLER, *Model Theory*, third ed., North-Holland, Amsterdam, New York, 1990; reprinted Dover Publications, Inc., Mineola, New York, 2012.
- [10] J. H. CONWAY, *On Numbers and Games*, Academic Press, London, 1976; *Second Edition*, A K Peters, Ltd., Natick, MA, 2001.
- [11] L. VAN DEN DRIES and P. EHRLICH, *Fields of Surreal Numbers and Exponentiation*. *Fundamenta Mathematicae*, vol. 167 (2001), pp. 173–188; erratum, *ibid.* 168 (2001), pp. 295–297.
- [12] P. EHRLICH, *An Alternative Construction of Conway's Ordered Field No*. *Algebra Universalis*, vol. 25 (1988), pp. 7–16; errata, *ibid.* 25 (1988), p. 233.
- [13] ———, *Absolutely Saturated Models*. *Fundamenta Mathematicae*, vol. 133 (1989), pp. 39–46.
- [14] ———, *Universally Extending Arithmetic Continua*, *Le Labyrinthe du Continu, Colloque de Cerisy* (H. Sinaceur and J.-M. Salanskis, editors), Springer-Verlag, France, Paris, 1992, pp. 168–177.
- [15] ———, *All Number Great and Small*, *Real Numbers, Generalizations of the Reals, and Theories of Continua* (P. Ehrlich, editor), Kluwer Academic Publishers, Dordrecht, Netherlands, 1994, pp. 239–258.
- [16] ———, *Number Systems with Simplicity Hierarchies: A Generalization of Conway's Theory of Surreal Numbers*, this JOURNAL, vol. 66 (2001), pp. 1231–1258.
- [17] ———, *Surreal Numbers: An Alternative Construction (Abstract)*. *Bulletin of Symbolic Logic*, vol. 8 (2002), p. 448.
- [18] ———, *Corrigendum to "Number Systems with Simplicity Hierarchies: A Generalization of Conway's Theory of Surreal Numbers"*, this JOURNAL, vol. 70 (2005), p. 1022.
- [19] ———, *Conway Names, the Simplicity Hierarchy and the Surreal Number Tree*. *Journal of Logic and Analysis*, vol. 3 (2011), no. 1, pp. 1–26.

- [20] ———. *The absolute arithmetic continuum and the unification of all numbers great and small*. *Bulletin of Symbolic Logic*, vol. 18 (2012), pp. 1–45.
- [21] H. GONSHOR, *An Introduction to the Theory of Surreal Numbers*, Cambridge University Press, Cambridge, 1986.
- [22] H. HAHN, *Über die nichtarchimedischen Grössensysteme*, *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften*, Wien, Mathematisch-Naturwissenschaftliche Klasse 116 (Abteilung IIa), 1907, pp. 601–655.
- [23] S. KUHLMANN and M. MATUSINSKI. *The Exponential-Logarithmic Equivalence Classes of Surreal Numbers*. *Order*, vol. 32 (2015), pp. 53–68.
- [24] E. MENDELSON, *An Introduction to Mathematical Logic*, fifth ed., CRC Press, Boca Raton, 2010.
- [25] D. SCHLEICHER and M. STOLL, *An introduction to Conway's games and numbers*. *Moscow Mathematical Journal*, vol. 6 (2006), pp. 359–358.

DEPARTMENT OF PHILOSOPHY
OHIO UNIVERSITY
ATHENS, OH 45701, USA
E-mail: ehrlich@ohio.edu

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN
URBANA, IL 61801, USA
E-mail: eakapla2@illinois.edu