

THE MULTILINEAR SPHERICAL MAXIMAL FUNCTION IN ONE DIMENSION

GEORGIOS DOSIDIS¹  AND JOÃO P. G. RAMOS² 

¹Department of Mathematical Analysis, Faculty of Mathematics and Physics, Charles University, Prague, Czechia

²Department of Mathematics, ETH Zürich, Zürich, Switzerland

Corresponding author: Georgios Dosidis, email: Dosidis@karlin.mff.cuni.cz

(Received 18 May 2022)

Abstract In dimension $n = 1$, we obtain $L^{p_1}(\mathbb{R}) \times \cdots \times L^{p_m}(\mathbb{R})$ to $L^p(\mathbb{R})$ boundedness for the multilinear spherical maximal function in the largest possible open set of indices and we provide counterexamples that indicate the optimality of our results.

Keywords: maximal functions; bilinear operators; sharp bounds

2020 Mathematics subject classification: 42B25; 39B72

1. Introduction

The Lebesgue differentiation theorem states that, for a function $f \in L^1_{loc}(\mathbb{R}^d)$, there is a null set $E \subset \mathbb{R}^d$ so that, if $x \in \mathbb{R}^d \setminus E$, then

$$\lim_{r \rightarrow 0} \frac{1}{|B_r(0)|} \int_{B_r(x)} f(y) dy = f(x).$$

A natural question, in that regard, is whether the same convergence holds if one replaces averages over balls by averages over *spheres*. In addition, the study of such spherical averages is deeply connected with the study of dimension-free bounds for the Hardy–Littlewood maximal function, as highlighted by Stein [31].

In this direction, such a theorem on spherical averages induces the study of the *spherical maximal function* defined by:

$$S(f)(x) := \sup_{t > 0} \left| \int_{\mathbb{S}^{n-1}} f(x - ty) d\sigma_{n-1}(y) \right|. \quad (1)$$

© The Author(s), 2024. Published by Cambridge University Press on Behalf of The Edinburgh Mathematical Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



The study of bounds for the spherical maximal function was initiated by Stein [30], who obtained its boundedness from $L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ when $n \geq 3$ and $p > \frac{n}{n-1}$ and showed that it is unbounded when $p \leq \frac{n}{n-1}$ and $n \geq 2$. The analogue of this result in dimension $n = 2$ was established later by Bourgain in [6], who also obtained a restricted weak type estimate in [5] in the case $n \geq 3$.

Further developments have been obtained by Seeger, Tao, and Wright, which, in [29], proved that the restricted weak type estimate does not hold in dimension $n = 2$. A number of other authors have also studied the spherical maximal function, among which we highlight [3, 9, 11, 25, 27, 28] and the references therein. Extensions of the spherical maximal function to different settings have also been established by several authors; for instance, see [8, 14, 20, 24].

The main object of this work is the m -linear analogue of the spherical maximal function, given by:

$$S^m(f_1, \dots, f_m)(x) := \sup_{t>0} \left| \int_{\mathbb{S}^{mn-1}} \prod_{j=1}^m f_j(x - ty^j) d\sigma_{mn-1}(y^1, \dots, y^m) \right|, \tag{2}$$

defined originally for Schwartz functions, where $d\sigma$ stands for the (normalized) surface measure of \mathbb{S}^{mn-1} .

The $m = 2$ case of (2) is called *the bi(sub)linear spherical maximal function*, and it was first introduced by Geba, Greenleaf, Iosevich, Palsson, and Sawyer [16], who obtained the first bounds for it. Later improved bounds were provided by [4, 18, 21, 22]. A multilinear (non-maximal) version of this operator when all input functions lie in the same space $L^p(\mathbb{R})$ was previously studied by Oberlin [26].

It was not until the work of Jeong and Lee [22] that the sharp open range of boundedness would be proved for the bilinear operator. Indeed, the authors proved in [22] that when $n \geq 2$, the bilinear maximal function is pointwise bounded by the product of the linear spherical maximal function and the Hardy–Littlewood maximal function, which implies boundedness in the optimal open set of exponents. This was generalized to the multilinear setting in [12]. See also [1, 2, 7, 13] for further developments.

The purpose of this work is to complement the results of [12, 22] in the $n = 1$ case. The spherical maximal operators are generally more singular when the dimension is smaller, which is reflected by the fact that the decay of the Fourier transform of the surface measure is smaller in low dimensions.

When $n \geq 2$, where the optimal boundedness range of the bilinear operator is $p > \frac{n}{2n-1}$. The optimality of the condition is found in [18] and yields the necessary condition $p > 1$ when $n = 1$. However, as was shown by Heo, Hong, and Yang in [21], when $n = 1$ the conditions $p_1, p_2 \geq 2$ are also necessary which further restricts the possible boundedness range.

Our first result establishes the $L^{p_1} \times L^{p_2} \rightarrow L^p$ boundedness of the one-dimensional bilinear operator ($m = 2$) in the region $p_1, p_2 > 2$ (see Figure 1). We show that this range is optimal via a modification of the counterexample in [21], which excludes the possibility of even a weak-type bound when p_1 or p_2 equals 2.

Theorem 1. *Let $p_1, p_2 > 1$ and $p = \frac{p_1 p_2}{p_1 + p_2}$. Then, there is a constant $C = C(p_1, p_2) < \infty$ such that:*

$$\|S^2(f, g)\|_{L^p} \leq C \|f\|_{L^{p_1}} \|g\|_{L^{p_2}}, \tag{3}$$




Figure 1. Range of $L^{p_1} \times L^{p_2} \rightarrow L^p$ boundedness of $S^2(f, g)$, when $n = 1$.

if and only if $p_1, p_2 > 2$. In this case S^2 admits a unique bounded extension from $L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R})$ to $L^p(\mathbb{R})$.

Moreover, for the end-point cases $p_1 = 2$ and $p_2 = 2$ the bilinear spherical maximal function S^2 fails to be weak type bounded. In particular, for any $1 \leq p_1, p_2 \leq \infty$, $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$, S^2 does not boundedly map $L^2 \times L^{p_2} \rightarrow L^{p, \infty}$ nor $L^{p_1} \times L^2 \rightarrow L^{p, \infty}$.

The boundedness result of Theorem 1 was also obtained independently by Christ and Zhou in [10], where the lacunary operator, with the supremum taken over the set $t \in \{2^k : k \in \mathbb{Z}\}$, is also treated.

The proof of this result is based on a decomposition of the circle into sectors, in which we may safely parametrize it. We then use the curvature of the circle in our favour, in order to show a different kind of pointwise domination with respect to the $n \geq 2$ case: instead of bounding pointwise by a product of the Hardy–Littlewood and spherical maximal functions, we obtain bounds with products of suitable p -maximal functions. In order to obtain these bounds, the curvature helps us by allowing us to insert power weights into the strategy, which effectively enable us to ‘transfer’ decay from one maximal function to the other.

Our second result deals with the multilinear case $m \geq 3$. Using the coarea formula (see [15, Theorem 3.2.22]), we see that the following pointwise bound holds, for fixed $t > 0$:

$$\begin{aligned}
 |S_t^m(f_1, \dots, f_m)(x)| &= \left| \int_{S^{m-1}} \prod_{k=1}^m f_k(x - ty_k) d\sigma(y) \right| \\
 &= \left| \int_{\mathbb{E}^{m-2}} \prod_{k=3}^m f_k(x - ty_k) \int_{r_y S^1} f_1(x - ty_1) f_2(x - ty_2) d\sigma(y_1, y_2) \frac{dy_3 \cdots dy_m}{r_y} \right| \\
 &\leq \left| \int_{\mathbb{E}^{m-2}} \prod_{k=3}^m f_k(x - ty_k) \int_{S^1} f_1(x - tr_y y_1) f_2(x - tr_y y_2) d\sigma(y_1, y_2) dy_3 \cdots dy_m \right| \\
 &\leq S^2(f_1, f_2)(x) \cdot M^{m-2}(f_3, \dots, f_m)(x),
 \end{aligned}$$

where \mathbb{B}^κ stands for the unit ball in \mathbb{R}^κ ,

$$M^m(f_1, \dots, f_m)(x) = \sup_{t>0} \int_{\mathbb{B}^m} \prod_{i=1}^m |f_i(x - ty_i)| dy_1 \cdots dy_m,$$

is the m -(sub)linear Hardy–Littlewood maximal function (first defined in [23]), and $r_y = \sqrt{1 - \sum_{k=3}^m y_k^2}$. Since M^m is pointwise bounded by the product of m Hardy–Littlewood maximal functions (denoted $Mf := M^1 f$), we arrive at the following estimates:

$$|S_t^m(f_1, \dots, f_m)(x)| \lesssim S^2(f_{i_1}, f_{i_2})(x) \prod_{j \neq i_1, i_2} M(f_j)(x), \tag{4}$$

using the fact that the operator S^m is symmetric with respect to permutations of the functions f_i . From these estimates and interpolation, we obtain $L^{p_1} \times \cdots \times L^{p_m} \rightarrow L^p$ boundedness for S^m in a certain range of exponents (see Figure 2). The range of exponents thus obtained turns out to be the optimal for the strong-type bounds. Unlike Theorem 1 our counterexamples here do not exclude the possibility of weak-type bounds on parts of the boundary; we discuss this point at the end of the section.

Theorem 2. *Let $n=1$, $m \geq 2$, $1 \leq p_i \leq \infty$ for $i = 1, \dots, m$, and $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}$. Then there is a constant $C < \infty$, only depending on p_1, \dots, p_m , such that*

$$\|S^m(f_1, \dots, f_m)\|_{L^p(\mathbb{R})} \leq C \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R})}, \tag{5}$$

for all Schwartz functions f_i , $i = 1, \dots, m$ if and only if all three of the following conditions hold:

- a) $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} < m - 1$,
- b) for every $i = 1, \dots, m$, $\sum_{j \neq i} \frac{1}{p_j} < m - \frac{3}{2}$,
- c) $(\frac{1}{p_1}, \dots, \frac{1}{p_m}) \notin \{0, 1\}^m \setminus \{(0, \dots, 0)\}$.

Additionally, if $(\frac{1}{p_1}, \dots, \frac{1}{p_m}) \in \{0, 1\}^m \setminus \{(0, \dots, 0)\}$, then we have the weak-type bound

$$\|S^m(f_1, \dots, f_m)\|_{L^{p, \infty}(\mathbb{R})} \leq C \prod_{i=1}^m \|f_i\|_{L^{p_i}(\mathbb{R})}, \tag{6}$$

for some constant $C = C(p_1, \dots, p_m)$ if and only if (a) and (b) both hold.

As an example we graph the region of boundedness for the trilinear spherical maximal function.




Figure 2. The $L^{p_1} \times L^{p_2} \times L^{p_3} \rightarrow L^p$ boundedness region of the trilinear spherical maximal operator ($n = 1$).

In order to prove the necessity of the conditions on the exponents in Theorem 2, we shall employ two different kinds of counterexamples: the first is where all functions involved are similarly concentrated around the origin, which gives us condition (a), and the second in which all but one function – at entry i – are similarly concentrated around the origin, whereas f_i is spread out; this gives us condition (b). A modification of such examples in the spirit of Stein’s original counterexample allows us to obtain condition (c) and the asserted lack of endpoint bounds.

Finally, let us mention for a brief moment the boundary case: for shortness of notation, define, for $i = 1, \dots, m$, the sets $\mathcal{H}, \mathcal{H}_i$ as:

$$\mathcal{H} := [0, 1]^m \cap \left\{ \sum_{j=1}^m x_j = m - 1 \right\} \cap \left[\bigcap_{i=1}^m \left\{ \sum_{j \neq i} x_j \leq m - \frac{3}{2} \right\} \right],$$

and

$$\mathcal{H}_i = [0, 1]^m \cap \left\{ \sum_{j \neq i} x_j = m - \frac{3}{2} \right\} \cap \left\{ \sum_{j=1}^m x_j \leq m - 1 \right\}.$$

In the diagram above, the set \mathcal{H} denotes the middle triangle in red, whereas each of the $\mathcal{H}_i, i = 1, 2, 3$, denote one of the red rectangles. In spite of Theorem 2 and the counterexamples it provides, the question of weak-type boundedness of S^m when $(\frac{1}{p_1}, \dots, \frac{1}{p_m})$ belongs in \mathcal{H} or \mathcal{H}_i remains open, as our counterexamples lie (sharply) in the corresponding Lebesgue spaces.

We would like to express our gratitude towards the anonymous referee for their helpful remarks that helped improve the exposition.

2. Boundedness of the multilinear spherical maximal function

Let $m \in \mathbb{N}$ be the index of multilinearity, and $t > 0$. Define for Schwartz functions f_1, \dots, f_m on the real line:

$$S_t^m(f_1, \dots, f_m)(x) = \int_{\mathbb{S}^{m-1}} \prod_{i=1}^m f_i(x - ty_i) d\sigma(y),$$

where \mathbb{S}^{m-1} is the unit sphere in \mathbb{R}^m , $y = (y_1, \dots, y_m) \in \mathbb{S}^{m-1}$, $y_i \in \mathbb{R}$ for $i = 1, \dots, m$, and $d\sigma$ is the (normalized) surface measure on \mathbb{S}^{m-1} . The multilinear spherical maximal operator is defined by:

$$S^m(f_1, \dots, f_m)(x) = \sup_{t>0} S_t^m(|f_1|, \dots, |f_m|)(x) = \sup_{t>0} \int_{\mathbb{S}^{m-1}} \prod_{i=1}^m |f_i(x - ty_i)| d\sigma(y).$$

Proof of Theorem 1, boundedness part. By sublinearity, we can assume without a loss of generality that $f, g \geq 0$. Fix then two indices $p_1, p_2 > 2$.

Decomposing the integral over \mathbb{S}^1 as the sum of the integrals over eight parts of the circle, we see that it is enough to deal with the integral over the set:

$$\left\{ (y_1, y_2) \in \mathbb{S}^1 : 0 \leq y_1 \leq \frac{1}{\sqrt{2}} \leq y_2 \leq 1 \right\},$$

as the treatment over the other sets is essentially equivalent. We then explicitly parametrize the circle over this arc, to obtain:

$$\begin{aligned} S_t^2(f, g)(x) &= \int_0^{1/\sqrt{2}} f(x - ty_1) g(x - t\sqrt{1 - y_1^2}) \frac{dy_1}{\sqrt{1 - y_1^2}} \\ &\leq \int_0^{1/\sqrt{2}} f(x - ty_1) g(x - t\sqrt{1 - y_1^2}) dy_1 \\ &= \int_0^{1/\sqrt{2}} f(x - ty_1) y_1^{-\frac{1-\varepsilon}{2}} g(x - t\sqrt{1 - y_1^2}) y_1^{\frac{1-\varepsilon}{2}} dy_1 \\ &\leq \left(\int_0^{1/\sqrt{2}} f^2(x - ty_1) y_1^{-1+\varepsilon} dy_1 \right)^{1/2} \left(\int_0^{1/\sqrt{2}} g^2(x - t\sqrt{1 - y_1^2}) y_1^{1-\varepsilon} dy_1 \right)^{1/2}, \end{aligned}$$

where $\varepsilon > 0$ small, to be chosen later. Since $y^{-1+\varepsilon} \chi_{0 \leq y \leq 1/\sqrt{2}} \in L^1$ and is decreasing for any $\varepsilon > 0$, the maximal function:

$$f \mapsto \sup_{t>0} \left(\int_0^{1/\sqrt{2}} f^2(x - ty_1) y_1^{-1+\varepsilon} dy_1 \right)^{1/2},$$

is bounded on L^{p_1} , since $p_1 > 2$. For the second term, we change variables by setting $z = \sqrt{1 - y_1^2}$ to get:

$$\begin{aligned} & \left(\int_0^{1/\sqrt{2}} g^2(x - t\sqrt{1 - y_1^2}) y_1^{1-\varepsilon} dy_1 \right)^{1/2} = \left(\int_{1/\sqrt{2}}^1 g^2(x - tz) (\sqrt{1 - z^2})^{-\varepsilon} z dz \right)^{1/2} \\ & \leq \left(\int_{1/\sqrt{2}}^1 g^2(x - tz) (\sqrt{1 - z^2})^{-\varepsilon} dz \right)^{1/2} \\ & \leq \left(\int_{1/\sqrt{2}}^1 g^{2q}(x - tz) dz \right)^{1/2q} \left(\int_{1/\sqrt{2}}^1 \frac{1}{\sqrt{1 - z^2}^{\varepsilon q'}} dz \right)^{1/2q'}, \end{aligned}$$

for any $1 \leq q, q' \leq \infty$ with $\frac{1}{q} + \frac{1}{q'} = 1$. We choose q sufficiently close to 1 so that $2 < 2q < p_2$ and then we choose ε to be sufficiently small so that $\varepsilon q' < 2$. In this way the second term in the above product is finite, and the maximal function:

$$g \mapsto \sup_{t>0} \left(\int_{1/\sqrt{2}}^1 g^{2q}(x - tz) dz \right)^{1/2q},$$

is bounded on L^{p_2} . Finally, taking supremum over $t > 0$ on both sides, we have:

$$\begin{aligned} & S^2(f, g)(x) \\ & \lesssim \left(\sup_{t>0} \left(\int_0^{1/\sqrt{2}} f^2(x - ty_1) y_1^{-1+\varepsilon} dy_1 \right)^{1/2} \right) \left(\sup_{t>0} \left(\int_{1/\sqrt{2}}^1 g^{2q}(x - tz) dz \right)^{1/2q} \right). \end{aligned}$$

Taking L^p norms on both sides and using Hölder’s inequality and the bounds discussed above completes the proof of Equation (3). □

Before moving on to the proof of the boundedness part of Theorem 2, we remark that the approach adopted below of using the Kolmogorov–Seliverstov–Plessner linearization and complex interpolation is by no means the only possible one; indeed, it has been brought to our attention by the anonymous referee that the results by Grafakos and Kalton in [19] may also be used to prove that part of Theorem 2.

Proof of Theorem 2, boundedness part. Again, by sublinearity, it is enough to assume that $f_i \geq 0$ for all $i = 1, \dots, m$. For $i_1, i_2 \in \{1, \dots, m\}$ we define the half-open tubes (see Figure 3):

$$T_{i_1, i_2} := \left\{ (y_1, \dots, y_m) \in [0, 1]^m : y_{i_1}, y_{i_2} < \frac{1}{2} \right\}.$$

The pointwise bound in Equation (4)

$$\sup_{t>0} |S_t^m(f_1, \dots, f_m)(x)| \lesssim S^2(f_{i_1}, f_{i_2})(x) \prod_{j \neq i_1, i_2} M(f_j)(x),$$




Figure 3. The tubes $T_{1,2}$, $T_{2,3}$ and $T_{1,3}$.

along with Theorem 1, the well-known bounds for the Hardy–Littlewood maximal function, and Hölder’s inequality yield strong-type bounds (5) for all:

$$\left(\frac{1}{p_1}, \dots, \frac{1}{p_m}\right) \in \left(T_{i_1, i_2} \setminus \{0, 1\}^m\right) \cup \{(0, \dots, 0)\},$$

and weak-type bounds (6) for

$$\left(\frac{1}{p_1}, \dots, \frac{1}{p_m}\right) \in (\{0, 1\}^m \setminus \{(0, \dots, 0)\}) \cap T_{i_1, i_2}.$$

We graph the tubes T_{i_1, i_2} in Figure 3 for $m = 3$. The multilinear Riesz–Thorin theorem [17, Corollary 7.2.11] states that, for a multilinear operator, strong-type bounds (5) on two points in $[0, 1]^m$ yield strong-type bounds on the line segment connecting them. The operator S^m is not linear, but we can use the Kolmogorov–Seliverstov–Plessner linearization (cf. [32, Chapter XIII]): Let $\tau : \mathbb{R} \rightarrow (0, \infty)$ be a measurable function and define:

$$S_\tau^m(f_1, \dots, f_m)(x) := \int_{\mathbb{S}^{m-1}} \prod_{j=1}^m f_j(x - \tau(x)y^j) d\sigma_{m-1}(y^1, \dots, y^m).$$

If S_τ^m is uniformly bounded from $L^{p_1} \times \dots \times L^{p_m}$ to L^p for all such measurable functions τ , then S^m is bounded on the same space. For any given measurable function τ , the operator S_τ^m is linear and we can thus use complex interpolation from the bounds on the T_{i_1, i_2} ’s. Since these bounds do not depend on τ , we also obtain them for S^m . Therefore, we conclude that S^m is strong-type bounded (5) for all $\left(\frac{1}{p_1}, \dots, \frac{1}{p_m}\right)$ that satisfy the conditions (a)–(c) in the statement of the theorem, which are precisely the points in the

convex hull of

$$\bigcup_{i_1, i_2} \left(T_{i_1, i_2} \setminus \{0, 1\}^m \right) \bigcup \{(0, \dots, 0)\}.$$

To confirm this, let $\frac{1}{p} = \left(\frac{1}{p_1}, \dots, \frac{1}{p_m}\right) \in [0, 1]^m$ be in the set of exponents such that (a), (b), and (c) are satisfied. If there at least 2 indices i_1, i_2 such that $p_{i_1}, p_{i_2} > 2$ then $\frac{1}{p}$ belongs in the tube T_{i_1, i_2} . If $p_i \leq 2$ for all $i = 1, \dots, m$, then the critical condition is (a) and $\frac{1}{p}$ belongs to the convex hull of $\mathcal{H} \cup \{0\}$. We interpolate the bounds on the tubes T_{i_1, i_2} to obtain strong-type bounds on the convex hull of $\mathcal{H} \cup \{0\}$, except for \mathcal{H} itself. Similarly, if $p_i > 2$ and $p_j \leq 2$ for all $j \neq i$, then the critical condition is:

$$\sum_{j \neq i} \frac{1}{p_j} < m - \frac{3}{2},$$

one of the conditions in (b). Then $\frac{1}{p}$ belongs in the convex hull of $\mathcal{H}_i \cup \{te_i, t \in (0, 1/2)\}$ and interpolation between points in

$$\bigcup_{j \neq i} T_{i, j}$$

yields strong-type bounds on this region minus \mathcal{H}_i itself. □

3. Counterexamples

Our starting point is Stein’s counterexample for the (sub)linear spherical maximal function in [30], which is the function $f(x) = |x|^{-n/p} (-\log(|x|))^{-\frac{1+\varepsilon}{p}} \chi_{|x| < 1/2}(x)$ for some $\varepsilon > 0$. Then $f \in L^p(\mathbb{R}^n)$, while $Sf \in L^p(\mathbb{R}^n)$ if and only if $p > \frac{n}{n-1}$.

For the bilinear case the authors in [18] use the same functions along with a geometric argument to ensure that in the diagonal $(y, y) \in \mathbb{R}^{2n}$, $y \in \mathbb{R}^n$, of the sphere S^{2n-1} the integral in the definition of $S^2(f_1, f_2)(x)$ is large enough to provide a counterexample. This was further expanded to the multilinear case in [12].

This counterexample is not optimal in one dimension, as was shown in [21], and a Knapp-type example further restricts the boundedness range from $p > 1$ (which the example in [18] implies) to $p_1, p_2 \geq 2$. Here we improve the example in [21], by introducing a blow-up function as in [30] in order to tackle the boundedness in the end-point cases $p_1 = 2$ and $p_2 = 2$.

Proof of Theorem 1, counterexample part. In [21] the authors showed that if the strong type bound Equation (3) holds, then $p_1, p_2 \geq 2$. Moreover, in [4] it was shown that $p = \frac{p_1 p_2}{p_1 + p_2} > 1$. A combination of the two examples shows that $p_1, p_2 > 2$ necessarily holds even for the weak type bound. We thus focus on this latter observation.

Let $g = \chi_{[-10, 10]}$ and $f(x) = |x|^{-1/2} (-\log(|x|))^{-1} \chi_{[-1/2, 1/2]}(x)$. Then $f \in L^2(\mathbb{R})$ and $g \in L^{p_2}(\mathbb{R})$ for any $p_2 \geq 1$. For any $1/4 \leq x \leq 1/2$ we choose $t = x$ in the definition of

$S^2(f, g)$ to estimate it from below by:

$$\begin{aligned} S^2(f, g)(x) &\geq \int_0^1 |x - xy|^{-1/2} (-\log(|x - xy|))^{-1} \frac{dy}{\sqrt{1 - y^2}} \\ &\geq \frac{\sqrt{x}}{\sqrt{2}} \int_0^1 (x - xy)^{-1} (-\log(x - xy))^{-1} dy \\ &\geq \frac{1}{\sqrt{2x}} \int_0^x u^{-1} (-\log(u))^{-1} du = +\infty, \end{aligned}$$

where we changed variables $u = x - xy$ in the passage from the second to the third line. Therefore $S^2(f, g)(x) = +\infty$ on a set of positive measure and the result follows for the $p_1 = 2$ case. Since the case $p_2 = 2$ is symmetric, this finishes our proof. \square

For the multilinear function we have two critical boundary cases: \mathcal{H} and \mathcal{H}_i , since all of the \mathcal{H}_i 's are similar by the symmetry of the operator. In the first case the counterexample is a characteristic function at the origin, similar to the functions in [12, 18, 30]. For the \mathcal{H}_i 's we use a Knapp-type example similar to the one in [21], a tube at the origin tangent to the sphere along the i th axis.

For the benefit of the reader, we first show that the open set of exponents is optimal. In this case characteristic functions suffice, which simplifies the computations and showcases the relevant geometry of the examples. We then include the appropriate blow-up in order to exclude strong-type bounds on \mathcal{H} and $\mathcal{H}_i, i = 1, \dots, m$.

Proof of Theorem 2, counterexample part. We start by showing that the open set of exponents in Theorem 2 is optimal.

Necessity of condition (a): If $f_1 = \dots = f_m = \chi_{[-\delta, \delta]}$, then for $1/2 \leq x \leq 1$ and $t = x\sqrt{m}$, we have:

$$\begin{aligned} S^m(f_1, \dots, f_m) &\geq \int_{\mathbb{S}^{m-1}} \prod_{i=1}^m \chi_{[-\delta, \delta]}(x(1 - \sqrt{m}y_j)) d\sigma(y_1, \dots, y_m) \\ &\geq \int_{\mathbb{S}^{m-1}} \prod_{j=1}^m \chi_{[-\frac{\delta}{x\sqrt{m}} - \frac{1}{\sqrt{m}}, \frac{\delta}{x\sqrt{m}} - \frac{1}{\sqrt{m}}]}(y_j) d\sigma(y_1, \dots, y_m) \\ &\gtrsim \int \prod_{j=1}^m \chi_{[-\frac{\delta}{x\sqrt{m}} - \frac{1}{\sqrt{m}}, \frac{\delta}{x\sqrt{m}} - \frac{1}{\sqrt{m}}]}(y_j) dy_1 \dots dy_{m-1} \gtrsim \delta^{m-1}, \end{aligned}$$

and thus, if S^m is bounded from $L^{p_1} \times \dots \times L^{p_m} \rightarrow L^p$, we should have

$$\delta^{m-1} \lesssim \|S^m(f_1, \dots, f_m)\|_{L^p} \lesssim \prod_{i=1}^m \|f_i\|_{L^{p_i}} \leq \delta^{\sum_{i=1}^m \frac{1}{p_i}},$$

and therefore $\sum_{i=1}^m \frac{1}{p_i} \leq m - 1$.

Necessity of condition (b): We set $f_1 = \chi_{[-10\sqrt{m}, 10\sqrt{m}]}$ and $f_2 = \dots = f_m = \chi_{[-\delta, \delta]}$. For $1/2 \leq x \leq 1$ we choose $t = x\sqrt{m-1}$ to estimate S^m from below. Analogously to the previous case, we have then:

$$\begin{aligned} S^m(f_1, \dots, f_m) &\geq \int_{\mathbb{S}^{m-1}} \chi_{[-10\sqrt{m}, 10\sqrt{m}]}(y_1) \prod_{j=2}^m \chi_{[-\delta, \delta]}(x(1 - \sqrt{m-1}y_j)) d\sigma(y_1, \dots, y_m) \\ &\geq \int_{\mathbb{S}^{m-1}} \prod_{j=2}^m \chi_{[-\frac{\delta}{x\sqrt{m-1}} - \frac{1}{\sqrt{m-1}}, \frac{\delta}{x\sqrt{m-1}} - \frac{1}{\sqrt{m-1}}]}(y_j) d\sigma(y_1, \dots, y_m) \\ &\gtrsim \int_{|y_1| \leq \sqrt{2\delta/x}} \left(\int \prod_{j=2}^m \chi_{[-\frac{\delta}{x\sqrt{m-1}} - \frac{1}{\sqrt{m-1}}, \frac{\delta}{x\sqrt{m-1}} - \frac{1}{\sqrt{m-1}}]}(y_j) \right. \\ &\quad \left. dy_2 \dots dy_{m-1} \right) dy_1 \\ &\gtrsim \delta^{m-3/2}. \end{aligned}$$

Thus, if S^m is bounded from $L^{p_1} \times \dots \times L^{p_m} \rightarrow L^p$,

$$\delta^{m-\frac{3}{2}} \lesssim \|S^m(f_1, \dots, f_m)\|_{L^p} \lesssim \prod_{i=1}^m \|f_i\|_{L^{p_i}} \lesssim_m \delta^{\sum_{i=2}^m \frac{1}{p_i}},$$

and therefore $\sum_{i=2}^m \frac{1}{p_i} \leq m - \frac{3}{2}$. □

We have thus showed that, in order for strong-type bounds to hold in Theorem 2, the set of exponents needs to be in the closure of the set defined by (a)–(c) in the statement of that result. With that proved, we move on to proving that the strong-type bounds *fail* also on the boundary sets \mathcal{H} and $\mathcal{H}_i, i = 1, \dots, m$.

First of all, we note the following calculus fact, which was also used in [18].

Lemma 1. *Let $r_1, r_2 > 0, t, s < e^{-\frac{r_2}{r_1}}$ and $t \leq Cs$ for some $C \geq 1$. Then, there exists an absolute constant C' (depending only on C, r_1, r_2) such that:*

$$s^{-r_1} \left(\log \frac{1}{s}\right)^{-r_2} \leq C' t^{-r_1} \left(\log \frac{1}{t}\right)^{-r_2}. \tag{7}$$

We then let $f_i = |x|^{-1/p_i} (-\log|x|)^{-2/p_i} \chi_{[-1/2, 1/2]}$ for $i = 1, \dots, m$, and note that $f_i \in L^{p_i}(\mathbb{R})$. For large $x > 0$, we choose $t = x\sqrt{m}$ to estimate $S^m(\vec{f})(x)$ from below by focusing on the region:

$$V_m(x) := \left\{ (y_1, \dots, y_m) \in \mathbb{S}^{m-1} : \left| \frac{1}{\sqrt{m}} - y_1 \right|, \dots, \left| \frac{1}{\sqrt{m}} - y_{m-1} \right| < \frac{1}{300m \cdot x\sqrt{m}} \right\}.$$

This yields the lower bound:

$$\begin{aligned}
 S^m(f_1, \dots, f_m)(x) &\geq \int_{V_m(x)} \prod_{i=1}^m f_i(x - \sqrt{m}xy_i) d\sigma(\vec{y}) \\
 &\geq \int_{V_m(x)} \prod_{i=1}^m |x - \sqrt{m}xy_i|^{-1/p_i} (-\log(|x - \sqrt{m}xy_i|))^{-2/p_i} d\sigma(\vec{y}).
 \end{aligned}$$

Notice now that, for $\vec{y} \in V_m^+(x) = \{\vec{y} \in V_m(x) : y_m > 0\}$, we have,

$$\begin{aligned}
 \left| \frac{1}{\sqrt{m}} - y_m \right| &= \frac{1}{\left| \frac{1}{\sqrt{m}} + y_m \right|} \left| \frac{1}{m} - y_m^2 \right| \leq \sqrt{m} \left| \frac{1}{m} - \left(1 - \sum_{j \leq m-1} y_j^2 \right) \right| \\
 &\leq 3 \sum_{j \leq m-1} \left| y_j - \frac{1}{\sqrt{m}} \right| < 3(m-1) \frac{1}{300m \cdot x\sqrt{m}} < \frac{1}{100x\sqrt{m}}.
 \end{aligned}$$

This in turn implies that the new variables $u_i := x - x\sqrt{m}y_i, i = 1, \dots, m$, satisfy $(\sum_{i \leq m-1} |u_i|^2)^{1/2}, |u_m| < e^{-2}$, which allows us to use Lemma 1 since $\max_{i,j} \frac{p_i}{p_j} = 2$ for indices in \mathcal{H} .

With this in mind, we locally parametrize $V_m^+(x)$ in terms of the first $(m-1)$ coordinates and use the aforementioned change of variables $\vec{y} \mapsto \tilde{u}$ in the lower bound above, noticing we are in a position to use Lemma 1, between $|u_i|$ and $|\tilde{u}|$, where $\tilde{u} := (u_1, \dots, u_{m-1})$. This implies, thus,

$$\begin{aligned}
 S^m(f_1, \dots, f_m)(x) &\geq C_m |x|^{1-m} \int_{B^{m-1}(0, \frac{1}{300m})} |\tilde{u}|^{-\frac{1}{p}} (-\log(|\tilde{u}|))^{-\frac{2}{p}} d\tilde{u} \\
 &\gtrsim \begin{cases} |x|^{1-m} & \text{if } \frac{1}{p} = m-1, \\ \infty & \text{if } \frac{1}{p} > m-1. \end{cases}
 \end{aligned}$$

This deals with the lack of strong-type bounds for the set \mathcal{H} .

We deal with the lack of strong-type bounds in each \mathcal{H}_i in a similar manner. Without loss of generality we focus on \mathcal{H}_m . Let then $f_i = |x|^{-1/p_i} (-\log|x|)^{-2/p_i} \chi_{[-1/2, 1/2]}$ for $i = 1, \dots, m-1$, and $f_m = |x|^{-1/p_m} (\log|x|)^{-2/p_m} \chi_{\mathbb{R} \setminus [-2, 2]}$. Note that $f_i \in L^{p_i}(\mathbb{R})$. For large $x > 0$, we choose $t = x\sqrt{m-1}$ to estimate $S^m(\vec{f})(x)$ from below by focusing on the region:

$$W_m(x) := \left\{ \vec{y} \in S^{m-1} : \left| \frac{1}{\sqrt{m-1}} - y_1 \right|, \dots, \left| \frac{1}{\sqrt{m-1}} - y_{m-1} \right| < \frac{10^{-4}}{mx\sqrt{m-1}} \right\},$$

over which $|1 - y_m\sqrt{m-1}| \approx 1$. Moreover, it can be seen that - by similar methods to the ones employed in the analysis of $V_m(x)$ above - for $\vec{y} \in W_m(x)$, we have $(1 - \sum_{i \leq m-1} y_i^2)^{-1/2} \geq c_m \left(\frac{x}{|\vec{v}|} \right)^{1/2}$, where $c_m > 0$ is a constant depending only on m ,

and $\tilde{v} = (v_1, \dots, v_{m-1})$, where $v_i = x - x\sqrt{m-1}y_i$. Parametrizing locally in terms of the first $(m-1)$ coordinates, changing variables $\vec{y} \mapsto \tilde{v}$ and using Lemma 1 again, we obtain:

$$\begin{aligned}
 S^m(f_1, \dots, f_m)(x) &\geq \int_{W_m(x)} \prod_{i=1}^m f_i(x - \sqrt{m-1}xy_i) d\sigma(\vec{y}) \\
 &\gtrsim x^{\frac{3}{2}-m-\frac{1}{pm}} (\log x)^{-\frac{2}{pm}} \int_{B^{m-1}(0, \frac{10^{-4}}{m})} |\tilde{v}|^{-\frac{1}{2}-\sum_{i=1}^{m-1} \frac{1}{p_i}} (-\log(|\tilde{v}|))^{-\sum_{i=1}^{m-1} \frac{2}{p_i}} d\tilde{v} \\
 &\gtrsim \begin{cases} x^{-\frac{1}{p}} (\log x)^{-\frac{2}{pm}} & \text{if } \sum_{i \leq m-1} \frac{1}{p_i} = m - \frac{3}{2}, \\ \infty & \text{if } \sum_{i \leq m-1} \frac{1}{p_i} > m - \frac{3}{2}. \end{cases}
 \end{aligned}$$

Thus, when $\sum_{i=1}^{m-1} \frac{1}{p_i} = m - \frac{3}{2}$, the above calculation shows that $S^m(\vec{f})(x) \gtrsim x^{-\frac{1}{p}} (\log \frac{1}{x})^{-\frac{2}{pm}}$ for x sufficiently large, and thus $S^m(\vec{f}) \notin L^p$, since $\frac{2p}{pm} < 1$. This completes the proof of the fact that no strong-type bounds can hold in the sets \mathcal{H}_i .

Finally, suppose that (c) is not satisfied. The counterexample in [12, Proposition 2] shows that the strong-type bound in Equation (5) cannot hold, since if, for instance, $p_1 = \dots = p_k = 1$ and $p_{k+1} = \dots = p_m = \infty$, we may take $f_1 = \dots = f_k = \chi_{(-1,1)}$ and $f_{k+1} = \dots = f_m \equiv 1$. Then, for large $x > 0$ and $t = x\sqrt{k}$:

$$\begin{aligned}
 S^m(f_1, \dots, f_m)(x) &\geq \int_{B^k(0,1)} \prod_{i=1}^k |f_i(x - x\sqrt{k}y_i)| dy_1 \dots dy_k \\
 &\gtrsim |x|^{-k},
 \end{aligned}$$

pointwise, which shows that Equation (5) cannot hold in this case. □

Funding statement. The first author was supported by the Primus research programme PRIMUS/21/SCI/002 of Charles University. The second author was supported by the European Research Council under the Grant Agreement No. 721675 ‘Regularity and Stability in Partial Differential Equations (RSPDE)’.

Competing interests. None declared.

References

- (1) T. Anderson and E. Palsson, Bounds for discrete multilinear spherical maximal functions, *Collect. Math.* **73**(1) (2022), 75–87.
- (2) T. Anderson and E. Palsson, Bounds for discrete multilinear spherical maximal functions in higher dimensions, *Bull. Lond. Math. Soc.* **53** (2021), 855–860.
- (3) J.-K. Bak, D. M. Oberlin and A. Seeger, Restriction of Fourier transforms to curves and related oscillatory integrals, *Amer. J. Math.* **131**(2) (2009), 277–311.

- (4) J. Barrionuevo, L. Grafakos, D. He, P. Honzík and L. Oliveira, Bilinear spherical maximal function, *Math. Res. Lett.* **25**(5) (2018), 1369–1388.
- (5) J. Bourgain, Estimations de certaines fonctions maximales. *C. R. Acad. Sci. Ser. I Math.* **301** (1985), 499–502.
- (6) J. Bourgain, Averages in the plane over convex curves and maximal operators, *J. Anal. Math.* **47**(1) (1986), 69–85.
- (7) T. Borges, B. Foster, Y. Ou, J. Pipher and Z. Zhou, Sparse bounds for the bilinear spherical maximal function, *J. Lond. Math. Soc.* **107**(4) (2023), 1409–1449.
- (8) C. P. Calderon, Lacunary spherical means, *Illinois J. Math.* **23**(3) (1979), 476–484.
- (9) A. Carbery, Radial Fourier multipliers and associated maximal functions, *North-Holland Math. Stud.* **111** (1985), 49–56.
- (10) M. Christ and Z. Zhou A class of singular bilinear maximal functions, (2022). Preprint, arXiv:2203.16725 [math.CA].
- (11) M. Cowling and G. Mauceri, On maximal functions, *Milan J. Math.* **49**(1) (1979), 79–87.
- (12) G. Dosidis, Multilinear spherical maximal function, *Proc. Amer. Math. Soc.* **149** (2021), 1471–1480.
- (13) G. Dosidis and L. Grafakos, On families between the Hardy–Littlewood and spherical maximal functions, *Arkiv Mat.* **59**(2) (2021), 323–343.
- (14) J. Duoandikoetxea and L. Vega, Spherical means and weighted inequalities, *J. Lond. Math. Soc.* **53**(2) (1996), 343–353.
- (15) H. Federer, *Geometric Measure Theory*, Die Grundlehren der mathematischen Wissenschaften, Band 153 (Springer-Verlag New York Inc., New York, 1969).
- (16) D. Geba, A. Greenleaf, A. Iosevich, E. Palsson and E. Sawyer, Restricted convolution inequalities, multilinear operators and applications, *Math. Res. Lett.* **20**(4) (2013), 675–694.
- (17) L. Grafakos, *Modern Fourier Analysis*, Graduate Texts in Mathematics, 3rd edn., vol. 250 (Springer, New York, 2014).
- (18) L. Grafakos, D. He and Honzík P., Maximal operators associated with bilinear multipliers of limited decay, *J. Anal. Math.* **143**(1) (2021), 231–251.
- (19) L. Grafakos and N. Kalton, Some remarks on multilinear maps and interpolation, *Math. Ann.* **319** (2001), 151–180.
- (20) A. Greenleaf, Principal curvature and harmonic-analysis, *Indiana Univ. Math. J.* **30**(4), (1981), 519–537.
- (21) Y. Heo, S. Hong and C. W. Yang, Improved bounds for the bilinear spherical maximal operators, *Math. Res. Lett.* **27**(2) (2020), 397–434.
- (22) E. Jeong and S. Lee, Maximal estimates for the bilinear spherical averages and the bilinear Bochner-Riesz, *J. Funct. Anal.* **279**(7) (2020), 108629.
- (23) A. Lerner, S. Ombrosi, C. Pérez, R. Torres and R. Trujillo-González, New maximal functions and multiple weights for the multilinear Calderón–Zygmund theory, *Adv. Math.* **220** (4) (2009), 1222–1264.
- (24) A. Magyar, E. Stein and S. Wainger, Discrete analogues in harmonic analysis: spherical averages, *Ann. Math. (2)* **155**(1) (2002), 189–208.
- (25) G. Mockenhaupt, A. Seeger and C. D. Sogge, Wave front sets, local smoothing and Bourgain’s circular maximal theorem, *Ann. Math. (2)* **136**(1) (1992), 207–218.
- (26) D. Oberlin, Multilinear convolutions defined by measures on spheres, *Trans. Amer. Math. Soc.* **310** (1988), 821–835.
- (27) J. L. Rubio de Francia, Maximal functions and Fourier transforms, *Duke Math. J.* **53**(2) (1986), 395–404.
- (28) W. Schlag, A geometric proof of the circular maximal theorem, *Duke Math. J.* **93** (1998), 505–534.

- (29) A. Seeger, T. Tao and J. Wright, Endpoint mapping properties of spherical maximal operators, *J. Inst. Math. Jussieu* **2**(1) (2003), 109–144.
- (30) E. M. Stein, Maximal functions: spherical means, *Proc. Nat. Acad. Sci.* **73**(7) (1976), 2174–2175.
- (31) E. M. Stein, The development of square functions in the work of A. Zygmund, *Bull. Amer. Math. Soc. (N. S.)* **7**(2) (1982), 359–376.
- (32) A. Zygmund, *Trigonometric Series*, 3rd edn., vols. I, II (Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2002).