

Brief Report

Cite this article: Ribeiro GG, Pessôa LR, de Abreu MDC, Corrêa LBNS, D'Avila Pereira A, Chagas MA, Brandão FZ, da Costa CAS, Boaventura GT. (2018) Taro flour (*Colocasia esculenta*) increases testosterone levels and gametogenic epithelium of *Wistar* rats. *Journal of Developmental Origins of Health and Disease* 9: 373–376. doi: 10.1017/S2040174418000120

Received: 10 November 2017
Revised: 29 January 2018
Accepted: 13 February 2018
First published online: 21 March 2018

Key words:

Colocasia esculenta; seminiferous tubules; *Wistar*

Address for correspondence:

G. G. Ribeiro, Laboratory of Experimental Nutrition, Department of Nutrition and Dietetics, College of Nutrition, Fluminense Federal University, Rua Mário Santos Braga, 30, Niterói, RJ, 24015-110, Brazil.

Taro flour (*Colocasia esculenta*) increases testosterone levels and gametogenic epithelium of *Wistar* rats

G. G. Ribeiro¹, L. R. Pessôa¹, M. D. C. de Abreu¹, L. B. N. S. Corrêa², A. D'Avila Pereira¹, M. A. Chagas², F. Z. Brandão³, C. A. S. da Costa¹ and G. T. Boaventura¹

¹The Laboratory of Experimental Nutrition, Department of Nutrition and Dietetics, College of Nutrition, Federal Fluminense University, Braga, Brazil, ²Department of Morphology, Biomedical Institute, Fluminense Federal University, Hernani Melo, Brazil and ³Laboratory of Hormonal Dosage, Faculty of Veterinary Medicine, Fluminense Federal University, Filho, Brazil

Abstract

This study evaluated the effects of diet containing taro flour on hormone levels and the seminiferous tubules morphology of rats. After weaning, the male rats were divided into two groups ($n = 12$ each): control group (CG) treated with control diet and taro group (TG), fed with 25% taro flour for 90 days. Food, caloric intake, mass and body length were evaluated at experiment end. Testis followed the standard histological processing. Immunostaining was performed using an anti-vimentin antibody to identify Sertoli cells. In histomorphometry, total diameter, total area, epithelial height, luminal height and luminal area were analyzed. The testosterone levels were performed using the radioimmunoassay method. Group TG presented ($P < 0.05$): increase in mass, body length, testicular weight, histomorphometric parameters and hormonal levels. Food intake, calorie and Sertoli cells not presented statistical differences. The taro promoted increase in the testicles parameters and hormones.

Introduction

Taro (*Colocasia esculenta* (L.) Schott) is a tropical tuber species of the Araceae family, and is widely consumed in Asia, Africa and South America.¹ This tuber is rich in starch and has several phytochemical compounds, such as polyphenols, tannins, flavonoids, alkaloids and saponins.^{1,2}

This tuber is commonly used in China as a drug because of its antioxidant and anti-inflammatory capacity and is widely used as therapeutic relief.³ In the literature, it was already reported the benefit that taro brings to women, especially with regard to postmenopausal health improvement, attenuating the symptoms relative to this stage, since it is a precursor in estrogen synthesis.^{3,4} In a study by Chiang *et al.*⁵ it was given a substrate of *Dioscorea (red mold dioscorea)* to ovariectomized rats with postmenopausal osteoporosis and the results were a decrease in estrogen-induced bone loss. Wu *et al.*⁴ observed that postmenopausal women, who received 390 g of yam (*Dioscorea alata*) for 30 days, had a higher increase of estrone, sex hormone-binding globulin and estradiol, decreased cholesterol and increased time of lipoprotein oxidation, thus decreasing the risk of cancer and cardiovascular diseases.

This performance is related to the androgenic action of these compounds present in taro. Saponin including diosgenin, has been investigated for its role in steroid production, this is because it binds to the receptors of these hormones or the enzymes that metabolize them.^{5–7} The alkaloids bind to the nervous system and modulate neurotransmitters in the gonadal tissue, there are still the hypothesis that polyphenols also have the ability to modulate neurotransmitters.⁸ It has also been found that flavonoids are capable of raising these hormone levels by serotonergic action or by preventing their metabolic degradation.⁶

Despite these reports, it has not yet been seen the performance of *Colocasia esculenta* in the hormonal levels of males and in the seminiferous tubules where this hormone is produced. So the goal of this study is to evaluate the effects of diet containing 25% of taro flour on hormone levels, seminiferous tubule morphology and number of Sertoli cells from *Wistar* rats at 90 days of age.

Materials and methods

The project was approved by the Ethics Commission on Animal Use (Ethics Committee on the Use of Animals – of the Pro-Rectorate of Research, Graduate and Innovation of the Federal Fluminense University (UFF) with registration number: 669/2015.

In the present study, rats *Wistar*, three months old, from the colonies of the Experimental Nutrition of Laboratory (LABNE) of the UFF were used. They were kept under temperature

control ($22 \pm 1^\circ\text{C}$), humidity ($60 \pm 10\%$) in an environment with artificial light–dark cycle (12–12 h).

Six nulliparous rats were used, ~3 months old. After mating, each female was relocated to individual cage with access to water *ad libitum* and commercial diet. After the pups birth litter were adjusted with six males per dams. At 21 days of lactation, the pups were weaned and randomly divided into two groups with 12 each until the 90 days of age. These two groups were divided as follows: the control group (CG), received control diet, and the experimental group, received diet containing 25% taro flour, according to Pessôa *et al.*⁹ Formulated according to the American Institute of Nutrition AIN-93G recommendations for rodent diets.¹⁰ The diet control is composed (g/100 g): 20 g casein, 52.9 g maize starch, 10 g sucrose, 7 g soy oil, 5 g fiber, 3.5 g mix mineral, 1 g mix vitamin, 0.25 g choline bitartrate, 0.3 g L-cystine, 0.0014 g tert-butylhydroquinone, 58 g carbohydrate, 7 g lipid, 17 g protein, 362 kcal energy. Diet experimental is added in 25 g taro flour (carbohydrate: 80 g/100 g; lipid: 0.4 g/100 g; protein: 7 g/100 g) and diverge in 28 g maize starch, 55 g carbohydrate, 7.4 g lipid, 17 g protein, 355 kcal energy, the other ingredients have the same amount as the control diet. The diets are similar with respect to the macronutrient content, however experimental diet content bioactive compounds present in taro.

The taros were bought at local market in Rio de Janeiro. They were washed and peeled, then cut into slices 1–3-cm thick called slats ‘chips’.¹¹ The chips were then immersed in water at 100°C and held for about 1 min in that water.¹² Thereafter, they will be frozen, and then dehydrated in a freeze dryer at 60°C until they acquire a constant weight.² After this process, the taros were made into a flour through an industrial mixer to be milled and then passed through a 35 mesh (0.5 mm mesh) sieve, where the ready meal was stored at -20°C until used.

Food intake was performed twice a week after weaning of the animals, controlling a supply of feed and rest ingestion. Body mass and length were measured at 90 days. The rats at 90 days of age will be fasted for ~8 h, then anesthetized by intraperitoneal injection of Thiopentax® at 5% (0.1 mg/100 g of body mass) being diluted with 50 ml of distilled water for each 1 g of anesthetic. After that, blood was collected by cardiac puncture. Samples were centrifuged, and serum was stored at -80°C for later analysis of testosterone (ng/dl) by radioimmunoassay method, through the Double Antibody RIA testosterone kit (MP Biomedicals, USA).

The testis were collected, weighed (g) and cleaved horizontally at about 1 cm. All material was processed following the standard laboratory protocol and then included in paraffin. The cuts were made with a thickness of 5- μm thick and then stained with hematoxylin and eosin. The images to perform the analyses were obtained under an Olympus BX-51 microscope, coupled to an Olympus video camera, the image being transferred to a LG monitor. In total, 30 photographs of two cuts of each animal were taken with a $20\times$ and $40\times$ fold objective of the rounded tubules, which were found in the spermatic cycle of 8–10.¹³ After the photos were taken, the seminiferous tubules were analyzed, with total diameter (μm), seminiferous epithelium height (μm), luminal height (μm), luminal (μm) and total (μm) area. These analyses were performed with ImageJ program with performed anti-vimentin immunohistochemistry technique to identify vimentin in Sertoli cells. All Sertoli cells were counted and, after that, divided by the number of tubules, expressed per unit.

Statistical analyses were performed using the statistical package Graph Pad Prism version 5.0, 2007 (San Diego, CA, USA). Data were analyzed by the Student's *t*-test method. All results were expressed as mean \pm S.E.M., considering the level of significance of $P < 0.05$.

Results

During the experiment, food intake and caloric intake were similar between groups. However, body mass and length and testicles mass were significantly higher ($P < 0.05$) in taro group (TG).

The interior of the testicles presented normal structure and well preserved. The organized germinative epithelium, with cells distributed in concentric layers from the base to the tubular lumen, and with well-defined tubular lumen (Fig. 1). Regarding at seminiferous tubules histomorphometric data, the total diameter, total area, seminiferous epithelium height, luminal height and luminal area showed a significant increase ($P < 0.05$) in TG (Fig. 1). The amount of Sertoli cells had no statistical differences between groups (Table 1). The levels of testosterone were significantly higher ($P < 0.05$) in TG.

Discussion

The taro is a food with high nutritional value and with great antioxidant capacity, to the 90 days, it was observed that the taro




Fig. 1. Photomicrography of the seminiferous tubules of Wistar rats (original magnification $20\times$), hematoxylin–eosin staining. (a) Group treated with a diet containing 250 g/1000 g cocoyam flour for 90 days. (b) Group treated with control diet.

Table 1. Food intake, body mass, length, histomorphometric analyses, testosterone levels and Sertoli cells

	CG (mean \pm S.E.M.)	TG (mean \pm S.E.M.)
Food intake (g/day)	54.80 \pm 3.58	63.98 \pm 3.65
Energy intake (kcal/day)	198.42 \pm 12.99	227.08 \pm 12.98
Body mass (g)	383.20 \pm 13.67	462.00 \pm 9.78*
Body length (cm)	41.72 \pm 0.33	42.92 \pm 0.25*
Testicular weight (g)	1.71 \pm 0.03	1.91 \pm 0.04*
Diameter total (μ m)	253.60 \pm 8.01	277.70 \pm 6.01*
Total area (μ m ²)	53.61 \pm 3.41	64.29 \pm 2.79*
Height epithelium (μ m)	40.89 \pm 0.82	44.21 \pm 0.95*
Luminal height (μ m)	162.00 \pm 6.65	180.90 \pm 4.33*
Luminal area (μ m ²)	21.89 \pm 1.89	27.25 \pm 1.25*
Sertoli cells (unit)	34.70 \pm 1.39	36.04 \pm 1.82
Testosterone (ng/ml)	1.60 \pm 0.49	4.27 \pm 1.03*

Control group (CG), fed with casein diet, Taro group (TG) diet containing 25 g/100 g taro flour for 90 days.

*The values are averages (Student's *t*-tests, $P < 0.05$).

had influenced the development of the seminiferous tubules. At the end of the experimental period, it was seen that food and caloric intake did not present significant differences. Body mass and length of the animals were significantly higher in the group fed with taro flour. These data differ from Chan *et al.* (2006), who found no differences in these criteria, after administration of yam flour (*Dioscorea alata*) for 90 days.¹¹ This increase in body development may be associated with elevated hormone levels, since testosterone assists muscle and bone development.⁷

Significant alterations in the testicular mass and seminiferous tubules morphology were observed in TG. About 85% of this organ is composed of the seminiferous tubules, testis weight is associated with sperm production.¹⁴ The tubule diameter is positively associated to spermatogenic activity and is in agreement with the value found for most amniotes (they are animals whose embryos are surrounded by an amniotic membrane) varying from 180 to 300 μ m,^{14,15} whereas the height of the seminiferous epithelium is related to the tubule functionality. These data were similar to those of Goel *et al.*¹⁶ which demonstrated that an extract of a plant (*Hippophae rhamnoides*) rich in flavonoids, polyphenols and tannins, resulted in increase in testicular mass, even if not significant and provided a significant increase in seminiferous tubule total diameter by 104% compared with the CG. This was also the case with an alcaloidal solution in which it promoted a significant increase in testicle weight, seminiferous tubule diameter and epithelial height. It was seen that flavonoids, polyphenols, tannins and alkaloids are capable of acting on testicular parameters exposed.^{17,18}

The number of Sertoli cells did not show statistical differences between the groups, however the TG had increase of 4%. The number of Sertoli cells correlates positively with testicular weight and with sperm production, moreover Wu *et al.*¹⁹ observed that cultures of Sertoli cells treated with diosgenin obtained a significant increase, this increase, although not significant may be due to the presence of saponin.

In this study, *Colocasia esculenta* helped increase the serum levels of testosterone and favored the seminiferous tubules

morphological alteration. Possibly, this hormonal increase may be related to gonadal performance in the pituitary stimulation of follicle stimulating hormone and luteinizing hormone, which was also suggested by Jianfeng *et al.* (2012). This author observed that in the extract of *Arctium lappa L.* rich in fitoquímicos, in the concentrations of 600 mg/kg and 1200 mg/kg promoted increase of more than 220% in testosterone concentrations and stimulated the sexual behavior of animals.⁶ In addition, Vyas and Raval (2016) observed that extract of a *Hygrophil aspinosa* plant, rich in alkaloids and saponin, in the concentration of 1000 μ g/ml promoted an increase in the levels of testosterone secreted by Leydig cells (*in vitro*) by more than 300%.¹⁸ Monet-Kuntz *et al.*²⁰ mentioned that elevations of testosterone levels lead to greater measures in testicular weights and seminiferous tubules dimensions, suggesting that testosterone exerts positive feedback on its own receptors acting as local action. It has already been seen that these actions may be due to the presence of polyphenols, flavonoids, alkaloids and saponins present in the taro.

Conclusion

Although the present study emphasizes that the consumption of taro flour, along with its compounds, are associated with higher testicular and hormonal parameters, later studies necessary to assess fertility in adult.

Acknowledgments. The authors are thankful for the College of Nutrition of the UFF.

Financial Support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest. None.

Ethical Standards. The protocol used to deal with experimental animals was approved by Ethics Committee on Animal Research of UFF, Niteroi-RJ, Brazil (protocol 669/2015). All procedures are in accordance with the provisions of Brazilian Society of Science and Laboratory's Animals and the Guide for the Care and Use of Laboratory Animals.

References

- Pereira FHF, Puiatti M, Miranda GV, *et al.* Genetic diversity in taro accessions. *Hortic Bras.* 2004; 22, 55–60.
- Chan YC, Hsu CK, Wang MF, Su TY. A diet containing yam reduces the cognitive deterioration and brain lipid peroxidation in mice with senescence accelerated. *Int J Food Sci Technol.* 2004; 39, 99–107.
- Depypere HT, Comhaire FH. Herbal preparations for the menopause: beyond isoflavones and black cohosh. *Maturitas.* 2014; 77, 191–194.
- Wu WH, Liu LY, Chung CJ, Jou HJ, Wang TA. Estrogenic effect of yam ingestion in healthy postmenopausal women. *J Am Coll Nutr.* 2005; 24, 235–243.
- Chiang SS, Chang SP, Pan TM. Osteoprotective effect of monascus-fermented dioscorea in ovariectomized rat model of postmenopausal osteoporosis. *J Agric Food Chem.* 2011; 59, 9150–9157.
- Jianfeng C, PengYing Z, ChengWei X, *et al.* Effect of aqueous extract of *Arctium lappa L.* (burdock) roots on the sexual behavior of male rats. *BMC Complement Altern Med.* 2012; 12, 1–8.
- Gauthaman K, Ganesan AP. The hormonal effects of Tribulus terrestris and its role in the management of male erectile dysfunction – an evaluation using primates, rabbit and rat. *Phytomedicine.* 2008; 15, 44–54.
- Schaffer S, Halliwell B. Do polyphenols enter the brain and does it matter? Some theoretical and practical considerations. *Genes Nutr.* 2012; 7, 99–109.
- Pessôa LR, Rêgo TS, Asht LS, *et al.* Serum and liver lipids distributions in streptozotocin induced diabetic rat treated with diet containing Yam (*Dioscorea bulbifera*) flour. *Nutr Hosp.* 2015; 31, 1647–1653.

10. Reeves PG. Components of the AIN-93 diets as improvements in the AIN-76A diet. *J Nutr.* 1997; 127, 838–841.
11. Chan YC, Hsu CK, Wang MF, *et al.* Beneficial effect of yam on the amyloid senescence β -protein, monoamine oxidase B and cognitive deficit in mice with accelerated. *J Sci Food Agric.* 2006; 86, 1517–1525.
12. Hashimoto N, Noda T, Kim SJ, *et al.* Yam contributes to improvement of glucose metabolism in rats. *Plant Foods Hum Nutr.* 2009; 64, 193–198.
13. Leblond CP, Clermont Y. Definition of the stages of the cycle of the seminiferous epithelium in the rat. *Ann N Y Acad Sci.* 1953; 55, 548–573.
14. Sinha-Hikin AP, Amador AG, Klemcke HG, *et al.* Correlative morphology and endocrinology of Sertoli cells in hamster testes in active and inactive states of spermatogenesis. *Endocrinology.* 1989; 125, 1829–1843.
15. Takashiba KS, Segatelli TM, Moraes SMF, *et al.* Testicular morphology in obese and sedentary Wistar rats submitted to physical training. *Acta Sci Health Sci.* 2011; 33, 25–33.
16. Goel HC, Samanta N, Kannan K, *et al.* Protection of spermatogenesis in mice against gamma ray induced damage by *Hippophae rhamnoides*. *Andrologia.* 2006; 38, 199–207.
17. Rafiee FM, Nejati V, Heidari R, *et al.* Protective effect of methanolic extract of *Berberis integerrima* Bunge. Root on carbon tetrachloride-induced testicular injury in Wistar rats. *Int J Reprod BioMed.* 2016; 14, 133–140.
18. Vyas NY, Raval MA. Aphrodisiac and spermatogenic potential of alkaloidal fraction of *Hygrophila spinosa* T. Ander in rats. *J Ethnopharmacol.* 2016; 194, 947–953.
19. Wu L, Dong H, Zhao J, *et al.* Diosgenin stimulates rat TM4 cell proliferation through activating plasma membrane translocation and transcriptional activity of estrogen receptors. *Biol Reprod.* 2015; 92, 1–10.
20. Monet-Kuntz C, Hochereau-de Reviers MT, Terqui M. . Variations in testicular androgen receptors and histology of the lamb testis from birth to puberty. *J Reprod Fertil.* 1984; 70, 203–210.