

# Self-harm in women with postpartum mental disorders

Benedicte Marie Johannsen<sup>1</sup>, Janne Tidselbak Larsen<sup>1,2</sup>,  
Thomas Munk Laursen<sup>1,2</sup>, Karyn Ayre<sup>3</sup>, Louise M. Howard<sup>3</sup>,  
Samantha Meltzer-Brody<sup>4</sup>, Bodil Hammer Bech<sup>5</sup> and Trine Munk-Olsen<sup>1</sup>

## Original Article

**Cite this article:** Johannsen BM, Larsen JT, Laursen TM, Ayre K, Howard LM, Meltzer-Brody S, Bech BH, Munk-Olsen T (2020). Self-harm in women with postpartum mental disorders. *Psychological Medicine* **50**, 1563–1569. <https://doi.org/10.1017/S0033291719001661>

Received: 5 December 2018

Revised: 11 June 2019

Accepted: 20 June 2019

First published online: 12 July 2019

### Key words:

Epidemiology; postpartum depression; postpartum mental disorders; self-harm; suicide; women's mental health

### Author for correspondence:

Benedicte Marie Johannsen, E-mail: [bjohannsen@econ.au.dk](mailto:bjohannsen@econ.au.dk)

<sup>1</sup>National Center for Register-based Research, Aarhus University, Fuglesangs Allé 26, 8210 Aarhus, Denmark; <sup>2</sup>CIRRAU, Centre for Integrated Register-based Research, Aarhus University, Fuglesangs Allé 26, 8210 Aarhus, Denmark; <sup>3</sup>Section of Women's Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; <sup>4</sup>Department of Psychiatry, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, USA and <sup>5</sup>Department of Public Health, Research Unit of Epidemiology, Aarhus University, Aarhus, Denmark

## Abstract

**Background.** Women suffering from first onset postpartum mental disorders (PPMD) have a highly elevated risk of suicide. The current study aimed to: (1) describe the risk of self-harm among women with PPMD and (2) investigate the extent to which self-harm is associated with later suicide.

**Methods.** We conducted a register-based cohort study linking national Danish registers. This identified women with any recorded first inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to their first child. The main outcome of interest was defined as the first hospital-registered episode of self-harm. Our cohort consisted of 1 202 292 women representing 24 053 543 person-years at risk.

**Results.** Among 1554 women with severe first onset PPMD, 64 had a first-ever hospital record of self-harm. Women with PPMD had a hazard ratio (HR) for self-harm of 6.2 (95% CI 4.9–8.0), compared to mothers without mental disorders; but self-harm risk was lower in PPMD women compared to mothers with non-PPMD [HR: 10.1, (95% CI 9.6–10.5)] and childless women with mental disorders [HR: 9.3 (95% CI 8.9–9.7)]. Women with PPMD and records of self-harm had a significantly greater risk for later suicide compared with all other groups of women in the cohort.

**Conclusions.** Women with PPMD had a high risk of self-harm, although lower than risks observed in other psychiatric patients. However, PPMD women who had self-harmed constituted a vulnerable group at significantly increased risk of later suicide.

## Background

The first 12 months after childbirth is a vulnerable time for any new mother, involving rapid biological, social, and emotional changes (Howard *et al.*, 2014; Jones *et al.*, 2014; Meltzer-Brody *et al.*, 2018). The incidence of new-onset mental disorder is high, particularly in the first 3 months postpartum (Wisner *et al.*, 2013; Jones *et al.*, 2014). Recent evidence further indicates variations in incidence, as moderate postpartum mental disorders (PPMD) (when defined as disorders treated with antidepressants and/or antipsychotics) occur in 7.72 cases per 1000 births, and moderate-severe disorders (when defined as disorders requiring psychiatric outpatient treatment) in 1.63 cases per 1000. In comparison, the most severe episodes requiring specialized inpatient psychiatric treatment, occur in 0.64 cases per 1000 births (Munk-Olsen *et al.*, 2016).

Negative mental health outcomes have been observed among women suffering from PPMD, including a concerning highly elevated risk of suicide within the first year after initial diagnosis (Appleby *et al.*, 1998; Johannsen *et al.*, 2016; Khalifeh *et al.*, 2016). This has been confirmed through inquiries into maternal deaths in the UK and other high-income countries which identified suicides as one of the leading causes of death among new mothers (Lewis, 2001; Cantwell, 2011; Thornton *et al.*, 2013).

Prior history of self-harm is an established and significant risk factor for later suicide amongst the general population (Cavanagh *et al.*, 2003; Carroll *et al.*, 2014; Runeson *et al.*, 2016; Olfson *et al.*, 2017). One small case-control study of female emergency department attendees showed that the rate of self-harm in the postpartum period was lower than expected given the high incidence of new-onset mental disorder on the postpartum period (Appleby and Turnbull, 1995). However, to our knowledge no previous study has investigated and quantified the risk of self-harm specifically among women with the first onset of PPMD, or whether self-harm is associated with later suicide in this population.

Consequently, the aims of this study were:

- To describe the risk of self-harm among women with a diagnosis of first-onset PPMD compared to other groups of women from the background population (both mothers and childless women, with and without non-postpartum onset of mental disorders).
- To investigate the extent to which self-harm in women with a diagnosis of first onset PPMD is associated with later suicide.

## Methods

### Study design and population

In order to investigate our aim, we conducted an epidemiological register-based cohort study using Danish population data.

For our main analysis, we included all women born in Denmark on 1 January 1963 or later. Follow-up started at the women's 15<sup>th</sup> birthday and was completed at date of first hospital-registered self-harm episode; date of death, emigration, or 31 December 2016, whichever came first. This allowed a maximum follow-up period of 39 years in the cohort comprising a total of 1 202 292 women.

The main exposure variable was the first onset of moderate-severe PPMD. Moderate-severe PPMD was defined as a woman experiencing any *first* inpatient or outpatient contact with a psychiatric facility within 90 days after giving birth to her first live-born child and is referred to as 'postpartum mental disorder' across this paper. In defining contacts and treatments at psychiatric treatment facilities we included all diagnostic codes in the F-chapter of mental and behavioral disorders in ICD-10 (F00–F99) and the corresponding diagnostic codes in ICD-8 (290.xx–315.xx) (Pedersen *et al.*, 2014). By defining PPMD as those occurring within a restricted interval of 3 months after giving birth, we assumed that childbirth was the specific triggering factor causing the onset of the disorder in the affected women.

The main outcome of interest was any first self-harm leading to a hospital-registered episode, as a recorded inpatient or outpatient contact of self-harm at either medical or psychiatric hospitals. This was defined by the women included in the cohort fulfilling at least *one* of the following five criteria (Helweg-Larsen *et al.*, 2005; Gasse *et al.*, 2018):

- (1) 'suicide attempt' as contact reason within the register,
- (2) an ICD-8 diagnosis within diagnostic codes: E9500–E9599,
- (3) contact with a main diagnosis within the ICD-10 F-chapter (mental disorder) and a secondary diagnosis of poisoning (ICD-10: T36–T50 and T52–T60) or lesions at the forearm, wrist, or hand (ICD-10: S51, S55, S59, S61, S65, and S69),
- (4) a main diagnosis of poisoning with weak analgesics, epileptic drugs, or carbon monoxide (ICD-10: T39, T42, T43, and T58) or
- (5) a diagnosis of suicide attempt or deliberate self-harm (ICD-10: X60–X84).

Please note, that we considered only the *first* recorded episode of self-harm from the list above.

### Data sources

Data for the study was based on a total of four national Danish registers:

- (1) The Danish Civil Registration System (CRS) (Pedersen, 2011) was initiated in 1968 and holds information on every individual in Denmark through a unique identification number (CRS number). This identification number allows linkage of information on individuals within and between registries. We included data on CRS number, gender, date of birth, vital status, data on all children and the women's parents from this register.
- (2) The Danish Psychiatric Central Register (PCR) (Mors *et al.*, 2011) includes data on all admissions to psychiatric inpatient facilities in Denmark since it was established in 1969. Information on all outpatient contacts to psychiatric facilities was added to the register from 1995. Therefore, we were able to identify information on psychiatric inpatient and outpatient contacts for the women included in the cohort. Additionally, we included data on any psychiatric contacts the women's parents might have had from 1969 and onwards, allowing us to adjust our analyses for confounding of family history of mental illness.
- (3) The Danish National Patient Register (NPR) (Lyngge *et al.*, 2011) provided data on all medical inpatient contacts to hospitals in Denmark from 1 January 1977 and onward. Information on outpatient contacts was added to the registry from 1 January 1995. This register allowed us to identify hospital records of self-harm among the women included in our cohort. Therefore, starting inclusion of women born from 1963 and onwards ensured complete follow-up information on the outcome of interest from the age of 15 years (start of follow-up) in 1978 for all women in the cohort. Furthermore, this register allowed us to draw information on parental self-harm and thereby the possibility to adjust for family history of self-harm in at least one parent.
- (4) The Danish Register of Causes of Death (Helweg-Larsen, 2011) was computerized in 1970 and provides data on all causes of death among Danish citizens dying in Denmark. This register contained the data necessary to define specifically suicides (ICD-10: X60–X84; ICD-8: E950–E959) as the cause of death among the women in our cohort as well as parental suicides. Hereby, we were able to adjust our analyses for suicide of at least one parent.

Until 31 December 1993 the diagnostic system used in Denmark was the ICD-8 (World Health Organization, 1976), and from 1 January 1994 the ICD-10 system was introduced (World Health Organization, 2016).

### Definition of reference and comparison groups

For our main analysis, the defined reference group consisted of mothers from the healthy background population, giving birth to their first child with no records of a psychiatric diagnosis either before or after childbirth.

In additional analyses, we compared the risk of self-harm among women with PPMD to other groups of women with varying status of mental health and motherhood. This addition of further comparison groups was done to examine how mental health and motherhood influences the risk of both self-harm and suicide (Appleby and Turnbull, 1995; Johannsen *et al.*, 2016). Consequently, all women in the cohort were classified into a total of five groups in a time-dependent variable depending on their mental health and parity status throughout the follow-up period:

- (1) Mothers with PPMD. This was defined as any first inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to their first child.
- (2) Mothers without mental disorders (reference group). This consisted of women with no registered history of contact to a psychiatric facility and who have given birth to at least one child.
- (3) Mothers with a diagnosis of mental disorder occurring outside the postpartum period. This was defined as women who have given birth to their first child and have a first inpatient or outpatient psychiatric contact outside the 90-day postpartum period (either before or after).
- (4) Childless women with mental disorders. This consisted of women with any first inpatient or outpatient contact to a psychiatric facility without any record of childbirth.
- (5) Childless women without mental disorders. This consisted of women without any record of contact to a psychiatric facility and without any record of childbirth.

When defining group 1 (mothers with PPMD) as the first-ever inpatient or outpatient contact within the short time-period of 90 days after giving birth, we assumed that childbirth was the specific trigger causing the psychiatric episode (Munk-Olsen *et al.*, 2006; Munk-Olsen *et al.*, 2014).

Please note that implementation of these five time-dependent variables enabled dynamic classifications where the women could transition between groups and contribute person-time to different comparison groups throughout the course of follow-up-time depending on when and if they had their first psychiatric contact and its possible relation to the potential birth of a first-born child (for further description see online Supplementary Appendix 1). Furthermore, our analysis built on the assumption that the specific dates of first psychiatric inpatient or outpatient contact and childbirth mark a change in group membership category. As a consequence, any potential prodromal phase leading up to the onset of mental disorders are not taken in to account, as is the case with any study design including time-dependent co-variables. Further, we conducted a sensitivity analysis of our main analysis expanding the onset of PPMD from the first 3 months (90 days) after giving birth to 6 months postpartum (180 days).

### Statistical analysis

We conducted survival analyses with hazard ratios (HRs) as the main outcome measure, using Cox regression in Stata, version 13 (StataCorp, College Station, Tex.). Age was chosen as the underlying time in all analyses as is customary in these types of studies with register-based Cox regression. Further, all analyses were adjusted for calendar time as a time-dependent variable. Further, we adjusted for the time-dependent variables of family history of mental disorder, parental self-harm and parental suicide defined as any psychiatric inpatient or outpatient contact, any hospital recorded episode of self-harm or death categorized as suicide among either of the women's parents.

### Analysis of short-term v. long-term risk of self-harm

In order to specifically study the short v. the long-term risk of self-harm following a diagnosis of PPMD, we conducted a sub-analysis calculating specific HRs for the risk of receiving a first-ever hospital record for self-harm up to 12 months after first inpatient or outpatient contact to a psychiatric facility as well as first-ever

self-harm episode beyond the first 12 months after diagnosis. These HRs were compared to the risk of first-ever self-harm episode of mothers without any record of self-harm in the entire time-period from 1963 to 2016.

### Analysis of suicide risk following a self-harm episode

In order to investigate our secondary aim of describing the risk of suicide among those with a history of self-harm, we conducted a sub-analysis on the sub-cohort of women registered with an episode of self-harm throughout the follow-up period. Here each woman was followed from the date of their first hospital-registered self-harm until the date of suicide, date of death by other causes or December 2016, whichever came first.

### Results

A total of 1 202 292 Danish women were included in our study cohort representing 24 053 543 person-years at risk throughout the entire follow-up period. We identified 1554 mothers with first onset PPMD (Table 1) i.e. experiencing first psychiatric outpatient or inpatient contact to a psychiatric facility within 90 days after giving birth to their first child. Among the mothers with PPMD, we identified 64 women who had a first-ever hospital record of self-harm following their diagnosis.

Among the 64 women with first onset PPMD and a hospital record of self-harm following their diagnosis, the majority were diagnosed with anxiety or stress-related mental disorders ( $n = 20$ , 31.3%) or affective mood disorders ( $n = 18$ , 28.1%) in the postpartum period. Six (9.4%) women had a schizophrenia or schizotypal disorder, and five (7.81%) women had diagnoses of mental and behavioral disorders associated with the puerperium. The remaining 15 (23.43%) women were recorded having other diagnoses including personality disorders.

Table 1 further shows the characteristics of the cohort in terms of age-distribution, parental history of mental illness, self-harm and/or suicide presented as person-years at risk and number of self-harm cases.

### Risk of self-harm

The adjusted HR of self-harm was 6.2 (95% CI 4.9–8.0) i.e. the risk was higher in women with first onset PPMD, compared to mothers from the healthy general population without mental disorders. However, when compared to both childless women and mothers with a mental disorder outside the postpartum period, women with PPMD had a lower risk of self-harm. More precisely, the risks for self-harm were: HR 10.1 (95% CI 9.6–10.5) for mothers with mental disorders occurring outside the postpartum period and 9.3 (95% CI 8.9–9.7) in childless women with mental disorders (see Table 2).

In our sensitivity analysis with the onset of PPMD expanded to 6 months (180 days) after childbirth, we found almost identical results (see online Supplementary Appendix 2).

### Timing of self-harm after diagnosis

Table 3 shows the short-term and long-term risk of self-harm among women with PPMD compared to the lifetime risk of self-harm in mothers from the background population without mental disorders. Overall, risk of self-harm was highest in close proximity to the time of diagnoses – the HR of self-harm within the first 12

**Table 1.** Characteristics of the cohort presented with person-years at risk and number of self-harm cases

	Person-years at risk	N (Self-harm cases)
<b>Exposure groups</b>		
Women with postpartum mental disorders <sup>a</sup> N = 1554	17 097	64
Mothers without mental disorders (reference) <sup>b</sup> N = 575 360	8 919 834	4950
Mothers with diagnosis of mental disorders outside the postpartum period <sup>c</sup> N = 80 622	675 005	4039
Childless women <i>with</i> mental disorders <sup>d</sup> N = 82 249	751 719	9837
Childless women <i>without</i> mental disorders <sup>e</sup> N = 462 507	13 689 888	18 441
<b>Age distribution in the cohort</b>		
15–25	10 563 262	25 005
25–35	7 384 391	7349
>35	6 105 888	4977
<b>Family history of mental illness in the cohort<sup>f</sup></b>		
Yes	3 553 229	10 938
No	20 500 312	26 393
<b>Family history of self-harm in the cohort<sup>g</sup></b>		
Yes	954 218	4380
No	23 099 323	32 951
<b>Family history of suicide in the cohort<sup>h</sup></b>		
Yes	256 265	663
No	23 797 276	36 668

<sup>a</sup>Women with any first psychiatric inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to her first child.

<sup>b</sup>Women with a record of childbirth but without any record of a psychiatric inpatient or outpatient contact.

<sup>c</sup>Women with a record of childbirth and an inpatient or outpatient contact to a psychiatric facility outside the 90-day postpartum period.

<sup>d</sup>Women without any record of childbirth but with a first inpatient or outpatient contact to a psychiatric facility.

<sup>e</sup>Women without any record of childbirth and without any record of an inpatient or outpatient contact to a psychiatric facility.

<sup>f</sup>At least one mother and/or father with a record of inpatient or outpatient psychiatric treatment.

<sup>g</sup>At least one mother and/or father with a hospital registered episode of self-harm.

<sup>h</sup>At least one mother and/or father dying from suicide.

months after PPMD diagnosis was 13.5 (95% CI 8.4–21.7) among women with PPMD ( $n = 17$  women). After the initial 12 months this risk decreased: HR 5.2 (95% CI 3.9–6.9) ( $n = 47$  women).

### Risk of recurrent episodes of self-harm and suicide

In total 33 (51.5%) of the 64 cases of self-harm in women with PPMD had at least one hospital record of subsequent self-harm during the follow-up period. 48.4% of the women self-harmed more than once, with eight individuals (12.5%) having more than four hospital registered episodes of self-harm (results not shown).

**Table 2.** Adjusted hazard ratios for self-harm in women with postpartum mental disorders and four comparison groups

	Hazard ratio (HR) <sup>a</sup>	95% CI
Women with postpartum mental disorders <sup>b</sup>	6.2	4.9–8.0
Mothers without mental disorders <sup>c</sup>	1	reference
Mothers with mental disorders diagnosed outside the postpartum period <sup>d</sup>	10.1	9.6–10.5
Childless women <i>with</i> mental disorders <sup>e</sup>	9.3	8.9–9.7
Childless women <i>without</i> mental disorders <sup>f</sup>	0.9	0.9–1.0

<sup>a</sup>Adjusted for calendar time and family history of mental illness, family history of self-harm and family history of suicide.

<sup>b</sup>Women with any first psychiatric inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to her first child.

<sup>c</sup>Women with a record of childbirth but without any record of a psychiatric inpatient or outpatient contact.

<sup>d</sup>Women with a record of childbirth and an inpatient or outpatient contact to a psychiatric facility outside the 90-day postpartum period.

<sup>e</sup>Women without any record of childbirth but with a first inpatient or outpatient contact to a psychiatric facility.

<sup>f</sup>Women without any record of childbirth and without any record of an inpatient or outpatient contact to a psychiatric facility.

The HR for suicide after an episode of self-harm in women with first onset PPMD compared to mothers without a mental disorders was 8.7 (95% CI 3.5–21.7) (see Table 4). Furthermore, women with PPMD had a significantly higher risk of suicide than the other groups of women with mental disorders – both when compared with mothers with non-PPMD (HR 2.7; 95% CI 1.9–3.7;  $p$  value = 0.0104) and childless women with mental disorders (HR 2.8; 95% CI 2.1–3.8;  $p$  value = 0.0125).

## Discussion

Using register data covering the entire Danish population, we found that women with first-onset PPMD are at increased risk of self-harm compared with mothers from the healthy background population. Nevertheless, their risk of self-harm was the lowest among all comparison groups of women with mental disorders – both mothers and non-mothers. Importantly, we further found that a record of self-harm among women with PPMD was associated with the significantly highest risk of later suicide among all the comparison groups.

### Risk of self-harm in mothers with PPMD

This present study is, the first population-based study to describe the incidence of self-harm among women with first-onset PPMD. Primiparous women experiencing PPMD within the first 90 days postpartum had an increased risk of hospital contact due to self-harm, which was approximately 6 times higher than in mothers without mental disorders.

Previous studies have reported increased rates of suicidal ideation among women in the postpartum period (Lindahl *et al.*, 2005; Wisner *et al.*, 2013), using data from a specific question relating to thoughts of self-harm (question 10) in the Edinburgh Postnatal Depression Scale (EPDS) in women with PPMD (Cox *et al.*, 1987; Lindahl *et al.*, 2005; Wisner *et al.*, 2013). While the EPDS is a highly valid tool in identifying high risk patients for postpartum depression and questions regarding suicidal behavior (thoughts of self-harm and/or suicide) are included (Adouard *et al.*, 2005; Rubertsson *et al.*, 2011), for this

**Table 3.** Short-term v. long-term risk of self-harm among women with postpartum mental disorders and four comparison groups

	Short-time risk of self-harm				Long-term risk of self-harm			
	Person years	N <sup>a</sup>	Short-term risk <sup>b</sup>	95% CI short-term risk	Person years	N <sup>c</sup>	Long-term risk <sup>b</sup>	95% CI long-term risk
Women with postpartum mental disorders <sup>d</sup>	1499	17	13.5	8.4–21.7	15 597	47	5.2	3.9–6.9
Mothers without mental disorders (reference) <sup>e</sup>					8 919 834	4950	1	(reference)
Mothers with diagnosis of mental disorder outside the postpartum period <sup>f</sup>	45 752	904	30.4	28.3–32.6	629 253	3135	8.4	8.0–8.7
Childless women <i>with</i> mental disorders <sup>g</sup>	91 258	3318	20.5	19.4–21.6	660 461	6519	7.7	7.4–8.1
Childless women <i>without</i> mental disorders <sup>h</sup>					13 689 887	18 441	1.0	0.9–1.0

<sup>a</sup>Number of women with the first record of self-harm within the first 12 months after first psychiatric inpatient or outpatient contact.

<sup>b</sup>Adjusted for age, calendar time, family history of mental illness, family history of self-harm and family history of suicide.

<sup>c</sup>Number of women with the first record of self-harm after 12 months following first psychiatric inpatient or outpatient contact.

<sup>d</sup>Women with any first psychiatric inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to her first child.

<sup>e</sup>Women with a record of childbirth but without any record of a psychiatric inpatient or outpatient contact.

<sup>f</sup>Women with a record of childbirth and an inpatient or outpatient contact to a psychiatric facility outside the 90-day postpartum period.

<sup>g</sup>Women without any record of childbirth but with a first inpatient or outpatient contact to a psychiatric facility.

<sup>h</sup>Women without any record of childbirth and without any record of an inpatient or outpatient contact to a psychiatric facility.

**Table 4.** Survival analysis showing the risk of suicide among the women with at least one hospital registered episode of self-harm in five comparison groups

	Person-years at risk	N <sup>a</sup>	Hazard ratio <sup>b</sup>	95% CI
Women with postpartum psychiatric disorders <sup>c</sup>	604	5	8.7	3.5–21.7
Mothers without psychiatric diagnosis (reference) <sup>d</sup>	59 913	61	1	reference
Mothers with psychiatric diagnosis outside the postpartum period <sup>e</sup>	35 752	90	2.7	1.9–3.7
Childless women <i>with</i> psychiatric diagnosis <sup>f</sup>	92 573	185	2.8	2.1–3.8
Childless women <i>without</i> psychiatric diagnosis <sup>g</sup>	264 007	165	0.7	0.5–0.9

<sup>a</sup>Women with at least one hospital record of self-harm who later died by completed suicide.

<sup>b</sup>Hazard ratio adjusted for age, calendar time, family history of mental illness, family history of self-harm and family history of suicide.

<sup>c</sup>Women with any first psychiatric inpatient or outpatient contact to a psychiatric facility within 90 days after giving birth to her first child.

<sup>d</sup>Women with a record of childbirth but without any record of a psychiatric inpatient or outpatient contact.

<sup>e</sup>Women with a record of childbirth and an inpatient or outpatient contact to a psychiatric facility outside the 90-day postpartum period.

<sup>f</sup>Women without any record of childbirth but with a first inpatient or outpatient contact to a psychiatric facility.

<sup>g</sup>Women without any record of childbirth and without any record of an inpatient or outpatient contact to a psychiatric facility.

study we used a much more specific instrument in evaluating suicidal behavior as we were able to identify women who had actually acted on potential self-harming impulses. One previous study has described an increased risk of self-harm among pregnant women with pre-existing mental disorders (Taylor *et al.*, 2016). However, so far no studies have described the extent to which women with first onset PPMD act on such thoughts of self-harm.

### Risk of self-harm compared to other groups of psychiatric patients

Motherhood, in general, appears to be a protective factor in terms of reducing self-harm, suicidal ideation, and mortality (Appleby and Turnbull, 1995; Johannsen *et al.*, 2016; Lysell *et al.*, 2018). However, mental disorders constitute a well-established major risk factor for self-harm (Christiansen *et al.*, 2013) and may, therefore, add to the complexity of the risk evaluation for suicidal and self-harming behavior in new mothers. In this present study, we considered various states of mental ill-health (absent, postpartum, non-postpartum) and also motherhood (mothers, non-mothers), by using multiple comparison groups of women

drawn from the background population. The risk of self-harm in women with PPMD was increased (HR 6.2; 95% CI 4.9–8.0) when compared to mothers without any record of psychiatric admissions. While our results confirmed a generally increased risk of self-harm in the three groups of women with mental disorders (both mothers and childless women), in our more detailed analyses we found that women with a diagnosis of PPMD had the lowest risk of hospital recorded contacts due to self-harm involving hospital contact when compared to groups of women with alternate onset of mental disorders (both mothers and childless women). These results indicate important clinical variations in self-harm risks, related to both mental health and motherhood.

### Self-harm as risk factor for later suicide in women with PPMD

Self-harm is one of the leading risk factors for later suicide (Cavanagh *et al.*, 2003; Carroll *et al.*, 2014; Runeson *et al.*, 2016; Olsson *et al.*, 2017). This was also confirmed in the present study, where women suffering from PPMD with records of self-harm had significantly highest risk of later suicide amongst all comparison groups (HR 8.7; 95% CI 3.5–21.7), including

women with onset of mental disorders at alternate time points (both mothers and childless women). These results add to the already worrying evidence suggesting that women with the most severe forms of mental disorders following childbirth are at significantly increased risk of dying by suicide, especially within the first year after diagnosis (Appleby *et al.*, 1998; Johannsen *et al.*, 2016; Khalifeh *et al.*, 2016; Lysell *et al.*, 2018). An additional aspect is that women who do exhibit suicidal behavior in the postpartum period are more likely to use violent methods when dying by suicide and thereby possibly indicating a higher suicidal intent (Appleby, 1991; Lindahl *et al.*, 2005; Esscher *et al.*, 2016). Unfortunately, we were not able to stratify our analysis on violent *v.* non-violent methods of self-harm and/or suicide due to lack of statistical power.

### Clinical perspectives

Our results demonstrate that women with PPMD who have a hospital record of self-harm following their psychiatric diagnosis, constitute a selected group of high-risk patients for later suicide. These results must, however, be considered alongside the observation that absolute number of women with first onset PPMD who self-harm and commit suicide are low (five women in the present study). Regardless, every single case represents a significantly increased relative risk among this specific group of vulnerable new mothers, which is clinically significant.

Our results also show that the first year following initial diagnosis carries the highest risk of self-harm for women with PPMD (see Table 3), which indicates a particular at-risk period. Hence, goals for future research projects should be aimed at tools for early identification of women: (1) who will develop the first onset PPMD following childbirth and (2) who are at risk of self-harm and suicidal behaviors, in order to identify them before they act on such impulses.

### Strengths and limitations

This study was based on information from the national Danish health registers. In Denmark there is free and equal access both to medical and mental health care and every individual treated is registered by CPR-number (Pedersen, 2011). This way we were able to ensure a representative study population, and thus minimize selection bias.

We aimed to investigate types of PPMD, and consequently defined these as disorders requiring inpatient and outpatient contacts to specialized psychiatric treatment facilities. While information of inpatient treatment was available throughout the entirety of the study period data on outpatient contact were added to the national Danish registers from 1995 and onwards. Some women from the comparison groups of women without mental disorders (mothers and non-mothers) might have received treatment for mild-moderate mental disorders at their general practitioners without treatment being registered in the Danish Psychiatric Central Register and this information would therefore not be captured in our data. However such possible misclassification would lead to an underestimation of the risk of self-harm and suicide, rather than dilute our findings.

As self-harm is often stigmatized all uncertainties regarding the correct coding of self-harm as a deliberate act *v.* an accident cannot be ruled out completely. We applied a Danish developed algorithm for defining self-harm (Gasse *et al.*, 2018) which

includes a more general spectrum of self-harm and could capture accidental injuries (e.g. overdose).

However, this potential misclassification bias would again lead to an underestimation of the prevalence of later suicide, as some deaths recorded as being accidental may have in fact been suicides, rather than dilute our findings.

The women included in this study were born from 1963 and followed until the year 2016. As a consequence, our cohort mainly consisting of younger women, illustrated with 17 947 653 (74.6%) of the person-years at risk within the age groups under 35 years (Table 1). This age distribution potentially limits generalizability of our results to older mothers. However, as it has been shown that self-harming behavior is especially prominent among younger individuals (Abdelraheem *et al.*, 2019) we do believe that our study population is suitable for answering the study questions proposed.


We adjusted for possible confounding from age, family history of mental disorders and family history of suicidal behavior from both parental self-harm and/or suicide in our analyses, however, as in any observational study, we cannot rule out possible residual confounding.

### Conclusion

Women with the most severe PPMD registered with a specialized psychiatric treatment have an increased risk of self-harm following their initial postpartum diagnosis when compared to mothers without a history of psychiatric diagnosis and treatment. However, among all groups of psychiatric patients (both mothers and childless women with alternate onset of mental disorders) women with PPMD have the lowest risk of self-harm.

Nevertheless, the particular concern should be focused on women with PPMD with a record of self-harm, as this group represents a clinical cohort with a significantly increased risk of subsequent suicide.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291719001661>

**Author ORCIDs.**  Trine Munk-Olsen, 0000-0002-0786-7147.

**Acknowledgements.** This study was supported by NIMH (National Institute of Mental Health) grant number: R01MH104468 and an unrestricted grant from the Lundbeck Foundation (grant number: R155-2012-11280)

### References

- Abdelraheem M, McAloon J and Shand F (2019) Mediating and moderating variables in the prediction of self-harm in young people: A systematic review of prospective longitudinal studies. *Journal of Affective Disorders* 246, 14–28.
- Adouard F, Glangeaud-Freudenthal NM and Golse B (2005) Validation of the Edinburgh postnatal depression scale (EPDS) in a sample of women with high-risk pregnancies in France. *Archives of Women's Mental Health* 8, 89–95.
- Appleby L (1991) Suicide during pregnancy and in the first postnatal year. *BMJ (Clinical Research Ed.)* 302, 137–140.
- Appleby L and Turnbull G (1995) Parasuicide in the first postnatal year. *Psychological Medicine* 25, 1087–1090.
- Appleby L, Mortensen PB and Faragher EB (1998) Suicide and other causes of mortality after post-partum psychiatric admission. *The British Journal of Psychiatry: The Journal of Mental Science* 173, 209–211.
- Cantwell R, Clutton-Brock T, Cooper G, Dawson A, Drife J, Garrod D, Harper A, Hulbert D, Lucas S, McClure J, Millward-Sadler H, Neilson J, Nelson-Piercy C, Norman J, O'Herlihy C, Oates M,

- Shakespeare J, de Swiet M, Williamson C, Beale V, Knight M, Lennox C, Miller A, Parmar D, Rogers J and Springett A (2011) Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006–2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG* **118**(Suppl 1), 1–203.
- Carroll R, Metcalfe C and Gunnell D (2014) Hospital presenting self-harm and risk of fatal and non-fatal repetition: systematic review and meta-analysis. *PLoS One* **9**, e89944.
- Cavanagh JT, Carson AJ, Sharpe M and Lawrie SM (2003) Psychological autopsy studies of suicide: a systematic review. *Psychological Medicine*, **33**, 395–405.
- Christiansen E, Larsen KJ, Agerbo E, Bilenberg N and Stenager E (2013) Incidence and risk factors for suicide attempts in a general population of young people: a Danish register-based study. *The Australian and New Zealand Journal of Psychiatry* **47**, 259–270.
- Cox JL, Holden JM and Sagovsky R (1987) Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry: The Journal of Mental Science* **150**, 782–786.
- Esscher A, Essen B, Innala E, Papadopoulos FC, Skalkidou A, Sundstrom-Poromaa I and Hogberg U (2016) Suicides during pregnancy and 1 year postpartum in Sweden, 1980–2007. *The British Journal of Psychiatry: The Journal of Mental Science* **208**, 462–469.
- Gasse C, Danielsen AA, Pedersen MG, Pedersen CB, Mors O and Christensen J (2018) Positive predictive value of a register-based algorithm using the Danish National Registries to identify suicidal events. *Pharmacoepidemiology and Drug Safety* **27**, 1131–1138.
- Helweg-Larsen K (2011) The Danish register of causes of death. *Scandinavian Journal of Public Health* **39**(Suppl), 26–29.
- Helweg-Larsen K, Kjoller M, Juel K, Sundaram V, Laursen B, Kruse M, Norlev J and Davidsen M (2005) *Selvmod I Danmark: Markant Fald I Selvmord, men Stigende Anatal Selvmordsforsog. Hvorfor? Statens Institut for Folkesundhed*. Copenhagen: Statens Institut for Folkesundhed.
- Howard LM, Molyneaux E, Dennis CL, Rochat T, Stein A and Milgrom J (2014) Non-psychotic mental disorders in the perinatal period. *Lancet* **384**, 1775–1788.
- Johannsen BM, Larsen JT, Laursen TM, Bergink V, Meltzer-Brody S and Munk-Olsen T (2016) All-cause mortality in women with severe postpartum psychiatric disorders. *The American Journal of Psychiatry* **173**, 635–642.
- Jones I, Chandra PS, Dazzan P and Howard LM (2014) Bipolar disorder, affective psychosis, and schizophrenia in pregnancy and the post-partum period. *Lancet* **384**, 1789–1799.
- Khalifeh H, Hunt IM, Appleby L and Howard LM (2016) Suicide in perinatal and non-perinatal women in contact with psychiatric services: 15 year findings from a UK national inquiry. *The Lancet Psychiatry* **3**, 233–242.
- Lewis G (2001) Chapter 1: introduction and key findings. In Lewis G (ed.), *Why Mothers Die 1997–1999 Confidential Enquiry Into Maternal and Child Health*. Cambridge: RCOG Press, p. 22.
- Lindahl V, Pearson JL and Colpe L (2005) Prevalence of suicidality during pregnancy and the postpartum. *Archives of Women's Mental Health* **8**, 77–87.
- Lynge E, Sandegaard JL and Rebolj M (2011) The Danish national patient register. *Scandinavian Journal of Public Health* **39**(Suppl), 30–33.
- Lysell H, Dahlin M, Viktorin A, Ljungberg E, D'onofrio BM, Dickman P and Runeson B (2018) Maternal suicide – Register based study of all suicides occurring after delivery in Sweden 1974–2009. *PLoS One* **13**, e0190133.
- Meltzer-Brody S, Howard LM, Bergink V, Vigod S, Jones I, Munk-Olsen T, Honikman S and Milgrom J (2018) Postpartum psychiatric disorders. *Nature Reviews Disease Primers* **4**, 18022.
- Mors O, Perto GP and Mortensen PB (2011) The Danish psychiatric central research register. *Scandinavian Journal of Public Health* **39**(Suppl), 54–57.
- Munk-Olsen T, Laursen TM, Pedersen CB, Mors O and Mortensen PB (2006) New parents and mental disorders: a population-based register study. *JAMA: The Journal of the American Medical Association*, **296**, 2582–2589.
- Munk-Olsen T, Jones I and Laursen TM (2014) Birth order and postpartum psychiatric disorders. *Bipolar Disorders* **16**, 300–307.
- Munk-Olsen T, Maegbaek ML, Johannsen BM, Liu X, Howard LM, Di Florio A, Bergink V and Meltzer-Brody S (2016) Perinatal psychiatric episodes: a population-based study on treatment incidence and prevalence. *Translational Psychiatry* **6**, e919.
- Olfson M, Wall M, Wang S, Crystal S, Gerhard T and Blanco C (2017) Suicide following deliberate self-harm. *The American Journal of Psychiatry* **174**, 765–774.
- Pedersen CB (2011) The Danish civil registration system. *Scandinavian Journal of Public Health* **39**(Suppl), 22–25.
- Pedersen CB, Mors O, Bertelsen A, Waltoft BL, Agerbo E, Mcgrath JJ, Mortensen PB and Eaton WW (2014) A comprehensive nationwide study of the incidence rate and lifetime risk for treated mental disorders. *JAMA Psychiatry* **71**, 573–581.
- Rubertsson C, Borjesson K, Berglund A, Josefsson A and Sydsjo G (2011) The Swedish validation of Edinburgh Postnatal Depression Scale (EPDS) during pregnancy. *Nordic Journal of Psychiatry* **65**, 414–418.
- Runeson B, Haglund A, Lichtenstein P and Tidemalm D (2016) Suicide risk after nonfatal self-harm: a national cohort study, 2000–2008. *Journal of Clinical Psychiatry* **77**, 240–246.
- Taylor CL, Van Ravesteyn LM, Van Denberg MP, Stewart RJ and Howard LM (2016) The prevalence and correlates of self-harm in pregnant women with psychotic disorder and bipolar disorder. *Archives of Women's Mental Health* **19**, 909–915.
- Thornton C, Schmied V, Dennis CL, Barnett B and Dahlen HG (2013) Maternal deaths in NSW (2000–2006) from nonmedical causes (suicide and trauma) in the first year following birth. *BioMed Research International* **2013**, 623743.
- Wisner KL, Sit DK, Mcshea MC, Rizzo DM, Zoretich RA, Hughes CL, Eng HF, Luther JF, Wisniewski SR, Costantino ML, Confer AL, Moses-Kolko EL, Famy CS and Hanusa BH (2013) Onset timing, thoughts of self-harm, and diagnoses in postpartum women with screen-positive depression findings. *JAMA Psychiatry (Chicago, Ill.)* **70**, 490–498.
- World Health Organization (1976) *Klassifikation af sygdomme: Udvidet dansk-latinsk udgave af verdenssundhedsorganisationens internationale klassifikation af sygdomme, skader og dødsårsager*. 8th revision ed. Aarhus: Danish National Board of Health.
- World Health Organization (2016) *International statistical classification of diseases and related health problems*. 10th Revision.