



## Climate Change and Agriculture Research Paper

**Cite this article:** Roman M, Hlisnikovský L, Menšík L, Zemanová V, Kunzová E (2025). Temporal trends in winter wheat yield: the role of NPK-fertilization and climate over decades of field experiments. *The Journal of Agricultural Science* 1–10. <https://doi.org/10.1017/S0021859625000024>

Received: 17 July 2024  
Revised: 17 November 2024  
Accepted: 8 December 2024

**Keywords:**  
grain yield; nitrogen optimization; precipitation; temperature; *Triticum aestivum* L

**Corresponding author:**  
Muhammad Roman;  
Email: [m.maana26@outlook.com](mailto:m.maana26@outlook.com)

# Temporal trends in winter wheat yield: the role of NPK-fertilization and climate over decades of field experiments

Muhammad Roman<sup>1</sup> , Lukáš Hlisnikovský<sup>2</sup> , Ladislav Menšík<sup>2</sup>,  
Veronika Zemanová<sup>2</sup> and Eva Kunzová<sup>2</sup>

<sup>1</sup>Department of Environment, Faculty of Environment, Jan Evangelista Purkyně University in Ústí nad Labem, Ústí nad Labem, Czech Republic and and <sup>2</sup>Department of Nutrition Management, Crop Research Institute, Prague, Czech Republic

### Abstract

Changes in climate patterns have a significant impact on agricultural production. A comprehensive understanding of weather changes in arable farming is essential to ensure practical and effective strategies for farmers. Our research aimed to investigate how different fertilization interacts with environmental factors, examine their effects on wheat yield and varietal response over time, minimize nitrogen (N) fertilizer using alfalfa as a preceding crop, and recommend an optimum N dose based on the latest weather conditions. A long-term experiment including 15 seasons (1961–2022) was studied, where a wheat crop followed alfalfa with different N applications. Our results indicated that the average temperature in the Caslav region has increased by 0.045°C per year, more significantly since 1987. Moreover, precipitation slightly decreased by 0.247 mm, but not significantly. The average November temperatures are gradually rising, positively affecting wheat grain yield. July precipitation negatively impacted grain yield only in years with extraordinary rainfall. Additionally, new wheat varieties (Contra, Mulan, Julie) yielded statistically more than the old variety (Slavia). Effectively managing nitrogen under various climate conditions is essential for promoting plant growth and reducing environmental N losses. The optimal N dosage was determined at 65 kg/ha N, resulting in an average yield of 9.1 t/ha following alfalfa as a preceding crop. Alfalfa reduces the need for N fertilization and contributes to sustainable conventional agriculture. Our findings will serve as a foundation for designing future climate change adaptation strategies to sustain wheat production.

### Introduction

Climate change significantly affects global agricultural production, including Europe and the Czech Republic. Adaptation methods are necessary to reduce the impacts of climate change, as it poses a substantial threat to agriculture (Zhu *et al.*, 2022; Grados *et al.*, 2024; Kamalova *et al.*, 2024). Furthermore, it is predicted that in the future, global temperatures will rise by about 2.5–3°C, and precipitation patterns and rates will vary on the location (Pielke *et al.*, 2022; IPCC, 2023). Efficient crop management practices can be developed by analysing how crop growth interacts with weather conditions and agricultural practices (Müller *et al.*, 2014). Rising temperatures are being documented globally, including in the Czech Republic (Zahradníček *et al.*, 2021; World Bank, Climate Change Knowledge Portal, Czech Republic, 2024), Poland (Kundzewicz and Matczak, 2012), Germany (Hemmerle and Bayer, 2020), Austria (Benz *et al.*, 2018), France (Ribes *et al.*, 2016), United Kingdom (Wreford and Topp, 2020), Europe (Twardosz *et al.*, 2021) and in Russia (Kamalova *et al.*, 2024).

Two key factors that significantly influence wheat yield are weather changes (temperature and precipitation) and fertilization. However, properly managing other factors (insects, diseases, water, etc.) also contributes to wheat yield. In recent years, particularly since 2005, temperatures have notably risen worldwide (Wójcik-Gront and Gozdowski, 2023). Recently, Donmez *et al.* (2024) found a significant correlation between temperature, precipitation and net agricultural productivity, which could affect agricultural yield by affecting photosynthesis and respiration. Average annual temperatures have shown a significant upward trend across all European sub-regions. The observed trend indicates a notable decline in winter rainfall, recorded at 1.3 mm in Eastern Europe. At the same time, in Northern Europe, there has been a significant increase of 1.5 mm per year in winter precipitation (Lopes, 2022). Increasing temperature has been documented as a crucial factor contributing to yield reduction. The average temperature annually increased by 0.05°C in the Caslav region of the Czech Republic (Hlisnikovský *et al.*, 2023b). We can conclude that temperature and

precipitation are essential factors in crop yield formation when evaluating the effects of climatic conditions.

In the Czech Republic, wheat is the primary cereal crop, covering approximately 32% of the total growing area between 2000 and 2022 and accounting for 57% of the total cereal cultivation area (Hlisnikovský *et al.*, 2023b). Wheat yield is affected in different ways, such as tillage (Peng *et al.*, 2020), variety selection (Morgounov *et al.*, 2014) and mainly fertilization and weather (Hlisnikovský *et al.*, 2023b). High temperatures can adversely impact photosynthesis efficiency, irrigation practices and plant respiration. The temperature increases observed in Eastern Europe have notably contributed to the rise in wheat yield (Lopes, 2022). In contrast, Asseng *et al.* (2015) analysed data from 30 wheat crop models from 1981 to 2010. They discovered that warming influences wheat yield in most growing areas, projecting a 6% decrease for each degree Celsius. Additionally, Moore and Lobell (2015) demonstrated that climate patterns could account for 10% of the deceleration in wheat and barley production in Europe, with changes in agriculture and environmental policies possibly responsible for the remainder. Furthermore, May and July temperatures have been linked to wheat yield in Northern Europe ( $-0.30$  t/ha °C), barley in Southern Europe ( $-0.14$  t/ha °C), and maize in Western and Southern Europe ( $-0.42$  and  $-0.39$  t/ha °C), respectively. As temperatures rise, it becomes imperative to extensively study how varietal response, crop rotation and sustainable fertilizer practices interact to enhance wheat yield in the Czech Republic and Europe.

The dynamic and interrelated effects of genotype, weather and management present obstacles to formulating practical agronomic guidelines. Potential alternates to this approach are to deal with management practices, such as yield stability (Lollato *et al.*, 2019), long-term experiment (LTE) outputs (Assefa *et al.*, 2016; Wójcik-Gront, 2018; Lollato *et al.*, 2019) or varietal response (Mourtzinis *et al.*, 2018; Wójcik-Gront, 2018). LTE studies demonstrate that integrated nutrient management is critical for sustaining crop yield. This approach offers the chance to observe prolonged fluctuations in crop yield and related factors (Donmez *et al.*, 2024; Walia *et al.*, 2024). Additionally, Liang *et al.* (2024) predicted that wheat yield would increase by 5.8–13.5% with fertilizer treatments under future climate scenarios. Among the primary nutrients for wheat and other crops, nitrogen plays a crucial role. There has been a global increase in awareness regarding the necessity to boost crop production while mitigating environmental concerns linked to nitrogen fertilizer (Dai *et al.*, 2023). The global use of nitrogen fertilizer has significantly increased, rising from 112.5 million tons in 2015 to approximately 118.2 million tons by 2019 (Sharma and Bali, 2017). Nitrate leaching and water pollution pose significant environmental challenges worldwide in Europe, the USA, China and other rainforest areas. These issues primarily result from the excessive application of nitrogen fertilizer (Zhu and Chen, 2002; Yang *et al.*, 2006; Hangs *et al.*, 2013). LTEs are conducted across approximately 700–800 locations worldwide, following Liebig's (1840–1845) formulation of the 'law of the minimum' and the principles of mineral nutrition (Rusu *et al.*, 2024). The LTEs on tillage, fertilizer and crop rotations have been conducted worldwide under uniform conditions to understand site-specific impacts. Climate change is anticipated to modify these conditions, leading to increased temperatures, more frequent droughts and intensified weather events that cause a risk to ecosystems and their functions. In light of climate change, it is crucial to know the effect of weather patterns on cropping systems and facilitate the

formulation of adaptation strategies to safeguard future productivity (Donmez *et al.*, 2024).

One approach to mitigate the impact of weather-induced variations in crop yield from year to year is through fertilization. Mineral fertilizer nutrients are readily available, exhibit uniformity and possess a well-defined composition, facilitating precise dosing. Implementing intensive fertilizer-based cultivation methods for wheat, especially during and after the Green Revolution, has been crucial in increasing yield levels and ensuring food security (Kardes and Gunes, 2024). Nevertheless, there needs to be more understanding of the effects of temperature and rainfall variations throughout the crop growth period on historical yield progress in Europe. It is imperative to evaluate yield progress periodically, typically every 10–20 years, to identify emerging techniques and policies beyond the decline in yield. The goal of this research was to analyse the following: (a) weather developments in the LTE region ( $H_0$ : there are no significant trends in temperature and precipitation;  $H_A$ : there are significant trends in temperature and precipitation), (b) the relationships between weather parameters and wheat yield ( $H_0$ : there is no relationship between weather parameters and winter wheat grain yield;  $H_A$ : the weather has an impact on winter wheat grain yield), (c) the effect of NPK fertilization on winter wheat grain yield ( $H_0$ : NPK fertilization has no significant effect on grain yield;  $H_A$ : NPK fertilization significantly affects grain yield), and (d) the optimal nitrogen dose for wheat cultivated under specific soil and climate conditions.

## Materials and methods

### Experiment layout

The LTE was located on the southern edge of Caslav city in the Czech Republic, Central Europe ( $49^{\circ}53.67547'N$ ,  $15^{\circ}23.73552'E$ ), and established in 1956. According to the Köppen–Geiger climate classification, the area belonged to the Cfb/Dfb zone (Tolasz *et al.*, 2007; Beck *et al.*, 2018). The soil type was Greyic Phaeozem (Schad, 2016). The long-term mean, minimal, maximal temperatures and precipitation were 9.1, 2.4, 15.8°C and 517 mm, respectively (Chotusice meteorological station, approximately 5 km away). The elevation is 263 m above sea level. In the LTE, there were four fields. Each field was divided into 48 plots, where 12 different fertilizer treatments with four replications were continuously analysed in a completely randomized design ( $12 \times 4 = 48$  plots). The size of an individual plot was  $9 \times 9$  m. Together 15 seasons were evaluated in this paper: 1980, 1981, 1982, 2003, 2004, 2005, 2006, 2011, 2012, 2013, 2014, 2019, 2020, 2021 and 2022. In these years, alfalfa was the preceding crop of winter wheat.

In this study, we evaluated a total of five out of 12 fertilization treatments: (1) control (unfertilized since 1956), (2) application of mineral phosphorus (P) and potassium (K) – PK treatment, (3) application of mineral nitrogen (N), P and K (NPK1), (4) NPK2, and (5) NPK3. The rates of mineral N in NPK1, NPK2 and NPK3 treatments were 40, 80 and 120 kg/ha between 1980 and 2014 (11 seasons). In 2018, along with introducing a new wheat variety (Julie), the methodology of the long-term trial was modified, and N rates were increased to 60, 100 and 140 kg/ha in NPK1, NPK2 and NPK3 treatments, respectively. The doses of mineral forms of P and K were adjusted during the experiment based on soil analyses. All treatments consistently received the same doses of mineral P and K fertilizers, with the variation between treatments lying in the different doses of mineral N. Mineral N was applied as lime ammonium nitrate, mineral

P as granulated superphosphate and K as potassium chloride. The mineral P and K nutrients were applied during the autumn. Mineral N was applied during different stages of wheat cultivation. Before sowing the wheat in autumn, 40 kg/ha of N was applied as part of the NPK1, NPK2 and NPK3 treatments. At the beginning of spring, for regeneration purposes (BBCH 21–29), an additional 40 kg/ha of N was applied as part of the NPK2 and NPK3 treatments. Finally, 40 kg/ha of N was applied to support grain production in May, specifically as part of the NPK3 treatment (BBCH 49–51). For the most recent evaluation period (4 seasons, 2019–2022), during which mineral N rates were increased, the distribution was as follows: 60 kg/ha N in autumn (NPK1, NPK2 and NPK3 treatments), 40 kg/ha N in spring (regeneration, NPK2 and NPK3 treatments) and 40 kg/ha N (NPK3 treatment). Spring applications were made at the same developmental stages as the previous fertilizer rates. Wheat was usually planted in October at a depth of 3–4 cm and a row spacing of 12.5 cm. The sowing rate was generally 400 seeds/m<sup>2</sup>. Pesticides were applied during the trial as needed, while growth regulators were not used.

### Statistical analyses

Trends in weather patterns (temperature and precipitation) were assessed using the Mann–Kendall trend test (Mann, 1945; Kendall, 1975), supplemented by Sen's slope estimation (Sen, 1968). The homogeneity of the weather data was assessed using the Pettitt's test (Pettitt, 1979). The relationship between weather and yield was analysed using correlation. Wheat yield data were analysed using the Shapiro–Wilk (Shapiro and Wilk, 1965) and Anderson–Darling (Anderson and Darling, 1954) tests for assessing the data distribution, followed by classical ANOVA (with Games–Howell post hoc test [Games and Howell, 1976]), or Kruskal–Wallis one-way ANOVA. A linear plateau response model was used for N optimization. The analyses and figures were performed using the Statistica 14.0 (Tibco Software, Palo Alto, CA, USA), SigmaPlot 14.5 (Systat Software Inc., San Jose, CA, USA) and XLStat software (Lumivero, Burlington, MA, USA).

## Results

### Climate changes

At the site of the LTE in Caslav, we observed an increasing trend in average, minimal and maximal temperatures from 1961 to 2022. All three trends are statistically significant. The average temperature increases by 0.045°C per year (Fig. 1(a)), and based on the homogeneity test, the year 1987 was identified as the breaking point (Fig. 1(b)). The minimal temperature increases by 0.047°C annually (Fig. 1(c)), with the breaking point occurring in 1987 (Fig. 1(d)). The maximal temperature increases by 0.053°C annually (Fig. 1(e)), and the breaking point occurred in 1988 (Fig. 1(f)). On the other hand, the total annual precipitation decreases slightly by 0.247 mm, but the trend is insignificant. Based on the results of the climate analysis, we can conclude that crops in Caslav are experiencing higher temperatures since 1987–1988. At the same time, they have a limited capacity to compensate for the impact of warmer air with precipitation that is more or less the same as in the past.

### Correlation between climate and wheat yield

According to the correlation analysis, only two out of 60 relationships between climate parameters and winter wheat grain yield were statistically significant. These were: (a) the average temperature in November (moderate and positive relationship,  $r = 0.6$ , Fig. 2(a)), and (b) the sum of precipitation in July (moderate and negative relationship,  $r = -0.6$ , Fig. 2(b)).

The relationship between the average temperature in November and wheat grain yield is positive. Higher average temperatures increase wheat yield, particularly when high mineral N inputs are applied. The development of average temperatures in November (1961–2022) shows an increasing and statistically significant trend (Fig. 3(a)). Moreover, the average temperatures in November are gradually rising, which currently acts and will continue to act as a beneficial weather factor, positively affecting wheat grain yield in Caslav.

A negative relationship between wheat yield and the sum of precipitation in July indicates that higher July rainfall significantly reduces wheat grain yield. High harvest can be expected if July rainfall ranges between zero and 120 mm, with the peak at 60 mm (Fig. 2(b)). This rainfall is close to the long-term average of 72 mm (1961–2022). The trend of July precipitation is slightly increasing and at the very limit of statistical significance ( $P = 0.076$ , Fig. 3(b)). Based on the data, July precipitation affects grain yield negatively, but only in years with exceptionally high rainfall. Such heavy rainfalls, exceeding 120 mm, only occurred in six out of 62 years (1961–2022).


### Fertilization and grain yield

The grain yield of winter wheat, following alfalfa in the crop rotation, has gradually increased in all analysed fertilization treatments since 1980 (Fig. 4). The average inter-annual grain yield increase was lowest in the PK treatment (80.3 kg/ha), followed by control (82.8 kg/ha), NPK1 (96.3 kg/ha), NPK3 (103.0 kg/ha) and NPK2 (105.6 kg/ha). The results show that wheat cultivation is sustainable even without applying mineral fertilizers if wheat follows alfalfa cultivation.

The main reason for the increasing yield trends is the utilization of new wheat varieties. The lowest average grain yield was provided by Slavia (4.8 t/ha), followed by Contra (7.4 t/ha), Mulan (7.9 t/ha) and Julie (8.3 t/ha). As mentioned in the 'Materials and methods' section, along with the introduction of the Julie variety, mineral N rates were also increased by 20 kg/ha in all treatments. Therefore, the Julie variety's average yield was higher than Contra due to increased N doses. However, comparing yield from the unfertilized control treatment helps eliminate the effect of fertilization. As a result, all the latest wheat varieties (Contra, Mulan, Julie) yielded statistically comparable results (Table 1).

The effect of each fertilization treatment was divided into two parts. The period from 1980 to 2014 ( $n = 11$  seasons) was evaluated in the first part. In this period, 40, 80 and 120 t/ha N were applied to wheat in treatments NPK1, NPK2 and NPK3, respectively. According to the ANOVA results, the fertilizer treatment significantly affected grain yield ( $P < 0.05$ ). The results are shown in Table 2.

The lowest average grain yield was recorded in the unfertilized control treatment, which provided statistically similar results to the PK treatment. Although this control treatment has not received fertilization since the experiment was established in




**Figure 1.** Development of the (a) average, (c) minimum and (e) maximum temperatures (°C) in Caslav between 1961 and 2022. Breaking points in (b) average, (d) minimum and (f) maximum temperatures are based on Pettitt's test. Red lines indicate temperatures before the breaking point, and green lines after the breaking point. Sen's slope is a non-parametric estimate of the slope of a trend.

1956, the yield was relatively high. This high yield is attributed to a suitable preceding crop, alfalfa, which can fix airborne N and supply this essential nutrient to the following crop. The PK treatment provided a statistically comparable yield to all NPK treatments. High yield in this treatment results from the preceding crop combined with an adequate supply of other essential macronutrients (P and K). Applying mineral N resulted in a significantly higher yield than the control, but there were insignificant differences between the NPK treatments. When comparing the impact of years and fertilizer treatments on wheat grain yield, it was found that the year factor had a dominant effect (74%), whereas the effect of the fertilizer treatment was marginal (25%). Applying mineral N thus gave the wheat a sufficient supply of this essential nutrient to support its yield potential. However,

the wheat varieties have reached their maximum yield potential under the current soil and climatic conditions.

In 2018, the LTE was modified. Mineral N rates were increased because the initial rates seemed to limit achieving higher yields that modern wheat varieties can produce. Thus, each treatment increased the mineral N rate by 20 kg/ha. Also, a modern and high-yielding wheat variety (var. Julie) was sown in 2018. The results are shown in Table 3.

During the modified period (2019–2022), the wheat grain yield was significantly influenced by the fertilizer treatment (by 52%), the year (by 40%) and their interaction (by 8%). Compared with the previous methodology (1980–2014), the impact of fertilizer treatment increased. The control treatment yielded the lowest grain yield, significantly lower than all other treatments. However,



**Figure 2.** The relationships between wheat yield (t/ha), N dose (kg/ha) and (a) average temperature in November (°C), and (b) sum of precipitation (mm) in July. The colour scales (right corners) represent winter wheat's yield rate (t/ha).

yield in the control treatment was exceptionally high compared to the average grain yield in the Czech Republic (5.9 t/ha, 2018–2022). A distinct position was held by the PK treatment,

which resulted in yield between those of the unfertilized control and the NPK treatments. Finally, the highest yield was achieved by NPK treatments. No differences were recorded among individual NPK treatments as in the previous evaluation.

Wheat (Julie variety) was used in the trial from 2019 to 2022. It is a modern and utilized variety known for its high yield in various soil and climatic conditions. This variety can maintain a high yield even after cereal crops and exhibits stable baking quality parameters and excellent hardiness. Thanks to its high protein content, this grain falls into the highest category for bakery use (class E). Given that it is an actively utilized variety, we used the trial results to determine the optimal dosage of mineral N considering the specific soil and climatic conditions. For this, we used a non-linear response model called the linear plateau model. According to the model, the optimal N dosage was determined at 65 kg t/ha N, corresponding with the average yield of 9.1 t/ha (Fig. 5).


## Discussion

### Climate change


Climate warming profoundly impacts agroclimatic resources and agricultural production. Recently, Kamalova *et al.* (2024) found that temperature and precipitation during the growing cycles significantly affect cereal production. Further, Zhang *et al.* (2022) highlighted that climate change aspects, such as rainfall and additional inputs, positively impact wheat production. The findings of this study suggest that the patterns of minimum, mean and maximum temperatures are statistically significant and show a clear upward trend (Fig. 1). Based on our data analysis, the temperature has been gradually increasing. The mean temperature increases by 0.045°C per year. In another research, statistically significant increases in mean, minimum and maximum temperatures have been found in the Czech Republic since 1961 (Zahradníček *et al.*, 2021; Brázdil *et al.*, 2022), which is aligned with our findings. The current study indicates that the mean annual temperature ranges from 8 to 10°C, and the average annual precipitation fluctuates between 550 and 1050 mm across Germany, the Czech Republic and Slovakia (Mozny *et al.*, 2023). Further, an increase in mean annual temperature (1.5°C) between 1961 and 2000 was observed in the Moravian region of Czech Republic (Dolák *et al.*, 2023). These findings are aligned with findings in the Calsav region, which shows that the temperature is increasing in the Czech Republic. Still, the increase in the Calsav region (0.045°C) is lower than in other regions of the Czech Republic.

In our study, precipitation slightly decreased (insignificant linear trend), which is similar to other experiments conducted in the Czech Republic (Lhotka *et al.*, 2018; Hlisenikovsky *et al.*, 2023a, 2024), Serbia (Gocic and Trajkovic, 2013) and Slovakia (Repel *et al.*, 2021). Extreme climate events are becoming more frequent in the Czech Republic and Europe (Lhotka *et al.*, 2018; Grillakis, 2019). An extreme situation happened in 2012 when a severe drought led to historically low yield in the South Moravian Region of the Czech Republic (Hlisenikovsky *et al.*, 2023a). Extreme climate events are rising in the Czech Republic and Europe (Ahmad *et al.*, 2021). In general, warming tends to intensify with altitude.

In contrast, the impact of altitude on changes in precipitation is comparatively minimal (Pernicová *et al.*, 2024), which can also be observed from our study where temperature significantly increased but precipitation did not decrease significantly. While



**Figure 3.** The development of (a) the average temperature (°C) in November and (b) the sum of precipitation (mm) in July at the Caslav trial station between 1961 and 2022.



**Figure 4.** The effect of wheat varieties and fertilization treatment on wheat grain yield (t/ha) in Caslav between 1980 and 2022 ( $n = 15$  seasons). Similar letters are not statistically significantly different ( $\alpha < 0.05$ ).

the trend in precipitation is minor and shows a slight decrease in Caslav, other researchers' findings underscore the fluctuating distribution of precipitation throughout the year within the climate change framework (Brázdil *et al.*, 2021). Rising temperatures result in elevated evapotranspiration rates (i.e. evaporation losses), which lead to soil moisture depletion, reduced yield due to water scarcity and heightened crop water requirements (Kirkegaard *et al.*, 2007).

#### Climate and wheat yield

Globally, wheat is the second most-produced cereal crop (Kardes and Gunes, 2024). It is important to know how we can overcome the effects of changing climate to optimize wheat yield. The ideal average temperature for optimum wheat yield ranges between 12 and 13°C in May and 16 and 17°C in June, the South Moravian Region, Czech Republic (Hlisnikovský *et al.*, 2023a). Another

study forecasted that the ongoing weather trend in the Czech Republic will positively affect wheat yield (Zahradníček *et al.*, 2021). According to our results, the average temperatures in November are gradually rising, positively affecting wheat grain yield in the Caslav region. Similarly, Hlisnikovský *et al.* (2023a) found that the average temperature in November is increasing, and there is a notable increase in wheat yield. Moreover, temperature increases significantly contribute to increasing wheat yield, mostly likely due to better temperatures for photosynthesis and crop development. From this perspective, these weather conditions appear advantageous for wheat cultivation. Moreover, LTE conducted in Prague also revealed a consistent correlation between November temperatures and winter wheat grain yield, which is aligned with our findings (Addy *et al.*, 2020; Vanongeval and Gobin, 2023; Hlisnikovský *et al.*, 2023b).

Our data showed that July precipitation negatively affects grain yield, but only in years with exceptionally high rainfall. Such

**Table 1.** Wheat grain yield (t/ha) as affected by the wheat variety in the unfertilized control treatment in Caslav between 1980 and 2022 ( $n = 15$  seasons)

Wheat variety	Grain yield (t/ha)
Slavia	$3.7 \pm 1.4^A$
Contra	$6.3 \pm 1.4^B$
Mulan	$6.9 \pm 1.3^B$
Julie	$6.7 \pm 0.8^B$

Note: The average grain yield ( $\pm$ standard deviation), followed by the same letter, are not statistically significantly different ( $\alpha < 0.05$ ). The comparison was made using the Kruskal–Wallis method, followed by the Conover–Iman procedure.

**Table 2.** Effect of fertilizer treatment on wheat grain yield (t/ha) in Caslav

Fertilizer treatment	Grain yield (t/ha)
Control	$5.8 \pm 1.9A$
PK	$6.5 \pm 1.6AB$
NPK1	$7.2 \pm 1.8B$
NPK2	$7.3 \pm 2.1B$
NPK3	$7.6 \pm 2.1B$

Results from 11 seasons between 1980 and 2014.

Note: The average grain yield ( $\pm$ standard deviation), followed by the same letter, is not statistically significantly different ( $\alpha < 0.05$ ). ANOVA was done using the Kruskal–Wallis method, followed by the Conover–Iman procedure.

**Table 3.** Effect of fertilizer treatment on grain yield (t/ha) of Julie variety in Caslav


Fertilizer treatment	Grain yield (t/ha)
Control	$6.7 \pm 0.2A$
PK	$7.6 \pm 0.2B$
NPK1	$9.0 \pm 0.3C$
NPK2	$9.0 \pm 0.3C$
NPK3	$9.3 \pm 0.4C$

Results from four seasons (2019–2022).

Note: The average grain yield ( $\pm$ standard error), followed by similar letters, are not statistically significantly different ( $\alpha < 0.05$ ). Results are based on ANOVA results, followed by the Games–Howell post hoc test.

heavy rainfalls, exceeding 120 mm, only occurred in six out of 62 years (1961–2022). The trend of July precipitation is slightly increasing and at the very limit of statistical significance ( $P = 0.076$ , Fig. 3(b)). High yield can be expected if July rainfall ranges between 0 and 120 mm, with the peak at 60 mm (Fig. 2(b)). In Sweden, De Toro *et al.* (2015) found that rainy harvesting period was sometimes connected with reduced yield of cereals, which is aligned with our findings. In another study, a negative correlation between July rainfall and grain yield was also recorded in the evaluation of Rothamsted LTE (Addy *et al.*, 2020).

Precipitation is positively correlated with the yield of most crops. However, changes in precipitation distribution may lead to reduced crop yield due to water shortages during the vegetation period, impacting plant growth and development (Wójcik-Gront and Gozdowski, 2023). Excessively wet harvest periods prevent the timely entry of harvesting equipment into the field, promote

**Figure 5.** The response of wheat variety Julie (grain yield, t/ha, black circles) to increasing application of mineral N in Caslav between 2019 and 2022 (four seasons). The data are presented in a linear plateau response model (black line).

root and wind lodging of the plants and promote the development of pathogenic organisms, ultimately reducing grain yield per hectare.

### Wheat varieties and N fertilization

European wheat varieties have shown the incapacity of current uniform cultivars to endure climate changes (Kahiluoto *et al.*, 2019). Hence, there is a need to establish sustainable systems capable of ensuring food security by stabilizing agricultural production (Frison *et al.*, 2011). Fertilizer application and the development of varieties are ways to deal with climate effects (Van Frank *et al.*, 2020). The rise in wheat production after the Green Revolution has been primarily attributed to advancements in management techniques, including nitrogen fertilization and the introduction of new cultivars (Wang and Frei, 2011; Holman *et al.*, 2016). Despite evidence of genetic advancements in recent decades, Europe still lacks a comprehensive understanding of genotype, environment and management condition interactions with the physiological characteristics behind genotypic adaptation (Senapati and Semenov, 2020). The significance of this issue cannot be overstated, particularly as climate change rapidly transforms global environmental conditions (Blanco *et al.*, 2017; Agovino *et al.*, 2019).

In our study, the yield variation of different wheat varieties can be noted within a period. Wheat yield increased with time and also varied with different varieties. The effect of nitrogen treatments was also significant (Table 1). However, the difference between the control and the maximum yielding treatment NPK3 was only 1.8 t/ha (Table 2). It is attributed to the cultivation of alfalfa, which was a preceding crop in the crop rotation. Moreover, preceding crops can affect yield more (1995–1998) than different wheat varieties (Hlisnikovský *et al.*, 2023a). Some other reasons may also affect yield, but with LTE analysis, we can get more evidence of the inadequate difference between the control and different N treatments. Legumes like alfalfa can fix atmospheric N when adequate soil N is unavailable. The fixation of atmospheric nitrogen ( $N_2$ ) by legumes per season ranges from 24 to 250 kg/ha N, with the highest rate observed in Alfalfa

(*Medicago sativa* L.) (Epstein and Bloom, 1853). Alfalfa is a superior rotation crop with deep roots that can absorb residual soil N from deeper soil layers and increase N availability to subsequent shallow-rooted crops. Ultimately, alfalfa can reduce farmer's economic expenses by reducing N fertilizer and minimizing environmental pollution. With evidence from our results, various studies have found that alfalfa cultivation can reduce N application for the following crops from 40 to 80 kg/ha (Thiessen Martens *et al.*, 2005; Ballesta and Lloveras, 2010; N'Dayegamiye *et al.*, 2015). Sometimes, N fertilization can be omitted altogether (Yost *et al.*, 2021). As a preceding crop, alfalfa also explains why wheat yield is statistically comparable in all fertilizer treatments. The crop rotation and proper fertilizer applications can be attributed to sustainable wheat production, even with low N applications, ultimately minimizing farmers' expenses. Further, in this experiment, different wheat varieties were also evaluated in different periods. The latest wheat variety (Julie) was analysed to recommend the current N dose for farmers in the Caslav region. According to the model, the optimal N dosage was determined at 65 kg/ha N, corresponding with the average yield of 9.1 t/ha at the Caslav region, when wheat follows alfalfa in the crop rotation.

## Conclusion

The study analyses the relation between winter wheat yield, weather changes and the role of alfalfa in minimizing N fertilizer applications. Alfalfa as a preceding crop reduces the need of N fertilization and contributes to sustainable conventional agriculture. Further, higher grain yield is associated with warmer November and July precipitation. Modern wheat varieties showed an upward yield trend even in the unfertilized control variant, indicating that wheat cultivation is sustainable even without applying fertilizers. Taking into account the above findings, it could be suggested that the farmers' community will be encouraged to utilize timely climate information issued from National Meteorological Departments for farm-level decision to enhance their crop production.

**Data.** Data will be made available on request.

**Author contributions.** Conceptualization, M. R. and L. H.; methodology, E. K.; validation, M. R., L. H. and E. K.; formal analysis, M. R., L. H. and E. K.; investigation, M. R. and V. Z.; resources, L. M. and E. K.; data curation, E. K.; writing original draft preparation, M. R., L. H. and V. Z.; writing review and edit, M. R. and L. H.; visualization, L. H.; supervision, E. K.; project administration, L. M. and E. K.; funding acquisition, L. M. and E. K. All authors have read and agreed to the published version of the manuscript.

**Funding statement.** This work was supported by the Ministry of Agriculture of the Czech Republic (grant numbers RO0423, QL24020149, QK22010251, QK23020056) and UJEP-SGS-2024-44-003-3.

**Competing interests.** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Ethical standards and consent to participate.** Not applicable.

## References

AdDY JW, Ellis RH, Macdonald AJ, Semenov MA and Mead A (2020) Investigating the effects of inter-annual weather variation (1968–2016) on the functional response of cereal grain yield to applied nitrogen, using data from the Rothamsted Long-Term Experiments. *Agricultural and Forest Meteorology* **284**, 107898.

- Agovino M, Casaccia M, Ciommi M, Ferrara M and Marchesano K (2019) Agriculture, climate change and sustainability: the case of EU-28. *Ecological Indicators* **105**, 525–543. <https://doi.org/10.1016/j.ecolind.2018.04.064>
- Ahmad MJ, Cho GH, Kim SH, Lee S, Adelodun B and Choi KS (2021) Influence mechanism of climate change over crop growth and water demands for wheat-rice system of Punjab, Pakistan. *Journal of Water and Climate Change* **12**, 1184–1202. <https://doi.org/10.2166/wcc.2020.009>.
- Anderson TW and Darling DA (1954) A test of goodness of fit. *Journal of the American Statistical Association* **49**, 765–769. doi: 10.1080/01621459.1954.10501232.
- Assefa Y, Vara Prasad PV, Carter P, Hinds M, Bhalla G, Schon R and Ciampitti IA (2016) Yield responses to planting density for US modern corn hybrids: a synthesis-analysis. *Crop Science* **56**, 2802–2817. <https://doi.org/10.2135/cropsci2016.04.0215>
- Asseng S, Ewert F, Martre P, Rötter RP, Lobell DB, Cammarano D and Zhu Y (2015) Rising temperatures reduce global wheat production. *Nature Climate Change* **5**, 143–147. <https://doi.org/10.1038/nclimate2470>
- Ballesta A and Lloveras J (2010) Nitrogen replacement value of alfalfa to corn and wheat. *Spanish Journal of Agricultural Research* **8**, 159–169. <https://doi.org/10.5424/sjar/2010081-1155>
- Beck HE, Zimmermann NE, McVicar TR, Vergopolan N, Berg A and Wood EF (2018) Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data* **5**, 1–12. <https://doi.org/10.1038/sdata.2018.214>
- Benz SA, Bayer P, Winkler G and Blum P (2018) Recent trends of groundwater temperatures in Austria. *Hydrology and Earth System Sciences* **22**, 3143–3154. <https://doi.org/10.5194/hess-22-3143-2018>
- Blanco M, Ramos F, Van Doorslaer B, Martínez P, Fumagalli D, Ceglar A and Fernández FJ (2017) Climate change impacts on EU agriculture: a regionalized perspective taking into account market-driven adjustments. *Agricultural Systems* **156**, 52–66. <https://doi.org/10.1016/j.agsy.2017.05.013>
- Brzdil R, Zahradníček P, Dobrovolný P, Štěpánek P and Trnka M (2021) Observed changes in precipitation during recent warming: The Czech Republic, 1961–2019. *International Journal of Climatology* **41**, 3881–3902. <https://doi.org/10.1002/joc.7048>
- Brzdil R, Zahradníček P, Dobrovolný P, Řehoř J, Trnka M, Lhotka O and Štěpánek P (2022) Circulation and climate variability in the Czech Republic between 1961 and 2020: a comparison of changes for two 'normal' periods. *Atmosphere* **13**, 137. <https://doi.org/10.3390/atmos13010137>
- Dai J, Gui H, Shen F, Liu Y, Bai M, Yang J and Siddique KH (2023) Fertilizer 15N balance in a soybean–maize–maize rotation system based on a 41-year long-term experiment in Northeast China. *Frontiers in Plant Science* **14**, 1105131. <https://doi.org/10.3389/fpls.2023.1105131>
- De Toro A, Eckersten H, Nkurunziza L and Von Rosen D (2015) Effects of extreme weather on yield of major arable crops in Sweden: Analysis of long-term experiment data. *Swedish University of Agricultural Sciences Aspects of Applied Biology* **128**, 165–172.
- Dolák L, Řehoř J, Lásková K, Štěpánek P and Zahradníček P (2023) Air temperature variability of the northern mountains in the Czech Republic. *Atmosphere* **14**, 1063. <https://doi.org/10.3390/atmos14071063>
- Donmez C, Sahingoz M, Paul C, Cilek A, Hoffmann C, Berberoglu S and Helming K (2024) Climate change causes spatial shifts in the productivity of agricultural long-term field experiments. *European Journal of Agronomy* **155**, 127121. <https://doi.org/10.1016/j.eja.2024.127121>
- Epstein E and Bloom AJ (1853) *Mineral Nutrition of Plants: Principles and Perspectives*. Sunderland: Sinauer Inc. Publishers.
- Frison EA, Cherfas J and Hodgkin T (2011) Agricultural biodiversity is essential for a sustainable improvement in food and nutrition security. *Sustainability* **3**, 238–253. <https://doi.org/10.3390/su3010238>
- Games PA and Howell JF (1976) Pairwise multiple comparison procedures with unequal n's and/or variances: a Monte Carlo study. *Journal of Educational Statistics* **1**, 113–125.
- Gocic M and Trajkovic S (2013) Analysis of changes in meteorological variables using Mann-Kendall and Sen's slope estimator statistical tests in Serbia. *Global and Planetary Change* **100**, 172–182. <https://doi.org/10.1016/j.gloplacha.2012.10.014>
- Grados D, Kraus D, Haas E, Butterbach-Bahl K, Olesen JE and Abalos D (2024) Common agronomic adaptation strategies to climate change may



- increase soil greenhouse gas emission in Northern Europe. *Agricultural and Forest Meteorology* **349**, 109966. <https://doi.org/10.1016/j.agrformet.2024.109966>
- Grillakis MG (2019) Increase in severe and extreme soil moisture droughts for Europe under climate change. *Science of the Total Environment* **660**, 1245–1255. <https://doi.org/10.1016/j.scitotenv.2019.01.001>
- Hangs RD, Schoenau JJ and Lafond GP (2013) The effect of nitrogen fertilization and no-till duration on soil nitrogen supply power and post-spring thaw greenhouse-gas emissions. *Journal of Plant Nutrition and Soil Science* **176**, 227–237. <https://doi.org/10.1002/jpln.201200242>
- Hemmerle H and Bayer P (2020) Climate change yields groundwater warming in Bavaria, Germany. *Frontiers in Earth Science* **8**, 575894. <https://doi.org/10.3389/feart.2020.575894>
- Hlisenikový L, Menšík L, Barlóg P and Kunzová E (2023a) How weather and fertilization affected grain yield and stability of winter wheat in a long-term trial in the south Moravian region, Czech Republic. *Agronomy* **13**, 2293. <https://doi.org/10.3390/agronomy13092293>
- Hlisenikový L, Menšík L and Kunzová E (2023b) Development and the effect of weather and mineral fertilization on grain yield and stability of winter wheat following alfalfa – analysis of long-term field trial. *Plants* **12**, 1392. <https://doi.org/10.3390/plants12061392>
- Hlisenikový L, Menšík L, Roman M and Kunzová E (2024) The evaluation of a long-term experiment on the relationships between weather, nitrogen fertilization, preceding crop, and winter wheat grain yield on Cambisol. *Plants* **13**, 802. <https://doi.org/10.3390/plants13060802>
- Holman FH, Riche AB, Michalski A, Castle M, Wooster MJ and Hawkesford MJ (2016) High throughput field phenotyping of wheat plant height and growth rate in field plot trials using UAV based remote sensing. *Remote Sensing* **8**, 1031. <https://doi.org/10.3390/rs8121031>
- Intergovernmental Panel on Climate (IPCC) Change (2023) *Climate Change 2021-The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Kahiluoto H, Kaseva J, Balek J, Olesen JE, Ruiz-Ramos M, Gobin A and Trnka M (2019) Decline in climate resilience of European wheat. *Proceedings of the National Academy of Sciences* **116**, 123–128. <https://doi.org/10.1073/pnas.1804387115>
- Kamalova R, Bogdan E, Belan L, Tuktarova I, Firstov A, Vildanov I and Saifullin I (2024) Assessment of changes in agroclimatic resources of the Republic of Bashkortostan (Russia) under the context of global warming. *Climate* **12**, 11. <https://doi.org/10.3390/cli12010011>
- Kardes TA and Gunes A (2024) Environmental and innovative fertilizer development strategies for wheat cultivation: urea-doped hydroxyapatite, biochar-coated diammonium phosphate, and biochar-coated urea in basal and top dressing. *Journal of Soil Science and Plant Nutrition* **24**, 1–16. <https://doi.org/10.1007/s42729-024-01737-6>
- Kendall MG (1975) *Rank Correlation Methods*, 4th Edn. London: Griffin.
- Kirkegaard JA, Lilley JM, Howe GN and Graham JM (2007) Impact of sub-soil water use on wheat yield. *Australian Journal of Agricultural Research* **58**, 303–315. <https://doi.org/10.1071/AR06285>
- Kundzewicz ZW and Matczak P (2012) Climate change regional review: Poland. *Wiley Interdisciplinary Reviews: Climate Change* **3**, 297–311. <https://doi.org/10.1002/wcc.175>
- Lhotka O, Kyselý J and Farda A (2018) Climate change scenarios of heat waves in Central Europe and their uncertainties. *Theoretical and Applied Climatology* **131**, 1043–1054. <https://doi.org/10.1007/s00704-016-2031-3>
- Liang S, Sun N, Meersmans J, Longdoz B, Colinet G, Xu M and Wu L (2024) Impacts of climate change on crop production and soil carbon stock in a continuous wheat cropping system in southeast England. *Agriculture, Ecosystems Environment* **365**, 108909. <https://doi.org/10.1016/j.agee.2024.108909>
- Lollato RP, Ochsner TE, Arnall DB, Griffin TW and Edwards JT (2019) From field experiments to regional forecasts: upscaling wheat grain and forage yield response to acidic soils. *Agronomy Journal* **111**, 287–302. <https://doi.org/10.2134/agronj2018.03.0206>
- Lopes MS (2022) Will temperature and rainfall changes prevent yield progress in Europe? *Food and Energy Security* **11**, e372. <https://doi.org/10.1002/fes3.372>
- Mann HB (1945) Nonparametric tests against trend. *Econometrica* **13**, 245. <https://doi.org/10.2307/1907187>
- Moore FC and Lobell DB (2015) The fingerprint of climate trends on European crop yields. *Proceedings of the National Academy of Sciences* **112**, 2670–2675. <https://doi.org/10.1073/pnas.1409606112>
- Morgounov A, Abugalieva A and Martynov S (2014) Effect of climate change and variety on long-term variation of grain yield and quality in winter wheat in Kazakhstan. *Cereal Research Communications* **42**, 163–172. <https://doi.org/10.1556/CRC.2013.0047>
- Mourtzinis S, Kaur G, Orłowski JM, Shapiro CA, Lee CD, Wortmann C and Conley SP (2018) Soybean response to nitrogen application across the United States: a synthesis-analysis. *Field Crops Research* **215**, 74–82. <https://doi.org/10.1016/j.fcr.2017.09.035>
- Mozny M, Trnka M, Vlach V, Zalud Z, Cejka T, Hajkova L, Potopova V, Semenov MA, Semeradova D and Büntgen U (2023) Climate-induced decline in the quality and quantity of European hops calls for immediate adaptation measures. *Nature Communications* **14**, 6028. <https://doi.org/10.1038/s41467-023-41474-5>
- Müller C, Waha K, Bondeau A and Heinke J (2014) Hotspots of climate change impacts in sub-Saharan Africa and implications for adaptation and development. *Global Change Biology* **20**, 2505–2517. <https://doi.org/10.1111/gcb.12586>
- N'Dayegamiye A, Whalen JK, Tremblay G, Nyiraneza J, Grenier M, Drapeau A and Bipfubusa M (2015) The benefits of legume crops on corn and wheat yield, nitrogen nutrition, and soil properties improvement. *Agronomy Journal* **107**, 1653–1665. <https://doi.org/10.2134/agronj14.0416>
- Peng Z, Wang L, Xie J, Li L, Coulter JA, Zhang R and Whitbread A (2020) Conservation tillage increases yield and precipitation use efficiency of wheat on the semi-arid Loess Plateau of China. *Agricultural Water Management* **231**, 106024. <https://doi.org/10.1016/j.agwat.2020.106024>
- Pernicová N, Urban O, Čáslavský J, Kolář T, Rybníček M, Sochová I and Trnka M (2024) Impacts of elevated CO<sub>2</sub> levels and temperature on photosynthesis and stomatal closure along an altitudinal gradient are counteracted by the rising atmospheric vapor pressure deficit. *Science of the Total Environment* **921**, 171173. <https://doi.org/10.1016/j.scitotenv.2024.171173>
- Pettitt AN (1979) A non-parametric approach to the change-point problem. *Applied Statistics* **28**, 126. <https://doi.org/10.2134/agronj14.0416>
- Pielke Jr. R, Burgess MG and Ritchie J (2022) Plausible 2005–2050 emissions scenarios project between 2 and 3 degrees C of warming by 2100. *Environmental Research Letters* **17**, 024027. <https://doi.org/10.1088/1748-9326/ac4ebf>
- Repel A, Zelenáková M, Jothiprakash V, Hlavatá H, Blištan P, Gargar I and Purcz P (2021) Long-term analysis of precipitation in Slovakia. *Water* **13**, 952. <https://doi.org/10.3390/w13070952>
- Ribes A, Corre L, Gibelin AL and Dubuisson B (2016) Issues in estimating observed change at the local scale – a case study: the recent warming over France. *International Journal of Climatology* **36**, 3794–3806. <https://doi.org/10.1002/joc.4593>
- Rusu M, Mihai M, Tritean N, Mihai V, Moldovan L, Ceclan OA and Toader C (2024) Results of long term fertiliser experiments-use in sustainable soil fertility protection measures. *Romanian Agricultural Research* **41**, 329–341. <https://doi.org/10.59665/rar4131>
- Schad P (2016) The international soil classification system WRB, third edition 2014. In *Novel Methods for Monitoring and Managing Land and Water Resources in Siberia*. Cham: Springer, pp. 563–571. <https://doi.org/10.1007/978-3-319-24409-9-25>
- Sen PK (1968) Estimates of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association* **63**, 1379–1389.
- Senapati N and Semenov MA (2020) Large genetic yield potential and genetic yield gap estimated for wheat in Europe. *Global Food Security* **24**, 100340. <https://doi.org/10.1016/j.gfs.2019.100340>
- Shapiro SS and Wilk MB (1965) An analysis of variance test for normality (complete samples). *Biometrika* **52**, 591.
- Sharma LK and Bali SK (2017) A review of methods to improve nitrogen use efficiency in agriculture. *Sustainability* **10**, 51. <https://doi.org/10.3390/su10010051>
- Thiessen Martens JR, Entz MH and Hoepfner JW (2005) Legume cover crops with winter cereals in southern Manitoba: fertilizer replacement values for oat. *Canadian Journal of Plant Science* **85**, 645–648.
- Tolasz R, Míková T, Valeriánová A and Voženílek V (2007) *Climate Atlas of Czechia*. Prague: Czech Hydrometeorological Institute, p. 256.

- Twardosz R, Walanus A and Guzik I** (2021) Warming in Europe: recent trends in annual and seasonal temperatures. *Pure and Applied Geophysics* **178**, 4021–4032. <https://doi.org/10.1007/s00024-021-02860-6>
- Van Frank G, Rivière P, Pin S, Baltassat R, Berthelot JF, Caizergues F and Goldringer I** (2020) Genetic diversity and stability of performance of wheat population varieties developed by participatory breeding. *Sustainability* **12**, 384. <https://doi.org/10.3390/su12010384>
- Vanongeval F and Gobin A** (2023) Adverse weather impacts on winter wheat, maize and potato yield gaps in northern Belgium. *Agronomy* **13**, 1104. <https://doi.org/10.3390/agronomy13041104>
- Walia SS, Dhaliwal SS, Gill RS, Kaur T, Kaur K, Randhawa MK and Hossain A** (2024) Improvement of soil health and nutrient transformations under balanced fertilization with integrated nutrient management in a rice-wheat system in Indo-Gangetic Plains—A 34-year research outcomes. *Heliyon* **10**, 4. <https://doi.org/10.1016/j.heliyon.2024.e25113>
- Wang Y and Frei M** (2011) Stressed food – the impact of abiotic environmental stresses on crop quality. *Agriculture, Ecosystems Environment* **141**, 271–286. <https://doi.org/10.1016/j.agee.2011.03.017>
- Wójcik-Gront E** (2018) Variables influencing yield-scaled Global Warming Potential and yield of winter wheat production. *Field Crops Research* **227**, 19–29. <https://doi.org/10.1016/j.fcr.2018.07.015>
- Wójcik-Gront E and Gozdowski D** (2023) Effect of climate change in years 2006–2019 on crop yields in Poland. *European Journal of Sustainable Development* **12**, 225–225. <https://doi.org/10.14207/ejsd.2023.v12n4p225>
- World Bank, Climate Change Knowledge Portal, Czech Republic** (2024) Date accessed: November 12, 2024, from <https://climateknowledgeportal.worldbank.org/country/czech-republic/climate-data-historical>
- Wreford A and Topp CF** (2020) Impacts of climate change on livestock and possible adaptations: a case study of the United Kingdom. *Agricultural Systems* **178**, 102737. <https://doi.org/10.1016/j.agsy.2019.102737>
- Yang SM, Malhi SS, Song JR, Xiong YC, Yue WY, Lu LL and Guo TW** (2006) Crop yield, nitrogen uptake and nitrate-nitrogen accumulation in soil as affected by 23 annual applications of fertilizer and manure in the rainfed region of Northwestern China. *Nutrient Cycling in Agroecosystems* **76**, 81–94. <https://doi.org/10.1007/s10705-006-9042-x>
- Yost MA, Pound CA, Creech JE, Cardon GE, Pace MG, Kitchen B and Russell K** (2021) Nitrogen requirements of first-year small grains after alfalfa. *Soil Science Society of America Journal* **85**, 1698–1709. <https://doi.org/10.1002/saj2.20269>
- Zahradníček P, Brázdil R, Štěpánek P and Trnka M** (2021) Reflections of global warming in trends of temperature characteristics in the Czech Republic, 1961–2019. *International Journal of Climatology* **41**, 1211–1229. <https://doi.org/10.1002/joc.6791>
- Zhang H, Tang Y, Chandio AA, Sargani GR and Ankrah Twumasi M** (2022) Measuring the effects of climate change on wheat production: evidence from Northern China. *International Journal of Environmental Research and Public Health* **19**, 12341. <https://doi.org/10.3390/ijerph191912341>
- Zhu ZL and Chen DL** (2002) Nitrogen fertilizer use in China – contributions to food production, impacts on the environment and best management strategies. *Nutrient cycling in Agroecosystems* **63**, 117–127. <https://doi.org/10.1023/A:1021107026067>
- Zhu P, Burney J, Chang J, Jin Z, Mueller ND, Xin Q and Ciais P** (2022) Warming reduces global agricultural production by decreasing cropping frequency and yields. *Nature Climate Change* **12**, 1016–1023. <https://doi.org/10.1038/s41558-022-01492-5>