

Immunodiagnostic monoclonal antibody-based sandwich ELISA of fasciolosis by detection of *Fasciola gigantica* circulating fatty acid binding protein

PANAT ANURACPREEDA^{1,2*}, RUNGLAWAN CHAWENGGKIRTTIKUL³ and PRASERT SOBHON²

¹ Division of Agricultural Science, Mahidol University, Kanchanaburi Campus, Saiyok, Kanchanaburi 71150, Thailand

² Department of Anatomy, Faculty of Science, Mahidol University, Rama VI Rd., Bangkok 10400, Thailand

³ Department of Microbiology, Faculty of Science, Mahidol University, Rama VI Rd., Bangkok 10400, Thailand

(Received 5 April 2016; revised 18 May 2016; accepted 22 May 2016; first published online 17 June 2016)

SUMMARY

Up to now, parasitological diagnosis of fasciolosis is often unreliable and possesses low sensitivity. Hence, the detection of circulating parasite antigens is thought to be a better alternative for diagnosis of fasciolosis, as it reflects the real parasite burden. In the present study, a monoclonal antibody (MoAb) against recombinant *Fasciola gigantica* fatty acid binding protein (rFgFABP) has been produced. As well, a reliable sandwich enzyme-linked immunosorbent assay (sandwich ELISA) has been developed for the detection of circulating FABP in the sera of mice experimentally and cattle naturally infected with *F. gigantica*. MoAb 3A3 and biotinylated rabbit anti-recombinant FABP antibody were selected due to their high reactivities and specificities. The lower detection limit of sandwich ELISA was 5 pg mL⁻¹, and no cross-reaction with other parasite antigens was observed. This assay could detect *F. gigantica* infection from day 1 post infection. In experimental mice, the sensitivity, specificity and accuracy of this assay were 93·3, 100 and 98·2%, while in natural cattle they were 96·7, 100 and 99·1%. Hence, this sandwich ELISA method showed high efficiencies and precisions for diagnosis of fasciolosis by *F. gigantica*.

Key words: *Fasciola gigantica*, fatty acid binding protein, monoclonal antibody, sandwich ELISA, Seroantigen, immunodiagnosis.

INTRODUCTION

Fasciolosis due to *Fasciola gigantica* is an important disease of ruminants in the tropical countries of Asia and Africa resulting in great economic losses to the livestock industry for more than US \$3·2 billion per annum (Sobhon *et al.* 1998; Spithill *et al.* 1999; Torgerson and Claxton, 1999; Mas-Coma *et al.* 2005, 2009). Furthermore, human infections with *F. gigantica* are also reported in many countries (Ashrafi *et al.* 2006a, b; Le *et al.* 2007). As a result infection cannot be examined by microscopic detection in the feces due to the absence of the parasite's eggs during the non-reproductive phase of the flukes (Ghosh *et al.* 2005; Kumar *et al.* 2008). Alternatively, diagnosis of fasciolosis in animals has been accomplished by two accesses of immunoassays, i.e. detection of antibody in the serum samples of infected animals (Zimmerman *et al.* 1985; Swarup *et al.* 1987; Fagbemi and Obarisiagbon, 1990; Guobadia and Fagbemi, 1995; Sriveny *et al.* 2006) and detection of circulating antigen (Langley and Hillyer, 1989; Fagbemi *et al.*

1995; Viyanant *et al.* 1997; Velusamy *et al.* 2004; Anuracpreeda *et al.* 2009a, b, 2013a, 2016b). The detection of antigen rather than antibodies is considered to be a more reliable method for identifying animals with pre-patent infection, which could not be detected by the usual parasitological test. Moreover, the antigen detection can evaluate the status of infection, which could be used to monitor the efficacy of treatment (Mbuh and Fagbemi, 1996; Hillyer, 1999).

In trematode species, fatty acid binding proteins (FABPs) play a vital role in the uptake and transport of lipid molecules, i.e. long chain fatty acids and cholesterol from the host (Meyer *et al.* 1970). These proteins were first isolated from *Fasciola hepatica* by affinity chromatography using rabbit antisera against somatic antigen of *Schistosoma mansoni* (Hillyer *et al.* 1977). In addition, a cDNA library has been made for these proteins from *F. hepatica* (Rodriguez-Perez *et al.* 1992; Chicz, 1994), and an almost identical group of proteins has been characterized and expressed in *Escherichia coli* (Smooker *et al.* 1997). In comparison with *F. hepatica*, Grams *et al.* (2000) had cloned FABP cDNA from *F. gigantica* by using real-time polymerase chain reaction (RT-PCR). In this study, we produced a monoclonal antibody (MoAb) against recombinant *F. gigantica* FABP, and use it in a sandwich

* Corresponding author: Division of Agricultural Science, Mahidol University, Kanchanaburi Campus, Saiyok, Kanchanaburi 71150, Thailand. Tel. +66 3458 5060. Fax. +66 3458 5077. E-mail: Panat1@yahoo.com; panat.anu@mahidol.ac.th

enzyme-linked immunosorbent assay (sandwich ELISA) for detection of circulating FABP antigen of *F. gigantica* in the sera of experimentally and naturally infected animals. The use of this MoAb could provide the immunodiagnosis of fasciolosis with high sensitivity, specificity and accuracy.

MATERIALS AND METHODS

Ethics statement

All animal experiments were approved by the Animal Care and Use Committee (SCMUACUC), Faculty of Science, Mahidol University, Thailand and were specifically used for this study. National Laboratory Animal Center, Nakorn Pathom, Mahidol University, Thailand has earned AAALAC international accreditation. On necropsy time, all animals (mice, hamsters and rabbits) were anaesthetized. The thoraco-abdominal cavity of mice and hamsters was opened, whereas the cardiac withdrawal was performed to collect all blood from the rabbits. The cattle sera were obtained from Department of Livestock, Ministry of Agriculture and Co-operatives, Bangkok, Thailand. Also, we have received consent to collect the parasite specimens from animals at the abattoir.

Collection of parasite specimens

Metacercariae of F. gigantica. A method described by Anuracpreeda *et al.* (2011, 2016b) was used to obtain *F. gigantica* metacercariae. Briefly, the metacercariae were obtained from experimentally infected snails, *Lymnaea ollula*. The snails were infected with miracidia, hatched from the mature eggs, and conceded to develop into sporocysts and cercariae. After 42–56 days, the cercariae were shed from these snails and attached to the cellophane papers and transformed into metacercariae. The metacercariae were brushed off and collected from cellophane papers and washed several times with Hank's balance salt (HBS) solution containing 100 U mL⁻¹ penicillin and 100 mg mL⁻¹ streptomycin and used immediately.

Newly excysted juveniles (NEJ) of F. gigantica.

Fasciola gigantica NEJ was obtained by the method according to Anuracpreeda *et al.* (2011, 2016b). To activate the excystment, the metacercariae were incubated in a solution containing 1% (w/v) pepsin (pepsin A from porcine gastric mucosa, P-7000, Sigma-Aldrich Co.) and 0.4% (v/v) HCl at 37 °C for 45 min. After washing with distilled water, they were incubated in a solution of 0.02 M sodium dithionite (Fluka Biochemika), 0.2% (w/v) taurocholic acid (T-4009, Sigma-Aldrich Co.), 1% (w/v) NaHCO₃, 0.8% (w/v) NaCl and 0.005% (v/v) HCl at 37 °C for 45 min and washed with distilled water. The activated metacercariae were excysted in fresh Roswell Park Memorial Institute (RPMI)-1640 medium (Sigma

Chemical Co., St. Louis, MO, USA) containing 10% fetal calf serum and 10 µg mL⁻¹ gentamycin at 37 °C overnight. On the following day, the NEJ were collected and washed several times with HBS solution and used immediately.

Juvenile parasites of F. gigantica. A method described by Anuracpreeda *et al.* (2011, 2014) was used to obtain juvenile parasites from male Golden Syrian hamsters were orally infected with metacercariae. At 1, 3 and 5 weeks after infections, the infected animals were sacrificed and the liver was teased to collect the parasites. They were washed several times with HBS solution and used immediately.

Adult stages of F. gigantica and other parasites.

For the cross-reactivity study, adult trematodes including *Fasciola hepatica*, *Eurytrema pancreaticum*, *Gigantocotyle explanatum*, *Cotylophoron cotylophorum*, *Paramphistomum cervi*, *Paramphistomum gracile*, *Fischoederius cobboldi*, *Gastrothylax crumenifer*, *Carmyerius spatiosus*, *Carmyerius gregarious* and *Schistosoma spindale*, adult cestodes including *Moniezia benedeni* and *Avitellina centripunctata*, as well as adult nematodes including *Haemonchus placei*, *Trichuris* sp. and *Setaria labiato-papillosa* were collected from the infected cattle or water buffaloes killed at local abattoirs. Adult *Opisthorchis viverrini* were obtained from hamsters infected orally with metacercariae. Adult *Schistosoma* sp. including *Schistosoma mansoni*, *Schistosoma japonicum* and *Schistosoma mekongi* were obtained by perfusing mice after being infected with schistosome cercariae. All parasite specimens were washed several times in HBS solution before being processed for further experiments (Anuracpreeda *et al.*, 2012, 2015, 2016b, c, d; Panyarachun *et al.*, 2010, 2013).

Preparations of parasite antigens

Whole body (WB) antigens of the parasites. WB antigens of the parasites were carried out according to the method of Anuracpreeda *et al.* (2008, 2016a, b). Briefly, the WB antigens were obtained by extracting whole parasites (metacercariae, NEJ, 1, 3, 5-week-old juveniles, adults of *F. gigantica* and other species) in a lysis buffer containing 10 mM Tris-HCl, pH 7.2, 150 mM NaCl, 0.5% (v/v) Triton X-100, 1 mM EDTA and 1 mM PMSF (P-7626, Sigma-Aldrich Co.). The extracted antigens were homogenized, sonicated and rotated for at 4 °C 1 h. Thereafter, the suspensions were centrifuged at 5000 × g, at 4 °C for 20 min to get rid of the parasites' eggs, and the supernatants were collected, lyophilized and kept at -70 °C until use in subsequent experiments.

Tegumental antigens (TA) of adult F. gigantica.

Fasciola gigantica TA was prepared as per the method of Anuracpreeda *et al.* (2006, 2016b).

Briefly, adult parasites were incubated in an extracting solution containing 1% Triton X-100 in 0.05 M Tris buffer, pH 8.0, 0.01 M EDTA, 0.15 M NaCl at room temperature for 20 min. Thereafter, the solution was collected and centrifuged at $5000 \times g$ at 4 °C for 20 min. The extracted TA was collected and equilibrated in 0.01 M phosphate buffered saline (PBS), pH 7.2, at 4 °C for 24 h, using Spectra/Por dialysis membrane (Spectrum Medical Industries, Los Angeles, California, USA). Then it was lyophilized, and stored at -70 °C until further use.

Excretory-secretory (ES) antigens of adult F. gigantica. The ES antigens were obtained according to the method of Anuracpreeda *et al.* (2013b, 2016b). Briefly, live adult worms were individually incubated at 37 °C for 3 h in the sterile RPMI-1640 medium (Gibco) [pH 7.4 with HEPES 20 mM, supplemented with penicillin (50 IU mL⁻¹), streptomycin (50 µg mL⁻¹) and gentamycin (50 µg mL⁻¹)]. Then, the medium was centrifuged at $5000 \times g$ at 4 °C for 20 min to get rid of the eggs. Thereafter, the supernatants containing ES antigens were collected and equilibrated in 0.01 M PBS, pH 7.2, at 4 °C for 24 h, using Spectra/Por dialysis membrane before it was lyophilized and stored at -70 °C until use.

Protein determinations. The protein contents in the parasites' extracts were determined using Lowry's method (Lowry *et al.* 1951). These extracts were stored at -70 °C until use.

Preparation of recombinant F. gigantica fatty acid binding protein (rFgFABP). The method described by Grams *et al.* (2000) and Sirisriro *et al.* (2002) was used to obtain rFgFABP. Briefly, the complementary DNA (cDNA) of adult FgFABP (AdFgFABP) gene was cloned and amplified from the adult *F. gigantica* cDNA library. A fragment of FgFABP was isolated by polymerase chain reaction (PCR) analysis and cloned into pGEM[®]-T easy vector (Promega, Madison, USA) and prepared for sequencing by Macrogen Inc. (South Korea). The full-length FgFABP cDNA was subcloned into the pET-30b vector (Novagen) and transformed into *Escherichia coli* BL21 (DE). The rFgFABP expression was induced with isopropyl-β-D-thiogalactoside (IPTG) to 1 mM final concentration at 37 °C. The rFgFABP was purified by using nickel-nitrilotriacetic acid (Ni-NTA) affinity chromatography (QIAGEN). The eluted protein fractions were analyzed by Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and FABP-containing fractions were pooled and dialysed against 0.01 M PBS buffer, pH 7.4, at 4 °C overnight. The rFgFABP was stored at -70 °C until used in further experiments.

Production of monoclonal antibodies (MoAbs) against rFgFABP. The method described by Anuracpreeda *et al.* (2014, 2016b) was used to obtain MoAbs against rFgFABP. Briefly, MoAbs against rFgFABP were produced by fusion of mouse myeloma cells (P3 × 63-Ag8.653) with the spleen cells from inbred BALB/c mice immunized with rFgFABP using polyethylene glycol (Sigma-Aldrich Inc., Saint Louis, MO, USA). MoAbs, expressed from the hybridoma cells, were screened for recognition of *F. gigantica* antigens by indirect enzyme-linked immunosorbent assay (indirect ELISA). The highly reactive hybridoma cells were cloned by limiting dilution methods using a feeder layer of spleen cells. The antibody isotypes were screened by a standard ELISA using the SBA Clonotyping[™] System/HRP (SouthernBiotech, USA). One hybridoma clone (3A3) producing a high titre of antibody against rFgFABP was selected.

Assessing the reactivity and specificity of MoAb 3A3. Indirect ELISA method described by Anuracpreeda *et al.* (2014, 2016b) was used for assessing the reactivity and specificity of MoAb 3A3. For reactivity study, 100 µL of 100 µg mL⁻¹ of WB, TA and ES of *F. gigantica* diluted in coating buffer (15 mM Na₂CO₃, 35 mM NaHCO₃, pH 9.6) was added into each well of a flat bottom F96 microtiter plate (Nunc A/S, Roskilde, Denmark) and incubated at 37 °C overnight. After washing three times with 0.05% Tween 20 in normal saline solution (NSST), the nonspecific binding was blocked by adding 100 µL well⁻¹ of a blocking solution containing 0.25% bovine serum albumin (BSA), 0.05% Tween 20 (Sigma) in 0.01 M PBS, pH 7.2 at 37 °C for 1 h. Thereafter, the plate was similarly washed, and 100 µL of undiluted MoAb 3A3 was added and incubated at 37 °C for 2 h. After washing again, the plate was incubated with 100 µL well⁻¹ of horseradish peroxidase conjugated goat anti-mouse immunoglobulin (Sigma-Aldrich Inc.) diluted in blocking solution at 1:6000 at 37 °C for 1 h. Subsequently, the plate was washed with the same buffer, and 100 µL well⁻¹ of 3,3',5,5'-tetramethyl benzidine (TMB) substrate (KPL, Gaithersburg, USA) was added and incubated for 10 min at room temperature. Then the enzymatic reaction was stopped by the addition of 100 µL 1N HCl. Finally, the optical density (OD) value at 450 nm was read in a microplate reader (Multiskan Ascent, Labsystems, Helsinki, Finland).

In the cross-reactivity study, a flat bottom F96 microtiter plate was coated with 10 µg mL⁻¹ of WB extracts from *F. gigantica* as well as WB from other trematode, cestode and nematode parasites in 100 µL well⁻¹ of coating buffer, and incubated at 37 °C overnight. The plate was then washed three times with 0.05% NSST, blocked with 100 µL well⁻¹ of a blocking solution at 37 °C for 1 h. After

washings of the plate, 100 μL of undiluted MoAb 3A3 and diluted MoAb 3A3 at 1:10, 1:100, 1:500, 1:1000 and 1:10 000 were added to the wells and incubated at 37 °C for 2 h. After washing again, 100 μL well⁻¹ of horseradish peroxidase conjugated goat anti-mouse immunoglobulin (Sigma-Aldrich Inc.) diluted 1:6000 in the blocking solution was added and incubated at 37 °C for 1 h. Thereafter, 100 μL of TMB substrate solution was added to each well, after 10 min the enzymatic reaction was stopped by the addition of 100 μL 1N HCl. Finally, the OD value was measured at 450 nm using a microplate reader (Multiskan Ascent, Labsystems, Helsinki, Finland).

For immunoblotting assay, proteins (10 μg) in WB from *F. gigantica* as well as WB from other trematode, cestode and nematode parasites were separated by 12.5% SDS-PAGE (Laemmli, 1970), and transferred to nitrocellulose (NC) membranes (Bio-Rad, Philadelphia, PA, USA) for immunoblotting (Towbin *et al.* 1979). Each NC membrane was blocked with a blocking solution (5% skimmed milk in Tris buffered saline (TBS) pH 7.4 containing 0.05% Tween 20) at room temperature for 2 h, and incubated in undiluted MoAb 3A3 and myeloma culture fluid (as the negative control) at room temperature for 2 h. After washing three times with TBS, bound MoAb 3A3 on the NC membranes was detected by incubation with the horseradish peroxidase-conjugated goat anti-mouse immunoglobulin (Sigma-Aldrich Inc.), diluted to 1:4000 with 1% skimmed milk in TBS, containing 0.05% Tween 20, pH 7.4, at room temperature for 1 h. Thereafter, the NC membranes were washed and developed with specific TMB substrate solution at room temperature for 3–5 min. Finally, the reaction was stopped with distilled water.

Production of polyclonal antibody (PoAb) against rFgFABP. PoAb against rFgFABP was prepared by immunizing New Zealand White rabbits with rFgFABP as per the method of Anuracpreeda *et al.* (2013a, 2016b). Briefly, 500 μg rFgFABP in 500 μL PBS solution was mixed with an equal volume of complete Freund's adjuvant (Sigma-Aldrich Inc.), and injected subcutaneously into the rabbits. Two boosters followed at 3-week intervals with 250 μg rFgFABP in PBS emulsified in incomplete Freund's adjuvant (Sigma-Aldrich Inc.) via the same route. Blood samples were collected 1-week after final boost and tested for the antibody titers in the antisera.

Purification of MoAb 3A3 and PoAb against rFgFABP. The IgG fraction of both MoAb and PoAb was purified by 50% saturated ammonium sulphate, dialyzed against an excess of PBS and applied to an affinity chromatography in a Mab trap protein G Sepharose column (Amersham Pharmacia Biotech

AB, Uppsala, Sweden). Thereafter, the purified IgG of PoAb was conjugated with biotin using N-hydroxysuccinimidobiotin (Sigma Co) as described earlier by Anuracpreeda *et al.* (2016b).

Animals

Experimental mice. One hundred and twenty 5-week-old outbred ICR mice, from National Laboratory Animal Center, Mahidol University, Nakorn Pathom, Thailand, were used in this study. All animals were subdivided into two groups: control and infected groups, 60 control and 60 infected mice were made up of 5 subgroups (12 mice per group). Mice were sacrificed at 1, 4, 7, 21 and 35 days post infection. Each mouse of infected groups was orally infected with 30 metacercariae of *F. gigantica*, while control groups received only 0.5 mL of 0.85% NaCl solution. At necropsy day, all infected mice were anaesthetized and their peritoneal cavities were opened. The carcasses and organs were carefully examined for any pathological alterations and the presence of worms. The livers were teased and submerged in HBS solution and the parasites were recovered. Blood samples of all animals were collected aseptically into tubes without anticoagulant, and allowed to clot at room temperature for 1 h before being centrifuged at 3500 $\times g$ for collection of the sera. For cross-reactivity study, 20 sera were collected from mice infected with *S. mansoni* and 20 sera from hamsters infected with *Opisthorchis viverrini*. In addition, 60 sera from non-infected hamsters were collected and used as the negative control.

Natural cattle. Serum samples from cattle with monoinfections of *F. gigantica*, other trematode, cestode, nematode parasites and non-infected cattle were obtained from fields in many zones of Thailand. Sixty fasciolosis sera were obtained from cattle with confirmed *F. gigantica* infection using a standard parasitological method. Fecal samples of the animals were examined for *Fasciola* eggs by sedimentation method (Soulsby, 1965) and for other parasite eggs by flotation method (Hammond and Sewell, 1972). Sera from *P. cervi* infection (paramphistomosis, $n = 50$), *M. benedeni* infection (monieziaiasis, $n = 10$), strongylid infections ($n = 10$), *Trichuris* sp. infection (trichuriasis, $n = 10$) and *Strongyloides* sp. infection (strongyloidiasis, $n = 10$) were tested for the cross reactivity study. Negative control sera were also collected from non-infected cattle ($n = 60$) whose stool samples at the time of blood collection contained no parasite eggs.

The lower detection limit and the specificity of sandwich ELISA. The recombinant FABP (rFABP) and WB antigens of metacercariae, NEJ, 1, 3, 5-week-old juveniles, adults and TA as well as ES antigens of adult *F. gigantica* were serially diluted in a

solution containing 1% BSA-0.05% PBST and tested to evaluate the lower detection limit of sandwich ELISA. The OD values were examined and correlated with the amounts of antigen. The end point of detection limit was judged to be the lowest amount of antigen still showing the positive OD values. To determine the specificity of ELISA, WB antigens from other trematode, cestode and nematode parasites were prepared at various concentrations and used to detect possible presence of FABP antigen.

Detection of circulating FABP antigen by sandwich ELISA. The sandwich ELISA was performed as per the method described previously by Anuracpreeda *et al.* (2016b). A 50 μL of rabbit anti mouse IgG (Dako A/S, Glostrup, Denmark) in coating buffer (10 $\mu\text{g mL}^{-1}$) was coated in each well of a flat bottom F96 micro-ELISA plate (Nunc A/S, Roskilde, Denmark), and incubated at 4 °C overnight. To remove excess antibody, the plate was washed with a washing buffer containing 0.05% Tween 20 in normal saline solution (NSST) at room temperature for 1 min. Thereafter, 50 μL of purified MoAb 3A3 diluted in 1% BSA in PBS pH 7.2 (10 $\mu\text{g mL}^{-1}$) was applied to each well of the precoated plate, and incubated at 37 °C for 3 h. After three washings with the same washing buffer, 150 μL of 5% skim milk in PBS was added and incubated at 37 °C for 1 h to block the unbound sites. The wells were washed three times with washing buffer, and 50 μL of reference antigens or samples in 1% BSA-0.05% Tween 20 in PBS (PBST) was added to triplicate wells, and then incubated at 4 °C overnight. After washing again, 50 μL of biotinylated rabbit IgG antibody against rFgFABP (2 $\mu\text{g mL}^{-1}$ of 1% BSA-0.05% Tween 20 in PBST) was added to the wells and allowed to react at 37 °C for 90 min. Afterward, the wells were washed and followed by the addition of 50 μL of streptavidin-conjugated peroxidase (Zymed Laboratory Inc.) diluted 1:6000 in 1% BSA-0.05% PBST. The plate was incubated at 37 °C for 1 h and washed as described earlier. A 50 μL of TMB substrate solution was added to each well, and after 10 min the reaction was stopped with 50 μL of 1 N HCl. Finally, the OD was measured at 450 nm using an ELISA reader (Multiskan Ascent, Labsystems, Helsinki, Finland).

Detection of the antibody against FABP by indirect ELISA. Indirect ELISA used for detection of the antibody against FABP in the sera of infected animals was followed according to the method described by Anuracpreeda *et al.* (2016b). ELISA plate was coated with 1 $\mu\text{g mL}^{-1}$ of rFgFABP (100 $\mu\text{L well}^{-1}$) in coating buffer, and incubated at 4 °C overnight. IgG antibodies were detected with horseradish peroxidase-conjugated goat anti-mouse immunoglobulin.

Assessment of results and statistical analysis. The ELISA cut-off value between negative and positive samples was calculated as the average of the OD of control non-infected sera plus three times the standard deviations (s.d.). Each serum sample was tested in triplicate and expressed as an individual mean OD. All data from the detection of FABP in the serum samples of experimentally infected mice and naturally infected cattle were calculated and analyzed with independent-sample *t*-test using SPSS for Windows program version 19.0 (SPSS Inc., Chicago, Illinois). The *P*-value greater than 0.05 was considered not significant and less than 0.05 and 0.01 was considered to be highly and very highly significant, respectively. The method of Galen (1980) was used to calculate the diagnostic sensitivity, specificity, positive and negative predictive values, false positive and negative rate, as well as accuracy. These values were calculated using the formulas as follows: sensitivity = [no. of true positives / (no. of true positives + no. of false negatives)] \times 100, specificity = [no. of true negatives / (no. of true negatives + no. of false positives)] \times 100, positive predictive value = [no. of true positives / (no. of true positives + no. of false positives)] \times 100, negative predictive value = [no. of true negatives / (no. of true negatives + no. of false negatives)] \times 100, false positive rate = [no. of false positives / (no. of false positives + no. of true negatives)] \times 100, false negative rate = [no. of false negatives / (no. of false negatives + no. of true positives)] \times 100, and accuracy = [all with true positives and negatives / all test done] \times 100. The primary data of the sandwich ELISA are as follows: true negative = number of control samples (other parasitosis and non-infected controls) that show negative result, true positive = number of proven *F. gigantica* infection samples that show positive result, false positive = number of control samples that show positive result and false negative = number of proven *F. gigantica* infection samples that show negative result.

RESULTS

Expression and purification of rFgFABP

The cDNA sequence encoding FgFABP was subcloned in the expression vector, pET-30b vector (Novagen) and transformed into *E. coli* BL21 (DE). The rFgFABP was purified by using Ni-NTA affinity chromatography (QIAGEN) and analyzed by SDS-PAGE. The calculated molecular weight (MW) of rFgFABP appeared at approximately 20 kDa (Supplementary Fig. S1).

MoAbs against rFgFABP

Six stable hybridoma clones of MoAbs against rFgFABP, designated 1B6, 2H8, 3A3, 4G9, 5C3, 6D1, were produced. They were selected and

expanded in culture flasks to obtain large volume of MoAb, which were then collected for further experiments. The immunoglobulin isotypes of all MoAbs were found to be IgG₁ and κ light chain. Clone 3A3 had the highest titre (up to 3.65 in ELISA OD reading at 450 nm with the cut off point at 0.15); hence, it was used in this study.

Reactivity and specificity of MoAb 3A3

The native FgFABP in WB, TA and ES fractions were reacted with MoAb 3A3 and the relative levels of FgFABP in each fraction were assessed by indirect ELISA. The levels of reactivity of FgFABP in WB, TA and ES with MoAb 3A3 were significantly higher when compared with control myeloma culture fluid (Fig. 1). In both indirect ELISA (Fig. 2) and immunoblotting assay (Fig. 3A and B) were used for studying the specificity of MoAb 3A3. MoAb 3A3 exhibited strong reaction with FABP antigen in WB of *F. gigantica* at MW of 14.5 kDa, while showing no cross-reaction with WB antigens from 15 trematodes (*F. hepatica*, *O. viverrini*, *E. pancreaticum*, *G. explanatum*, *S. spindale*, *S. mansoni*, *S. japonicum*, *S. mekongi*, *P. cervi*, *P. gracile*, *F. cobboldi*, *G. crumenifer*, *C. spatiosus*, *C. gregarious* and *C. cotylophorum*), from two cestodes (*M. benedeni* and *A. centripunctata*), as well as from three nematodes (*Trichuris* sp., *H. placei* and *S. labiato-papillosa*).

The lower detection limit and the specificity of sandwich ELISA

The ELISA cut-off value was 0.15. The lower detection limit of this assay was evaluated from the different concentrations of rFABP, WB, TA and ES antigens of adult *F. gigantica*. Based on the lowest concentrations of antigen that still exhibited the sensitivity of test, this test could detect rFABP and FABP in WB, TA and ES fractions of *F. gigantica* at the lowest concentrations of 5, 50, 100 and 200 pg mL⁻¹, respectively (Fig. 4A). Likewise, the lowest dilution of WB in Met and NEJ antigens was at 400 pg mL⁻¹, and for 1-, 3-, 5-week-old juveniles and adult antigens was at 50 pg mL⁻¹ (Fig. 4B). Moreover, this sandwich ELISA described herein was highly specific for FABP antigen, as no cross-reactivity was demonstrated when the assay was employed to detect this antigen at various concentrations of other parasite antigens (Table 1).

Application of sandwich ELISA for detection of circulating FABP antigen in sera from infected animals

Experimentally infected animals. At day 35 post infection, the morbidity rate of infected mice was 100% and no mortality has occurred throughout




Fig. 1. The levels of the native FABP in WB, TA and ES fractions of adult *F. gigantica* was evaluated by its reactivity with MoAb 3A3 using indirect ELISA. Significant increase of OD values representing FABP levels were shown in WB, TA and ES antigens (red column) when compared with the control myeloma culture fluid (green column). The result was considered significant if *P*-value is less than 0.05 ($P < 0.05$) represented in asterisk (*). ELISA, enzyme-linked immunosorbent assay; FABP, fatty acid binding protein; OD, optical density; MoAb, monoclonal antibody; WB, whole body; TA, tegumental antigens; ES, excretory-secretory.

this study. The parasite recoveries varied from 5 to 19 worms per mouse with an average of 10.92 ± 4.17 . For liver pathology, the liver were swollen and covered with large patches of fibrinopurulent exudates that later caused adhesion of all of the liver lobes with adjacent organs in the abdominal cavity. Likewise, numerous white tracks and foci were seen and appeared as irregular lines that scattered throughout the centre of the hepatic lobes. The mean ODs for antigen detection were significantly different from those of control sera at day 1, 4 ($P < 0.05$), 7, 21 and 35 ($P < 0.01$) post infection. The levels of detectable antigen in infected sera peaked at day 21 to day 35 post infection (Fig. 5A). The total numbers of *F. gigantica* metacercariae-infected mice sera and the numbers as well as the percentages of positive sera at day 1, 4, 7, 21 and 35 post infection were 83.33% (10 out of 12), 83.33% (10 out of 12), 100% (12 out of 12), 100% (12 out of 12) and 100% (12 out of 12), respectively (Fig. 5B).

Additionally, sera from, 60 fasciolosis mice, 20 schistosomiasis mice, 20 sera opisthorchiasis hamsters, as well as 60 and 60 non-infected healthy mice and hamsters, were examined. The results revealed that 56 of 60 (93.3%) fasciolosis sera were positive, whereas all 160 (100%) sera from those with other infections and non-infected healthy animals were negative (Fig. 6A). Therefore, this




Fig. 2. Mean ELISA OD values of cross-reactivities studies between MoAb 3A3 and the panel of antigen from various trematode, cestode and nematode parasites. Fg, *F. gigantica*; Fh, *F. hepatica*; Ov, *O. viverrini*; Ep, *E. pancreaticum*; Ge, *G. explanatum*; Ss, *S. spindale*; Sm, *S. mansoni*; Sj, *S. japonicum*; Sme, *S. mekongi*; Pc, *P. cervi*; Pg, *P. gracile*; Fc, *F. cobboldi*; Gc, *G. crumenifer*; Cs, *C. spatiosus*; Cg, *C. gregarious*; Cc, *C. cotylophorum*; Mb, *M. benedeni*; Ac, *A. centripunctata*; Ts, *Trichuris* sp.; Hp, *H. placei*; Sp, *S. labiato-papillosa*. These parasites' antigens were allowed to react with undiluted MoAb culture fluid (Undil) and MoAb at dilution 1:10, 1:100, 1:500, 1:1000 and 1:10 000. Horizontal dotted line is the cut off level, i.e. the mean OD of control sera plus 3 s.d. The OD values greater than this cut-off value are considered to be positive detection. ELISA, enzyme-linked immunosorbent assay; OD, optical density; MoAb, monoclonal antibody.

assay showed a sensitivity and specificity of 93.3 and 100%, respectively, with a positive predictive value of 100%, a negative predictive value of 97.6%, false positive rate of 0%, false negative rate of 6.7% and an accuracy of 98.2% (Table 2).

Naturally infected animals. Two hundred and ten sera from cattle were collected and tested by sandwich ELISA. Sixty sera from cattle infected with fasciolosis, 50 from cattle infected with paramphistomosis, 10 from cattle infected with moniezia, 10 from cattle infected with strongylid infections, 10 from cattle infected with trichuriasis, 10 from cattle infected with strongyloidiasis and 60 from non-infected cattle were examined. The results exhibited that 58 of 60 (96.7%) fasciolosis sera were positive, while all 90 sera from those infected with other parasites, and all 60 sera from non-infected controls were negative (Fig. 6B). Hence, the method showed a sensitivity and specificity of 96.7 and 100%, respectively, with a positive predictive value of 100%, a negative predictive value of 98.7%, a false positive rate of 0%, a false negative rate of 3.3% and an accuracy of 99.1% (Table 2).

Indirect ELISA for the detection of antibody against *F. gigantica* FABP. The numbers of serum samples from infected mice being tested and percentages of positive detection during the period of 1–35 days post infection were 8.3% (1 out of 12) at day 1,

16% (2 out of 12) at day 4, 33.3% (4 out of 12) at day 7, 33.3% (4 out of 12) at day 21 and 75% (9 out of 12) at day 35 (Fig. 5B). The mean OD for mouse antibody diagnosed fasciolosis at 35 days post infection is significantly different from that of the control sera OD ($P < 0.05$).

DISCUSSION

The problem of precisely diagnosing fasciolosis using the conventional parasitological tests, including microscopic detection of the fluke's eggs in the feces and detection of serum antibodies, has led to develop a more accurate diagnostic method. In this study, we have detected circulating *F. gigantica* FABP antigen in the sera from both *F. gigantica* experimentally infected mice and naturally infected cattle using a MoAb-based sandwich ELISA. The MoAb 3A3 specific to FgFABP exhibited no cross-reactivities with antigens of other adult parasites, including *F. hepatica*, *O. viverrini*, *E. pancreaticum*, *G. explanatum*, *S. spindale*, *S. mansoni*, *S. japonicum*, *S. mekongi*, *P. cervi*, *P. gracile*, *F. cobboldi*, *G. crumenifer*, *C. spatiosus*, *C. gregarious*, *C. cotylophorum*, *M. benedeni*, *A. centripunctata*, *Trichuris* sp., *H. placei* and *S. labiato-papillosa*. Hence, we could use this MoAb 3A3 to detect the circulating FABP antigen in both early and late stages of infection. In our earlier study, we have also produced MoAb specific to




Fig. 3. The specificity of MoAb 3A3 is tested with WB antigens from *F. gigantica*, other trematode, cestode and nematode parasites by using immunoblotting technique. (A) Immunoblot analysis showing the cross-reactivities of MoAb clone 3A3 with WB antigens from Fg, *F. gigantica* (lane 2); Fh, *F. hepatica* (lane 3); Ov, *O. viverrini* (lane 4); Ep, *E. pancreaticum* (lane 5); Ge, *G. explanatum* (lane 6); Ss, *S. spindale* (lane 7); Sm, *S. mansoni* (lane 8); Sj, *S. japonicum* (lane 9); Sme, *S. mekongi* (lane 10); Pc, *P. cervi* (lane 11); Pg, *P. gracile* (lane 12). (B) Immunoblot analysis of the cross-reactivities of MoAb 3A3 with WB antigens from *F. gigantica* (lane 2), Fc, *F. cobboldi* (lane 3); Gc, *G. crumenifer* (lane 4); Cs, *C. spatiosus* (lane 5); Cg, *C. gregarious* (lane 6); Cc, *C. cotylophorum* (lane 7); Mb, *M. benedeni* (lane 8); Ac, *A. centripunctata* (lane 9); Ts, *Trichuris* sp. (lane 10); Hp, *H. placei* (lane 11); Sp, *S. labiato-papillosa* (lane 12). Lane 1 of (A) and (B) is WB antigen from *F. gigantica* (Fg) blotted with the MCF, which is used as the negative control. STD is the lane containing standard protein molecular weight markers. MCF, myeloma culture fluid; MoAb, monoclonal antibody; WB, whole body.

28.5 kDa tegumental antigen (28.5 kDa TA), recombinant cathepsin B3 (rCatB3) and recombinant cathepsin L1 (rCatL1) of adult *F. gigantica* as well as developed reliable MoAb-based sandwich ELISA for diagnosis of fasciolosis by *F. gigantica* in experimentally infected mice and naturally infected cattle sera (Anuracpreeda *et al.* 2009a, 2013a, 2016b). Close to our findings, Viyanant *et al.* (1997) produced MoAb against TA of *F. gigantica* whose target antigen was recognized at MW of 66 kDa. However, they reported that cross reactivity




Fig. 4. The lowest concentrations of rFABP and FABP in WB of various stages of *F. gigantica*, TA and ES antigens of *F. gigantica* as examined by the sandwich ELISA assay. (A) Lines with black circle, white circle, black triangle and white triangle indicate the concentration levels of rFABP and FABP in WB, TA and ES antigens of adult *F. gigantica*. The arrows indicate the lowest concentrations of FABP that could still be detected. (B) Lines with black circle, white triangle, black square, white diamond, black triangle and white hexagon exhibit the concentration levels of FABP in WB fractions of adult, 5-, 3-, 1-week-old juveniles, NEJ and metacercariae (Met) of *F. gigantica*, respectively. The arrows indicate the lowest concentrations that FABP could still be detected in each sample. ELISA, enzyme-linked immunosorbent assay; rFABP, recombinant fatty acid binding protein; MoAb, monoclonal antibody; WB, whole body; TA, tegumental antigens; ES, excretory-secretory; NEJ, Newly excysted juveniles.

with other parasite antigens from *Paramphistomum* sp., *S. spindale*, *S. mansoni* and *S. mekongi* was occurred and the MoAb clone was not stable. Fagbemi *et al.* (1997) used MoAbs against whole worm antigens of *F. gigantica* for the diagnosis of fasciolosis in cattle; however, this MoAb exhibited more cross-reactivities with other antigens from *P. microbothrium*, *Dicrocoelium hospes* and *Schistosoma bovis*. Likewise, Arafa *et al.* (1999) reported that the diagnosis of human fasciolosis was developed using MoAbs against *F. gigantica* ES

Table 1. Specificity testing of a sandwich ELISA to various crude preparations from trematode, cestode and nematode parasite antigens

Antigens ^a	OD at 450 nm ^b
Trematodes	
<i>F. gigantica</i> , rFABP	3.511
<i>F. gigantica</i>	3.121
<i>F. hepatica</i>	0.089
<i>O. viverrini</i>	0.081
<i>E. pancreaticum</i>	0.071
<i>G. explanatum</i>	0.076
<i>S. spindale</i>	0.081
<i>S. mansoni</i>	0.079
<i>S. japonicum</i>	0.085
<i>S. mekongi</i>	0.083
<i>P. cervi</i>	0.074
<i>P. gracile</i>	0.073
<i>F. cobboldi</i>	0.086
<i>C. spatiosus</i>	0.069
<i>C. gregarious</i>	0.071
<i>C. cotylophorum</i>	0.078
Cestodes	
<i>M. benedeni</i>	0.069
<i>A. centripunctata</i>	0.067
Nematodes	
<i>Trichuris</i> sp.	0.077
<i>H. placei</i>	0.068
<i>S. labiato-papillosa</i>	0.071
Reagent blank	0.159

^a The protein content of each parasite antigen preparation was adjusted to 20 µg mL⁻¹, and a 50-µL volume was used for analysis.
^b Mean OD was determined in triplicates performed on three separate occasions.

antigens at MW of 49.5 kDa. It was found that cross-reactivity with *S. mansoni* antigen occurred by using these MoAbs. Another study of *F. gigantica* revealed that MoAb against isoforms of cathepsin Ls of juvenile and adult fluke was used for diagnosis of animal fasciolosis (Wongwairoot *et al.* 2015).

In this study, the results indicated the advantage of using the rabbit anti-mouse IgG to precoat the plate, which helped to enhance the binding of MoAb 3A3, and hence also the binding to antigen. Consequently, the assay was able to detect rFABP and FABP antigens in WB, TA and ES fractions of *F. gigantica* at the concentrations as low as 5, 50, 100 and 200 pg mL⁻¹, respectively. In addition, the FABP antigen could also be detected in WB of Met and NEJ at 400 pg mL⁻¹, and in WB of 1-, 3-, 5-week-old juveniles and adult antigens 50 pg mL⁻¹. Furthermore, the use of MoAb as the antigen-capturing antibody increased the specific binding of the PoAb, which could help to reduce the background and yielded very low detection limits of the assay. The detection limits of our assay are lower than those of Langley and Hillyer (1989) who detected *F. hepatica* ES antigens in serum samples of experimentally infected mice at a concentration 0.25 ng mL⁻¹. Also, the detection




Fig. 5. Detection of circulating FABP antigens and antibody against FABP antigens of *F. gigantica* is evaluated by ELISA assay. (A) Detection of circulating FABP antigens of *F. gigantica* in the serum samples of the infected mice as compared with non-infected mice by sandwich ELISA. Black stars represent OD values of individual mouse serum; white squares indicate mean OD values of all mice in the experimental group. The horizontal dotted line denotes the cut-off level. Any values above this line are considered positive detection. (B) Comparison between detection of circulating FABP antigens using sandwich ELISA and antibody against FABP antigens using indirect ELISA in the sera of mice infected with *F. gigantica*. ELISA, enzyme-linked immunosorbent assay; FABP, fatty acid binding protein; OD, optical density.

limits are better than the results reported by Mezo *et al.* (2004) who developed the capture ELISA for the detection of *F. hepatica* ES antigens in fecal supernatants of infected animals, and found that the detection limit was 0.3 (for sheep) and 0.6 ng mL⁻¹ (for cattle). In our previous study, we have developed a sandwich ELISA for the detection of circulating 28.5 kDa TA in the serum samples of mice experimentally infected with *F. gigantica*. The result revealed that the lower detection limit was 600 pg mL⁻¹ (for TA), 16 ng mL⁻¹ (for WB antigen), and 60 ng mL⁻¹ (for ES antigen), which are considerably higher than in the present study (Anuracpreeda *et al.* 2009a). Likewise, a sandwich




Fig. 6. Sandwich ELISA reactivity pattern denotes (A) the relative levels of circulating FABP antigens (OD values) in sera from mice infected with *F. gigantica* (fasciolosis), *S. mansoni* (schistosomiasis) and hamster infected with *O. viverrini* (opisthorchiasis). Serum samples from non-infected mice and hamsters were used as controls. The horizontal dotted line represents the cut-off level for a positive detection. (B) The relative levels of circulating FABP antigens (OD values) in sera from cattle naturally infected with *F. gigantica* (fasciolosis), *P. cervi* (paramphistomosis), *M. benedeni* (Moniezia), Strongylids (strongylid infection), *Trichuris* sp. (trichuriasis), and *Strongyloides* sp. (strongyloidiasis). Sera from non-infected cattle were used as controls. The horizontal dotted line exhibits the cut-off level for a positive detection. ELISA, enzyme-linked immunosorbent assay; FABP, fatty acid binding protein; OD, optical density.

ELISA was used for detection of circulating CatB3 antigen in the sera of mice and cattle infected with *F. gigantica* with the lower detection limit at 10 (for rCatB3 antigen), 100 (for Met antigen) and 400 pg mL^{-1} (for NEJ antigen), which were still higher than in this study (Anuracpreeda *et al.* 2013a). Similarly, we have developed and used a sandwich ELISA for detection of rCatL1 and CatL1 in WB and ES fractions of *F. gigantica* with

the lower detection limit at 3, 50 and 100 pg mL^{-1} , respectively. The lowest detection limit for WB in Met, NEJ, 1-week-old juvenile antigens was at 100 pg mL^{-1} , and for 3-, 5-week-old juveniles and adult antigens was at 50 pg mL^{-1} (Anuracpreeda *et al.* 2016b). For human fasciolosis, the lowest detection limits reported herein are lower than that reported by Espino and Finlay (1994) who used a sandwich ELISA to detect the *F. hepatica* ES

Table 2. Calculation of diagnostic values of the sandwich ELISA for FABP antigen detection in sera of mice experimentally and cattle naturally infected with *F. gigantica*

Calculations	Experimentally infected mice (%)	Naturally infected cattle (%)
1. Sensitivity	93.3	96.7
2. Specificity	100	100
3. Positive predictive value	100	100
4. Negative predictive value	97.6	98.7
5. False positive rate	0	0
6. False negative rate	6.7	3.3
7. Accuracy	98.2	99.1

antigens in stool samples of patients, and found that the lower detection limit was 15 ng mL⁻¹, and that of Demerdash *et al.* (2011) who developed a sandwich ELISA to detect the *F. gigantica* ES antigens in both serum and stool samples of patients at a concentration level 3 ng mL⁻¹. In the present study, the sensitivity and specificity of this MoAb-based sandwich ELISA were also considered very high as in the sera of experimentally infected mouse at 93.3 and 100%, while in the sera of naturally infected cattle they were 96.7 and 100%, respectively. In addition, the accuracy of this assay in the sera of naturally infected cattle was 99.1%, which is comparable with that of experimentally infected mouse at 98.2%. Our findings indicated that this sandwich ELISA could be successfully applied to naturally infected cattle with a large body size.

In the present study, we have shown that the circulating FABP antigen in the sera of mice experimentally infected with *F. gigantica* was detectable as early as the first day after infection, with the peak levels occurred during the day 21 and day 35. This data herein is correlated with the reports of our previous studies (Anuracpreeda *et al.* 2016b). It is possible that the levels of the circulating antigen in infected serum samples were corresponded with the pattern of the parasites' migration in the host. During the first few weeks post infection, the antigen was slowly and continually released into the hosts' circulation. Once the worms reached and became mature stage in the biliary system of the liver, they might release a large amount of antigens into the hosts' circulation. On the other hand, Langley and Hillyer (1989) who used rabbit hyper-immune serum to detect circulating *F. hepatica* ES antigen in the sera from experimentally infected mouse at the first week after infection. As well, Viyanant *et al.* (1997) who developed a sandwich ELISA for detection of circulating 66 kDa TA in the sera of cattle experimentally infected with *F. gigantica* at the first week post infection. Another detections were also reported by Fagbemi *et al.* (1995) who developed and utilized a sandwich

ELISA for detection of circulating 88 kDa *F. gigantica* antigen in the sera of experimentally infected cattle at the second and third weeks post infection, and Velusamy *et al.* (2004) reported that the circulating 54 kDa *F. gigantica* antigen was detected in the sera of experimentally infected cattle at the second week post infection.

Concluding remarks

The results obtained in this study clearly indicated that a reliable MoAb-based sandwich ELISA showed high efficiencies and precisions. This assay method could be used as an important diagnostic tool not only for both early and late detections of fasciolosis but also for the seroepidemiological screening of animals, which in turn could contribute to the monitoring and control of the disease from different areas.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/S0031182016001104>.

ACKNOWLEDGEMENTS

We are grateful to Assoc. Prof. Dr. David Piedrafita, Federation University, Australia, for providing *F. hepatica* antigen.

FINANCIAL SUPPORT

This research was financially supported by Research Grants from The Thailand Research Fund and Mahidol University to Panat Anuracpreeda.

REFERENCES

- Anuracpreeda, P., Wanichanon, C., Chaithirayanon, K., Preyavichyapugdee, N. and Sobhon, P. (2006). Distribution of 28.5 kDa antigen in tegument of adult *Fasciola gigantica*. *Acta Tropica* **100**, 31–40.
- Anuracpreeda, P., Wanichanon, C. and Sobhon, P. (2008). *Paramphistomum cervi*: antigenic profile of adults as recognized by infected cattle sera. *Experimental Parasitology* **118**, 203–207.
- Anuracpreeda, P., Wanichanon, C., Chawengkirtikul, R., Chaithirayanon, K. and Sobhon, P. (2009a). *Fasciola gigantica*: immunodiagnosis of fasciolosis by detection of circulating 28.5 kDa tegumental antigen. *Experimental Parasitology* **123**, 334–340.
- Anuracpreeda, P., Wanichanon, C. and Sobhon, P. (2009b). *Fasciola gigantica*: immunolocalization of 28.5 kDa antigen in the tegument of metacercaria and juvenile fluke. *Experimental Parasitology* **122**, 75–83.
- Anuracpreeda, P., Songkoomkroong, S., Sethadavit, M., Chotwiwatthanakun, C., Tinikul, Y. and Sobhon, P. (2011). *Fasciola gigantica*: production and characterization of a monoclonal antibody against recombinant cathepsin B3. *Experimental Parasitology* **127**, 340–345.
- Anuracpreeda, P., Panyarachun, B., Ngamniyom, A., Tinikul, Y., Chotwiwatthanakun, C., Poljaroen, J. and Sobhon, P. (2012). *Fischoederius cobboldi*: a scanning electronmicroscopy investigation of surface morphology of adult rumen fluke. *Experimental Parasitology* **130**, 400–407.
- Anuracpreeda, P., Chawengkirtikul, R., Tinikul, Y., Poljaroen, J., Chotwiwatthanakun, C. and Sobhon, P. (2013a). Diagnosis of *Fasciola gigantica* infection using a monoclonal antibody-based sandwich ELISA for detection of circulating cathepsin B3 protease. *Acta Tropica* **127**, 38–45.

- Anuracpreeda, P., Poljaroen, J., Chotwiwatthanakun, C., Tinikul, Y. and Sobhon, P. (2013b). Antigenic components, isolation and partial characterization of excretion-secretion fraction of *Paramphistomum cervi*. *Experimental Parasitology* **133**, 327–333.
- Anuracpreeda, P., Srirakam, T., Pandonlan, S., Changklungmoa, N., Chotwiwatthanakun, C., Tinikul, Y., Poljaroen, J., Meemon, K. and Sobhon, P. (2014). Production and characterization of a monoclonal antibody against recombinant cathepsin L1 of *Fasciola gigantica*. *Acta Tropica* **135**, 1–9.
- Anuracpreeda, P., Phutong, S., Ngamniyom, A., Panyarachun, B. and Sobhon, P. (2015). Surface topography and ultrastructural architecture of the tegument of adult *Carnyverius spatiosus* Brandes, 1898. *Acta Tropica* **143**, 18–28.
- Anuracpreeda, P., Chawengkirttikul, R. and Sobhon, P. (2016a). Antigenic profile, isolation and characterization of whole body extract of *Paramphistomum gracile*. *Parasite Immunology* (In Press) doi: 10.1111/pim.12330.
- Anuracpreeda, P., Chawengkirttikul, R. and Sobhon, P. (2016b). Immunodiagnosis of *Fasciola gigantica* infection using monoclonal antibody-based sandwich ELISA and immunochromatographic assay for detection of circulating cathepsin L1 protease. *PLoS ONE* **11**, 1–22. e0145650.
- Anuracpreeda, P., Chawengkirttikul, R. and Sobhon, P. (2016c). Surface histology, topography, and ultrastructure of the tegument of adult *Orthocoleium parvipapillatum* (Stiles & Goldberger, 1910). *Parasitology Research* (In Press) doi: 10.1007/s00436-016-5024-3.
- Anuracpreeda, P., Chankaew, K., Puttarak, P., Koedrih, P., Chawengkirttikul, R., Panyarachun, B., Ngamniyom, A., Chanchai, S. and Sobhon, P. (2016d). The anthelmintic effects of the ethanol extract of *Terminalia catappa* L. leaves against the ruminant gut parasite, *Fischoederius cobboldi*. *Parasitology* **143**, 421–433.
- Arafa, M. S., Abaza, S. M., El-Shewy, K. A., Mohareb, E. W. and El-Moamly, A. A. (1999). Detection of *Fasciola*-specific excretory/secretory (E/S) protein fraction band (49.5 kDa) and its utilization in diagnosis of early fascioliasis using different diagnostic techniques. *Journal of the Egyptian Society of Parasitology* **29**, 911–926.
- Ashrafi, K., Valero, M. A., Panova, M., Periago, M. V., Massoud, J. and Mas-Coma, S. (2006a). Phenotypic analysis of adults of *Fasciola hepatica*, *Fasciola gigantica* and intermediate forms from the endemic region of Gilan. *Parasitology International* **55**, 249–260.
- Ashrafi, K., Valero, M. A., Massoud, J., Sobhani, A., Solaymani-Mohammadi, S., Conde, P., Khoubbane, M., Bargues, M. D. and Mas-Coma, S. (2006b). Plant-borne human contamination by fascioliasis. *American Society of Tropical Medicine and Hygiene* **75**, 295–302.
- Chicz, R. M. (1994). Submitted to the Protein Sequence Database, August 1994. Accession No. A44638.
- Demerdash, Z. A., Diab, T. M., Aly, I. R., Mohamed, S. H., Mahmoud, F. S., Zoheiry, M. K., Mansour, W. A., Attia, M. E. and El-Bassiouny, A. E. (2011). Diagnostic efficacy of monoclonal antibody based sandwich enzyme linked immunosorbent assay (ELISA) for detection of *Fasciola gigantica* excretory/secretory antigens in both serum and stool. *Parasites and Vectors* **4**, 176.
- Espino, A. and Finlay, C. (1994). Sandwich enzyme-linked immunosorbent assay for detection of excretory/secretory antigens in humans with fascioliasis. *Journal of Clinical Microbiology* **32**, 190–193.
- Fagbemi, B. O. and Obarisiagbon, I. O. (1990). Comparative evaluation of the enzyme-linked immunosorbent assay (ELISA) in the diagnosis of natural *Fasciola gigantica* infection in cattle. *Veterinary Quarterly* **12**, 35–38.
- Fagbemi, B. O., Obarisiagbon, I. O. and Mbuh, J. V. (1995). Detection of circulating antigen in sera of *Fasciola gigantica*-infected cattle with antibodies reactive with a *Fasciola*-specific 88-kDa antigen. *Veterinary Parasitology* **58**, 235–246.
- Fagbemi, B. O., Aderibigbe, O. A. and Guobadia, E. E. (1997). The use of monoclonal antibody for the immunodiagnosis of *Fasciola gigantica* infection in cattle. *Veterinary Parasitology* **69**, 231–240.
- Galen, R. S. (1980). Predictive value and efficiency of laboratory testing. *Pediatric Clinics of North America* **27**, 861–869.
- Ghosh, S., Rawat, P., Velusamy, R., Joseph, D., Gupta, S. C. and Singh, B. P. (2005). 27 kDa *Fasciola gigantica* glycoprotein for the diagnosis of prepatent fasciolosis in cattle. *Veterinary Research Communications* **29**, 123–135.
- Grams, R., Vichasri-Grams, S., Sobhon, P., Upatham, E. S. and Viyanant, V. (2000). Molecular cloning and characterization of antigen encoding genes from *Fasciola gigantica*. In *International Proceedings of the Second Congress of the Federation of Immunological Societies of Asia-Oceania* (ed. Sirisinha, S., Chaiyaroj, S. C. and Tapchaisri, P.), pp. 39–43. Monduzzi Editore, Bangkok, Thailand, January 23–27. Bologna, Italy.
- Guobadia, E. E., and Fagbemi, B. O. (1995). Immunodiagnosis of fasciolosis in ruminants using a 28-kDa cysteine protease of *Fasciola gigantica* adult worms. *Veterinary Parasitology* **57**, 309–318.
- Hammond, J. A. and Sewell, M. M. H. (1972). Flotation on the Sellotape (demonstration). *Transactions of the Royal Society of Tropical Medicine and Hygiene* **66**, 547.
- Hillyer, G. V. (1999). Immunodiagnosis of human and animal fasciolosis. In *Fasciolosis* (ed. Dalton, J. P.), pp. 435–447. CABI Publishing, Oxon, UK.
- Hillyer, G. V., del Llano de Diaz, A. and Reyes, C. N. (1977). *Schistosoma mansoni*: acquired immunity in mice and hamsters using antigens of *Fasciola hepatica*. *Experimental Parasitology* **42**, 348–355.
- Kumar, N., Ghosh, S. and Gupta, S. C. (2008). Early detection of *Fasciola gigantica* infection in buffaloes by enzyme-linked immunosorbent assay and dot enzyme-linked immunosorbent assay. *Parasitology Research* **103**, 141–150.
- Langley, R. J. and Hillyer, G. V. (1989). Detection of circulating parasite antigen in murine fascioliasis by two-site enzyme-linked immunosorbent assays. *American Journal of Tropical Medicine and Hygiene* **41**, 472–478.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Le, T. H., De, N. V., Agatsuma, T., Blair, D., Vercruysse, J., Dorny, P., Nguyen, T. G. and McManus, D. P. (2007). Molecular confirmation that *Fasciola gigantica* can undertake aberrant migrations in human hosts. *Journal of Clinical Microbiology* **45**, 648–650.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. and Randal, R. J. (1951). Protein measurement with the Folin phenol reagent. *Journal of Chemical Biology* **193**, 265–275.
- Mas-Coma, S., Bargues, M. D. and Valero, M. A. (2005). Fascioliasis and other plant-borne trematode zoonoses. *International Journal for Parasitology* **35**, 1255–1278.
- Mas-Coma, S., Valero, M. A. and Bargues, M. D. (2009). Chapter 2. *Fasciola*, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Advances in Parasitology* **69**, 41–146.
- Mbuh, J. V. and Fagbemi, B. O. (1996). Antibody and circulating antigen profiles before and after chemotherapy in goats infected with *Fasciola gigantica*. *Veterinary Parasitology* **66**, 171–179.
- Meyer, F., Myer, H., and Bueding, E. (1970). Lipid metabolism in the parasitic and free-living flatworms, *Schistosoma mansoni* and *Dugesia dorotocephala*. *Biochimica et Biophysica Acta*. **210**, 257–266.
- Mezo, M., Gonzalez-Warleta, M. and Ubeira, F. M. (2004). An ultra-sensitive capture ELISA for detection of *Fasciola hepatica* coproantigens in sheep and cattle using a new monoclonal antibody (MM3). *Journal of Parasitology* **90**, 845–852.
- Panyarachun, B., Sobhon, P., Yotsawan, T., Chotwiwatthanakun, C., Anunpinit, V. and Anuracpreeda, P. (2010). *Paramphistomum cervi*: surface topography of the tegument of adult fluke. *Experimental Parasitology* **125**, 95–99.
- Panyarachun, B., Ngamniyom, A., Sobhon, P. and Anuracpreeda, P. (2013). Morphology and histology of the adult *Paramphistomum gracile* Fischöder, 1901. *Journal of Veterinary Science* **14**, 425–432.
- Rodriguez-Perez, J., Rodriguez-Medina, J. R., Garcia-Blanco, M. A. and Hillyer, G. V. (1992). *Fasciola hepatica*: molecular cloning, nucleotide sequence, and expression of a gene encoding a polypeptide homologous to a *Schistosoma mansoni* fatty acid binding protein. *Experimental Parasitology* **74**, 400–407.
- Sirisiro, A., Grams, R., Vichasri-Grams, S., Ardseungneon, P., Pankao, V., Meepool, A., Chaithirayanon, K., Viyanant, V., Tan-Ariya, P., Upatham, E. S. and Sobhon, P. (2002). Production and characterization of a monoclonal antibody against recombinant fatty acid binding protein of *Fasciola gigantica*. *Veterinary Parasitology* **105**, 119–129.
- Smooker, P. M., Hickford, D. E., Vaiano, S. A., and Spithill, T. W. (1997). Isolation, cloning and expression of fatty acid binding proteins from *Fasciola gigantica*. *Experimental Parasitology* **85**, 89–91.
- Sobhon, P., Anantavara, S., Dangprasert, T., Viyanant, V., Krailas, D., Upatham, E. S., Wanichanon, C., Kusamran, T. (1998). *Fasciola gigantica*: studies of the tegument as a basis for the developments of immunodiagnosis and vaccine. *Southeast Asian Journal of Tropical Medicine and Public Health* **29**, 387–400.
- Soulsby, E. J. L. (1965). *Textbook of Veterinary Clinical Parasitology*. I. Helminths. Blackwell Scientific Publications, Oxford, p. 1120.
- Spithill, T. W., Smooker, P. M. and Copeman, D. B. (1999). *Fasciola gigantica*: epidemiology, control, immunology and molecular biology. In *Fasciolosis* (ed. Dalton, J. P.), pp. 465–525. CABI Publishing, Oxon.
- Sriveny, D., Raina, O. K., Yadav, S. C., Chandra, D., Jayraw, A. K., Singh, M., Velusamy, R., and Singh, B. P. (2006). Cathepsin L cysteine

proteinase in the diagnosis of bovine *Fasciola gigantica* infection. *Veterinary Parasitology* **135**, 25–31.

Swarup, D., Pachauri, S. P., Sharma, B., and Bandhopadhyay, S. K. (1987). Serodiagnosis of *Fasciola gigantica* infection in buffaloes. *Veterinary Parasitology* **24**, 67–74.

Torgerson, P. and Claxton, J. (1999). Epidemiology and control. In *In Fasciolosis* (ed. Dalton, J. P.), pp. 113–149. CABI Publishing, Oxon.

Towbin, H., Staehelin, T. and Gordon, J. (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheet: procedure and some applications. *Proceedings of the National Academy of Sciences of the United States of America* **76**, 4350–4354.

Velusamy, R., Singh, B. P., Sharma, R. L. and Chanda, D. (2004). Detection of circulating 54 kDa antigen in sera of bovine calves experimentally infected with *F. gigantica*. *Veterinary Parasitology* **119**, 187–195.

Viyanant, V., Krailas, D., Sobhon, P., Upatham, E. S., Kusamran, T., Chompoochan, T., Thammasart, S. and Prasittirat, P. (1997). Diagnosis of cattle fasciolosis by the detection of a circulating antigen using a monoclonal antibody. *Asian Pacific Journal of Allergy and Immunology* **15**, 153–159.

Wongwairo, S., Kueakhai, P., Changklungmoa, N., Jaikua, W., Sansri, V., Meemon, K., Songkoomkroong, S., Riengrojpitak, S. and Sobhon, P. (2015). Monoclonal antibody against recombinant *Fasciola gigantica* cathepsin L1H could detect juvenile and adult cathepsin Ls of *Fasciola gigantica*. *Parasitology Research* **114**, 133–140.

Zimmerman, G. L., Nelson, M. J. and Clark, C. R. (1985). Diagnosis of ovine fascioliasis by a dot enzyme-linked immunosorbent assay: a rapid microdiagnostic technique. *American Journal of Veterinary Research* **46**, 1513–1515.