

Pandora's Box highlights research findings, news and matters of interest from around the world. Most of these article types are relatively brief, typically up to 1500 words with up to 12 references. In addition to printed content, *BJPsych International* has increased its online visibility through the creation of videos and podcasts (<https://www.cambridge.org/core/journals/bjpsych-international/videos-and-podcasts#videos>) and its popular arts blog *Muses* (<https://www.cambridge.org/core/blog/tag/muses-the-arts-blog/>).

The time has now come for *BJPsych International* to expand into publishing original research and systematic reviews and meta-analyses, directly contributing to building an evidence base for good practice and policy worldwide, with a special focus on countries that have historically contributed little to the international scientific literature. It has been recognised that a significant increase in the word count for these papers would be necessary in order for the journal to attract top-quality submissions. To that end, two new article types will be introduced: Original Articles, up to 4000 words with up to 40 essential references; and Reviews, up to 6000 words with up to 150 essential references. Together with the fact that *BJPsych International* is open access, not charging a publication fee to the authors of accepted papers or to its readers, this will further elevate the journal to a key

publication in the field of international psychiatry and mental health, facilitating cross-fertilisation without barriers across all cultures and continents.

Summary

Capitalising on its strengths, *BJPsych International* continues its exciting journey of growth and development. With a prominent online presence, a large readership, clinically relevant content and a high commitment to making a difference, the journal is placed in an excellent position to serve its mission. The inclusion of original research and systematic reviews and meta-analyses will further advance its impact; we welcome submissions of such papers from around the world!

Declaration of interest




M.K. is Editor in Chief of *BJPsych International*.

References

- 1 World Health Organization. *World Mental Health Report: Transforming Mental Health for All*. WHO, 2022 (<https://iris.who.int/bitstream/handle/10665/356119/9789240049338-eng.pdf?sequence=1>) [cited 17 Feb 2024].
- 2 World Health Organization. *Comprehensive Mental Health Action Plan 2013–2030*. WHO, 2021 (<https://iris.who.int/bitstream/handle/10665/345301/9789240031029-eng.pdf?sequence=1>) [cited 17 Feb 2024].



The development of the Helping your Anxious Child programme: a parent-mediated group intervention for parents of autistic children in South Asia

Caitlin Kittridge,¹  Priyanka Rob,² Amy Fisher-Rogers,³ Tarana Anis,⁴ Udena Attygalle,⁵ Farzana Islam,⁶ Aditya Narain Sharma⁷  and Jacqui Rodgers⁸ 

¹Research Assistant, School of Psychology, Newcastle University, Newcastle Upon Tyne, UK

²Research Assistant, School of Psychology, Newcastle University, Newcastle Upon Tyne, UK

³Research Assistant, School of Psychology, Newcastle University, Newcastle Upon Tyne, UK

⁴Senior Child Psychologist, Child Development Centre, Evercare Hospital, Dhaka, Bangladesh

⁵Consultant Child and Adolescent Psychiatrist, Sirimavo Bandaranayake Children's Hospital, Peradeniya, Sri Lanka

⁶Specialist, Child Development Centre, Evercare Hospital, Dhaka, Bangladesh

Autistic children are at increased risk of experiencing a range of mental health difficulties, including anxiety. A number of intervention programmes are now available in high-income countries to support autistic children. However, to date there are no evidence-based interventions to support families of such children in South Asia. Based on consultations with clinicians, researchers and parents in Bangladesh and Sri Lanka, we developed a culturally tailored two-session skills-based group programme for parents whose autistic children present with anxiety. This paper describes the process of creating this programme, to be delivered by mental health professionals.

Autism spectrum disorder (ASD) is a lifelong neurodevelopmental condition characterised by difficulties with social communication and social interaction, and restrictive and repetitive behaviours, interests or activities. Autistic children are at increased risk of experiencing a range of mental health difficulties, including anxiety, which has been found to affect around 40%.¹ Anxiety can have a negative impact on a child's social development² and academic attainment.³

A number of parent-led cognitive-behavioural therapy (CBT) programmes for typically developing children experiencing anxiety are available⁴ and in recent years programmes have been developed in high-income countries to address anxiety in autistic children (for a recent review see⁵).

⁷Clinical Senior Lecturer and Honorary Consultant in Child and Adolescent Psychiatry, Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

⁸Professor of Psychology and Mental Health, Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK. Email jacqui.rodgers@newcastle.ac.uk

Keywords. Anxiety or fear-related disorders; autism spectrum disorders; low- and middle-income countries; neurodevelopmental disorders; psychosocial interventions.

First received 9 Jun 2023
Final revision 17 Oct 2023
Accepted 30 Oct 2023

doi:10.1192/bji.2023.37

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Anxiety in autism is particularly complex and may include autism-related features such as anxiety related to sensory differences, restrictive and repetitive behaviours, uncertainty and unusual phobias or idiosyncratic fears.⁶ Consequently, anxiety interventions for autistic children need to be tailored to meet their needs and may include modifications such as visual aids, significant behavioural components, emotional literacy work and incorporation of interests.⁷ Furthermore, parent-mediated approaches may be helpful in supporting generalisation of strategies across a range of contexts outside a clinical setting. Over the past 15 years or so a range of manualised interventions for autistic children experiencing anxiety have been developed in high-income countries.⁸ Anxiety in autism has also been identified as a research priority for the autism community globally,⁹ further highlighting the significant international importance. The high rates of anxiety in autistic children, limited availability of trained professionals in low- and middle-income countries (LMICs) and lack of an evidence-based intervention specifically to address anxiety experienced by children on the spectrum in LMICs all point towards a need for an intervention to be developed. The Helping your Anxious Child (HAC) programme was created to fill this treatment gap.

The HAC programme is underpinned by pre-existing frameworks from parent groups for anxious children developed in the West. HAC is designed to suit the needs of parents in South Asia, with adaptations designed in co-production with clinicians, researchers and parents of autistic children to ensure suitability and cultural appropriateness.¹⁰

The Helping your Anxious Child programme

Helping your Anxious Child was developed through collaborations via the North East England South Asia Mental health Alliance (NEESAMA; www.NEESAMA.org). NEESAMA seeks to bring together international partners in the UK and South East Asia to improve research, training and clinical service delivery within mental health services. At an international summit in Dhaka in 2019 an unmet need for an intervention to support autistic children experiencing anxiety was identified. This resulted in the Helping your Anxious Child (HAC) programme being developed through partnerships from Sri Lanka, Bangladesh and the UK.

The HAC programme is a manualised parent-mediated group intervention to address anxiety experienced by autistic children, specifically in LMICs. Based on CBT techniques, HAC aims to increase understanding of anxiety symptoms and how they may relate to autism, to identify helpful and less helpful strategies in managing anxiety and, if developmentally appropriate, to promote the use of helpful strategies to increase children's ability to manage their anxiety and experience of stressful situations on their own. In

addition to being cost-effective, a parent group programme was considered as it allows parents to teach children coping strategies across a range of settings and it has the potential to improve family and child outcomes.^{11,12}

The HAC programme was designed following consultation with parents, researchers and clinicians in Sri Lanka and Bangladesh to discuss potential benefits and challenges of delivering a parent group programme in those countries. The collaboration also ensured that the programme was potentially feasible for delivery within existing services. Following consultation with clinicians in Bangladesh and Sri Lanka, examples of specific adaptations to the programme included the following: (a) delivering fewer, longer sessions than the number typically included in such programmes (this is because parents advised that it would be very challenging to attend for more than two sessions); (b) incorporating stress management techniques using methods and techniques that parents and clinicians advised that families may be familiar with, for example butterfly hugs and pranayama; (c) not including the term autism in the title of the programme, owing to concerns regarding stigma; (d) providing a range of culturally adapted case examples; (e) providing opportunity for parents to practise their skills on a variety of culturally appropriate case examples; (f) reducing the emphasis on homework, to reduce its burden; (g) running smaller parent groups such as groups of three; and (h) increasing the amount of interactive activities/experiential exercises.

The HAC programme comprises two 4-h parent group sessions with a recommended 5–10 parents per group, guided by two facilitators. The sessions provide a range of strategies for managing anxious and stressful situations and are individualised to meet the needs of each parent and their child. Materials for the delivery of the programme include a presentation for each of the sessions, a therapist manual and a parent handout. The programme components include: (a) psychoeducation on cognitive, behavioural, emotional and physical presentations of anxiety; (b) psychoeducation on parent-reported common anxiety triggers and on recognising anxiety triggers using the STAR framework ('situation, trigger, anxiety, response'); (c) supporting children to develop emotional literacy; (d) working towards SMART ('specific, measurable, achievable, relevant and time-bound') goals; (e) drawing on a toolbox of strategies (physical/relaxation/sensory tools, social tools and thinking tools) to apply in example case studies, and reflecting on how parents can apply these in real-life situations; and (f) managing setbacks.

Results of pilot HAC programmes and next steps

Pilot HAC programmes have now been delivered to 14 parents of autistic children in Sri Lanka and

Bangladesh. Ethical permission was granted in Bangladesh and was not deemed necessary in Sri Lanka as the programme was conducted as part of routine clinical work. Written consent was obtained from parents to allow their experiences and feedback to be used for this research. Parents and clinicians also shared some reflections on their experiences and thoughts on the HAC programme.

Initial interviews and focus groups with parents suggest that the programme was well received by families in Sri Lanka and Bangladesh. Clinicians and group facilitators in Sri Lanka reflected that seven families participated despite a national fuel shortage and difficulties with transportation in 2022. Further, reflections from clinicians and group facilitators in Sri Lanka suggest that the structure was acceptable to parents as they found it helpful to describe their child's anxiety triggers in this way and valued the framework. All parents identified a SMART goal to work on, and by the end of the programme, parents were able to modify their SMART goal to include strategies such as 'labelling my own emotions with my child to enable him to go to the grocery store three times a week during less crowded times'. This suggests that the structure of the groups was acceptable and the experiential exercises enabled parents to adopt strategies and amend their SMART goals. Further, in Bangladesh, parents reflected that they valued sharing their own methods and techniques with the group during the experiential exercises. Clinicians reflected that this supported the therapeutic process within the groups. Lastly, parents in Bangladesh reflected that being aware of their own emotion regulation strategies during the experiential exercises enabled them to consider how this affected their child's coping strategies.

Above all, preliminary findings suggest possible beneficial outcomes following feedback from parents who took part as well as the professionals who delivered the training. Parents and facilitators also reported the programme to be feasible and acceptable. The next steps include the completion of a larger trial of the programme. In addition to this we aim to create an adaptation of HAC to provide a psychoeducation programme for teachers in LMICs supporting autistic children experiencing anxiety.

Data availability

The data are not publicly available due to their containing information that could compromise the privacy of research participants.

Author contributions

All authors contributed to the writing of this article.

Funding

The NEESAMA meeting that led to this work, held in Kathmandu Nepal in 2018 and Dhaka Bangladesh in 2019, received funding from the British Council UK, Cumbria Northumberland Tyne and Wear NHS Foundation Trust and Newcastle University.

Declaration of interest

None.

References

- 1 van Steensel FJ, Bögels SM, Perrin S. Anxiety disorders in children and adolescents with autistic spectrum disorders: a meta-analysis. *Clin Child Fam Psychol Rev* 2011; 14: 302–17.
- 2 de Lijster JM, Dieleman GC, Utens E, Dierckx B, Wierenga M, Verhulst FC, et al Social and academic functioning in adolescents with anxiety disorders: a systematic review. *J Affect Disord* 2018; 230: 108–17.
- 3 Woodward LJ, Fergusson DM. Life course outcomes of young people with anxiety disorders in adolescence. *J Am Acad Child Adolesc Psychiatry* 2001; 40: 1086–93.
- 4 Creswell C, Chessell C, Halliday G. Parent-led cognitive behaviour therapy for child anxiety problems: overcoming challenges to increase access to effective treatment. *Behav Cogn Psychother* [Epub ahead of print] 2 Dec 2022. Available from: <https://doi.org/10.1017/S1352465822000546>.
- 5 Vasa RA, Keefer A, McDonald RG, Hunsche MC, Kerns CM. A scoping review of anxiety in young children with autism spectrum disorder. *Autism Res* 2020; 13: 2038–57.
- 6 Wigham S, Rodgers J, South M, McConachie H, Freeston M. The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in autism spectrum disorder. *J Autism Dev Disord* 2015; 45: 943–52.
- 7 Winch AT, Klaver SJ, Simpson SD, Calub CA, Alexander K. Identifying and treating anxiety in youth with autism spectrum disorder/intellectual developmental disorder: a review. *Evid-Based Pract Child Adolesc Ment Health* 2021; 7: 160–72.
- 8 Perihan C, Burke M, Bowman-Perrott L, Bicer A, Gallup J, Thompson J, et al Effects of cognitive behavioral therapy for reducing anxiety in children with high functioning ASD: a systematic review and meta-analysis. *J Autism Dev Disord* 2020; 50: 1958–72.
- 9 Lord C, Charman T, Havdahl A, Carbone P, Anagnostou E, Boyd B, et al The Lancet Commission on the future of care and clinical research in autism. *Lancet* 2022; 399: 271–334.
- 10 Pervin M, Hagmayer Y. Attitudes towards evidence-based practice of professionals working with children and adolescents with autism spectrum disorder in Bangladesh. *Adm Policy Ment Health* 2022; 49: 861–80.
- 11 Bearss K, Johnson C, Smith T, Lecavalier L, Swiezy N, Aman M, et al Effect of parent training vs parent education on behavioral problems in children with autism spectrum disorder: a randomized clinical trial. *JAMA* 2015; 313: 1524–33.
- 12 Rodgers J, Goodwin J, Garland D, Grahame V, Isard L, Kernohan A, et al Coping with Uncertainty in Everyday Situations (CUES[®]) to address intolerance of uncertainty in autistic children: an intervention feasibility trial. *J Autism Dev Disord* 2023; 53: 3460–74.