

A new cheilostome bryozoan from a dinosaur site in the Upper Cretaceous (Campanian) Judith River Formation of Montana

Paul D. Taylor¹*  and Raymond R. Rogers²

¹Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom <p.taylor@nhm.ac.uk>

²Geology Department, Macalester College, 1600 Grand Ave., Saint Paul, MN 55105, USA <rogers@macalester.edu>

Abstract.—Few bryozoans have been described from the Cretaceous Western Interior Seaway (WIS), which is consistent with the low diversity of other typically stenohaline groups in this large expanse of relatively shallow marine water. Here we describe a new cheilostome bryozoan, *Conopeum flumineum* n. sp., based on well-preserved material from the Campanian Judith River Formation of the Upper Missouri River Breaks National Monument in north-central Montana. The new species shows strong morphological similarities with *Conopeum seurati*, a Recent species that is often categorized as brackish, but which is euryhaline and can also be found in marine and stenohaline environments. The new Campanian bryozoan species was found in a locality also containing fragmentary remains of dinosaurs and other terrestrial vertebrates, as well freshwater mollusks and terrestrial plant debris. The sedimentology and facies associations of the fossil-bearing site suggest that the depositional setting was a swamp or tidally influenced fluvial backwater on the Judith River coastal plain. The proximity of the site to the western shoreline of the WIS presumably made it susceptible to occasional marine flooding during storms or extreme tides. Previous occurrences of *Conopeum* in the Cretaceous of the Western Interior have also been associated with dinosaur remains, corroborating the very nearshore and at times even ‘upstream’ distribution of this euryhaline genus.

UUID: <http://zoobank.org/bb1fdc8a-5017-44c5-9251-9e24ef3995e3>.

Introduction

The Cretaceous Western Interior Seaway (WIS; Fig. 1.1) is characterized by a depauperate fauna in which some marine invertebrate groups are rare or absent (Hay et al., 1993). For instance, compared to rocks of similar age elsewhere, few brachiopods, corals, sponges, or echinoids have been recorded from the WIS. Bryozoans are another group seldom recorded from this vast expanse of shallow sea (Cuffey, 1994). Only a small number of papers have described or mentioned Cretaceous bryozoans from the WIS, and the total known diversity of these colonial invertebrates is at least an order of magnitude less than that found in sediments deposited at the same time along the Gulf and Atlantic coasts of North America (Taylor and McKinney, 2006), as well as in the Upper Cretaceous of Europe and elsewhere (see Taylor, 2019). Most of the WIS bryozoans are cheilostomes; cyclostome bryozoans have been recorded only in the Turonian of the WIS (Hattin, 1986), while an immured ctenostome *Pierrella larsoni* Wilson and Taylor, 2013, is known from the Pierre Shale of South Dakota and Colorado. Apart from the ascophoran cheilostome *Dysnoetopora demissa* (White, 1879) from the Mesaverde Formation of Wyoming (Toots and Cutler, 1962), the WIS cheilostomes are of malacostegan grade. Unlike the majority of cheilostome species, these lack ovicells and are thought to have possessed long-lived planktotrophic larvae like their living

relatives (Taylor, 1988). The WIS malacostegans include: (1) *Pyrripora shawi* Frey and Larwood, 1971 (= *Herpetopora anglica* Lang, 1914) from the Niobrara Chalk of Kansas; (2) *Eokotosokum bicystosum* (Allan and Sanderson, 1945) from the Edmonton Group of Alberta (Taylor and Cuffey, 1992); *Villicharixa lintonensis* (Cuffey et al., 1981) from the Fox Hills Sandstone of North Dakota and the Edmonton Group of Alberta (Taylor and Cuffey, 1992); (3) *Heteroconopeum ramosa* (Toots and Cutler, 1962) from the Mesaverde Formation of Wyoming; and (4) un-named species attributed to *Conopeum* sp. from the Fruitland Formation of New Mexico (Kues, 1983) and the Kaiparowits Formation of southern Utah (Roberts et al., 2008). Here, we describe a new WIS species of *Conopeum* that is notable in having a very similar morphology to *Conopeum seurati* Canu, 1928, an extant species unusual in being able to live in extremely low salinity, brackish waters (Winston, 1977).

The *Conopeum* specimens described in this report were collected from the Upper Cretaceous Judith River Formation in north-central Montana, within the confines of the Upper Missouri River Breaks National Monument (Fig. 1). The Judith River Formation is noteworthy because the first skeletal remains of dinosaurs described from North America were collected in 1855 from Judith River strata near the confluence of the Judith and Missouri rivers (Leidy, 1856; Thomson, 2006). To this day, the Judith River Formation and age-equivalent units elsewhere in the Western Interior Basin (e.g., Two Medicine, Oldman, Dinosaur Park, Kaiparowits, Fruitland, and Aguja formations) remain the focus of intense sampling and study in

*Corresponding author




Figure 1. Location and stratigraphy of the locality (site UC-8315) containing *Conopeum flumineum* n. sp. in the Campanian Judith River Formation, Montana, USA. (1) Regional setting in relation to Campanian shoreline of Western Interior Seaway (WIS). Gray-shaded area represents approximate distribution of terrestrial landscapes in the Western Interior of North America during the Campanian, west of the WIS (based on Blakey, 2013 and Slattery et al., 2015, modified from Rogers et al., 2020). (2) Location of site UC-8315 in Upper Missouri River Breaks National Monument. (3) Stratigraphy of site UC-8315 in Coal Ridge Member of Judith River Formation, in close proximity to transgressing shoreline of marine Woodhawk Member. (4) Field view of site UC-8315. Dashed line marks base of the bed containing *Conopeum flumineum* n. sp. Arrow points to 65 cm hoe pick resting on fossil-bearing stratum.

relation to their vertebrate fossils, and together these Campanian units yield some of the richest records of non-avian dinosaurs known from the entire Mesozoic (Weishampel et al., 2004). In contrast, invertebrate fossils in the Judith River Formation remain relatively undocumented and understudied, despite their abundance and common occurrence alongside vertebrates in numerous localities. Here we describe the first bryozoans recovered from the Judith River Formation, and as clarified above, one of very few bryozoans known from the entire Western Interior Cretaceous record. Interestingly, specimens of this new cheilostome bryozoan were recovered in direct association with vertebrate, invertebrate, and plant fossils with indisputable terrestrial and freshwater affinities.

Geological setting

The Judith River Formation in its type area in the Upper Missouri River Breaks National Monument is represented by ~180 m of terrestrial, paralic, and shallow-marine strata that range in age from middle to late Campanian (Goodwin and Deino, 1989; Rogers et al., 2016). The formation correlates to the west (updip) with alluvial facies of the middle and upper Two Medicine Formation, and is bound above and below by open-marine shales of the Claggett and Bearpaw formations, respectively (Fig. 1). A recent reappraisal of the stratigraphy of the Judith River Formation by Rogers et al. (2016) advanced three new formal members and linked them to distinct accommodation regimes in both terrestrial and marine settings. The specimens described in this report were collected from the new Coal Ridge Member (Fig. 1.3), a succession of coastal-plain alluvial/paralic facies that accumulated in a high accommodation setting landward of backstepping shorelines during the initial transgression of the Bearpaw Sea (Rogers, 1994, 1998; Rogers and Kidwell, 2000; Rogers et al., 2016). The Coal Ridge Member spans approximately the upper half of the Judith River Formation (up to 90 m) in the type area, and consists of fine-grained sandstone bodies of fluvial origin (often with indication of tidal influence) interstratified with overbank deposits, including beds of carbonaceous shale and lignite, that represent hydromorphic floodplains and expansive coastal swamps and mires. Recent $^{40}\text{Ar}/^{39}\text{Ar}$ analyses of sanidine crystals extracted from bentonite beds indicate that the Coal Ridge Member accumulated between 76.2–75.2 Ma (Rogers et al., 2016), which renders it late Campanian in age.

Fossil vertebrates, invertebrates, and plants are well represented in the coastal paralic facies of the Coal Ridge Member. These fossils are often found concentrated in vertebrate microfossil bonebeds (also known as vertebrate microfossil assemblages or ‘microsites,’ sensu Brinkman, 1990; Eberth et al., 2007; Rogers and Kidwell, 2007; Sankey and Baszio, 2008; Rogers and Brady, 2010), which are localized accumulations of small bones and teeth, including a variety of skeletal components and fragmentary remains from small animals such as frogs, salamanders, fish, and mammals, and small skeletal components or broken skeletal fragments from larger animals, including turtles, crocodiles, and dinosaurs. These localized multitaxic concentrations of small disarticulated and dissociated vertebrate fossils also commonly include shells and shell fragments of freshwater mollusks (bivalves and gastropods), and degraded (coalified) plant debris. In fact, many of the known vertebrate fossil sites in the Coal Ridge Member were initially discovered due to their conspicuous veneers of mollusk shell debris on outcrop.

The vertebrate microfossil bonebed that yields the *Conopeum* specimens was discovered in 1983 by J.C. Harkson during a government-funded survey commissioned to identify fossil resources along the Missouri River corridor, and is designated site UC-8315 (Fig. 1). This well-documented locality has been featured in three previous reports that detail stratigraphic and taphonomic attributes of the Judith River fossil record (Rogers and Kidwell, 2000; Rogers and Brady, 2010; Rogers et al., 2017). The matrix of site UC-8315 consists of light olive-gray (Munsell 5Y 6/2) mudstone. The tabular bed of mudstone that

hosts the fossil concentration is ~60 cm thick, and can be traced laterally for 40+ m to the limits of exposure (due to modern-day erosion). The site yields abundant fossil bone in association with copious remains of freshwater mollusks and carbonaceous plant debris over its entire extent. Vertebrate taxa represented in the UC-8315 collection include hadrosaurian, ceratopsian, and theropod dinosaurs, crocodylians, champsosaurs, turtles, small teleosts, gar, amphibians (salamanders and frogs), lizards, and mammals. The molluscan sample includes unionid and sphaeriid bivalves and small viviparid gastropods. Trematode flatworm parasites, diagnosed by distinctive igloo-shaped reaction traces on sphaeriid shell fragments, are the most recent addition to the UC-8315 faunal list (Rogers et al., 2018).

The sedimentology, fossil content, and facies associations of site UC-8315 are consistent with deposition in a low-energy aqueous setting, such as a lowland swamp or fluvial backwater, on the Judith River coastal plain. Based on regional mapping and thickness trends of the correlative marine Woodhawk Member of the Judith River Formation (Rogers et al., 2016), which consists of three back-stepping, shallow-marine sequences composed of sandy shoreface and deltaic strata (Fig. 1), site UC-8315 was located very close to the western margin of the WIS, perhaps no more than 1 or 2 km inland from the contemporaneous marine strand.

Materials and methods

Bulk samples of fossiliferous matrix excavated from locality UC-8315 were processed using an automated sieve system that washed sets of nested sieve pairs with openings of 500 µm and 2 mm concurrently. The unconsolidated mudstone matrix of UC-8315 readily disaggregated in a water bath, and remnant bioclasts in the sieves were dried under heat lamps and sorted into vertebrate, invertebrate, and plant fractions under light microscopes.

Approximately 50, mm-scale specimens of *Conopeum flumineum* n. sp. were isolated from the UC-8315 bioclast fraction, and selected well-preserved specimens were imaged using back-scattered electrons with a LEO ABT-55 SEM at the Natural History Museum in London.

Repositories and institutional abbreviations.—Fossil material collected from locality UC-8315 is currently curated at the Science Museum of Minnesota (SMM) and the National Museum of Natural History, Smithsonian Institution (USNM). The type material described in this report is deposited at the USNM. Comparative material of the modern species *Conopeum seurati* is deposited in the Department of Life Sciences, Natural History Museum, London (NHMUK).

Systematic paleontology

Phylum Bryozoa Ehrenberg, 1831

Order Cheilostomata Busk, 1852

Suborder Membraniporina Ortmann, 1890 (= Malacostegina Levensen, 1902)

Superfamily Membraniporoidea Busk, 1854

Family Electridae Stach, 1937

Genus *Conopeum* Gray, 1848

Type species.—*Millepora reticulum* Linnaeus, 1767, by monotypy.

Conopeum flumineum new species

Figures 2, 3

Holotype.—USNM PAL 771789 (Fig. 2.1–2.3). Cretaceous, upper Campanian, Judith River Formation, Coal Ridge Member; Upper Missouri River Breaks National Monument (locality UC-8315 of Rogers et al., 2017, fig. 1), Montana, USA.

Paratypes.—USNM PAL 771790–771795. Same details as for holotype.

Diagnosis.—*Conopeum* with multilamellar colonies, small- to intermediate-sized autozooids, narrow gymnocyst not expanded proximally, no gymnocystal spines, circumopesia cryptocyst, granular and with an outer groove, closure plates not observed.

Occurrence.—Upper Campanian, Judith River Formation, Coal Ridge Member; Upper Missouri River Breaks National Monument, Montana, USA.

Description.—Colony encrusting, multiserial, multilamellar (Fig. 3.6), in some places with zooids arranged in well-defined longitudinal rows (Fig. 3.2, 3.3), but elsewhere chaotically arranged (Figs 2.1, 3.1). Possible pore chambers (Fig. 2.4) narrow and contained mainly within the thickness of the zooidal walls, not indenting the zooidal cavities. Basal zooidal walls convex on colony underside, without an uncalcified window. Ancestrula and early astogeny unknown.

Autozooids variable in size, small to intermediate, ovoidal (Fig. 2.3) to rounded rectangular (Fig. 3.4) in outline shape, on average 1.5x longer than wide, surrounded by a thin fissure (Fig. 3.4). Gymnocyst narrow, broadening slightly at the corners of the zooids, but not proximally expanded. Cryptocyst moderately broad, circumopesia, inwardly sloping, not shelf-like, a groove present near the outer circumference (Fig. 2.3), surface texture pustulose, the pustules sometimes radially aligned but usually evenly distributed. Spines and their bases not seen. Opesia occupying most of the frontal surface, ovoidal to rounded rectangular, on average about twice as long as wide. Intramural buds occasionally developed (Figs. 2.2, 3.5). Closure plates not observed. Ovicells not observed, presumed absent.

Kenozooids variable in size and shape, the smallest examples adventitious and seemingly lacking a cryptocyst (Fig. 2.4), but most interzooidal, infilling irregular spaces between the larger autozooids, with a pustulose cryptocyst similar to that of the autozooids (Fig. 2.3).

Avicularia not observed, presumed absent.

Etymology.—*Flumen* is the Latin translation of river, referring to the Judith River Formation in which this species occurs.




Figure 2. *Conopeum flumineum* n. sp. from the Campanian Judith River Formation of Upper Missouri River Breaks National Monument, Montana, USA. (1–3) Holotype, USNM PAL 771789; (1) general view of the colony fragment with irregularly arranged zooids (scale bar = 500 μ m); (2) detail, including intramural bud indicated by an arrow (scale bar = 200 μ m); (3) kenozooid (left) and autozooid (right) with cryptocyst groove arrowed (scale bar = 100 μ m). (4) Paratype, USNM PAL 771790; broken edge of fragment (probably upside down with distal at the bottom) showing apparent pore chamber (arrow) and small kenozooid (center) (scale bar = 200 μ m).

Dimensions.—Based on 5 zooids from each of 4 colony fragments. Zooid length: mean (\bar{x}) 444 μ m, standard deviation (SD) 89 μ m, range 291–629 μ m. Zooid width: \bar{x} 295 μ m, SD 45 μ m, range 200–359 μ m. Opesia length: \bar{x} 326 μ m, SD 67 μ m, range 200–457 μ m. Opesia width: \bar{x} 166 μ m, SD 30 μ m, range 109–215 μ m.

Remarks.—More than 50 Recent and fossil species have been assigned to *Conopeum*, including at least 11 from the Cretaceous. However, the generic attribution of many of these species warrants re-evaluation, although this is hampered by the simple skeletal morphology of the genus, which provides few taxonomic characters. Among the key features of *Conopeum* are the absence of ovicells and avicularia (but see Grischenko et al., 2007 for a possible exception), and the poor development on the surface of the autozooids of the gymnocyst relative to the cryptocyst. Establishing the absence of both ovicells and avicularia depends on the availability of a large sample of specimens, which is often not the case for

fossils. The numerous and well-preserved specimens from the Judith River Formation include neither ovicells nor avicularia, and the close similarity of this species to the extant *Conopeum seurati* (Fig. 4) leaves little doubt that it belongs to *Conopeum*.

Some of the Cretaceous species placed in *Conopeum* have since been transferred to *Heteroconopeum* (see Taylor and McKinney, 2006), whereas others were provisionally reassigned by Martha et al. (2019) to their new Albian–Maastrichtian genus *Iyarisipora*. The latter genus differs from the type and other Recent species of *Conopeum*, including *C. flumineum* n. sp., in having closure plates that are calcified across the entire frontal surface of the zooid and which contain pores. Many putative species of *Conopeum* described from the Cretaceous are difficult to evaluate because they lack adequate illustration. A bryozoan identified as ?*Conopeum* sp. from the late Campanian Fruitland Formation of New Mexico encrusting a dinosaur bone (Kues, 1983) differs from *C. flumineum* n. sp. in having small distal spine bases. To judge from their figures, the *Conopeum* sp. described by Roberts et al. (2008) from the upper Campanian




Figure 3. *Conopeum flumineum* n. sp. from the Campanian Judith River Formation of Upper Missouri River Breaks National Monument, Montana, USA. (1) Paratype, USNM PAL 771791, colony fragment with autozooids ranging from ovoidal to rounded rectangular in outline shape (scale bar = 500 µm). (2) Paratype, USNM PAL 771792, colony fragment consisting of rounded rectangular zooids (scale bar = 500 µm). (3, 4) Paratype, USNM PAL 771793; (3) general view (scale bar = 500 µm); (4) detail showing almost rectangular zooids with a narrow peripheral gymnocyst and a broad cryptocyst (scale bar = 200 µm). (5) Paratype, USNM PAL 771794, damaged intramural bud (scale bar = 200 µm). (6) Paratype, USNM PAL 771795, part of a larger fragment showing multilamellar colony with areas of autozooids aligned in rows and others where their arrangement is more chaotic (scale bar = 1 mm).

Kaiparowits Formation of southern Utah has zooids averaging ~700 µm in length compared to the mean value of 444 µm for *C. flumineum* n. sp.

As already noted, this new species most closely resembles *Conopeum seurati* Canu, 1928, an extant species revised recently by Gordon et al. (2020) (Fig. 4). Autozooids in both




Figure 4. Comparative images of *Conopeum seurati* from modern lagoons in Tunisia. (1–3) Topotypes from Wadi Bezirk, NHMUK 2020.1.3.1; (1) lightly bleached zooids with cryptocystal groove arrowed (scale bar = 200 μ m); (2) dried, unbleached zooids showing two autozooids with flap-like opercula visible just above tears in their frontal membranes, and several small kenozooids, one of which is indicated by an arrow (scale bar = 200 μ m); (3) raised intramural bud (arrowed) potentially forming the origin of an overgrowing layer of zooids (scale bar = 500 μ m). (4) NHMUK 2020.1.3.2, Wadi Tinja; early astogeny with the ancestrula labelled 'A' and three succeeding generations of budded zooids labelled I, II, and III (scale bar = 500 μ m).

the Cretaceous and Recent species have a groove around the outer edge of the cryptocyst (compare Figs. 2.3 and 4.1), a feature lacking in other species assigned to *Conopeum*. Differences between the two species include the appreciably broader cryptocyst and presence of a narrow gymnocyst in *C. flumineum* n. sp., as well as the occasional development of spines in *C. seurati*, which have not been seen in the Judith River Formation material. Closure plates occur in *C. seurati*, where they were described as 'kleistozooids' by Poluzzi and Sabelli (1985), but have not been observed in *C. flumineum* n. sp. Autozooid size is similar in *C. flumineum* n. sp. and *C. seurati*: autozooid length ranges from 291–629 μ m in *C. flumineum* n. sp., compared with 342–722 μ m in *C. seurati* (Gordon et al., 2020); autozooid width ranges from 200–359 μ m in *C. flumineum* n. sp. compared with 150–501 μ m in *C. seurati* (Gordon et al., 2020).

The origin of multilamellar growth in *C. flumineum* n. sp. is not apparent from the available material. However, given that intramural buds are present in this species (Fig. 2.2), it is possible that multilamellar growth was achieved through eruptive overgrowths originating from these intramural buds, as is observed to occur in *C. seurati* (Fig. 4.3).

Fissures surrounding the zooids point to the past presence of organic cuticle of the outer layer forming the exterior walls that bound each zooid. Decay of the cuticle produces lines of weakness allowing taphonomic separation of the zooids. Parts of colonies with rectangular autozooids in longitudinal rows often have autozooids in adjacent rows at the same level (Fig. 3.2, 3.3) rather than in a brickwall-like alternation more typically found among cheilostomes. The difference between fragments with almost rectangular zooids aligned in rows (Fig. 3.2–3.4) and those with more ovoidal zooids arranged irregularly (Figs. 2.1, 2.2, 3.1) can be interpreted as reflecting growth freely across the substrate and confined growth, respectively.

Discussion

Terrestrial depositional environments in general, and dinosaur-bearing facies in particular, are not the most obvious places to expect to find fossil bryozoans, a phylum usually characterized as being stenohaline and fully marine. Admittedly, phylactolaemates and a few species of ctenostome bryozoans inhabit

freshwater environments today, but both of these groups lack mineralized skeletons, in contrast to the stenolaemates and cheilostomes that constitute the overwhelming majority of bryozoans found in the fossil record. Yet, with the addition of the Judith River Formation example described here, three terrestrial localities associated with dinosaurs along the margins of the Cretaceous WIS are now known to yield cheilostome bryozoans (Kues, 1983; Roberts et al., 2008; this report). Such apparently anomalous occurrences can be understood in the context of the small number of bryozoans with calcite skeletons that flourish in very low salinity environments at the present-day. Most of these euryhaline species are malacostegan cheilostomes and are usually identified as species of either *Conopeum* or *Einhornia* (*Electra* in older publications).

Examples of present-day occurrences of brackish water malacostegans have been described from inland waterways of Holland (Bijma and Boekschoten, 1985), the Coorong Lagoon of South Australia (Palinska et al., 1999), and the Albufeira coastal lagoon of Portugal (Freitas et al., 1994). Somewhat counterintuitively, bryozoan growth in the Portuguese example was observed to flourish after heavy rainfall caused salinity levels to drop to 6.5–9‰. It would be interesting to know whether this event also introduced terrestrial nutrients leading to an increase in phytoplankton that could trigger rapid bryozoan growth.

Some euryhaline cheilostomes are very fast-growing. For example, Dudley (1973) reported that colonies of *Conopeum tenuissimum* (Canu, 1908) could bud 150 zooids in the first week of their life. Kues (1983) noted that because modern *Conopeum* colonies are fast-growing, the *Conopeum?* sp. colonies he described encrusting Campanian dinosaur bones from the Fruitland Formation of New Mexico might conceivably have grown during a single season. If the Judith River species *Conopeum flumineum* n. sp. grew at a similar rate, suitable growing conditions would have been needed for only a brief amount of time to allow for the formation of mineralized colonies of the size found in this deposit.

Opportunistic growth of *C. flumineum* n. sp. during one or more short-lived episodes of elevated salinity is consistent with our current understanding of the paleoenvironment. Sedimentologic and paleontological evidence suggest that the aquatic setting represented by site UC-8315 was predominantly freshwater in nature, and was populated by a wide range of terrestrial, semi-aquatic, and aquatic animals, including groups that are generally salt intolerant, such as amphibians, sphaeriid bivalves, and viviparid gastropods. Taphonomic evidence suggests that the fossil assemblage preserved in site UC-8315 is time-averaged, with the accumulation of bioclastic debris transpiring over decades to centuries in a long-lived aquatic basin on the Judith River coastal plain (Rogers, 1993; Rogers and Brady, 2010; Rogers et al., 2017). Facies mapping indicates that site UC-8315 was located in very close proximity to the western shoreline of the WIS, being at most a few km inland, and thus occasional marine influence during deposition is certainly plausible, and arguably even likely given the low topographic gradient of the Cretaceous coastal plain in this region (Eberth, 1996). Sporadic storm surges linked to tropical cyclones (e.g., Eberth, 2015) and extreme tides may have been capable of transporting planktotrophic cheilostome larvae inland to the site, and resultant elevated salinities may have been established long enough for the development

of mature colonies, especially if *C. flumineum* n. sp. was indeed fast-growing, like living representatives of the genus. Lastly, the preservational quality of the *C. flumineum* n. sp. specimens is consistent with an in situ interpretation, because the fragile well-preserved colonies would have been susceptible to abrasion and fragmentation during long-distance transport.

The close morphological similarity between the Campanian species *Conopeum flumineum* n. sp. and the living *C. seurati*, imply that *C. seurati* may be the best ecological analogue for the Campanian species. *Conopeum seurati* is generally characterized as a brackish-water species (e.g., Koletic et al., 2014). According to Ryland (1970), European populations of *C. seurati* can tolerate salinities down to 1‰. The species is widespread in the Baltic Sea, for example, encrusting rotten wood in the Kiel Canal where salinities are 12‰ (Nikulina and Schäfer, 2006). However, *C. seurati* also occurs in higher salinity settings. At Avonmouth Dock on the Bristol Channel, UK, *C. seurati* has been found living in salinities of 18.5–27.6‰ (O’Dea and Okamura, 1999), and it is found in salinities of 30–38‰ at Ria de Aveiro, Portugal (Marchini et al., 2007). It has also been recorded from hypersaline lagoons, such as the Karvasta lagoon system of western Albania (Nonnis Marzano et al., 2010). The broad range of salinities over which *C. seurati* occurs at the present-day allows this species to be classified as euryhaline. If the morphologically similar Campanian species *C. flumineum* n. sp. was similarly euryhaline, its occurrence alone is uninformative about the paleosalinity of the Judith River Formation site UC-8315. That said, the association with unequivocal freshwater taxa is consistent with the site being characterized by freshwater conditions in general, with only occasional influxes of marine waters prompting higher salinities suitable for colonization by *C. flumineum* n. sp. (see above). The window for *C. flumineum* n. sp. to flourish may have been fleeting, and its fossils are decidedly rare in comparison with associated freshwater forms (e.g., sphaeriids, viviparids). It is also important to point out that *C. flumineum* n. sp. has only been found in this one Judith River site, despite the fact that several other similar fossil localities in the Coal Ridge Member have been collected and processed in identical fashion (Rogers et al., 2017).

Finally, from a phylogenetic standpoint, the recognition of a Campanian cheilostome species closely resembling the extant *Conopeum seurati* is significant. A molecular phylogeny (Gordon et al., 2020, fig. 8) recovered *C. seurati* as the basal species in a *Conopeum* clade forming the sister genus of the neocheilostomes with ‘conventional ovicells’ (Ostrovsky, 2021), crownward of malacostegan and some other more minor groups of cheilostomes (see Taylor, 2020, fig. 8.8). The *Conopeum* clade has particular importance in understanding the origin of neocheilostomes, the dominant bryozoans living today, which began an explosive radiation in the mid Cretaceous (Taylor, 1988). *Conopeum seurati* has a planktotrophic larva, which is shell-less and less flattened than those typical of ‘malacostegines,’ and probably has a shorter duration before settlement and metamorphosis (Cook, 1962). Such larvae are, in terms of morphology and time spent in the plankton, transitional to the non-planktotrophic coronate larvae that are brooded in the ovicells of neocheilostomes. *Conopeum flumineum* n. sp. from the Campanian may be an early example of a cheilostome possessing these traits.

Acknowledgments

We thank K. Curry Rogers, Z. Fulbright, J. Harkson, S. Kidwell, G. Liggett, K. Moffett, L. Rogers, B. Severy-Hoven, J. Thies, and J. Thole for insights, permits, and assistance in the lab and field. We also thank the many Macalester students who have worked to sieve and sort tiny fossils from various Judith River sites. Funding was provided by the National Science Foundation (grant EAR-1052673), the Bureau of Land Management (grant L10AC16281), Macalester College, and the Leverhulme Trust Research Project Grant (RPG-2016-429) ‘Molecules meet fossils—an integrated approach to studying palaeodiversity,’ awarded to A. Waeschenbach.

References

- Allan, J.A., and Sanderson, J.O.G., 1945, Geology of the Red Deer and Rosebud sheets, Alberta: Report of the Research Council of Alberta, v. 13, p. 1–109.
- Bijma, J., and Boekschoten, G.J., 1985, Recent bryozoan reefs and stromatolite development in brackish inland lakes, SW Netherlands: *Senckenbergiana Maritima*, v. 17, p. 163–185.
- Blakey, R., 2013, Regional paleogeographic views of earth history: Colorado Plateau Geosystems, Inc., <http://cpgeosystems.com>.
- Brinkman, D.B., 1990, Paleocology of the Judith River Formation (Campaian) of Dinosaur Provincial Park, Alberta, Canada: evidence from vertebrate microfossil localities: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 78, p. 37–54.
- Busk, G., 1852, An account of the Polyzoa, and sertularian zoophytes, collected in the voyage of the Rattlesnake, on the coasts of Australia and the Louisiade Archipelago &c, in MacGillivray, J., ed., *Narrative of the Voyage of H.M.S. Rattlesnake, commanded by the late Captain Owen Stanley ... 1846–1850; including discoveries and surveys in New Guinea, the Louisiade Archipelago, etc., to which is added the account of Mr E. B. Kennedy's expedition for the exploration of the Cape York Peninsula [including Mr W. Carron's narrative]*. Vol. 1: London, T.W. Boone, p. 343–402.
- Busk, G., 1854, Catalogue of Marine Polyzoa in the Collection of the British Museum. Part II. Cheilostomata (part.): London, Trustees of the British Museum, 66 p.
- Canu, F., 1908, Iconographie des bryozoaires fossiles de l'Argentine. Première Partie: *Anales del Museo Nacional de Buenos Aires*, v. 10, p. 245–341.
- Canu, F., 1928, Trois nouveaux bryozoaires d'eau douce: *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, v. 19, p. 262–264.
- Cook, P.L., 1962, The early larval development of *Membranipora seurati* (Canu) and *Electra crustulenta* (Pallas), (Polyzoa Ectoprocta): *Cahiers de Biologie Marine*, v. 3, p. 57–60.
- Cuffey, R.J., 1994, Cretaceous bryozoan faunas of North America—preliminary generalizations, in Hayward, P.J., Ryland, J.S., and Taylor, P.D., eds., *Biology and Palaeobiology of Bryozoans*: Fredensborg, Olsen and Olsen, p. 55–56.
- Cuffey, R.J., Feldmann, R.M. and Pohlable, K.E., 1981, New Bryozoa from the Fox Hills Sandstone (Upper Cretaceous, Maestrichtian) of North Dakota: *Journal of Paleontology*, v. 55, p. 401–409.
- Dudley, J.E., 1973, Observations on the reproduction, early larval development, and colony astogeny of *Conopeum tenuissimum* (Canu): *Chesapeake Science*, v. 14, p. 270–278.
- Eberth, D.A., 1996, Origin and significance of mud-filled incised valleys (Upper Cretaceous) in southern Alberta, Canada: *Sedimentology*, v. 43, p. 459–477.
- Eberth, D.A., 2015, Origins of dinosaur bonebeds in the Cretaceous of Alberta, Canada: *Canadian Journal of Earth Sciences*, v. 52, p. 655–681.
- Eberth, D.A., Rogers, R.R., and Fiorillo, A.R., 2007, A practical approach to the study of bonebeds, in Rogers, R., Eberth, D., and Fiorillo, A., eds., *Bonebed: Genesis, Analysis, and Paleobiological Significance*: Chicago, The University of Chicago Press, p. 265–331.
- Ehrenberg, C.G., 1831, *Symbolae physicae, seu icones et descriptiones animalium evertibratorum, sepositis insectis, quae ex itineribus per Libyam, Aegyptum, Nubiam, Dongalam, Syriam, Arabiam et Habessiniam public institutis sumptu Friderici Guilelmi Hemprich et Christiano Godofredi Ehrenberg medicinae et chirurgiae doctorum, studio annis MDCCCXX–MDCCCXXV redierunt*: Berlin, G. Reimer, 128 p.
- Freitas, M.C., Cacao, M., Cancela da Fonseca, L., Caroca, C., and Galopim de Carvalho, A.M., 1994, Unusual co-occurrence of serpulids and Bryozoa in a lagoonal system (Albufeira coastal lagoon—Portugal): *Gaia*, v. 8, p. 39–46.
- Frey, R.W. and Larwood, G.P., 1971, *Pyripora shawi*; new bryozoan from the Upper Cretaceous of Kansas (Niobrara Chalk) and Arkansas (Brownstown Marl): *Journal of Paleontology*, v. 45, p. 969–976.
- Goodwin, M.B., and Deino, A.L., 1989, The first radiometric ages from the Judith River Formation (Late Cretaceous), Hill County, Montana: *Canadian Journal of Earth Sciences*, v. 26, p. 1384–1391.
- Gordon, D.P., Sutherland, J.E., Perez, B.A., Waeschenbach, A., Taylor, P.D., and Di Martino, E., 2020, The bryozoan genus *Conopeum* (Electridae) in New Zealand, with description of a new species and discussion of the morphological and genetic characters of *Conopeum seurati* (Canu, 1928): *Journal of Natural History*, v. 54, p. 947–970.
- Gray, J.E., 1848, List of the Specimens of British Animals in the Collections of the British Museum. Part I. Centrifonae or Radiated Animals: London, Trustees of the British Museum, 173 p.
- Grischenko, A.V., Dick, M.H. and Mawatari, S.F., 2007, Diversity and taxonomy of intertidal Bryozoa (Cheilostomata) at Akkeshi Bay, Hokkaido, Japan: *Journal of Natural History*, v. 41, p. 1047–1161.
- Hattin, D.E., 1986, Carbonate substrates of the Late Cretaceous Sea, Central Great Plains and Southern Rocky Mountains: *Palaios*, v. 1, p. 347–367.
- Hay, W.W., Eicher, D.L., and Diner, R., 1993, Physical oceanography and water masses in the Cretaceous Western Interior Seaway, in Caldwell, W.G.E., and Kauffman, E.G., eds., *Evolution of the Western Interior Basin: Geological Association of Canada Special Paper*, v. 39, p. 297–318.
- Koletic, N., Novosel, M., Rajevic, N., and Franjevic, D., 2014, Bryozoans are returning home: recolonization of freshwater ecosystems inferred from phylogenetic relationships: *Ecology and Evolution*, v. 5, p. 255–264.
- Kues, B.S., 1983, Bryozoan and crustacean remains from Fruitland Formation (Upper Cretaceous) of New Mexico: *New Mexico Geology*, August 1983, p. 52–56.
- Lang, W.D., 1914, Some new genera and species of Cretaceous cheilostome Polyzoa: *Geological Magazine*, decade 6, v. 1, p. 436–444.
- Leidy, J., 1856, Notices of the remains of extinct reptiles and fishes discovered by Dr. F.V. Hayden in the bad lands of the Judith River, Nebraska Territory: *Proceedings of the Academy of Natural Sciences*, v. 8, p. 72–73.
- Levensen, G.M.R., 1902, *Studies on Bryozoa: Videnskabelige Meddelelser fra den naturhistoriske Forening i Kjøbenhavn*, v. 54, p. 1–31.
- Linnaeus, C., 1767, *Systema Naturae*. 12th edition: Holmiae, Laurentii Salvii, 1327 p.
- Marchini, A., Cunha, M.R., and Occhipinti-Ambrogi, A., 2007, First observations on bryozoans and entoprocts in the Ria de Aveiro (NW Portugal) including the first record of the Pacific invasive cheilostome *Tricellaria inopinata*: *Marine Ecology*, v. 28, suppl. 1, p. 154–160.
- Martha, S.O., Taylor, P.D., and Rader, W.L., 2019, Early Cretaceous gymnoleamete bryozoans from the early to middle Albian of the Glen Rose and Walnut formations of Texas, USA: *Journal of Paleontology*, v. 93, p. 260–277.
- Nonnis Marzano, C., Gravina, M.F., Fianchini, A., Portacci, G., Gherardi, M., Tursi, A., and Corriero, G., 2010, The macrozoobenthos of the Karavasta lagoon system (Albania): local assemblages and geographical comparisons: *Marine Ecology*, v. 31, p. 622–632.
- Nikulina, E.A., and Schäfer, P., 2006, Bryozoans of the Baltic Sea: *Meyniana*, v. 58, p. 75–95.
- O'Dea, A., and Okamura, B., 1999, Influence of seasonal variation in temperature, salinity and food availability on module size and colony growth of the estuarine bryozoan *Conopeum seurati*: *Marine Biology*, v. 135, p. 581–588.
- Ortmann, A.E., 1890, Die japanische Bryozoenfauna (Bericht über die von Herrn Dr. L. Döderlein im Jahre 1880–81 gemachten Sammlungen.): *Arch Naturgeschichte*, v. 54, p. 1–72.
- Ostrovsky, A.N., 2021, Sexual reproduction in Bryozoa, in Schwaha, T., ed., *Handbook of Zoology. Phylum Bryozoa*: Berlin, De Gruyter, p. 101–121.
- Palinska, K.A., Scholz, J., Sterflinger, K., Gerdes, G., and Bone, Y., 1999, Microbial mats associated with bryozoans (Coorong Lagoon, South Australia): *Facies*, v. 41, p. 1–14.
- Poluzzi, A., and Sabelli, B., 1985, Polymorphic zooids in deltaic species populations of *Conopeum seurati* (Canu, 1928) (Bryozoa, Cheilostomata): *Marine Ecology*, v. 6, p. 265–284.
- Roberts, E.M., Tapanila, L., and Mijal, B., 2008, Taphonomy and sedimentology of storm-generated continental shell beds: a case example from the Cretaceous Western Interior Basin: *The Journal of Geology*, v. 116, p. 462–479.
- Rogers, R.R., 1993, Systematic patterns of time-averaging in the terrestrial vertebrate record: a Cretaceous case study, in Kidwell, S.M., and Behrens-meyer, A.K., eds., *Taphonomic Approaches to Time Resolution in Fossil Assemblages: Short Courses in Paleontology (The Paleontological Society)*, no. 6, p. 228–249.
- Rogers, R.R., 1994, Nature and origin of through-going discontinuities in non-marine foreland basin deposits, Upper Cretaceous, Montana: implications for sequence analysis: *Geology*, v. 22, p. 1119–1122.

- Rogers, R.R. 1998, Sequence analysis of the Upper Cretaceous Two Medicine and Judith River formations, Montana: nonmarine response to the Claggett and Bearpaw marine cycles: *Journal of Sedimentary Research*, v. 68, p. 615–631.
- Rogers, R.R., and Brady, M.E., 2010, Origins of microfossil bonebeds: insights from the Upper Cretaceous Judith River Formation of north-central Montana: *Paleobiology*, v. 36, p. 80–112.
- Rogers, R.R., and Kidwell, S.M., 2000, Associations of vertebrate skeletal concentrations and discontinuity surfaces in nonmarine and shallow marine records: a test in the Cretaceous of Montana: *The Journal of Geology*, v. 108, p. 131–154.
- Rogers, R., and Kidwell, S., 2007, A conceptual framework for the genesis and analysis of vertebrate skeletal concentrations, *in* Rogers, R., Eberth, D., and Fiorillo, A., eds., *Bonebed: Genesis, Analysis, and Paleobiological Significance*: Chicago, The University of Chicago Press, p. 1–63.
- Rogers, R.R., Kidwell, S.M., Deino, A., Mitchell, J.P., and Nelson, K., 2016, Age, correlation, and lithostratigraphic revision of the Upper Cretaceous (Campanian) Judith River Formation in its type area (north-central Montana), with a comparison of low- and high-accommodation alluvial records: *The Journal of Geology*, v. 124, p. 99–135.
- Rogers, R.R., Carrano, M.T., Curry Rogers, K.A., Perez, M., and Regan, A.K., 2017, Isotaphonomy in concept and practice: an exploration of vertebrate microfossil bonebeds in the Upper Cretaceous (Campanian) Judith River Formation, north-central Montana: *Paleobiology*, v. 43, p. 248–273.
- Rogers, R.R., Curry Rogers, K.A., Bagley, B.C., Goodin, J.J., Hartman, J.H., Thole, J.T., and Zatoń, M., 2018, Pushing the record of trematode parasitism of bivalves upstream and back to the Cretaceous: *Geology*, v. 46, p. 431–434.
- Rogers, R.R., Regan, A.K., Weaver, L.N., Thole, J.T., and Fricke, H.C., 2020, Tracking authigenic mineral cements in fossil bones from the Upper Cretaceous (Campanian) Two Medicine and Judith River formations of Montana: *Palaaios*, v. 35, p. 135–150.
- Ryland, J.S., 1970, *Bryozoans*: London, Hutchinson, 175 p.
- Sankey, J.T., and Baszio, S., eds., 2008, *Vertebrate Microfossil Assemblages: Their Role in Paleocology and Paleobiogeography*: Bloomington, Indiana University Press, 278 p.
- Slattery, J.S., Cobban, W.A., McKinney, K.C., Harries, P.J., and Sandness, A.L., 2015, Early Cretaceous to Paleocene paleogeography of the Western Interior Seaway: the interaction of eustasy and tectonism: *Wyoming Geological Association Guidebook 2015*, p. 22–60.
- Stach, L.W., 1937, Bryozoa of Lady Julia Percy Island: *Proceedings of the Royal Society of Victoria*, v. 49, p. 374–384.
- Taylor, P.D., 1988, Major radiation of cheilostome bryozoans: triggered by the evolution of a new larval type?: *Historical Biology*, v. 1, p. 45–64.
- Taylor, P.D., 2019, A brief review of the scanty fossil record of Cretaceous bryozoans from Gondwana: *Australasian Palaeontological Memoirs*, v. 52, p. 147–154.
- Taylor, P.D., 2020, *Bryozoan Paleobiology*: Oxford, Wiley Blackwell, 320 p.
- Taylor, P.D., and Cuffey, R.J. 1992, Cheilostome bryozoans from the Upper Cretaceous of the Drumheller area, Alberta, Canada: *Bulletin of the British Museum (Natural History), Geology Series*, v. 48, p. 13–24.
- Taylor, P.D., and McKinney, F.K., 2006, Cretaceous Bryozoa from the Campanian and Maastrichtian of the Atlantic and Gulf Coastal plains, United States: *Scripta Geologica*, v. 132, p. 1–346.
- Thomson, K. S., 2006, Marginalia: American dinosaurs: Who and what was first?: *American Scientist*, v. 94, p. 209–211.
- Toots, H., and Cutler, J.F., 1962, Bryozoa from the 'Mesaverde' Formation (Upper Cretaceous) of southeastern Wyoming: *Journal of Paleontology*, v. 36, p. 81–86.
- Weishampel, D.B., Barrett, P.M., Coria, R.A., Le Loeuff, J., Xu, X., Zhao, X.J., Sahni, A., Gomani, E.M.P., and Noto, C.R., 2004, *Dinosaur distribution*, *in* Weishampel, D.B., Dodson, P., Osmolska, H., eds., *The Dinosauria*, 2nd edition: Los Angeles, University of California Press, p. 517–606.
- White, C.A., 1879, Paleontological papers no. 11: remarks upon certain Carboniferous fossils: *Bulletin of the United States Geological and Geographical Survey of the Territories*, v. 5, no. 2, p. 209–221.
- Wilson, M.A., and Taylor, P.D., 2013, Palaeoecology, preservation and taxonomy of encrusting ctenostome bryozoans inhabiting ammonite body chambers in the Late Cretaceous Pierre Shale of Wyoming and South Dakota, USA: *Lecture Notes in Earth System Sciences*, v. 143, p. 419–433.
- Winston, J.E., 1977, Distribution and ecology of estuarine ectoprocts: a critical review: *Chesapeake Science*, v. 18, p. 34–57.

Accepted: 5 April 2021