

Hexane–ethanol extract of *Glycyrrhiza uralensis* containing licoricidin inhibits the metastatic capacity of DU145 human prostate cancer cells

So Young Park¹, Soon Sung Lim¹, Jin Kyu Kim², Il-Jun Kang¹, Jong-Sang Kim³, Choonghwan Lee⁴, Jongdai Kim^{5,6} and Jung Han Yoon Park^{1,5*}

¹Department of Food Science and Nutrition, Hallym University, 39 Hallymdaehak-gil, Chuncheon 200-702, South Korea

²Institute of Natural Medicine, Hallym University, Chuncheon 200-702, South Korea

³Department of Animal Science and Biotechnology, Kyungpook National University, Daegu 702-701, South Korea

⁴Division of Bioscience and Biotechnology, Konkuk University, Seoul 143-701, South Korea

⁵Medical and Bio-Materials Research Center, Kangwon National University, Chuncheon 200-701, South Korea

⁶Department of Food Science and Biotechnology, Kangwon National University, Chuncheon 200-701, South Korea

(Received 30 November 2009 – Revised 18 March 2010 – Accepted 13 April 2010 – First published online 21 May 2010)

Licorice extracts are known to exhibit anti-carcinogenic activities. However, chronic licorice consumption can lead to serious side effects due to the presence of considerable quantities of glycyrrhizin, which causes severe hypokalaemia and hypertension. In the present study, we evaluated the effects of a hexane–ethanol extract of *Glycyrrhiza uralensis* (HEGU), which lacks glycyrrhizin, on the metastatic characteristics of DU145 prostate cancer cells. HEGU inhibited basal and epidermal growth factor-induced cell migration, invasion and adhesion in a dose-dependent fashion. HEGU significantly suppressed the secretion and activation of the matrix metalloproteinase (MMP)-2 and MMP-9. The secretion of tissue inhibitor of metalloproteinase (TIMP)-1 was reduced, but that of TIMP-2 was increased in HEGU-treated cells. HEGU reduced the protein levels of integrin- α 2, the intercellular adhesion molecule, and the vascular cell adhesion molecule. An active fraction of HEGU was separated via column chromatography, and the structure of the active component, licoricidin, was identified via ¹H NMR and ¹³C NMR. The treatment of DU145 cells with licoricidin induced a reduction in cell migration and the secretion of MMP-9, TIMP-1, urokinase-type plasminogen activator and vascular endothelial growth factor, as well as in the expression of adhesion molecules. These results indicate that HEGU, which contains licoricidin, is a potent anti-metastatic agent, which can markedly inhibit the metastatic and invasive capacity of malignant prostate cancer cells. The observed reductions in the activation of proteases and the levels of adhesion molecules may constitute a component of the mechanisms by which HEGU inhibits the migration and adhesion of prostate cancer cells.

***Glycyrrhiza uralensis*: Licoricidin: Metastasis: Prostate cancer: Matrix metalloproteinase: Adhesion molecules**

Licorice root is one of the oldest botanicals in traditional Oriental medicine. Its extract has been utilised worldwide as an additive for flavouring and sweetening of tobacco, candies, chewing gum, toothpaste and beverages. Additionally, it has been demonstrated to perform a variety of pharmaceutical functions, including detoxification, anti-ulcer, anti-inflammatory, anti-viral, anti-atherogenic and anti-carcinogenic activities. Licorice extracts have been shown to evidence anti-carcinogenic activities in both *in vivo* and *in vitro* animal models. For example, in a mouse xenograft model, licorice extract has been demonstrated to inhibit the growth of colon tumours⁽¹⁾. *In vitro* studies have shown that licorice extract inhibits the proliferation of prostate cancer cells⁽²⁾, and induces apoptosis in breast⁽³⁾ and gastric⁽⁴⁾ cancer cells. These results show that licorice extract has the potential to be developed into a cancer chemopreventive agent.

Licorice contains a considerable quantity of glycyrrhizin, a triterpene compound, which is normally regarded to be its principal biologically active component (reviewed in Wang & Nixon⁽⁵⁾ and Asl & Hosseinzadeh⁽⁶⁾). The results of previous *in vivo* studies have shown that glycyrrhizin inhibits pulmonary metastasis in mice injected with B16 melanoma cells⁽⁷⁾. The results of previously conducted *in vitro* cell culture studies have demonstrated that glycyrrhizin induces apoptosis in several human cancer cell lines^(8–10). However, glycyrrhizin is converted by human intestinal bacteria to glycyrrhetic acid⁽¹¹⁾, which has been reported to induce severe hypertension and hypokalaemia^(12,13). Therefore, our research group has attempted to prepare a licorice extract which lacks glycyrrhizin. In order to eliminate glycyrrhizin from the licorice extract, we had previously prepared a hexane–ethanol (9:1, v/v) extract of *Glycyrrhiza uralensis* (HEGU), which harbours no detectable glycyrrhizin⁽¹⁴⁾.

Abbreviations: DMEM, Dulbecco's modified Eagle's medium; EGF, epidermal growth factor; FBS, fetal bovine serum; HEGU, hexane–ethanol extract of *Glycyrrhiza uralensis*; ICAM, intercellular adhesion molecule; MMP, matrix metalloproteinase; M_r , molecular weight; TIMP, tissue inhibitor of metalloproteinase; uPA, urokinase-type plasminogen activator; VCAM, vascular cell adhesion molecule; VEGF, vascular endothelial growth factor.

* **Corresponding author:** J. H. Y. Park, fax +82 33 256 0199, email jyoona@hallym.ac.kr

Prostate cancer is one of the most common malignancies in men in the USA⁽¹⁵⁾, with an increasing incidence worldwide. Morbidity and mortality due to prostate cancer result primarily from metastasis to distant organs. The process of metastasis involves a series of separate steps, which include the separation of tumour cells in the primary tumor, local invasion, intravasation of invading cells into the vasculature or lymphatic systems, extravasations, angiogenesis and proliferation at the new site (reviewed in Steeg⁽¹⁶⁾). Matrix-degrading proteases, adhesion molecules and motility factors perform an important function in cancer invasion and metastasis (reviewed in Bogenrieder & Herlyn⁽¹⁷⁾). Matrix metalloproteinase (MMP), a zinc-dependent proteinase, is one of the most potent extracellular matrix-degrading enzymes⁽¹⁸⁾. MMP-2 (72 kDa gelatinase A) and MMP-9 (92 kDa gelatinase B) appear to be involved in the hydrolysis of basal membrane Type IV collagen, and have also been frequently associated with the invasive metastatic potential of tumour cells, including prostate cancer⁽¹⁹⁾. In addition to MMP, urokinase-type plasminogen activator (uPA) is a serine protease which is involved in tissue remodelling and cell migration⁽²⁰⁾. Cellular adhesion molecules are critical participants in cell–cell interactions, and interactions between cells and components of the extracellular matrix⁽²¹⁾. These molecules have been implicated in a broad variety of cellular functions, including apoptosis and metastasis. Integrins, another important group of cellular adhesion molecules, are involved in the process of local stromal invasion, as well as in invasion at the metastatic site⁽²²⁾.

The principal objective of the present study was to characterise the effects of HEGU lacking glycyrrhizin on the metastatic capacity of androgen-insensitive DU145 prostate cancer cells, and to identify the active component of HEGU. We demonstrated that HEGU inhibits the migration, invasion and adhesion of DU145 cells as well as MMP activities and the expression of adhesion molecules. Additionally, we identified licoricidin as the active component of HEGU.

Experimental methods

Materials

Reagents were purchased from the following suppliers: horseradish peroxidase-conjugated anti-rabbit and anti-mouse IgG from Amersham (Arlington Heights, IL, USA); antibodies against MMP-9, integrin- α 2, tissue inhibitor of metalloproteinase-1 (TIMP-1), TIMP-2, intercellular adhesion molecule (ICAM), vascular cell adhesion molecule (VCAM) and vascular endothelial growth factor (VEGF) from Santa Cruz Biotechnology (Santa Cruz, CA, USA); anti-uPA antibody from Calbiochem (Darmstadt, Germany); epidermal growth factor (EGF) from R&D Systems (Minneapolis, MN, USA); Adhesion Assay kit from Chemicon International (Temecula, CA, USA); and Centricon Plus-20 from Millipore Corporation (Billerica, MA, USA). Unless otherwise specified, all other materials were purchased from Sigma (St Louis, MO, USA).

Preparation and fractionation of hexane–ethanol extract of *Glycyrrhiza uralensis*, and identification of an active component (licoricidin)

Dried *G. uralensis* Fischer (Leguminosae) was purchased from Dae Kwang Company (Chuncheon, South Korea). A voucher

specimen is deposited in the Department of Food Science and Nutrition, Hallym University. HEGU was prepared as described previously⁽¹⁴⁾. HEGU (88.0 g) was subjected to flash column chromatography over silica gel and eluted with a gradient system of *n*-hexane–ethyl acetate (10:0–5:5, v/v) to yield 125 fractions that were pooled on the basis of their TLC patterns. Analytical silica gel (Kieselgel 60 F254, 0.25 mm; Merck, Darmstadt, Germany) plates were utilised for the determination of TLC patterns. The ability of the pooled fractions to inhibit DU145 cell migration was assessed, and the most active G89 (fractions 86–89, 2.0 g) was purified via column chromatography over a Cosmosil C18 eluting with a gradient of methanol–H₂O (20:80–10:90) to yield twenty subfractions. The ability of these twenty subfractions to inhibit DU145 cell migration was assessed, and the most efficacious fraction, F7 (5.4 mg), was purified further via recrystallisation. The compound in F7 was structurally identified via ¹H NMR and ¹³C NMR. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DPX 400 (400 MHz) spectrometer using tetramethylsilane as an internal standard.

Quantitative analysis of licoricidin in hexane–ethanol extract of *Glycyrrhiza uralensis*

Licoricidin contents in the HEGU were estimated via HPLC using a Thermo HPLC with a P2000 Gradient Pump (Thermo Separation Products, San Jose, CA, USA), an AS3000-021 thermostated autosampler, a SCM 1000-022 degasser and an Agilent Zorbax SB-C18 reversed-phase column (150 mm × 4.6 mm, inner diameter 5 μ m) coupled with an Agilent Zorbax Extend C18 guard column (10 mm × 4.6 mm inner diameter 5 μ m). The mobile phase consisted of 0.01% trifluoroacetic acid in water (A) and methanol (B). The gradient program was as follows: 60% B in 0–10 min, 60–100% B in 10–30 min at a flow rate of 0.8 ml/min using a Thermo 6000 LP detector at a wavelength of 254 nm. The standard was dissolved in methanol, and the standard curve was plotted using concentrations of 0.01–1 mg/ml, which is almost linear (R^2 0.99). The concentration of licoricidin was calculated via comparison of the peak area of the sample with that of the standard.

Cell culture

DU145 human prostate cancer cells (the American Type Culture Collection, Manassas, VA, USA) were grown in Dulbecco's modified Eagle's medium (DMEM)/F12 containing 100 ml/l of fetal bovine serum (FBS), with 62.5 mg/l of penicillin and 100 mg/l of streptomycin (Gibco BRL, Gaithersburg, MD, USA).

Migration, invasion and adhesion assays

For the transwell migration assay, DU145 cells were serum deprived in DMEM/F12 supplemented for 24 h with 1% charcoal-stripped FBS. Transwell filters (8 μ m pore size) were precoated with 10 μ g of Type IV collagen. The lower chambers of the wells of twenty-four-well plates were filled with DMEM/F12 containing 1% charcoal-stripped FBS and 0.1% bovine serum albumin in the absence or presence of

10 ng/ml EGF. The cells were then plated onto the filter in the transwell inserts at 2.5×10^4 cells/filter, and treated with the indicated concentrations of HEGU, fractions and licoricidin. The cells were then incubated for 4 h. After incubation, the migrated cells were stained with haematoxylin and eosin. For the invasion assay, procedures same as those described for the migration assay were conducted, except that the cells were plated on a matrigel-coated transwell filter (BD Biosciences, San Jose, CA, USA).

For the adhesion assay, DU145 cells were plated in human collagen Type I-coated CytoMatrix Cell Adhesion Strips. The cells were incubated for 45 min in DMEM/F12 containing 1 % charcoal-stripped FBS with various concentrations of HEGU with or without 10 ng/ml of EGF. The strips were rinsed twice with $\text{Ca}^{2+}/\text{Mg}^{2+}$ -containing PBS, and stained for 5 min with 0.2 % crystal violet in 10 % ethanol. After staining, the cell-bound stains were quantified by determining the absorbance at 570 nm.

Gelatin zymography

Cells were serum starved in DMEM/F12 and treated with 10 ng/ml EGF for 18 h in the absence or presence of HEGU. Conditioned media were collected, and the proteins were concentrated via centrifugal ultrafiltration using a 10 000 molecular weight (M_r) cut-off Centricon Plus-20 filter. Proteins in the conditioned media were separated via SDS-PAGE with 7.5 % acrylamide gel containing 1 % gelatin. The volumes of media loaded onto the gel were adjusted for equivalent protein levels. After electrophoresis, the gels were washed twice in 2.5 % Triton X-100 for 30 min to completely eliminate SDS. The gels were then rinsed twice with zymogen activation buffer (50 mM-Tris-HCl, 0.02 % Brij-35, 5 mM- CaCl_2 and 0.2 mM-NaCl), and were incubated for 48 h at 37°C in the same buffer. After incubation, the gels were stained for 2 h with 0.25 % Coomassie Blue solution and destained. Conditioned medium (40 μl) from the human fibrosarcoma cell line HT1080, which secretes MMP-9⁽²³⁾ and MMP-2⁽²⁴⁾, was used as a control.

Western blot analyses

Total cell lysates were prepared as described previously⁽²⁵⁾, and the protein contents were determined using a bicinchoninic acid protein assay kit (Pierce, Rockford, IL, USA). The total cell lysates (50 μg protein) and concentrated conditioned media (50 or 80 μg protein) were utilised for the Western blot analyses under reducing conditions as described previously⁽²⁵⁾. The relative abundance of each band was quantified using the Bio-profile Bio-ID application (Vilber-Lourmat, Marine La Vallée, France).

Statistical analysis

The data were expressed as the means with their standard errors and analysed via ANOVA. Differences among the treatment groups were evaluated by Duncan's multiple-range test, using the SAS system for Windows version 8.1 (SAS Institute, Cary, NC, USA).

Results

Hexane-ethanol extract of Glycyrrhiza uralensis inhibits the migration and invasion of DU145 cells

We have demonstrated previously that EGF stimulates the migration, invasion and adhesion of DU145 cells⁽²⁶⁾. In the present study, we cultured the cells in the absence or presence of EGF to determine whether HEGU inhibits the cell migration and invasion of DU145 cells. As anticipated, EGF significantly enhanced the migration of DU145 cells. In the absence of EGF, 7.5 $\mu\text{g}/\text{ml}$ of HEGU drastically inhibited the migration of DU145 cells. In the presence of EGF, 2.5–7.5 $\mu\text{g}/\text{ml}$ of HEGU inhibited the migration of DU145 cells in a dose-dependent manner (Fig. 1). The transwell invasion assay was also conducted using matrigel-coated filters, which demonstrated that HEGU markedly inhibited the basal and EGF-stimulated invasion of DU145 cells (Fig. 2). We reported previously that HEGU moderately inhibited DU145 cell growth when the cells were treated for 48 h with 5 $\mu\text{g}/\text{ml}$ of HEGU⁽¹⁴⁾. However, 2.5–7.5 $\mu\text{g}/\text{ml}$ of HEGU did not influence cell viability when the cells were treated for 24 h with HEGU (Mira Son & JHY Park, unpublished results), and therefore we utilised 2.5–7.5 $\mu\text{g}/\text{ml}$ of HEGU in the experiments conducted in the present study.

Hexane-ethanol extract of Glycyrrhiza uralensis alters the secretion of matrix metalloproteinase, tissue inhibitor of metalloproteinase and urokinase-type plasminogen activator by DU145 cells

In order to evaluate the possible anti-metastatic mechanisms of HEGU, we evaluated the effects of HEGU on the secretion of MMP using gelatin zymography. EGF stimulated the activity of pro-MMP-9, and HEGU inhibited the activity of both pro-MMP-9 and active MMP-9, whether the cells were treated with EGF or not (Fig. 3(A)). Additionally, the activity of pro-MMP-2 (72 kDa) was increased by EGF treatment and reduced by HEGU, whether the cells were treated with EGF or not. We detected another band with an apparent M_r of 43 000, and this band was increased by EGF treatment and reduced by HEGU treatment in a dose-dependent fashion (Fig. 3(A)). Because it has been reported previously that the MMP-2 proenzyme is converted to a 42.5 kDa stable active enzyme⁽²⁷⁾, we tentatively concluded that the 43 000 M_r species is a product of MMP-2 activation. Western blot analysis demonstrated that the secretion of pro-MMP-9 was significantly increased by EGF treatment, and that HEGU treatment effected a reduction in the protein levels of MMP-9 under both basal and EGF-stimulated conditions (Fig. 3(B)).

Because HEGU inhibited the expression and activation of MMP, we subsequently examined the effects of HEGU on the secretion of TIMP, natural inhibitors of MMP⁽¹⁹⁾. The results of Western blot analysis showed that EGF stimulated the secretion of TIMP-1, and that HEGU inhibited the secretion of this protein in the absence or presence of EGF (Fig. 3(B)). However, the secretion of TIMP-2 was not affected by EGF treatment, but was significantly increased as the result of treatment with 7.5 $\mu\text{g}/\text{ml}$ of HEGU (Fig. 3(B)).

The uPA system is believed to perform a pivotal function in tissue degradation, cell migration, angiogenesis, and cancer invasion and metastasis⁽²⁸⁾. Thus, we quantified uPA secretion




Fig. 1. Effect of hexane–ethanol extract of *Glycyrrhiza uralensis* (HEGU) on the epidermal growth factor (EGF)-induced migration of DU145 cells. DU145 cells were serum deprived for 24 h in Dulbecco's modified Eagle's medium (DMEM)/F12 containing 1% charcoal-stripped fetal bovine serum (FBS). The lower side of a 6.5 mm transwell filter was precoated with Type IV collagen. Cells were plated into the upper compartment at 25 000 cells/well, and were treated with 0–7.5 µg/ml of HEGU. The lower compartment was then filled with DMEM/F12 with 1% charcoal-stripped FBS and 0.1% bovine serum albumin with or without EGF. The cells were incubated for 4 h, and the migrated cells were stained with haematoxylin and eosin (H&E). (A) Photographs of H&E-stained cells (100×). (B) Quantitative analysis of the migrated cells. Each bar represents the means with their standard errors ($n = 3$). ^{a,b,c,d,e} Mean values with unlike letters were significantly different ($P < 0.05$).

levels via Western blotting. EGF increased the secretion of uPA (53 000 M_r pro-form and 33 000 M_r active form), and HEGU inhibited the secretion of these proteins. In addition to the pro- and active forms of uPA, we detected another band with a slightly higher M_r than that of pro-uPA only in the HEGU-treated cells (Fig. 3(B)). Western blot analysis with a VEGF antibody demonstrated that VEGF secretion was increased as the result of EGF treatment, and also that HEGU inhibited the secretion of VEGF under both basal and EGF-stimulated conditions (Fig. 3(B)).

Hexane–ethanol extract of Glycyrrhiza uralensis inhibits the adhesion of DU145 cells

In order to evaluate the effects of EGF/HEGU on the adhesion of DU145 cells, we used human collagen Type I-coated strips. EGF stimulated DU145 cell adhesion to Type I collagen, and HEGU inhibited basal and EGF-stimulated cell adhesion (Fig. 4(A)). The expression of a variety of adhesion molecules was detected via Western blotting. When the cells were treated with EGF, the levels of integrin- $\alpha 2$ protein were increased

by 170%, an increase which was significantly inhibited by treatment with 5 µg/ml of HEGU. The levels of ICAM and VCAM were not affected significantly by EGF. However, they were lower in cells treated with 7.5 µg/ml of HEGU than in cells treated with EGF only (Fig. 4(B)).

Licoricidin is an active component of hexane–ethanol extract of Glycyrrhiza uralensis

For the identification of active component(s), HEGU (88.0 g) was fractionated with a gradient system of *n*-hexane–ethyl acetate on flash column chromatography, and the resulting 125 fractions were pooled into twenty-nine groups on the basis of their TLC patterns. The ability of each group to inhibit DU145 cell migration was determined. Among the twenty-nine groups, G89 most potently inhibited DU145 cell migration (Fig. 5(A)). G89 (2.0 g) was fractionated further by column chromatography over a Cosmosil C18 eluted with a gradient of methanol–H₂O to yield twenty subfractions. Among these twenty subfractions, the F7 subfraction (5.4 mg) evidenced the highest inhibitory effect on DU145 cell




Fig. 2. Effect of hexane–ethanol extract of *Glycyrrhiza uralensis* (HEGU) on epidermal growth factor (EGF)-induced invasion of DU145 cells. DU145 cells were serum deprived for 24 h in Dulbecco's modified Eagle's medium (DMEM)/F12 containing 1% charcoal-stripped fetal bovine serum (FBS). The cells were plated in 6.5 mm matrigel-coated transwells at 25 000 cells/well, and were treated with 0–7.5 µg/ml of HEGU. The lower compartment was filled with DMEM/F12 with 1% charcoal-stripped FBS and 0.1% bovine serum albumin with or without EGF. The cells were incubated for 14 h, and the invaded cells were stained with haematoxylin and eosin (H&E). (A) Photographs of the H&E-stained cells (100 ×). (B) Quantitative analysis of the invaded cells. Each bar represents the means with their standard errors ($n = 3$). ^{a,b,c,d,e,f} Mean values with unlike letters were significantly different ($P < 0.05$).

migration (Fig. 5(B)). After further purification, the structure of the resultant pure compound in F7 was identified as licoricidin (Fig. 5(C)) via comparisons of its ^1H , ^{13}C NMR and MS spectra with those reported previously in the literature⁽²⁹⁾. The licoricidin content of the HEGU was 0.35%.

Licoricidin inhibits the migration of DU145 cells

When DU145 cells were treated with 0–5 µg/ml of licoricidin in the absence or presence of EGF, licoricidin inhibited the migration of DU145 cells in a dose-dependent fashion, whether the cells were treated with EGF or not (Fig. 6). However, 1–5 µg/ml of licoricidin did not affect cell viability when the cells were treated for 24 h with licoricidin (data not shown).

Licoricidin alters the expression of proteins involved in the regulation of metastasis in DU145 cells

As HEGU inhibited the secretion of MMP-9, TIMP-1, uPA and VEGF, we evaluated the effects of licoricidin on the secretion of these proteins. Western blot analysis showed that the secretion of MMP-9, TIMP-1, uPA and VEGF was reduced, and that of TIMP-2 was significantly increased in the licoricidin-treated cells (Fig. 7(A)). Similar to the HEGU-treated cells (Fig. 3(B)), a band with a M_r slightly higher than that of pro-uPA was detected in the cells treated with 5 µg/ml licoricidin, whether the cells were treated with EGF or not (Fig. 7(A)).

The expression of the adhesion molecules integrin- $\alpha 2$ and ICAM was reduced in a dose-dependent manner in licoricidin-treated cells. The levels of VCAM were significantly reduced in cells treated with 5 µg/ml of licoricidin (Fig. 7(B)).




Fig. 3. Effect of hexane-ethanol extract of *Glycyrrhiza uralensis* (HEGU) on epidermal growth factor (EGF)-induced secretion of matrix metalloproteinase (MMP), tissue inhibitor of metalloproteinase (TIMP), urokinase-type plasminogen activator (uPA) and vascular endothelial growth factor (VEGF) by DU145 cells. DU145 cells were plated in 100 mm dishes at 2×10^6 cells/dish in Dulbecco's modified Eagle's medium (DMEM)/F12 supplemented with 10% fetal bovine serum. One day later, the monolayers were serum starved for 24 h with serum-free DMEM/F12. The cells were incubated for 18 h with 0–7.5 $\mu\text{g/ml}$ of HEGU in serum-free media with or without EGF. The media were collected and concentrated for gelatin zymography (A) and Western blotting (B) after 18 h of conditioning. The volumes of media loaded onto the gel were adjusted for equivalent proteins. Photographs of a Coomassie Blue-stained gel (A) and chemiluminescent detection of the blots (B), which are representative of three independent experiments, are shown. In the first lane of (A), serum-free HT1080 cell-conditioned medium was loaded. The relative abundance of each band (B) was estimated via densitometric scanning of the exposed films. The control levels were set to one. The adjusted means with their standard errors (n 3) of each band are shown above each blot. ^{a,b,c,d} Mean values with unlike letters were significantly different ($P < 0.05$).

Discussion

In a previous study, we demonstrated that HEGU, which harbours no glycyrrhizin, suppressed doxorubicin-induced apoptosis in H9c2 rat cardiac myoblasts and induced reductions in the viable cell numbers of a variety of cancer cell types⁽¹⁴⁾. We have also observed previously that HEGU exerts potent anti-inflammatory effects in murine macrophages and skin⁽³⁰⁾. As we determined previously that EGF augments the migration, invasion and adhesion of androgen-independent DU145 prostate cancer cells⁽²⁶⁾, in the present study, an attempt was made to evaluate the effects of HEGU on the basal

and EGF-stimulated metastatic potential of DU145 cells. We observed that HEGU at concentrations of 2.5–7.5 $\mu\text{g/ml}$ ⁽¹⁾ markedly inhibits cell migration, invasion and adhesion⁽²⁾; reduces the secretion of MMP-2, MMP-9, TIMP-1, uPA and VEGF⁽³⁾; stimulates the secretion of TIMP-2; and⁽⁴⁾ reduces the expression levels of integrin- α 2, ICAM and VCAM proteins. These findings indicate that HEGU may be developed into an agent for the prevention of cancer metastasis.

The constituents of licorice include triterpene saponins, flavonoids, isoflavonoids and chalcones, with glycyrrhizin generally being considered the primary biologically active component (reviewed in Wang & Nixon⁽⁵⁾ and Asl & Hosseinzadeh⁽⁶⁾). It has also been demonstrated that glycyrrheticin




Fig. 4. Effect of hexane-ethanol extract of *Glycyrrhiza uralensis* (HEGU) on epidermal growth factor (EGF)-induced cell adhesion in DU145 cells. (A) DU145 cells were serum deprived for 24 h in Dulbecco's modified Eagle's medium (DMEM)/F12 containing 1% charcoal-stripped fetal bovine serum (FBS). Cells were plated in CytoMatrix™ human collagen I cell adhesion strips. Cells were incubated for 45 min in DMEM/F12 containing 1% charcoal-stripped FBS medium with 0–7.5 $\mu\text{g/ml}$ HEGU with or without EGF. Cells were stained with 0.2% crystal violet, and the cell-bound stains were colorimetrically quantified. Each bar represents the means with their standard errors (n 4). (B) DU145 cells were plated, serum starved and treated with HEGU as described in Fig. 3. Total cell lysates were subjected to immunoblotting with antibodies raised against integrin- α 2, intercellular adhesion molecule (ICAM) or vascular cell adhesion molecule (VCAM). Photographs of chemiluminescent detection of the blots, which are representative of three independent experiments, are shown. The relative abundance of each band was estimated via densitometric scanning of the exposed films, and the expression levels were normalised to β -actin. The adjusted means with their standard errors (n 3) of each band are shown above each blot. ^{a,b,c,d,e,f} Mean values with unlike letters were significantly different ($P < 0.05$).

acid exerts an inhibitory effect on the proliferation of LNCaP androgen-dependent prostate cancer cells⁽³¹⁾. Other components that are present in licorice extracts have also been demonstrated to exert anti-carcinogenic activities. For example, licochalcone A has been shown to inhibit inflammatory ear oedema formation and tumour promotion in mice⁽³²⁾. Isoliquiritigenin inhibited pulmonary metastasis⁽³³⁾,

the migration and invasion of prostate cancer cells⁽²⁶⁾, the pulmonary metastasis of mouse renal cell carcinoma⁽³³⁾, and 7,12-dimethylbenz(*a*)anthracene/12-*O*-tetradecanoylphorbol-13-acetate-induced cutaneous oxidative stress and tumour promotion⁽³⁴⁾. Because glycyrrhizin concentration in dried licorice root ranges between 3.63 and 13.06 g/100 g⁽⁵⁾, the chronic consumption of licorice in candies and beverages




Fig. 5. Purification and characterisation of the active compound licoricidin. DU145 cells were plated, serum starved as described in Fig. 1 and treated with 5 μ g/ml of each fraction of hexane-ethyl acetate extract of *Glycyrrhiza uralensis* (HEGU). The lower compartment of the matrigel-coated transwells was filled with Dulbecco's modified Eagle's medium/F12 with 1% charcoal-stripped fetal bovine serum and 0.1% bovine serum albumin with or without epidermal growth factor. The cells were incubated for 4 h, and the migrated cells were stained with haematoxylin and eosin. Quantitative analyses of the migrated cells are shown. (A) *n*-Hexane-ethyl acetate fractions of HEGU were pooled into twenty-nine groups on the basis of their TLC patterns. Groups (pooled fractions) exhibiting sufficient activity are shown: 64 (fractions 61–65), 76 (fractions 75–78), 80 (fractions 79–82), 89 (fractions 86–89), 91 (fractions 90–93), 96 (fractions 94–96) and 105 (fractions 102–105). Each bar represents the means with their standard errors (n 3). * Mean values were significantly different from HEGU ($P < 0.05$). (B) Methanol-H₂O fractions of G89. (C) Chemical structure of licoricidin. ¹H NMR (400 MHz, CD₃OD): δ 1.66, 1.67, 1.76, 1.77 (each 3H, s, CH₃), 2.72 (1H, dd, 16.0 and 11.0 Hz, H-4_{ax}), 2.90 (1H, ddd, 16.0, 5.1, and 1.8 Hz, H-4_{eq}), 3.23 (2H, t, $J = 7.2$ Hz, H-9), 3.36 (2H, brd, $J = 7.0$ Hz, H-7'), 3.32 (1H, m, H-3), 3.67 (3H, s, OCH₃), 3.89 (1H, t, $J = 10.2$ Hz, H-2_{ax}), 4.18 (2H, ddd, $J = 2.0, 3.1,$ and 10.2 Hz, H-2_{eq}), 5.19 (2H, m, H-10 and H-8'), 6.09 (1H, s, H-8), 6.35 (1H, d, $J = 8.4$ Hz, H-5'), 6.75 (1H, d, $J = 8.4$ Hz, H-6'). ¹³C NMR (100 MHz, CD₃OD): 158.47 (C-5), 155.90 (C-7), 155.71 (C-4'), 154.78 (C-8a), 154.31 (C-2'), 132.07 (C-9'), 130.84 (C-11), 125.61 (C-8'), 125.27 (C-10), 124.33 (C-6'), 121.62 (C-1'), 117.60 (C-3'), 115.12 (C-4a), 108.72 (C-5'), 108.61 (C-6), 100.05 (C-8), 71.26 (C-2), 60.95 (OCH₃), 32.79 (C-3), 27.43 (C-4), 26.01 (C-11'), 25.96 (C-13), 23.63 (C-7', 9), 18.02 (C-10'), 17.97 (C-12).




Fig. 6. Effect of licoricidin on epidermal growth factor (EGF)-induced migration of DU145 cells. DU145 cells were plated, serum starved as described in Fig. 1 and treated with 0–5 µg/ml of licoricidin. The lower compartment of the matrigel-coated transwells was filled with Dulbecco's modified Eagle's medium/F12 with 1% charcoal-stripped fetal bovine serum and 0.1% bovine serum albumin with or without EGF. The cells were incubated for 4 h, and the migrated cells were stained with haematoxylin and eosin (H&E). (A) Photographs of H&E-stained cells (100 ×). (B) Quantitative analysis of the migrated cells. Each bar represents the means with their standard errors (n 3). ^{a,b,c,d,e} Mean values with unlike letters were significantly different ($P < 0.05$).

is not recommended. However, licorice contains many bioactive components other than glycyrrhizin, and the identification and characterisation of such compounds may facilitate the development of effective new chemopreventive and/or chemotherapeutic agent(s).

In the present study, we demonstrated that licoricidin is an active component of HEGU, which is responsible, at least in part, for the inhibition of the metastatic potential of prostate cancer cells. To the best of our knowledge, the biological effects of licoricidin have not been abundantly reported, except that licoricidin exerts anti-bacterial effects against *Staphylococcus aureus* and *Helicobacter pylori* ^(35,36). This is the first study in which the anti-metastatic effect of licoricidin has been detected and described. However, the absorption and metabolism characteristics of licoricidin from HEGU were not determined in the present study. In order to determine whether the licoricidin concentrations used in the cell culture studies are relevant in an *in vivo* context, more studies will be required to determine the concentrations of licoricidin in the blood of animals and human subjects after the administration of licoricidin or HEGU.

In the present study, HEGU was demonstrated to markedly inhibit not only the secretion of MMP-2 and MMP-9, but also the activation of these enzymes (Fig. 3). TIMP are natural inhibitors of the MMP found in the majority of tissues and body fluids. By inhibiting the activities of MMP, they participate in the remodelling of ECM tissue ⁽³⁷⁾. Although the over-expression of TIMP-1 and TIMP-2 has been demonstrated to inhibit tumour growth, invasion and metastasis, conflicting results have also been reported ⁽³⁸⁾. In our previous study, we observed that EGF increased TIMP-1 secretion without any detectable changes in TIMP-2 ⁽²⁶⁾, a finding which is consistent with the findings of the present study. Herein, we observed that HEGU and licoricidin inhibited both basal and EGF-stimulated TIMP-1 secretion and increased TIMP-2 secretion (Figs. 3 and 7). Consistent with our findings, it has also been reported in previous studies that external stimuli including growth factors (basic fibroblast growth factor, platelet-derived growth factor and EGF), phorbol esters, serum and cytokines (IL-6, IL-1 and IL-1 β) induce TIMP-1 expression in a variety of cell types, including FTC-133 thyroid carcinoma cells, whereas TIMP-2 is constitutive ^(37,39).




Fig. 7. Effect of licoricidin on proteins involved in the regulation of metastasis in DU145 cells. DU145 cells were plated and serum starved as described in Fig. 3. The cells were incubated with 0–5 μg/ml of licoricidin in serum-free media with or without epidermal growth factor for 18 h. (A) The 18 h conditioned media were collected and concentrated for Western blotting. The volumes of media loaded onto the gel were adjusted for equivalent proteins. The relative abundance of each band was estimated via the densitometric scanning of the exposed films, and the adjusted means with their standard errors (*n* 3) of each band are shown above each blot. (B) Total cell lysates were subjected to immunoblotting with antibodies raised against integrin-α2, intercellular adhesion molecule (ICAM) or vascular cell adhesion molecule (VCAM). Photographs of chemiluminescent detection of the blots, which are representative of three independent experiments, are shown. The relative abundance of each band was estimated via densitometric scanning of the exposed films, and the expression levels were normalised to β-actin. The adjusted means with their standard errors (*n* 3) of each band are shown above each blot. ^{a,b,c,d} Mean values with unlike letters were significantly different (*P* < 0.05).

Furthermore, high serum TIMP-1 levels have recently been associated with adverse prognoses in endometrial carcinoma⁽⁴⁰⁾. Collectively, these findings are reflective of a more complex role for TIMP-1 during tumourigenesis than in the regulation of MMP activity. The observed increase in

the secretion of TIMP-2 may have contributed to reduced MMP activation in HEGU-treated cells.

The uPA/uPA receptor system is profoundly associated with prostate cancer metastasis⁽²⁸⁾. In the present study, EGF increased the protein levels of both 53 000 *M_r* pro-uPA and 33 000 *M_r* active uPA, and HEGU inhibited both the basal and EGF-stimulated secretion of these proteins. Additionally, we detected another band that migrated at a slightly slower rate than the 53 000 *M_r* band, and this band was induced by HEGU/licoricidin treatment (Fig. 3(B)). uPA is secreted as a 411-amino acid inactive proenzyme (pro-uPA), which has an apparent *M_r* of 53 kDa and undergoes several post-translational modifications. The zymogen is cleaved by plasmin or other proteases at Lys158–Ile159 to produce the active two-chain form held together by a single disulphide bond⁽²⁰⁾. It has also been reported in previous studies that the primary translation product is the 431-residue pre-pro-uPA which, via the removal of a twenty-residue-long signal peptide, gives rise to the secreted pro-uPA⁽⁴¹⁾. Thus, we tentatively concluded that the slightly higher *M_r* band observed in HEGU or licoricidin-treated cells may be attributable to pre-pro-uPA. HEGU may have prevented the removal of the signal peptide, and thus resulted in the accumulation of the higher *M_r* band. These results demonstrate that reduced uPA secretion contributes to reductions in the invasion and migration of HEGU-treated DU145 cells.

The formation of a new tumour vasculature is a critical process supporting tumour growth⁽⁴²⁾. VEGF promotes angiogenesis via the activation of a group of signalling pathways that stimulate the growth, migration, and survival of endothelial cells, vasculature permeability and the mobilisation of endothelial precursor cells from the bone marrow into the general circulation⁽⁴³⁾. In the present study, HEGU and licoricidin significantly inhibited basal and EGF-induced VEGF production in DU145 cells (Figs. 3(B) and 7(A)). Because VEGF plays a primary role in both tumour progression and angiogenesis⁽⁴⁴⁾, the HEGU-induced inhibition of VEGF production may constitute a promising strategy for the treatment of prostate cancer.

In addition to the inhibition of migration and invasion, we observed that HEGU inhibited the adhesion of DU145 cells and reduced the levels of the adhesion molecules, integrin-α2, ICAM and VCAM. Among the many changes inherent to the tumour progression process, alterations in cell–cell and cell–matrix adhesions appear to have a crucial role in the promotion of tumour cell invasion, migration and metastatic propagation⁽⁴⁵⁾. Our results show that HEGU inhibits cell adhesion, which is associated with reductions in the expression of integrin-α2, ICAM and VCAM.

In conclusion, in the present study, we have demonstrated that HEGU lacking glycyrrhizin inhibits the migration, invasion and adhesion of DU145 cells, and that this can be associated with reductions in the levels of MMP-9, MMP-2, uPA and VEGF secretion, as well as in the expression levels of adhesion molecules, and increases in the levels of TIMP-2 secretion. We also showed that licoricidin is an active component that is present in HEGU. Because metastasis is the primary cause of cancer deaths and the incidence of hypertension increases as men age, the HEGU-mediated inhibition of prostate cancer cell metastasis may have important preventive and therapeutic benefits. Future *in vivo* efficacy studies are warranted in light of these findings.

Acknowledgements

This work was supported by a grant (code no. 20070301034039) from the BioGreen 21 Program, Rural Development Administration, and by the grant of the Korean Ministry of Education, Science and Technology (The Regional Core Research Program/Medical and Bio-Materials Research Center). The authors declare that they have no conflict of interest. The authors' contributions are as follows: S. Y. P., I.-J. K., J.-S. K., C. L., J. K. and J. H. Y. P. designed the experiments. S. Y. P. conducted the biochemical analyses and analysed data. S. S. L. and J. K. K. prepared fractions of HEGU and identified the active component. S. Y. P. wrote the first draft, and J. H. Y. P. revised the paper. All the authors read and approved the final manuscript.

References

- Lee CK, Park KK, Lim SS, *et al.* (2007) Effects of the licorice extract against tumor growth and cisplatin-induced toxicity in a mouse xenograft model of colon cancer. *Biol Pharm Bull* **30**, 2191–2195.
- Adams LS, Seeram NP, Hardy ML, *et al.* (2006) Analysis of the interactions of botanical extract combinations against the viability of prostate cancer cell lines. *Evid Based Complement Alternat Med* **3**, 117–124.
- Jo EH, Kim SH, Ra JC, *et al.* (2005) Chemopreventive properties of the ethanol extract of chinese licorice (*Glycyrrhiza uralensis*) root: induction of apoptosis and G1 cell cycle arrest in MCF-7 human breast cancer cells. *Cancer Lett* **230**, 239–247.
- Ma J, Peng W & Liang D (2000) Apoptosis of human gastric cancer cell line MGC-803 induced by *Glycyrrhiza uralensis* extract. *Zhongguo Zhong Xi Yi Jie He Za Zhi* **20**, 928–930.
- Wang ZY & Nixon DW (2001) Licorice and cancer. *Nutr Cancer* **39**, 1–11.
- Asl MN & Hosseinzadeh H (2008) Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phytother Res* **22**, 709–724.
- Kobayashi M, Fujita K, Katakura T, *et al.* (2002) Inhibitory effect of glycyrrhizin on experimental pulmonary metastasis in mice inoculated with B16 melanoma. *Anticancer Res* **22**, 4053–4058.
- Hibasami H, Iwase H, Yoshioka K, *et al.* (2005) Glycyrrhizin induces apoptosis in human stomach cancer KATO III and human promyelotic leukemia HL-60 cells. *Int J Mol Med* **16**, 233–236.
- Hibasami H, Iwase H, Yoshioka K, *et al.* (2006) Glycyrrhetic acid (a metabolic substance and aglycon of glycyrrhizin) induces apoptosis in human hepatoma, promyelotic leukemia and stomach cancer cells. *Int J Mol Med* **17**, 215–219.
- Thirugnanam S, Xu L, Ramaswamy K, *et al.* (2008) Glycyrrhizin induces apoptosis in prostate cancer cell lines DU-145 and LNCaP. *Oncol Rep* **20**, 1387–1392.
- Hattori M, Sakamoto T, Kobashi K, *et al.* (1983) Metabolism of glycyrrhizin by human intestinal flora. *Planta Med* **48**, 38–42.
- Stormer FC, Reistad R & Alexander J (1993) Glycyrrhizic acid in liquorice – evaluation of health hazard. *Food Chem Toxicol* **31**, 303–312.
- van Uum SH (2005) Liquorice and hypertension. *Neth J Med* **63**, 119–120.
- Choi HJ, Seon MR, Lim SS, *et al.* (2008) Hexane/ethanol extract of *Glycyrrhiza uralensis* licorice suppresses doxorubicin-induced apoptosis in H9c2 rat cardiac myoblasts. *Exp Biol Med* **233**, 1554–1560.
- Jemal A, Siegel R, Ward E, *et al.* (2008) Cancer statistics, 2008. *CA Cancer J Clin* **58**, 71–96.
- Steeg PS (2006) Tumor metastasis: mechanistic insights and clinical challenges. *Nat Med* **12**, 895–904.
- Bogenrieder T & Herlyn M (2003) Axis of evil: molecular mechanisms of cancer metastasis. *Oncogene* **22**, 6524–6536.
- Yoon SO, Park SJ, Yun CH, *et al.* (2003) Roles of matrix metalloproteinases in tumor metastasis and angiogenesis. *J Biochem Mol Biol* **36**, 128–137.
- Zhang L, Shi J, Feng J, *et al.* (2004) Type IV collagenase (matrix metalloproteinase-2 and -9) in prostate cancer. *Prostate Cancer Prostatic Dis* **7**, 327–332.
- Crippa MP (2007) Urokinase-type plasminogen activator. *Int J Biochem Cell Biol* **39**, 690–694.
- Orian-Rousseau V & Ponta H (2008) Adhesion proteins meet receptors: a common theme? *Adv Cancer Res* **101**, 63–92.
- Okegawa T, Pong RC, Li Y, *et al.* (2004) The role of cell adhesion molecule in cancer progression and its application in cancer therapy. *Acta Biochim Pol* **51**, 445–457.
- Moll UM, Youngleib GL, Rosinski KB, *et al.* (1990) Tumor promoter-stimulated Mr 92,000 gelatinase secreted by normal and malignant human cells: isolation and characterization of the enzyme from HT1080 tumor cells. *Cancer Res* **50**, 6162–6170.
- Azzam HS & Thompson EW (1992) Collagen-induced activation of the M(r) 72,000 type IV collagenase in normal and malignant human fibroblastoid cells. *Cancer Res* **52**, 4540–4544.
- Cho HJ, Kim WK, Kim EJ, *et al.* (2003) Conjugated linoleic acid inhibits cell proliferation and ErbB3 signaling in HT-29 human colon cell line. *Am J Physiol Gastrointest Liver Physiol* **284**, G996–1005.
- Kwon GT, Cho HJ, Chung WY, *et al.* (2009) Isoliquiritigenin inhibits migration and invasion of prostate cancer cells: possible mediation by decreased JNK/AP-1 signaling. *J Nutr Biochem* **20**, 663–676.
- Howard EW, Bullen EC & Banda MJ (1991) Regulation of the autoactivation of human 72-kDa progelatinase by tissue inhibitor of metalloproteinases-2. *J Biol Chem* **266**, 13064–13069.
- Li Y & Cozzi PJ (2007) Targeting uPA/uPAR in prostate cancer. *Cancer Treat Rev* **33**, 521–527.
- Fukai T, Toyono M & Nomura T (1988) On the structure of licoricidin. *Heterocycles* **27**, 2309–2313.
- Cho HJ, Lim SS, Lee YS, *et al.* (2010) Hexane/ethanol extract of *Glycyrrhiza uralensis* licorice exerts potent anti-inflammatory effects in murine macrophages and in mouse skin. *Food Chem* **121**, 959–966.
- Hawthorne S & Gallagher S (2008) Effects of glycyrrhetic acid and liquorice extract on cell proliferation and prostate-specific antigen secretion in LNCaP prostate cancer cells. *J Pharm Pharmacol* **60**, 661–666.
- Shibata S, Inoue H, Iwata S, *et al.* (1991) Inhibitory effects of licochalcone A isolated from *Glycyrrhiza inflata* root on inflammatory ear edema and tumour promotion in mice. *Planta Med* **57**, 221–224.
- Yamazaki S, Morita T, Endo H, *et al.* (2002) Isoliquiritigenin suppresses pulmonary metastasis of mouse renal cell carcinoma. *Cancer Lett* **183**, 23–30.
- Agarwal MK, Iqbal M & Athar M (2005) Inhibitory effect of 18beta-glycyrrhetic acid on 12-O-tetradecanoyl phorbol-13-acetate-induced cutaneous oxidative stress and tumor promotion in mice. *Redox Rep* **10**, 151–157.
- Fukai T, Marumo A, Kaitou K, *et al.* (2002) Anti-*Helicobacter pylori* flavonoids from licorice extract. *Life Sci* **71**, 1449–1463.
- Hatano T, Kusuda M, Inada K, *et al.* (2005) Effects of tannins and related polyphenols on methicillin-resistant *Staphylococcus aureus*. *Phytochemistry* **66**, 2047–2055.

37. Lambert E, Dasse E, Haye B, *et al.* (2004) TIMPs as multifacial proteins. *Crit Rev Oncol Hematol* **49**, 187–198.
38. Jiang Y, Goldberg ID & Shi YE (2002) Complex roles of tissue inhibitors of metalloproteinases in cancer. *Oncogene* **21**, 2245–2252.
39. Soula-Rothhut M, Coissard C, Sartelet H, *et al.* (2005) The tumor suppressor PTEN inhibits EGF-induced TSP-1 and TIMP-1 expression in FTC-133 thyroid carcinoma cells. *Exp Cell Res* **304**, 187–201.
40. Honkavuori M, Talvensaaari-Mattila A, Puistola U, *et al.* (2008) High serum TIMP-1 is associated with adverse prognosis in endometrial carcinoma. *Anticancer Res* **28**, 2715–2719.
41. Blasi F & Verde P (1990) Urokinase-dependent cell surface proteolysis and cancer. *Semin Cancer Biol* **1**, 117–126.
42. Folkman J (2002) Role of angiogenesis in tumor growth and metastasis. *Semin Oncol* **29**, 15–18.
43. Hicklin DJ & Ellis LM (2005) Role of the vascular endothelial growth factor pathway in tumor growth and angiogenesis. *J Clin Oncol* **23**, 1011–1027.
44. Goh PP, Sze DM & Roufogalis BD (2007) Molecular and cellular regulators of cancer angiogenesis. *Curr Cancer Drug Targets* **7**, 743–758.
45. Christofori G (2006) New signals from the invasive front. *Nature* **441**, 444–450.