

Polygenic risk for schizophrenia, transition and cortical gyrification: a high-risk study

E. Neilson¹, C. Bois¹, T.-K. Clarke¹, L. Hall², E. C. Johnstone¹, D. G. C. Owens¹, H. C. Whalley¹, A. M. McIntosh¹ and S. M. Lawrie¹

Original Article

Cite this article: Neilson E, Bois C, Clarke T-K, Hall L, Johnstone EC, Owens DGC, Whalley HC, McIntosh AM, Lawrie SM (2018). Polygenic risk for schizophrenia, transition and cortical gyrification: a high-risk study. *Psychological Medicine* **48**, 1532–1539. <https://doi.org/10.1017/S0033291717003087>

Received: 7 February 2017
Revised: 20 September 2017
Accepted: 21 September 2017
First published online: 25 October 2017

Key words:

Familial high risk; polygenic risk scores; structural MRI

Author for correspondence:

E Neilson, E-mail: s0830415@sms.ed.ac.uk

¹Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Kennedy Tower, Edinburgh, UK and ²International Centre for Life, Institute of Genetic Medicine, Newcastle University, Central Parkway, Newcastle upon Tyne, UK

Abstract

Background. Schizophrenia is a highly heritable disorder, linked to several structural abnormalities of the brain. More specifically, previous findings have suggested that increased gyrification in frontal and temporal regions are implicated in the pathogenesis of schizophrenia. **Methods.** The current study included participants at high familial risk of schizophrenia who remained well ($n = 31$), who developed sub-diagnostic symptoms ($n = 28$) and who developed schizophrenia ($n = 9$) as well as healthy controls (HC) ($n = 16$). We first tested whether individuals at high familial risk of schizophrenia carried an increased burden of trait-associated alleles using polygenic risk score analysis. We then assessed the extent to which polygenic risk was associated with gyral folding in the frontal and temporal lobes. **Results.** We found that individuals at high familial risk of schizophrenia who developed schizophrenia carried a significantly greater burden of risk-conferring variants for the disorder compared to those at high risk (HR) who developed sub-diagnostic symptoms or remained well and HC. Furthermore, within the HR cohort, there was a significant and positive association between schizophrenia polygenic risk score and bilateral frontal gyrification. **Conclusions.** These results suggest that polygenic risk for schizophrenia impacts upon early neurodevelopment to confer greater gyral folding in adulthood and an increased risk of developing the disorder.

Introduction

Schizophrenia is a highly heritable disorder, which affects approximately 1% of the general population (Harrison & Weinberger, 2005). The underlying genetic architecture of schizophrenia has been hard to define, partly because it is a pathophysiologically complex, clinically heterogeneous disorder. Increasingly, however, evidence points to a substantial proportion of trait variation being polygenic; consisting of many common single-nucleotide polymorphisms (SNPs), each of small effect (Purcell *et al.* 2009; Lee *et al.* 2012; Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). At present, the diagnosis of schizophrenia is based solely on the clinical picture, and no objective biological marker exists for predicting which individuals are more likely than others to transition to psychosis. Genetic stratification could provide a means by which individuals who are most likely to transition to psychosis are identified pre-morbidly. Recently, there has been success in predicting whether first-episode patients (FEPs) will develop schizophrenia, as opposed to any other psychoses, through the use of polygenic risk scores for schizophrenia (PGRS-SCZ) (Vassos *et al.* 2017). The accuracy of these scores is continually improving as the Genome-Wide Association Study (GWAS) discovery sample grows (Vassos *et al.* 2017). Early detection has important clinical implications, as the duration of untreated psychosis has been shown to be associated with worse clinical outcome (Fraguas *et al.* 2014), and early intervention may improve a patient's prognosis, or even prevent the onset of psychotic disorders (Heinssen & Insel, 2015).

Schizophrenia is also characterized by a host of structural brain alterations, frequently in frontal and temporal cortex (Lawrie & Abukmeil, 1998; Fornito *et al.* 2012; Haijma *et al.* 2013). These findings have also been shown in some studies of unaffected relatives and in well individuals who later go on to develop schizophrenia (Boos *et al.* 2007; Fusar-Poli *et al.* 2011; Cooper *et al.* 2014). Brain changes may provide crucial insights into the genetic mechanisms of schizophrenia, increasing our understanding of the disorder's underlying aetiology (Bois *et al.* 2015). In examining the relationship between structural brain alterations and genetic risk, prospective studies of currently unaffected individuals at familial high risk (HR) of developing schizophrenia allow researchers to assess whether brain alterations are specifically associated with the disorder itself, or more accurately characterized as trait or vulnerability markers for schizophrenia (Johnstone *et al.* 2005; McIntosh *et al.* 2011), whilst avoiding

potential confounding influences; such as anti-psychotic medication and effects of adaptation to chronic illness.

Brain structural findings in schizophrenia are widespread and may reflect dysconnectivity between diverse regions, particularly between and within frontal and temporal regions (White & Hilgetag, 2011). These differences may be revealed in measures of cortical complexity, such as the gyrification index (GI) (Zilles *et al.* 1988), a measure of cortical folding thought to reflect underlying neuronal connectivity (White & Hilgetag, 2011). A temporal relationship between neuronal migration and patterns of gyrification has also been shown, with disorders that affect early migration having more alterations in the resulting gyrification patterns (White & Hilgetag, 2011). Developmental studies of gyrification demonstrate that gyrification remains relatively stable following birth (Zilles *et al.* 1988), and is associated with genetic factors (Harris *et al.* 2007; Nanda *et al.* 2014). As such, patient-control differences may reflect changes that take place early in neurodevelopment.

Studies have shown increased gyrification in the frontal lobes of people with schizophrenia (Falkai *et al.* 2007; Palaniyappan *et al.* 2011), their unaffected relatives (Falkai *et al.* 2007), participants with an at-risk mental state (Sasabayashi *et al.* 2017) and those at familial HR who later developed schizophrenia (Harris *et al.* 2007). Increases in temporal lobe gyrification have been found in FEPs when compared with controls (Harris *et al.* 2004) as well as in those with an at-risk mental state (Sasabayashi *et al.* 2017); however, this was not reported within those at familial HR (Harris *et al.* 2007).

It is important to note that decreases in gyrification, within these regions, have also been reported within schizophrenia patients (Sallet *et al.* 2003; White *et al.* 2003; Bonnici *et al.* 2007; McIntosh *et al.* 2009; Palaniyappan *et al.* 2011) but not in those at HR. It may be that these differential results are due to the developmental stage (Bonnici *et al.* 2007) and duration of the illness (McIntosh *et al.* 2009; Mancini-Marie *et al.* 2015). Overall, it appears that the link between gyrification and schizophrenia is not completely clear. However, whilst the evidence is limited, the pattern of increased gyrification in relation to both frontal and temporal regions has been previously reported within those with, and at risk of, developing schizophrenia. Nevertheless, further investigation into the association between gyrification and schizophrenia, particularly in relation to familial HR samples, is required.

Despite previously mixed findings, gyrification has been identified as a potential endophenotype for schizophrenia (White & Gottesman, 2012) and has been linked to genetic loading for the disorder. Increased PGRS-SCZ was associated with decreased gyrification in fronto-parietal regions in two independent populations (Liu *et al.* 2016), but further investigation is needed to determine if PGRS-SCZ can be linked to gyrification within a schizophrenia patient or HR sample.

We have therefore constructed a PGRS-SCZ, obtaining marker weights from the most recent Psychiatric Genomics Consortium-2 (PGC2) GWAS data (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014) in participants from the Edinburgh High Risk Study (EHRS). We then tested for an association between PGRS-SCZ and the transition to frank schizophrenia in those at familial HR for the disorder. We hypothesized that there would be a greater burden of common genetic risk variants in those at HR who subsequently made the transition, compared with those at HR who remained well and controls. We also assessed whether there was an association

between PGRS-SCZ and frontal and temporal indices of gyrification within the HR cohort, as well as their subsequent clinical outcomes, with the expectation that PGRS-SCZ would be associated with altered gyrification.

Methods and materials

Participants

The present analysis was conducted on data from a prospective longitudinal cohort study of people at high familial risk of schizophrenia, conducted over 10 years (Johnstone *et al.* 2005). Participants came from families multiply affected with schizophrenia. The recruitment and clinical assessment process for the EHRS have been described in detail elsewhere (Hodges *et al.* 1999; Johnstone *et al.* 2000, 2005). In summary, at baseline/recruitment HR individuals aged 16–25 years, with no personal history of psychiatric disorder, were contacted throughout Scotland based on the criteria that they had at least two first- and/or second-degree relatives with a diagnosis of schizophrenia. Healthy controls (HC) without personal or family history of major psychiatric disorder were recruited from the same social and geographical networks as the HR subjects in order to minimize potential confounding environmental influences. The groups were similar in sex distribution, paternal social class and education, with the vast majority of HC and HR individuals being either in full-time employment or education at entry into the study; however, genotyping was not conducted until the end of the study.

Informed consent was obtained from all participants, as approved by the Psychiatry and Clinical Psychology subcommittee of the Multi-Centre Research Ethics Committee for Scotland. All applications for continuation and amendment to this study have been filed appropriately with the Scotland Research Ethics Committee, and conformed to the Declaration of Helsinki.

The presence and absence of psychotic and other symptoms was established by Present State Examinations at 18-month intervals on up to five occasions, which was the main clinical assessment used for the study (Wing *et al.* 1974). During the course of the study, 21 HR individuals developed International Classification of Diseases, 10th Edition (ICD-10) schizophrenia, HR[ill], 19 of whom had full clinical assessments and 17 had a usable structural magnetic resonance imaging (MRI) scan before disorder onset. Only a subset of these ($n = 9$) were genotyped and are included in the current analysis. Those in the HR[ill] group were formally diagnosed after an average of 929 days (s.d. = 138) and were not offered re-scanning or re-assessment once the diagnosis had been made. In contrast, some individuals had psychotic symptoms at one or more assessments of the study, but were never ill enough to meet criteria for diagnosis, as they either had only one key symptom, or their symptoms were too transient or mild to satisfy diagnostic criteria – these individuals were designated HR[symp] ($n = 60$). The remaining HR subjects, who never had any psychotic symptoms throughout the course of the study were designated HR[well] ($n = 66$). None of the HR individuals were on any form of anti-psychotic medication at the time of scanning. Participants for the HC group ($n = 36$) were recruited from the social network of the HR individuals and had no personal or family history of psychotic illness, but could have a family history of other psychiatric illness. HCs were otherwise as similar to the HR participants as possible.

Numbers of participants who had both structural MRI and genotyped data amounted to 46% of the initial sample and were designated to sample groups as follows: HC ($n = 16$), HR[well] ($n = 31$), HR[symp] ($n = 28$), HR[ill] ($n = 9$). More detailed demographic information is presented in Table 1.

Genotyping and derivation of PGRS

Information on the genotyping process can be found in online Supplementary Data.

Polygenic profile scores were generated using imputed genotype data. Imputation was performed in accordance with the ENIGMA2 1000 genomes protocol (ENIGMA2 Genetics Support Team, 2012). SNPs with an imputation R^2 quality score of >0.3 were retained for further analysis resulting in 6 145 246 SNPs. All subsequent analyses were performed in PLINK (Purcell *et al.* 2007). Further QC criteria were applied to imputed data. Individuals with missingness $>2\%$ were excluded, as were SNPs with a genotype call rate of $<98\%$, Hardy-Weinberg equilibrium p value $<1 \times 10^{-6}$, a minor allele frequency of $<5\%$, or those that were strand ambiguous. Clump-based linkage disequilibrium pruning (r^2 0.2, 300 kb window) was

performed to create a SNP set in approximate linkage equilibrium. Marker weights (logarithm of the odds ratio) and p value association statistics for SNPs were derived from the most recent PGC GWAS of schizophrenia (9.8 million autosomal SNPs) (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). PGRS were generated at five p value thresholds by multiplying the number of copies of the reference allele by its marker weight for each SNP, and summing across all SNPs, as described in Purcell *et al.* (2009).

Imaging parameters

Structural brain images were acquired using a 42 SPE Siemens (Erlangen, Germany) Magnetom scanner operating at 1.0 T. The sequence was a three-dimensional (3D) magnetization prepared rapid acquisition gradient echo sequence consisting of a 180° inversion pulse followed by a fast low-angle shot collection (flip angle 12° , repetition time 10 ms, echo time 4 ms, inversion time 200 ms, relaxation delay time 500 ms, field of view 250 mm), giving 128 contiguous slices with a thickness of 1.88 mm. The sequence was selected in order to obtain optimal grey and white matter contrast.

Table 1. Demographic information for healthy controls and the high-risk sub-groups

	HC ($n = 16$)	HR[well] ($n = 31$)	HR[symp] ($n = 28$)	HR[ill] ($n = 9$)	Statistics
Age	21.46	22.17	20.96	19.33	$F = 2.78, p = 0.047^*$
Mean (s.d.)	(2.46)	(2.62)	(3.01)	(2.60)	
Gender					$\chi^2 = 6.13, p = 0.105$
Male	9	18	8	5	
Female	7	14	20	4	
Handedness					$\chi^2 = 3.52, p = 0.741$
Right	9	27	21	9	
Left	2	2	4	0	
Mixed	0	1	2	0	
NA	0	1	1	0	
NART IQ	100.5	102.0	98.1	101.7	$F = 1.23, p = 0.306$
Mean (s.d.)	(8.43)	(8.72)	(9.77)	(11.20)	
Employment and education					$\chi^2 = 1.56, p = 0.957$
In education	5	7	7	2	
Unemployed	1	1	1	1	
Employed	6	11	9	2	
NA	4	12	11	4	
Paternal social class					$\chi^2 = 15.64, p = 0.618$
Non-manual	3	3	2	1	
Manual skilled	4	13	16	6	
Manual semi-skilled	3	2	3	1	
Manual unskilled	1	3	3	0	
NA	5	10	4	1	

* $p \leq 0.05$.

Significant effect of age was due to the HR[well] cohort being significantly older than the HR[ill] group.

All demographics were measured at the time of scan.

Handedness was measured using the Annett Hand Preference Questionnaire.

FreeSurfer analysis

Cortical reconstructions were generated using the surface-based stream of the software FreeSurfer, version 5.3.0 (<http://surfer.nmr.mgh.harvard.edu/fswiki/recon-all/>). Briefly, this processing includes motion correction and averaging of T1-weighted images, removal of non-brain tissue using a hybrid watershed/surface deformation procedure, automated Talairach transformation, intensity normalization (Ségonne *et al.* 2007), tessellation of the grey matter–white matter boundary, automated topology correction (Sled *et al.* 1998; Fischl *et al.* 2001) and surface deformation to place the grey/white and grey/cerebrospinal fluid borders optimally (Fischl *et al.* 1999). This method uses both intensity and continuity information from the entire 3D MR volume in segmentation and deformation procedures to produce representations of cortical thickness and surface area (Fischl & Dale, 2000). The maps are created using spatial intensity gradients across tissue classes and are therefore not simply reliant on absolute signal intensity. Reconstructed data sets were visually inspected for accuracy, and all segmentation errors were corrected by a trained, group-blinded rater (CB).

Gyrification indices

Local gyrification indices (LGI) were obtained with the method of Schaer *et al.* (2008), which operates on the images reconstructed through the FreeSurfer pipeline described above. Schaer *et al.*'s method is a vertex-wise extension of Zilles' original 2D GI, which is a measure that gives an estimate of the inner folded contour compared with the outer perimeter of the cortex (Zilles *et al.* 1988). A detailed description of Schaer *et al.*'s method can be found in online Supplementary Data.

Statistical analysis

All statistical analyses were computed with R version 3.2.2 (<http://www.r-project.org>). Analyses of covariance (ANCOVAs) were used to implement mixed-model linear analyses. A genetic relationship matrix (GRM) was created using autosomal SNPs that passed QC (Gorjanc *et al.* 2007). The ASReml-R (www.vsnr.com.au/software/asreml) software package was used to implement mixed linear model analyses with an inverse relationship matrix derived from the GRM fit as a random effect. These analyses were performed to determine whether genetic relatedness within the sample impacted on the associations reported, a method described previously (McIntosh *et al.* 2016).

To account for potential confounding due to population stratification, multi-dimensional scaling (MDS) components were created following the ENIGMA 1000 genomes protocol (ENIGMA2 Genetics Support Team, 2012), using the software package, PLINK (Purcell *et al.* 2007). Four MDS components were included in subsequent analyses models within this study, consistent with previous publications (McIntosh *et al.* 2013; Whalley *et al.* 2015). Results presented here primarily refer to PGRS-SCZ at the $p \leq 0.1$ threshold, as this threshold was shown to explain the most phenotypic variance in the discovery cohort (Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014). Results at the four other thresholds ($p \leq 0.01, 0.05, 0.5$ and 1) are presented in online Supplementary Data. All analyses are subject to a statistical significance level of $p < 0.05$.

Group comparisons in mean PGRS

Gender, age (mean centred), MDS components 1–4 and group status (HR[well], HR[symp], HR[ill] and HC) were entered as fixed effects in the ANCOVA models, with family structure fitted as a random effect in subsequent ASReml-R models. Family structure was inferred using genomic data from 61 268 SNPs across the genome and an inverse G matrix created. PGRS-SCZ was entered as the dependent variable in both models.

For the ASReml-R models, Wald's conditional *F*-test was used to calculate *p* values for fixed effects.

All ANCOVA and ASReml-R models were corrected for multiple comparisons using false discovery rate (FDR) correction, with a rate of $p = 0.05$ (Genovese *et al.* 2002), with pairwise comparisons adjusted using Tukey from the R package 'lsmeans'.

PGRS and frontal and temporal gyrification

The covariates were entered as in the above analyses, with the addition of global gyrification (left/right) as a covariate, to assess localized, as opposed to global changes in gyrification. Frontal/temporal (left/right) gyrification was entered as the dependent variable, with PGRS-SCZ as the predictor variable of interest. These models were conducted for controls, the HR cohort as a whole, and then with the HR group split by subsequent clinical outcome. For the latter model, a group by PGRS-SCZ interaction was used as the predictor of interest with further analyses of individual sub-groups, FDR corrected, to explore significant relationships.

Mean gyrification

Comparisons of mean gyrification utilized the means of left and right, frontal and temporal gyrifications within the sample.

Results

Demographics

Statistical analysis of demographic variables revealed a significant difference in age between groups, with pairwise comparisons suggesting this was due to the HR[well] cohort being significantly older than the HR[ill] group ($p = 0.044$). However, all statistical models controlled for age within this study.

No other significant differences were found for any other demographics of the sample (Table 1).

Group comparisons in mean polygenic scores

Mean PGRS-SCZ for each group, at all five thresholds, are reported in online Supplementary Data, Table S1.

The main effect of group on PGRS-SCZ was significant at the $p \leq 0.1$ threshold ($F_{3,74} = 2.91, p = 0.040$). Differences at other thresholds can be seen in online Supplementary Data, Table S1.

Subsequent pairwise comparisons, adjusted using Tukey, revealed that this was because the HR[ill] cohort had significantly increased PGRS-SCZ compared with controls ($p = 0.037$). These results are illustrated in Fig. 1. No other contrasts reached significance (online Supplementary Data, Table S2).

Importantly, this main effect was also significant when adjusted for relatedness in ASReml-R ($F_{3,74} = 3.22, p = 0.028$).




Fig. 1. Boxplot illustrating differences in mean PGRS-SCZ by group, where mean polygenic loading for the HR[ill] group ($n=9$) was significantly different to the HC group ($n=16$). Y-axis represents PGRS-SCZ. X-axis are the four groups: HR+ = HR[symp], HR = HR[well], LL = HR[ill] and CON = healthy controls. Two outliers are highlighted in this figure; however, these individuals were <3 standard deviations away from the mean and the decision was therefore made to include them within the analysis.

Associations between polygenic scores and frontal and temporal gyrification

Frontal lobe

For the HR cohort, as a whole, there was a significant effect of PGRS-SCZ on left frontal ($F_{1,59} = 7.37$, $p = 0.010$) and right frontal gyrifications ($F_{1,59} = 5.38$, $p = 0.024$), where an increase in genetic loading was associated with bilateral hypergyrification. These findings survived FDR correction (left frontal: $p_{\text{corrected}} = 0.035$, right frontal: $p_{\text{corrected}} = 0.048$) and are illustrated in Figs 2a and b. These results were also apparent when controlling for

relatedness: left frontal gyrification ($F_{1,58,9} = 7.12$, $p = 0.010$, $p_{\text{corrected}} = 0.039$), right frontal gyrification ($F_{1,59} = 5.38$, $p = 0.024$, $p_{\text{corrected}} = 0.048$). No effect was found for HC (online Supplementary Data, Table S3).

Analysis of the differences between the HR sub-groups revealed a significant interaction of group and PGRS-SCZ on the right frontal lobe ($F_{2,56} = 4.78$, $p = 0.012$), which survived FDR correction ($p_{\text{corrected}} = 0.050$).

Further exploration of the HR sub-groups individually revealed a significant positive relationship with polygenic loading and right frontal gyrification for the HR[symp] group ($F_{1,20} = 10.63$, $p = 0.004$, $p_{\text{corrected}} = 0.012$). Only a trend-level effect, in the same direction, was found for the HR[ill] group, although a large effect size ($\eta^2 = 0.98$) suggests that, had this cohort included a larger number of participants, this result may have reached significance (Table 2). No effect was found within the HR[well] group (online Supplementary Data, Table S4).

There was also a positive effect of PGRS-SCZ within the left frontal lobe for the HR[symp] sub-group ($F_{1,20} = 6.21$, $p = 0.022$), but this did not survive FDR correction ($p_{\text{corrected}} = 0.065$). No significant effects were found for the remaining HR sub-groups in this region (Table 2).

Temporal lobe

No associations between PGRS-SCZ and temporal gyrification reached significance in the temporal lobe for either the HR cohort, HR cohort split by subsequent clinical outcome or in controls (online Supplementary Data, Tables S3 and S4).

Group comparisons in mean gyrification

Models comparing mean gyrification between groups failed to reach significance within any of the areas of interest (online Supplementary Data, Table S5). However, inspection of the means suggested that there was a pattern of incremental increase in gyrification for participants at HR who went on to develop symptoms and transition into schizophrenia (Table 3).




Fig. 2. Scatter plots showing a positive relationship between PGRS-SCZ and frontal gyrification in the (a) left and (b) right hemisphere. Y-axis represents (a) left and (b) right frontal gyrifications. X-axis represents PGRS-SCZ.

Table 2. Significance and effect size values for the effect of polygenic risk for schizophrenia on the left and right frontal lobes, within the high-risk sub-groups

	HR[well] (n = 31)		HR[symp] (n = 28)		HR[ill] (n = 9)	
	Effect size	p (P _{corrected})	Effect size	p (P _{corrected})	Effect size	p (P _{corrected})
Left frontal	0.037	0.360 (0.383)	0.237	0.022 (0.065)	0.680	0.383 (0.383)
Right frontal	0.002	0.818 (0.818)	0.347	0.004 (0.012)	0.983	0.084 (0.126)

Effect size = partial η^2 .

Table 3. Mean gyrification for each group within the regions of interest, along with their standard deviations

	HC (n = 16) Mean (s.d.)	HR[well] (n = 31) Mean (s.d.)	HR[symp] (n = 28) Mean (s.d.)	HR[ill] (n = 9) Mean (s.d.)
Left frontal	2.69 (0.13)	2.67 (0.11)	2.69 (0.13)	2.75 (0.11)
Right frontal	2.69 (0.12)	2.67 (0.12)	2.70 (0.16)	2.75 (0.11)
Left temporal	3.25 (0.22)	3.24 (0.15)	3.27 (0.17)	3.34 (0.15)
Right temporal	3.20 (0.18)	3.18 (0.15)	3.23 (0.18)	3.30 (0.12)

Discussion

To our knowledge, the present study is the first to examine PGRS-SCZ in a prospectively ascertained familial HR cohort, and to show that such scores are associated with transition to schizophrenia. The current results, along with the recent finding that the PGRS-SCZ can be used to predict between FEPs who go on to develop schizophrenia or other psychoses (Vassos *et al.* 2017), have important implications for our understanding of the aetiology of schizophrenia. It is likely, for example, that genetic and environmental factors, and their interactions, determine who of those at elevated genetic risk go on to get a broader phenotype of psychotic symptoms or the narrower phenotype of schizophrenia. Although PGRS are currently underpowered to be of clinical utility in a predictive capacity, they are sufficiently well-powered to test for association between different phenotypes and the PGRS (Dudbridge, 2013). Therefore, using neuroimaging phenotypes in combination with PGRS could become a useful tool in clinical settings in the future. The utility of both measures will increase as collaborative efforts yield ever-larger sample sizes; however, due to the current sample size, the results reported here should be interpreted with caution.

This study is also the first to examine the relationship between PGRS-SCZ and gyrification within a HR sample. We found that within the HR cohort, but not controls, an increased genetic load for schizophrenia was positively and significantly associated with increased bilateral frontal gyrification. As there is a temporal relationship between patterns of neuronal migration and patterns of gyrification, with disorders that affect migration early having more pronounced alterations in the pattern of gyrification, our results suggest that a mechanism by which an increased genetic load for schizophrenia increases frontal gyrification may be involved in the pathogenesis of the disorder. However, the current results require replication in a large data set to provide further evidence for this theory. Nevertheless, our current findings are in keeping with our own and other earlier studies, using less automated measures of GI which found an association with familial

risk for schizophrenia and propensity to develop the disorder (Vogelely *et al.* 2000; Harris *et al.* 2004, 2007).

We also found that increasing PGRS-SCZ was related to right frontal gyrification in the HR[symp] and HR[ill] sub-groups of the HR cohort. Hence, those at HR who developed schizophrenia had a significantly greater proportion of risk alleles for schizophrenia which, for the HR[symp] cohort and, at trend level for the HR[ill] cohort, correlated with gyrification estimates. Although, this finding was only at trend level for the HR[ill] sub-group, the result was also associated with a large effect size and it is suggested that the lack of significance was a result of low numbers within this group. The interaction effect of group and PGRS-SCZ was not found within the left frontal lobe; however, a trend-level effect was found within the HR[symp] group individually.

It is also important to note that, analysis of the HR group as a whole, provided significant effects bilaterally. Thus, it may be that this polygenic risk-influenced gyrification is a 'trait' marker of schizophrenia, and provides a putative mechanism by which genetic influences affect subsequent neurodevelopment and increase an individual's risk of developing schizophrenia. Further analysis of cohorts, using a larger sample size, are required to determine if this result is related to the HR status as a whole or can be predicted by subsequent clinical outcome. As schizophrenia is such a complex and heterogeneous disorder, however, it is unlikely to be fully mediated by this one mechanism. It is likely that other genetic and environmental risks, through development and nearer to the time of onset of the disorder, interact with the polygenic risk-influenced gyrification differences which we have found.

Lack of an effect within the temporal lobe was not entirely unexpected. Previous research has linked temporal lobe abnormalities in gyrification to participants with developed schizophrenia (White *et al.* 2003; Harris *et al.* 2004) rather than those at HR (Harris *et al.* 2007), suggesting that this deficit may be more related to the impact of the disorder rather than its genetic architecture. It would be of interest for future studies to further investigate this by replicating the current result within an independent

HR sample as well as to conduct these analyses within a patient population.

Contradictory to our previous findings within this sample (Harris *et al.* 2007), we did not find that gyrification predicted group status within the present study. However, investigation of the means suggests that there is an incremental increase in both the frontal and temporal lobes in those that develop symptoms and subsequently transition to schizophrenia. As outlined in the introduction, it is important to note here that the research into the link between gyrification and schizophrenia has been fairly inconsistent. However, within the current study, it is suggested that the lack of a significant effect may be due to the small sample size; as only a sub-section of the previous sample had genotyped information that could be used in the current analyses. Future studies should further investigate the relationship between transition to schizophrenia and gyrification and, in turn, the association with PGRS-SCZ.

The main limitation within this study is the small sample size, particularly in relation to the HR[ill] group, which may impede the generalizability of the results. Nevertheless, the incremental deviation from HC in gyrification estimates across the HR subgroups (Table 3) along with the case-control differences in PGRS-SCZ (Fig. 1) found within the current study, which are in keeping with previous findings (Harris *et al.* 2007; Falkai *et al.* 2007; Palaniyappan *et al.* 2011; Vassos *et al.* 2017), lend credibility to the results.

Another limitation relates to the parameters of the MRI scans for use with FreeSurfer. The recommended guidelines suggest 1.5–3 T with 1 mm³ isotropic resolution; however, the current MR images were collected at 1 T with a voxel size of 1.8 mm³ (<https://surfer.nmr.mgh.harvard.edu/fswiki/>). Although we are working with slightly lower spatial resolution parameters than those recommended by FreeSurfer, we have attempted to overcome this potential issue by manually editing all of the data and checking the fidelity of all of the imaging reconstructions. Moreover, by quantifying large-scale shape parameters such as LGI, which is essentially calculated over 25 mm of the cortical surface (Schaer *et al.* 2008), we are less likely to suffer from voxel-based inaccuracies than, for example, a per-vertex thickness measure that depends on grey-white contrast ratio. Furthermore, the current results rely on a comparison between subjects acquired on the same scanner, and the fact that we only see results in the frontal (and not temporal) regions suggests that the lower field strength does not hamper such comparisons. Thus, despite utilising slightly lower scan resolution than recommended, these results still provide meaningful insight into the relationship between polygenic loading for schizophrenia and frontal gyrification.

In conclusion, we have found that an increased genetic loading for schizophrenia was present in those who developed schizophrenia, compared with those at HR who developed symptoms, remained well or controls. This increased loading was associated with altered bilateral frontal gyrification within the HR cohort, but this association was not present in the control group. Limited evidence for this effect being dependent on subsequent clinical outcome was found within the right frontal lobe; however, small sample sizes impeded our ability to investigate this fully. Understanding the genetic architecture of schizophrenia and polygenic effects on brain structure could have critical implications for our knowledge of the aetiology and pathogenesis of the disorder and thus provide us with a means to provide more targeted interventions for people at an elevated familial risk of developing schizophrenia.

Supplementary Material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717003087>.

Acknowledgements. The authors would like to thank all the participants and their families for taking part in this study and the radiographers who acquired the MRI scans at the City Hospital in Edinburgh. The authors would also like to extend their thanks to both Lauren Navrady and Simon R. Cox (University of Edinburgh) for additional help with the statistical analyses. Medical Research Council (MRC) [Grant Numbers: G9226254 and G9825423] and the Dr Mortimer and Theresa Sackler Foundation provided financial support.

Declaration of Interest. The author SML has received financial support for research, in the past 3 years, from Roche Abbvie, Sunovion and Janssen, in relation to therapeutic studies of people with schizophrenia. He has also received personal payments for advisory panels and/or educational meetings from Janssen, Forum and Otsuka. AMM is supported by the Health Foundation through a Clinician Scientist Fellowship (Ref: 2268/4295) and by the National Alliance for Research on Schizophrenia and Depression (NARSAD) through an Independent Investigator Award. AMM has previously received financial support from Janssen and Lilly. HCW is supported by a JMAS SIM fellowship from the Royal College of Physicians of Edinburgh and an ESAT College Fellowship from the University of Edinburgh. The authors AMM, HCW and SML have received financial support from Pfizer (formerly Wyeth) in relation to imaging studies of people with schizophrenia and bipolar disorder. ECJ, DGCO, EN, CB, T-KC and LH report no financial conflicts of interest.

Ethical Standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Bois C *et al.* (2015) Structural magnetic resonance imaging markers of susceptibility and transition to schizophrenia: a review of familial and clinical high risk population studies. *Journal of Psychopharmacology* **29**, 144–154.
- Bonnici HM *et al.* (2007) Pre-frontal lobe gyrification index in schizophrenia, mental retardation and comorbid groups: an automated study. *NeuroImage* **35**, 648–654.
- Boos HB *et al.* (2007) Brain volumes in relatives of patients with schizophrenia: a meta-analysis. *Archives of General Psychiatry* **64**, 297–304.
- Cooper D *et al.* (2014) Multimodal voxel-based meta-analysis of structural and functional magnetic resonance imaging studies in those at elevated genetic risk of developing schizophrenia. *Psychiatry Research* **221**, 69–77.
- Dudbridge F (2013) Power and predictive accuracy of polygenic risk scores. *PLoS Genetics* **9**, e1003348.
- ENIGMA2 Genetics Support Team (2012) *ENIGMA2 1KGP Cookbook (v3). The Enhancing Neuroimaging Genetics through Meta-Analysis (ENIGMA) Consortium*. Viewed 27th July 2017 (http://enigma.ini.usc.edu/wp-content/uploads/2012/07/ENIGMA2_1KGP_cookbook_v3.pdf).
- Falkai P *et al.* (2007) Disturbed frontal gyrification within families affected with schizophrenia. *Journal of Psychiatric Research* **41**, 805–813.
- Fischl B and Dale AM (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences* **97**, 11050–5.
- Fischl B, Liu A and Dale AM (2001) Automated manifold surgery: constructing geometrically accurate and topologically correct models of the human cerebral cortex. *IEEE Transactions of Medical Imaging* **20**, 70–80.
- Fischl B, Sereno MI and Dale AM (1999) Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *NeuroImage* **9**, 195–207.
- Fornito A *et al.* (2012) Schizophrenia, neuroimaging and connectomics. *NeuroImage* **62**, 2296–2314.
- Fraguas D *et al.* (2014) Duration of untreated psychosis predicts functional and clinical outcome in children and adolescents with first-episode psychosis: a 2-year longitudinal study. *Schizophrenia Research* **152**, 130–138.

- Fusar-Poli P et al.** (2011) Neuroanatomy of vulnerability to psychosis: a voxel-based meta-analysis. *Neuroscience & Biobehavioural Reviews* **35**, 1175–1185.
- Genovese CR, Lazar NA and Nichols T** (2002) Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *NeuroImage* **15**, 870–878.
- Gorjanc G, Henderson DA and Kinghorn B, Percy A** (2007) *GeneticsPed: Pedigree and Genetic Relationship Functions*. R package version 1.22.0 (<http://rgenetics.org>).
- Haijma SV et al.** (2013) Brain volumes in schizophrenia: a meta-analysis in over 18 000 subjects. *Schizophrenia Bulletin* **39**, 1129–1138.
- Harris JM et al.** (2007) Increased prefrontal gyrification in a large high-risk cohort characterizes those who develop schizophrenia and reflects abnormal prefrontal development. *Biological Psychiatry* **62**, 722–729.
- Harris JM et al.** (2004) Gyrification in first-episode schizophrenia: a morphometric study. *Biological Psychiatry* **55**, 141–147.
- Harrison PJ and Weinberger DR** (2005) Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Molecular Psychiatry* **10**, 40–68.
- Heinssen RK and Insel TR** (2015) Preventing the onset of psychosis: not quite there yet. *Schizophrenia Bulletin* **41**, 28–29.
- Hodges A et al.** (1999) People at risk of schizophrenia. Sample characteristics of the first 100 cases in the Edinburgh High-Risk Study. *British Journal of Psychiatry* **174**, 547–553.
- Johnstone EC et al.** (2000) Edinburgh high risk study – findings after four years: demographic, attainment and psychopathological issues. *Schizophrenia Research* **46**, 1–15.
- Johnstone EC et al.** (2005) Predicting schizophrenia: findings from the Edinburgh High-Risk Study. *British Journal of Psychiatry* **186**, 18–25.
- Lawrie SM and Abukmeil SS** (1998) Brain abnormality in schizophrenia. A systematic and quantitative review of volumetric magnetic resonance imaging studies. *British Journal of Psychiatry* **172**, 110–120.
- Lee SH et al.** (2012) Estimating the proportion of variation in susceptibility to schizophrenia captured by common SNPs. *Nature Genetics* **44**, 247–250.
- Liu B et al.** (2016) Polygenic risk for schizophrenia influences cortical gyrification in 2 independent general populations. *Schizophrenia Bulletin* **43**, 673–680.
- Mancini-Marie A et al.** (2015) Sex, age, symptoms and illness duration and their relation with gyrification index in schizophrenia. *Clinical Schizophrenia & Related Psychoses* <https://doi.org/10.3371/CSRP.MAYO.070415>.
- McIntosh AM et al.** (2013) Polygenic risk for schizophrenia is associated with cognitive change between childhood and old age. *Biological Psychiatry* **73**, 938–943.
- McIntosh AM et al.** (2016) Genetic and environmental risk for chronic pain and the contribution of risk variants for major depressive disorder: a family-based mixed-model analysis. *PLoS Medicine* **13**, e1002090.
- McIntosh AM et al.** (2009) Prefrontal gyral folding and its cognitive correlates in bipolar disorder and schizophrenia. *Acta Psychiatrica Scandinavica* **119**, 192–198.
- McIntosh AM et al.** (2011) Longitudinal volume reductions in people at high genetic risk of schizophrenia as they develop psychosis. *Biological Psychiatry* **69**, 953–958.
- Nanda P et al.** (2014) Local gyrification index in probands with psychotic disorders and their first-degree relatives. *Biological Psychiatry* **76**, 447–455.
- Palaniyappan L et al.** (2011) Folding of the prefrontal cortex in schizophrenia: regional differences in gyrification. *Biological Psychiatry* **69**, 974–979.
- Purcell S et al.** (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics* **81**, 559–575.
- Purcell SM et al.** (2009) Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* **460**, 748–752.
- Sallet PC et al.** (2003) Reduced cortical folding in schizophrenia: an MRI morphometric study. *American Journal of Psychiatry* **160**, 1606–1613.
- Sasabayashi D et al.** (2017) Increased occipital gyrification and development of psychotic disorders in individuals with an at-risk mental state: a multicenter study. *Biological Psychiatry* <http://dx.doi.org/10.1016/j.biopsych.2017.05.018>.
- Schaer M et al.** (2008) A surface-based approach to quantify local cortical gyrification. *IEEE Transactions on Medical Imaging* **27**, 161–170.
- Schizophrenia Working Group of the Psychiatric Genomics Consortium** (2014) Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421–427.
- Ségonne F, Pacheco J and Fischl B** (2007) Geometrically accurate topology-correction of cortical surfaces using nonseparating loops. *IEEE Transactions on Medical Imaging* **26**, 518–529.
- Sled JG, Zijdenbos AP and Evans AC** (1998) A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE Transactions on Medical Imaging* **17**, 87–97.
- Vassos E et al.** (2017) An examination of polygenic score risk prediction in individuals with first-episode psychosis. *Biological Psychiatry* **81**, 470–477.
- Vogele K et al.** (2000) Disturbed gyrification of the prefrontal region in male schizophrenic patients: a morphometric postmortem study. *American Journal of Psychiatry* **157**, 34–39.
- Whalley HC et al.** (2015) Impact of cross-disorder polygenic risk on frontal brain activation with specific effect of schizophrenia risk. *Schizophrenia Research* **161**, 484–489.
- White T et al.** (2003) Gyrification abnormalities in childhood- and adolescent-onset schizophrenia. *Biological Psychiatry* **54**, 418–426.
- White T and Hilgetag CC** (2011) Gyrification and neural connectivity in schizophrenia. *Development and Psychopathology* **23**, 339–352.
- White T and Gottesman I** (2012) Brain connectivity and gyrification as endophenotypes for schizophrenia: weight of the evidence. *Current Topics in Medicinal Chemistry* **12**, 2393–2403.
- Wing JK, Cooper JE and Sartorius N** (1974) *The Description Classification of Psychiatric Symptoms: An Instruction Manual for the PSE and CATEGO System*. Cambridge University: London.
- Zilles K et al.** (1988) The human pattern of gyrification in the cerebral cortex. *Anatomy and Embryology* **179**, 173–179.