

# New exact multiplicity results with an application to a population model

**Philip Korman**

Department of Mathematical Sciences, University of Cincinnati,  
Cincinnati, OH 45221-0025, USA

**Junping Shi\***

Department of Mathematics, Tulane University, New Orleans,  
LA 70118, USA

(MS received 17 December 1999; accepted 14 September 2000)

We obtain some new exact multiplicity results for the Dirichlet boundary-value problem

$$\Delta u + \lambda f(u) = 0 \quad \text{for } x \in B^n, \quad u = 0 \quad \text{for } x \in \partial B^n$$

on a unit ball  $B^n$  in  $\mathbb{R}^n$ . We consider several classes of nonlinearities  $f(u)$ , including both positive and sign-changing cases. A crucial part of the proof is to establish positivity of solutions for the corresponding linearized problem. As an application we obtain exact multiplicity results for the Holling–Tanner population model.

## 1. Introduction

We study positive solutions of the Dirichlet boundary-value problem depending on a positive parameter  $\lambda$ ,

$$\Delta u + \lambda f(u) = 0 \quad \text{for } x \in B^n, \quad u = 0 \quad \text{for } x \in \partial B^n \quad (1.1)$$

on the unit ball  $B^n$  in  $\mathbb{R}^n$ ,  $n \geq 2$ . From the well-known result of Gidas *et al.* [6], any positive solution of (1.1) is radially symmetric, i.e.  $u = u(|x|)$ , and hence (1.1) reduces to an ordinary differential equation of the form (2.2) below.

The set of positive solutions of (1.1) (or, equivalently, of (2.2)) consists of simple curves in the function space  $\mathbb{R}^+ \times X = \{(\lambda, u)\}$ , where  $X = C^{2,\alpha}(\bar{B}^n)$ , and each solution curve can be parametrized by the maximal value of the solution (see [12] or [18]). To determine the exact shape of the solution curve, and to count the number of solutions for each given  $\lambda > 0$ , it is important to study the singular points on the solution curve. The singular points are where the curve makes a turn (so we called them the *turning points*), and at a turning point  $(\lambda, u)$ , the linearized equation

$$\Delta z + \lambda f'(u)z = 0 \quad \text{for } x \in B^n, \quad z = 0 \quad \text{for } x \in \partial B^n \quad (1.2)$$

has a non-trivial solution  $z$ . A bifurcation theorem of Crandall and Rabinowitz [2] can be applied to (1.1) near a turning point, providing a bifurcation theory approach

\*Present address: Department of Mathematics, College of William and Mary, Williamsburg, VA 23187, USA.

to the exact multiplicity results, which was developed and refined in [9,11,12,18,19]. The key to the bifurcation theory approach is

- (1) prove, for any turning point  $(\lambda, u)$ ,  $z$  can be chosen to be positive; and
- (2) prove that if  $z$  is positive, then there is at most one turning point.

The second part involves an estimate of the integral  $\int_{B^n} f''(u)z^3 dx$ , so both the positivity of  $z$  and the convexity properties of  $f$  are relevant (see lemma 2.5). The first part usually involves comparison arguments of Sturm’s type, and the choice of suitable comparison functions is critical and delicate.

In this paper, we mainly use a comparison function  $v(r) = ru_r(r) + (n-2)u(r) + \mu$  for some unspecified constant  $\mu$ , and it results in a condition on the nonlinearity  $f(u)$ ,

$$2[f'(u)]^2 - nf(u)f''(u) \geq 0 \text{ for all } u > 0 \quad (\text{or } c > u > 0). \tag{1.3}$$

Under (1.3) and some other conditions on  $f$ , we are able to show the positivity of  $z$ , and consequently we obtain some new exact multiplicity results for (1.1). The test function  $v$  and the condition (1.3) also appeared in [10], where the problem (1.1) with a nonlinearity  $f$  satisfying  $f(0) < 0$  was studied. We will consider the nonlinearities  $f$  satisfying (a)  $f(0) \geq 0$ , and  $f$  is asymptotically superlinear; (b) there is a constant  $c > 0$  such that  $f(u) > 0$  for  $u \in (0, c)$  and  $f(c) = 0$ ; and (c) there exists  $c > b > 0$  such that  $f(u) < 0$  for  $u \in (0, b)$ ,  $f(u) > 0$  for  $u \in (b, c)$  and  $f(0) = f(b) = f(c)$ . In all of these cases, we assume  $f(u)$  to satisfy (1.3).

In §3, we apply the exact multiplicity results to the Holling–Tanner population model,

$$\Delta u + \lambda \left( mu - u^2 - \frac{ku}{1 + u} \right) = 0 \quad \text{for } x \in B^n, \quad u = 0 \quad \text{for } x \in B^n, \tag{1.4}$$

with constants  $k, m > 0$ . The solutions of (1.4) are the steady-state solutions of the corresponding reaction–diffusion equation,

$$\left. \begin{aligned} u_t &= \Delta u + \lambda \left( mu - u^2 - \frac{ku}{1 + u} \right) && \text{for } x \in B^n, \\ u(x, t) &= 0 && \text{for } x \in \partial B^n, \\ u(x, 0) &= u_0(x) && \text{for } x \in B^n. \end{aligned} \right\} \tag{1.5}$$

When  $k = 0$ ,  $f_0(u) = mu - u^2$  and (1.4) is the well-known diffusive logistic equation, which has wide applications in population biology modelling. When  $k > 0$ , the term  $-ku/(1 + u)$  is one example of a predation term. Here,  $u$  is considered to be a population of prey, whose growth rate is decreased because of the existence of some predators. The predation term  $-ku/(1 + u)$  was introduced by Holling and Tanner (see, for example, [7]). We prove that condition (1.3) is satisfied for nonlinearity here when  $1 \leq n \leq 4$  and we completely classify the bifurcation diagrams for all parameters  $k > 0, m > 0$  and  $1 \leq n \leq 4$ . Our results allow us to characterize the attractor of (1.5) for any possible combination of parameters.

We mention that Du and Lou [5] have studied the bifurcation curve of solutions of an elliptic system with Holling–Tanner-type interaction of two species, and, for

some parameters, they obtained an exact S-shaped solution curve. The test function  $v(r) = ru_r(r) + (n - 2)u(r) + \mu$  was also used in [10]. But the special case of  $v(r) = ru_r(r) + (n - 2)u(r)$  was well known (see [13, 17, 19]), and, on the other hand, when  $n = 2$ , Du and Lou [4] used a test function  $v(r) = ru_r(r) + \mu$  in studying a problem from combustion theory. When  $n = 1$ , Schaaf [20] also obtained some exact multiplicity results, with  $f$  satisfying similar properties as (1.3). More previous results on using test functions to obtain the exact multiplicity results for (1.1) can be found in [9, 11, 12, 14, 18, 19].

Throughout the paper,  $\lambda_1$  is the principal eigenvalue of the Laplacian on  $B^n$  and  $\phi(x) > 0$  the corresponding eigenfunction. Also, a solution of (1.1) is always meant a positive solution of (1.1).

## 2. Exact multiplicity results

We consider the positive solutions of the Dirichlet problem

$$\Delta u + \lambda f(u) = 0 \quad \text{for } x \in B^n, \quad u = 0 \quad \text{for } x \in \partial B^n \tag{2.1}$$

on the unit ball  $B^n$  in  $\mathbb{R}^n$ ,  $n \geq 2$ . By the well-known result of Gidas *et al.* [6], any positive solution of (2.1) is radially symmetric, and hence it satisfies

$$u'' + \frac{n-1}{r}u' + \lambda f(u) = 0 \quad \text{for } 0 < r < 1, \quad u'(0) = u(1) = 0, \tag{2.2}$$

where  $r = |x|$ . The corresponding linearized problem is

$$\Delta z + \lambda f'(u)z = 0 \quad \text{for } x \in B^n, \quad z = 0 \quad \text{for } x \in \partial B^n, \tag{2.3}$$

and if  $u$  is a positive radially symmetric solution of (2.1), then it was shown by Lin and Ni [15] that any solution of (2.3) is also radially symmetric, and thus it satisfies

$$L[z] \equiv z'' + \frac{n-1}{r}z' + \lambda f'(u)z = 0 \quad \text{for } r < 1, \quad z'(0) = z(1) = 0. \tag{2.4}$$

Recall that a solution  $u$  of (2.2) is called *unstable* if the principle eigenvalue  $\mu_1$  of  $L[z] + \mu z = 0$ ,  $z'(0) = z(1) = 0$  is negative, otherwise we say it is *stable*. When  $\mu_1 < 0$ , the number of negative eigenvalues is called the *Morse index* of  $u$ . When  $\mu_1 = 0$ , the solution is *degenerate*, that is, equation (2.4) has a non-trivial solution, the corresponding eigenfunction  $z$  is of one sign and it can be chosen to be positive. For determining the precise bifurcation diagram of (2.2), it has been shown (see [9, 11, 12, 18, 19]) the importance of proving  $\mu_1 = 0$  at a degenerate solution (instead of  $\mu_k = 0$  for  $k > 1$ ). This is equivalent to excluding the possibility of a higher Morse index solution of (2.2).

The main tool for proving  $z > 0$  is a comparison lemma.

LEMMA 2.1. *Suppose  $L[z](t) = 0$ ,  $z \not\equiv 0$ . If there exists  $v \in C^2[a, b]$  such that  $L[v](t) \cdot v(t) \leq (\not\equiv) 0$ , then  $z$  has at most one zero in  $[a, b]$ . In addition, if  $z'(a) = 0$ , then  $z$  does not have any zero in  $[a, b]$ .*

For the proof, we refer to [10, 18]. Our first result in proving  $\mu_1 = 0$  (or  $z > 0$ ) is the following.

LEMMA 2.2. Assume that the function  $f(u) \in C^2[0, \infty)$  satisfies  $f(0) \geq 0$ , condition (1.3) for all  $u > 0$  and

$$f'(u) > 0 \quad \text{for all } u > 0. \tag{2.5}$$

Then any non-trivial solution  $z$  of the linearized problem (2.4) is of one sign, i.e. we may assume that  $z(r) > 0$  for all  $r \in [0, 1]$ .

*Proof.* We consider a test function  $v = ru'(r) + (n - 2)u(r) + \alpha$  with a constant  $\alpha$  to be specified. Recall that we denote the left-hand side of the linearized equation for (2.4) by  $L[z]$ . Compute

$$L[v] = \lambda[(n - 2)uf'(u) - nf(u) + \alpha f'(u)] \equiv \lambda g_\alpha(u(r)). \tag{2.6}$$

The sign of the test function  $v(r)$  is governed by the function

$$\alpha = h(r) \equiv -ru'(r) - (n - 2)u(r).$$

Indeed,  $v > 0$  ( $< 0$ ) when  $h(r) < \alpha$  ( $> \alpha$ ). Similarly, the sign of  $g_\alpha(u)$  is governed by

$$\alpha = j(r) \equiv \frac{nf(u(r)) - (n - 2)uf'(u(r))}{f'(u(r))}.$$

This time, in view of (2.5),  $g_\alpha(u) > 0$  ( $< 0$ ), provided  $\alpha > j(r)$  ( $\alpha < j(r)$ ). Notice that  $h(0) = -(n - 2)u(0) \leq 0$ ,  $h(1) = -u'(1) > 0$  and

$$h'(r) = -ru'' - (n - 1)u' = \lambda r f(u) > 0, \tag{2.7}$$

since, by our conditions,  $f(u)$  is positive. Also,  $j(1) \geq 0$  and

$$j'(r) \leq 0, \tag{2.8}$$

in view of the condition (1.3). It follows that only two cases are possible.

CASE 1. The functions  $h(r)$  and  $j(r)$  intersect exactly once on  $[0, 1]$ , say at  $r = r_0$ . We select  $\alpha = h(r_0) = j(r_0)$ . Then, on the interval  $[0, r_0)$ , we have  $v > 0$  and  $L[v] < 0$ , while on the interval  $(r_0, 1)$  the opposite inequalities hold. Lemma 2.1 implies that  $z(r)$  cannot have any roots and hence is of one sign on  $[0, 1]$ .

CASE 2.  $j(r) \geq h(r)$  for all  $r \in [0, 1]$ . This time we select  $\alpha = h(1) = \max_{r \in [0, 1]} h(r)$  to obtain  $v > 0$  and  $L[v] < 0$  on the entire interval  $[0, 1]$ . We again conclude that  $z(r)$  is of one sign. □

REMARK 2.3. In several previous works on the exact multiplicity results for (2.2), the function  $K_f(u) = uf'(u)/f(u)$  also played an important role when proving results of lemma 2.2 type.  $K_f(u)$  can be thought of as an indicator of the growth rate of  $f(u)$ , for example,  $K_f(u) = p$  if  $f(u) = u^p$ . It was shown in [19] that if  $f(u) > 0$  for  $u > 0$ ,  $K_f(u)$  is decreasing or  $K_f(u) \leq n/(n - 2)$ , then any non-trivial solution  $z(r)$  of (2.4) is of one sign. We point out that the condition (1.3) is also related to  $K_f$ . In fact, condition (1.3) is equivalent to saying that the function

$$A(u) = \frac{nf(u) - (n - 2)uf'(u)}{f'(u)}$$

is increasing and, since  $A(u) = u[nK_f^{-1}(u) - (n - 2)]$ , we have

$$A'(u) = \frac{n - 2}{K_f(u)} \left[ \frac{n}{n - 2} - K_f(u) \right] - \frac{nuK'_f(u)}{K_f^2(u)}.$$

So (1.3) holds if  $K_f(u)$  is decreasing and  $K_f(u) \leq n/(n - 2)$ . Also, if (1.3) holds, then, for any  $u > 0$ , either  $K_f(u)$  is decreasing or  $K_f(u) \leq n/(n - 2)$ . This observation will be useful in the proof of lemma 2.15 later.

REMARK 2.4. Condition (1.3) also implies some asymptotic growth restriction on  $f$ . In fact, condition (1.3) is equivalent to

$$\left( \frac{f}{f'} \right)' - \frac{n - 2}{n} \geq 0, \tag{2.9}$$

and, by integration, we obtain

$$f(u) \leq ae^{bu} \quad (n = 2) \quad \text{and} \quad f(u) \leq a(u + b)^{n/(n-2)} \quad (n \geq 3) \tag{2.10}$$

for all  $u \geq 0$  and some  $a, b > 0$ .

If the solution  $z$  of the linearized equation (2.4) is shown to be of one sign, there is a well-established theory on the set of positive solutions of (2.2), which we briefly review here. The details (and also the proof of all quoted facts) can be found in [9, 11, 12, 18, 19].

From the uniqueness of ordinary differential equations, for any  $s > 0$ , there is at most one  $\lambda(s) > 0$  such that (2.2) has a positive solution  $u(\cdot, s)$  with  $\lambda = \lambda(s)$  and  $u(0) = s$ . So the set of positive solutions of (2.2) can be globally parametrized by  $s = u(0)$ , thus the solution set is a curve of the form  $\{(\lambda(s), s)\}$ , where  $s > 0$  belongs to a certain admissible set. If  $\lambda'(s) \neq 0$ , then the corresponding solution  $u(\cdot, s)$  is non-degenerate, while if  $\lambda'(s) = 0$ , then the solution is degenerate. At a degenerate solution, we can show that

$$\lambda''(s) = \frac{-\lambda(s) \int_0^1 r^{n-1} f''(u(r)) z^3(r) dr}{\int_0^1 r^{n-1} f(u(r)) z(r) dr}, \tag{2.11}$$

where  $z$  is a non-trivial solution of (2.4). Here we assume that  $z(r) > 0$  for  $r \in [0, 1]$ . For the denominator in (2.11), we can show (see [18] or [9]) that

$$\int_0^1 r^{n-1} f(u(r)) z(r) dr = \frac{1}{2\lambda(s)} u'(1) z'(1) > 0 \tag{2.12}$$

if  $u'(1) < 0$  and  $z'(1) < 0$ , which are both true if  $f(0) \geq 0$ . So the direction of the turn of the bifurcation curve is mainly determined by the integral  $\int_0^1 r^{n-1} f''(u(r)) z^3(r) dr$ . Here we recall the following results from [19].

LEMMA 2.5. *Suppose that  $(\lambda(s), u(s))$  is a degenerate solution of (2.2),  $z$  is the corresponding solution of linearized equation (2.4) and  $z(r) > 0$  for  $r \in [0, 1]$ .*

- (1) *If  $f''(u) > 0$  for  $u > 0$ , then  $\lambda''(s) < 0$ .*
- (2) *If  $f''(u) < 0$  for  $u > 0$ , then  $\lambda''(s) > 0$ .*




Figure 1. Theorem 2.6.

- (3) If  $f(0) \geq 0$  and there exists  $\beta > 0$  such that  $f''(u) < 0$  for  $u$  in  $(0, \beta)$ ,  $f''(u) > 0$  for  $u$  in  $(\beta, \infty)$  and  $u(0) > \beta$ , then  $\lambda''(s) > 0$ .
- (4) If  $f(0) \leq 0$ , and there exists  $\beta > 0$  such that  $f''(u) > 0$  for  $u$  in  $(0, \beta)$ ,  $f''(u) < 0$  for  $u$  in  $(\beta, \infty)$  and  $u(0) > \beta$ , then  $\lambda''(s) < 0$ .

Combining lemmas 2.2 and 2.5, we have the following exact multiplicity result.

**THEOREM 2.6.** Assume  $f(u) > 0$  for  $u \geq 0$ ,  $f''(u) \geq 0$ ,  $\lim_{u \rightarrow \infty} f(u)/u = \infty$  and  $f$  satisfies (1.3) and (2.5). Then there exists a constant  $\lambda_0 > 0$  such that problem (2.2) has no solution for  $\lambda > \lambda_0$ , exactly two solutions for  $\lambda < \lambda_0$  and exactly one solution for  $\lambda = \lambda_0$ . Moreover, all solutions lie on a unique smooth solution curve (see figure 1).

*Proof.* Since  $f(0) > 0$ , then  $u = 0$  is a non-degenerate solution of (2.2) when  $\lambda = 0$  and, by the implicit function theorem, for any small  $\lambda > 0$ , there is a unique solution  $u(\lambda, \cdot)$  of (2.2) near  $u = 0$ . Furthermore,  $u(\lambda, \cdot)$  is positive by the maximum principle. From the remarks before lemma 2.5,  $u(\lambda, \cdot)$  can also be parametrized by  $u(\lambda, 0)$ . So a solution curve  $\Sigma = (\lambda(s), u(s))$  emerges from  $(\lambda, u) = (0, 0)$  and moves to the right as  $s$  increases. On the other hand, problem (1.1) has no positive solution for  $\lambda > 0$  large. In fact, if we multiply (1.1) by  $\phi$ , the principal eigenfunction of  $\Delta\phi + \lambda\phi = 0$ ,  $\phi(x) = 0$  for  $x \in \partial B^n$ , then we have

$$\lambda_1 \int_{B^n} u\phi \, dx = \lambda \int_{B^n} f(u)\phi \, dx \geq \lambda a \int_{B^n} u\phi \, dx, \tag{2.13}$$

where  $a = \min_{u \geq 0} f(u)/u > 0$ . Thus (1.1) has a positive solution only if  $\lambda \leq \lambda_1 a^{-1}$ . Thus the solution curve cannot continue to  $\lambda = \infty$ . Let

$$\lambda_* = \sup\{\lambda > 0 : \text{problem (2.2) has a positive solution with this } \lambda\}.$$

Then  $0 < \lambda_* \leq \lambda_1 a^{-1}$ .

We claim that there is a turning point (degenerate solution) on the solution curve  $\Sigma$ . Suppose there are no turning points, then  $\lambda'(s) > 0$  for all  $s > 0$  and  $\lim_{s \rightarrow \infty} \lambda(s) = \lambda_*$ . Since  $s = u(0)$  is a global parameter for all positive solutions, it follows that there are no positive solutions other than those on  $\Sigma$ . In particular, problem (2.2) has a unique positive solution for  $\lambda \in (0, \lambda_*)$  and no positive solution for  $\lambda \geq \lambda_*$ . However, since  $f$  satisfies the growth condition (2.10) and  $f(0) > 0$ , by a result of Lions [16, theorem 2.1 and remark 1.1], for the case  $n > 2$ , the problem (1.1) has at least two positive solutions for  $0 < \lambda < \lambda^*$ , which is a contradiction. For




Figure 2. Theorem 2.9.

the case  $n = 2$ , we use theorem 2.3 in [3] to conclude the existence of at least two positive solutions for  $0 < \lambda < \lambda^*$ . (Since in [3] it was assumed that  $f(0) = 0$ , the following modification is needed: to prove existence of the second solution, we apply the mountain pass lemma, using the solution bifurcating from zero (rather than the trivial solution) as a minimizer.) Thus there is a turning point on  $\Sigma$ .

At any turning point, by lemma 2.2,  $z$  can be chosen as positive and hence, by lemma 2.5,  $\lambda''(s) < 0$ , so the curve turns to the left and, after passing the turning point, the curve travels to the left. There are no other turning points, since at any turning point, the solution curve turns to the left, but when  $\Sigma$  moves to the left, it has to approach a possible turning point from the right. Therefore,  $\Sigma$  is monotone ( $\lambda'(s) < 0$ ) above the turning point. Let  $\bar{\lambda} = \lim_{s \rightarrow \infty} \lambda(s)$ . Then  $\bar{\lambda} \geq 0$ . Since, from [16], problem (2.2) has at least two solutions for all  $0 < \lambda < \lambda_*$ , then  $\bar{\lambda} = 0$ . □

EXAMPLE 2.7. As a very particular case, we recover the well-known result of Joseph and Lundgren [8] for the case  $f(u) = e^u$  and  $n = 2$  (actually, we get some extra information even in this case).

EXAMPLE 2.8. We have another application of lemma 2.2. In [1], Adimurthi gives an ingenious proof that the problem

$$\Delta u + \lambda u e^u = 0 \quad \text{for } B^2, \quad u = 0 \quad \text{for } x \in \partial B^2 \tag{2.14}$$

on a ball  $B^2$  in  $\mathbb{R}^2$  has at most one positive solution. We have a more detailed result. Let  $\lambda_1$  denote the principal eigenvalue of the Laplacian on the unit ball in  $\mathbb{R}^2$ .

THEOREM 2.9. *The set of positive solutions of (2.14) consists of a smooth solution curve, which bifurcates (to the left) at  $\lambda = \lambda_1$  from the trivial solution  $u = 0$ , and continues for all  $0 < \lambda < \lambda_1$ , tending to infinity as  $\lambda \rightarrow 0$  (see figure 2).*

*Proof.* It is well known that the bifurcation at  $\lambda = \lambda_1$  occurs. First we claim that the bifurcation is subcritical. In fact, we can compute  $\lambda'(0)$  and

$$\lambda'(0) = \frac{-\lambda_0 f''(0) \int_0^1 r^{n-1} \phi^3(r) \, dr}{2f'(0) \int_0^1 \phi^2(r) \, dr}, \tag{2.15}$$

so  $\lambda'(0) < 0$  as  $f''(0) > 0$ ,  $f'(0) > 0$  and  $\phi > 0$ . Thus  $\Sigma$  travels to the left initially near  $\lambda = \lambda_1$ . Since the nonlinearity is convex, while, by lemma 2.2, any non-trivial solution of the linearized problem is of one sign, it follows by lemma 2.5 that the

solution curve continues to the left without any turning point (degenerate solution). It follows from theorem 2.3 of [3] that the solution curve cannot go to infinity at a positive  $\lambda$ , while for  $\lambda = 0$  infinity is the only place the curve can go. (The result in [3] implies that for any sufficiently small  $\lambda$ , problem (2.14) has a positive solution. If the solution curve were to go to infinity at some positive  $\lambda_0$ , then all possible values of  $u(0, \lambda)$  would have been taken, and hence no positive solutions could exist for  $0 < \lambda < \lambda_0$ , which is a contradiction.) □

**EXAMPLE 2.10.** As our last example for lemma 2.2, we point out that for  $f(u) = e^{-1/(u+\varepsilon)}$  and  $\varepsilon > 0$ , problem (1.3) is satisfied for  $n = 2$ . In fact, Du and Lou [4] used the test function  $v(r) = ru_r(r) + \alpha$  to prove  $z > 0$  for that special example.

Next, we turn to the results for  $f$  not always increasing. Lemma 2.2 can be modified as follows.

**LEMMA 2.11.** *Assume that the function  $f(u) \in C^2[0, \infty)$  satisfies  $f(0) \geq 0$ ,  $f(u) > 0$  for  $0 < u < c$  and there exists  $b \in (0, c)$  such that*

$$f'(u) > 0 \quad \text{for } 0 < u < b, \quad f'(u) < 0 \quad \text{for } b < u < c, \tag{2.16}$$

*and  $f$  satisfies (1.3). Then any non-trivial solution  $z$  of the linearized problem (2.4) is of one sign, i.e. we may assume that  $z(r) > 0$  for all  $r \in [0, 1)$ .*

*Proof.* Let  $u$  be a degenerate solution of (2.2). If  $0 < u(0) < b$ , then  $f'(u(r)) > 0$  for  $r \in (0, 1)$ , and we can just use the proof of lemma 2.2. So we assume that  $u(0) > b$ , and so there exists  $r_0 \in (0, 1)$  such that  $u(r_0) = b$ . Then, as in lemma 2.2,  $h(0) \leq 0$ ,  $h(1) > 0$  and  $h'(r) > 0$ . On the other hand, for  $r \in [0, r_0)$ ,  $L[v(r)] = \lambda g_\alpha(u(r)) < 0$  for any  $\alpha > 0$  since  $f'(u) < 0$  for  $u \in (b, c)$ . For  $r \in (r_0, 1)$ , from (1.3), we still have  $j'(r) \leq 0$  and  $j(1) \geq 0$ . Observe also that  $j(r) \rightarrow +\infty$  as  $r \downarrow r_0$ . So only the following two cases are possible.

**CASE 1.** The functions  $h(r)$  and  $j(r)$  intersect exactly once on  $(r_0, 1)$ , say at  $r = r_1$ . We select  $\alpha = h(r_1) = j(r_1) > 0$ . Then on the interval  $[0, r_1)$  we have  $v > 0$  and  $L[v] < 0$ , while on the interval  $(r_0, 1)$  the opposite inequalities hold. So we can proceed as in lemma 2.2.

**CASE 2.**  $j(r) \geq h(r)$  for all  $r \in (r_0, 1)$ . Again, we select

$$\alpha = h(1) = \max_{r \in [0, 1]} h(r) > 0$$

to obtain  $v > 0$  and  $L[v] < 0$  on the entire interval  $[0, 1)$ . Thus we have  $z \neq 0$  for  $r \in [0, 1)$ . □

Applying lemma 2.11, we obtain two exact multiplicity results as follows.

**THEOREM 2.12.** *Assume  $f(0) = 0$ ,  $f'(0) > 0$  and  $f(u) > 0$  for  $u \in (0, c)$ , where  $0 \leq c \leq \infty$ . Assume  $f$  satisfies (1.3) and, for some  $c > \beta > 0$ , we have*

$$f''(u) > 0 \quad \text{for } 0 \leq u < \beta, \quad f''(u) < 0 \quad \text{for } \beta < u < c. \tag{2.17}$$

*If  $c = \infty$ , we also assume  $\lim_{u \rightarrow \infty} f(u)/u = 0$ . Define  $\lambda_0 = \lambda_1/f'(0)$ . Then there exists a constant  $\lambda_* < \lambda_0$  such that problem (2.2) has no solution for  $\lambda < \lambda_*$ , exactly*






Figure 3. Theorem 2.12.

two solutions for  $\lambda_0 > \lambda > \lambda_*$  and exactly one solution for  $\lambda = \lambda_*$  and  $\lambda \geq \lambda_0$ . Moreover, all solutions lie on a unique smooth solution curve (see figure 3).

*Proof.* It is well known that if  $f(0) = 0$  and  $f'(0) > 0$ , then  $\lambda_0 = \lambda_1/f'(0)$  is a bifurcation point for (1.1) and there is a solution curve bifurcating from  $(\lambda, u) = (\lambda_0, 0)$ . Similar to theorem 2.6, the solution curve  $\Sigma = (\lambda(s), u(s))$  has a global parameter  $s = u(0)$ . And, similar to theorem 2.9,  $\lambda'(0) < 0$  and  $\Sigma$  travels to the left initially near  $\lambda = \lambda_0$ .

Next we prove that (1.1) has no positive solution for small  $\lambda > 0$ . In fact, from the conditions on  $f$ , there exists  $a > 0$  such that  $f(u) \leq au$  for all  $u > 0$ . If we multiply (1.1) by  $u$  and integrate, then

$$\lambda_1 \int_{B^n} u^2 \, dx \leq \int_{B^n} |\nabla u|^2 \, dx = \lambda \int_{B^n} f(u)u \, dx \leq \lambda a \int_{B^n} u^2 \, dx, \tag{2.18}$$

so  $\lambda \geq \lambda_1 a^{-1}$ . Hence  $\Sigma$  cannot continue left to  $\lambda = 0$ ; it cannot blow up to  $s = \infty$  either, since  $s = u(0) < c$ , as follows from the maximum principle for the case  $c < \infty$ , and by an easy *a priori* estimate for the case  $c = \infty$ . So it has to bend back at some turning point.

By the assumptions,  $f$  satisfies either (2.5) or (2.16), so  $z > 0$  by lemmas 2.2 or 2.11. Similar to theorem 2.6, there is a unique turning point on  $\Sigma$  from lemma 2.5 (the fourth case) and, at the turning point, the curve bends to the right. Above the turning point,  $\lambda'(s) > 0$ , and so  $\Sigma$  moves to the right monotonously. By an argument in [11] (see also [19]), we can show that there are no other branches and, for the solutions on  $\Sigma$ , when  $\lambda \rightarrow \infty$ , the solution uniformly converges to  $c$  for any compact subset of  $(0, 1)$ . □

**EXAMPLE 2.13.** An example for theorem 2.12 is  $f(u) = -u(u - 2)(u + 1)$  and  $n = 2, 3$ . It is easy to check that  $f$  satisfies (2.16) and (2.17) for  $u > 0$  with  $b = \frac{1}{3}(1 + \sqrt{7})$ ,  $c = 2$  and  $\beta = \frac{1}{3}$ . For (1.3), we obtain

$$\begin{aligned} 2[f']^2 - nff'' &\equiv B(u) \\ &= 2[(9 - 3n)u^4 + (4n - 12)u^3 + (5n - 8)u^2 + (8 - 2n)u + 4]. \end{aligned} \tag{2.19}$$

When  $n = 3$ , we have  $B(u) = 22u^2 + 2u + 4 > 0$  for all  $u > 0$ , and when  $n = 2$ , we have  $B(u) = 6u^4 - 8u^3 + 24u^2 + 8u + 4 > 0$  for all  $u > 0$ .




Figure 4. Theorem 2.14.

**THEOREM 2.14.** *Assume  $f(0) = 0$ ,  $f'(0) = 0$  and  $f(u) > 0$  for  $u \in (0, c)$ , where  $c \leq \infty$ . Assume  $f$  satisfies (1.3) and (2.17). If  $c = \infty$ , we also assume that  $\lim_{u \rightarrow \infty} f(u)/u = 0$ . Then there exists a constant  $\lambda_* > 0$  such that problem (2.2) has no solution for  $\lambda < \lambda_*$ , exactly two solutions for  $\lambda > \lambda_*$  and exactly one solution for  $\lambda = \lambda_*$ . Moreover, all solutions lie on a unique smooth solution curve (see figure 4).*

The proof of this result is same as that of theorem 1.3 of [18], except here we use a different way to prove the positivity of  $z$ , as in lemma 2.11. So we refer the reader to [18] and omit the proof here.

Finally, we consider a problem with sign-changing  $f$ .

**LEMMA 2.15.** *Assume that the function  $f(u) \in C^2[0, \infty)$  satisfies  $f(0) = 0$ , and there exists  $b \in (0, c)$  such that*

$$f(u) < 0 \quad \text{for } 0 < u < b, \quad f(u) > 0 \quad \text{for } b < u < c, \tag{2.20}$$

*$\int_0^c f(u) du > 0$  and  $f$  satisfies (1.3) and (2.17). In addition, we assume that  $n \geq 3$ . Then any non-trivial solution  $z$  of the linearized problem (2.4) is of one sign, i.e. we may assume that  $z(r) > 0$  for all  $r \in [0, 1]$ .*

*Proof.* Since  $\int_0^c f(u) du > 0$ , there exists a unique  $\theta \in (b, c)$  such that

$$F(\theta) = \int_0^\theta f(u) du = 0.$$

For any solution  $u$  of (2.2), we have  $u(0) > \theta$ . In fact, multiplying (2.2) by  $u'$  and integrating over  $(0, 1)$ , we obtain

$$\frac{1}{2}[u'(1)]^2 + (n - 1) \int_0^1 \frac{[u'(r)]^2}{r} dr - \lambda F(u(0)) = 0, \tag{2.21}$$

where  $F(u) = \int_0^u f(t) dt$ . Thus  $F(u(0)) > 0$ , and so there exists  $r_1 \in (0, 1)$  such that  $u(r_1) = \theta$ . From the result of [14] (see also lemma 4.9 of [19]),  $z(r) \neq 0$  for  $r \in [r_1, 1)$ . Next we show that  $z$  has at most one zero on  $(0, r_1]$ . For  $r \in (0, r_1)$ ,  $f(u(r)) > 0$ , so the proof of lemma 2.11 can be carried over to here without changes. But we can only conclude that  $z$  has at most one zero in  $(0, r_1]$ , since in the case of lemma 2.11 we obtain that  $z$  has at most one zero in  $(0, 1]$ , while  $z(1) = 0$ .

Suppose that  $z$  has exactly one zero at some  $r = r_2$ . We exclude this possibility by several steps. Let  $u(r_2) = u_2$ . First we prove  $u_2 < \beta$ . If this is not true, then

$u(r_2) \geq \beta$ , and there exists  $r_3 \geq r_2$  such that  $u(r_3) = \beta$ . Consider the comparison function  $v_1(r) = f(u(r))$ . It is easy to verify that  $L[v_1] = f''(u(r))u_r^2(r)$ . So, in  $[0, r_3]$ ,  $v_1(r) > 0$  and  $L[v_1](r) \leq 0$ , and, by lemma 2.1,  $z$  has no zero in  $[0, r_3]$ , which contradicts with  $r_2 \in [0, r_3]$ .

Next we prove that  $K_f(u_2) > n/(n - 2)$ . Suppose this is not true. Then  $K_f(u_2) \leq n/(n - 2)$ . We claim that  $K_f(u) \leq n/(n - 2)$  for all  $u \in [u_2, u(0)]$ . In fact, from the remark after the proof of lemma 2.2, since  $f$  satisfies (1.3), then, for any  $u \in (0, c)$ , either  $K_f(u) \leq n/(n - 2)$  or  $K'_f(u) \leq 0$ . So, if there exists  $u_3 \in [u_2, u(0)]$  such that  $K_f(u_3) > n/(n - 2)$ , then, for some  $u_4 \in (u_2, u_3)$ ,  $K_f(u_4) > n/(n - 2)$  and  $K'_f(u_4) > 0$ , which contradicts with (1.3). Thus  $K_f(u) \leq n/(n - 2)$  for all  $u \in [u_2, u(0)]$ . Define  $v_2(r) = ru_r(r) + (n - 2)u(r)$  and  $v_3(r) = r^{-1}v_2(r)$ . We calculate that

$$L[v_2] = \lambda[(n - 2)uf'(u) - nf(u)] = \lambda f(u)(n - 2) \left[ K(u) - \frac{n}{n - 2} \right],$$

$$L[v_3] = \lambda \left[ \frac{n - 2}{r}(uf'(u) - f(u)) - (n - 3)\frac{v_2}{r^3} \right].$$

Since  $v'_2(r) = ru''(r) + (n - 1)u'(r) = -\lambda r f(u) < 0$  and  $v_2(0) > 0$ ,  $v_2(1) < 0$ , there exists  $r_4 \in (0, 1)$  such that  $v_2(r) > 0$  in  $(0, r_4)$  and  $v_2(r) < 0$  in  $(r_4, 1)$ . There are two cases to consider.

CASE 1.  $r_4 \geq r_2$ . On  $[0, r_2]$ ,  $v_2(r) \geq 0$  and  $L[v_2](r) \leq 0$ , since  $K_f(u) \leq n/(n - 2)$  for all  $u \in [u_2, u(0)]$ . But this implies that  $z$  has no zero in  $[0, r_2]$ , which contradicts with  $z(r_2) = 0$ .

CASE 2.  $r_4 < r_2$ . On  $[r_2, 1]$ ,  $v_3(r) < 0$  and  $L[v_3](r) \geq 0$ , since  $u_2 = u(r_2) < \beta$ , then  $f''(u(r)) \geq 0$ , which implies  $uf'(u) - f(u) \geq 0$  for  $r \in [r_2, 1]$ . This implies that  $z$  has at most one zero in  $[r_2, 1]$ , which contradicts with  $z(r_2) = z(1) = 0$ .

Therefore,  $K_f(u_2) > n/(n - 2)$ , and then  $K'_f(u_2) \leq 0$  by problem (1.3). Let  $\gamma = K_f(u_2)$ . Then, for any  $u > u_2$ ,  $K_f(u) < \gamma$ , otherwise  $K_f(u)$  would have to be increasing somewhere above the  $\gamma$  level, and hence there is  $u_4 > u_2$  such that  $K_f(u_4) > n/(n - 2)$  and  $K'_f(u_4) > 0$ , which contradicts with (1.3). This implies  $uf'(u) - \gamma f(u) \leq 0$  for  $u > u_2$ . Similarly,  $uf'(u) - \gamma f(u) \geq 0$  for  $u \in [b, u_2]$ . Furthermore, for  $u \in [0, b]$ , we also have  $uf'(u) - \gamma f(u) \geq 0$ , since  $\gamma > 1$  and  $uf'(u) - \gamma f(u) \geq uf'(u) - f(u) \geq 0$ . (The latter inequality holds since  $f''(u) > 0$ , for the case  $\beta > b$ . For the case  $\beta \leq b$ , the lemma follows easily by the remarks in the second paragraph of the present proof.) Hence we obtain that

$$u(r)f'(u(r)) - \gamma f(u(r)) \leq 0 \quad \text{for } r \in [0, r_2], \tag{2.22}$$

$$u(r)f'(u(r)) - \gamma f(u(r)) \geq 0 \quad \text{for } r \in [r_2, 1] \tag{2.23}$$

and  $z(r) > 0$  in  $[0, r_2]$ ,  $z(r) < 0$  in  $(r_2, 1)$ . Combining, we obtain

$$\int_0^1 r^{n-1}[u(r)f'(u(r)) - \gamma f(u(r))]z(r) dr < 0. \tag{2.24}$$

However, on the other hand, from a calculation in [18, lemma 2.3], we have

$$\int_0^1 r^{n-1}[u(r)f'(u(r)) - \gamma f(u(r))]z(r) dr = \frac{1}{2\lambda}(1 - \gamma)u'(1)z'(1) > 0, \tag{2.25}$$




Figure 5. Theorem 2.16.

since  $u'(1) < 0$ ,  $z'(1) > 0$  and  $1 - \gamma < 0$ . This is a contradiction, and  $z$  cannot have exactly one zero in  $[0, 1)$ . Therefore,  $z$  must be of one sign in  $[0, 1)$ .  $\square$

**THEOREM 2.16.** *Suppose that  $f$  satisfies the conditions of lemma 2.15. If  $c = \infty$ , we also assume  $\lim_{u \rightarrow \infty} f(u)/u = 0$ . Then there exists a constant  $\lambda_* > 0$  such that problem (2.2) has no solution for  $\lambda < \lambda_*$ , exactly two solutions for  $\lambda > \lambda_*$  and exactly one solution for  $\lambda = \lambda_*$ . Moreover, all solutions lie on a unique smooth solution curve (see figure 5).*

The proof of theorem 2.16 is same as the results in [11, 12, 18], except that now we use lemma 2.15 to prove  $z > 0$ . So we omit the proof. Some arguments in the proof of lemma 2.15 are similar to the proof in [14, 18].

### 3. Applications to a population growth model

We now apply our exact multiplicity results to the Holling–Tanner population model (1.4). The lines  $m = k$  and  $k = 1$  divide the parameter plane ( $m > 0, k > 0$ ) into four regions. For three of these regions and a part of the fourth we are able to give exact multiplicity results for  $1 \leq n \leq 4$ . Notice that the roots of  $f(u)$  are

$$0 \quad \text{and} \quad \frac{1}{2}[(m - 1) \pm \sqrt{(m - 1)^2 + 4(m - k)}],$$

so that the line  $m = k$  separates the regions where  $f(u)$  has one or two positive roots. As an application of theorems 2.12, 2.14 and 2.16, our result for the population model (1.4) is as follows.

**THEOREM 3.1.** *Suppose that  $m > 0, k > 0$  and  $1 \leq n \leq 4$ , and define  $\lambda_1^0 = \lambda_1/(m - k)$  for the case  $m - k > 0$ .*

- (i) *If  $m > k > 0$  and  $1 \geq k$ , then (1.4) has a unique solution for  $\lambda > \lambda_1^0$  and has no solution for  $\lambda_1^0 \geq \lambda > 0$  (see figure 6).*
- (ii) *If  $m > k > 1$ , then there exists  $\lambda_* \in (0, \lambda_1^0)$  such that (1.4) has exactly two solutions for  $\lambda_1^0 > \lambda > \lambda_*$ , has exactly one solution for  $\lambda = \lambda_*$  and  $\lambda \geq \lambda_1^0$  and has no solution for  $\lambda_* > \lambda > 0$  (see figure 3).*
- (iii) *If  $n \neq 2, k \geq m > 0, (\frac{1}{4})(m + 1)^2 > k > 1$  and*

$$F(u_+) > 0 \quad \text{for } u_+ = \frac{1}{2}[(m - 1) + \sqrt{(m + 1)^2 - 4k}], \tag{3.1}$$




Figure 6. Theorem 3.1(i).




Figure 7. Theorem 3.1.

where

$$F(u) = \int_0^u f(t) dt = \frac{1}{2}mu^2 - \frac{1}{3}u^3 - ku + k \ln(1 + u),$$

then (1.4) has exactly two solutions for  $\lambda > \lambda_*$ , has exactly one solution for  $\lambda = \lambda_*$  and has no solution for  $0 < \lambda < \lambda_*$  (see figures 4 and 5).

(iv) If  $(k, m) \in \mathbb{R}^+ \times \mathbb{R}^+$ , but not in the regions described above, then (1.4) has no solution for  $\lambda > 0$ .

We begin with a lemma.

LEMMA 3.2. Assume that  $1 \leq n \leq 4$ ,  $\frac{1}{4}(m + 1)^2 > k > 1$  and  $f(u) = mu - u^2 - ku/(1 + u)$ . Then  $2[f'(u)]^2 - nf(u)f''(u) \geq 0$  for  $0 \leq u \leq u_+$  if  $u_+ > 0$ , where  $u_+$  is defined in (3.1).

Proof. Note that  $u_+$  is the largest zero of  $f(u)$  if  $\frac{1}{4}(m + 1)^2 > k > 1$ , so  $f(u) < 0$  for  $u > u_+$ . First we compute

$$f'(u) = m - 2u - \frac{k}{(1 + u)^2}, \quad f''(u) = -2 + \frac{2k}{(1 + u)^3}, \quad f'''(u) = -\frac{6k}{(1 + u)^4}.$$

If  $m \geq k$ , then  $f'(0) = m - k \geq 0$  and so  $f(u) > 0$  for  $0 < u < u_+$ . If  $k \leq 1$ , then  $f''(u) \leq 0$  for all  $0 < u < u_+$  and  $M(u) \equiv 2[f'(u)]^2 - nf(u)f''(u) \geq 0$  for all  $0 < u < u_+$ . If  $k > 1$ ,  $f''(u) > 0$  for  $0 < u < \sqrt[3]{k} - 1$  and  $f''(u) < 0$  for

$\sqrt[3]{k} - 1 < u < u_+$ , then again  $M(u) \geq 0$  for  $\sqrt[3]{k} - 1 < u < u_+$ . For  $0 < u \leq \sqrt[3]{k} - 1$ ,  $M'(u) = (4 - n)f'f'' - nff''' > 0$  and  $M(0) = 2[f'(0)]^2 \geq 0$ , then  $M(u) > 0$  for  $0 < u \leq \sqrt[3]{k} - 1$ .

If  $m < k$ , then  $f'(0) < 0$ , and we have  $u_+ > 0$  only if  $\frac{1}{4}(m + 1)^2 > k > 1$ . In such a case, there exists  $u_1 \in (0, u_+)$ , such that  $f(u) < 0$  for  $0 < u < u_1$  and  $f(u) > 0$  for  $u_1 < u < u_+$ . For  $u_1 < u < u_+$ , we can proceed the same way as in the last paragraph to show that  $M(u) > 0$ . For  $0 < u \leq u_1$ , if  $f''(u) > 0$  for all  $u \in (0, u_1)$ , then  $M(u) > 0$ . So we assume that there exists  $\beta \in (0, u_1)$  such that  $f''(u) > 0$  for  $0 < u < \beta$  and  $f''(u) < 0$  for  $\beta < u < u_1$ . For  $0 < u < \beta$ ,  $M(u) > 0$ , since  $f < 0$  and  $f'' > 0$ . For  $\beta < u < u_1$ ,  $M'(u) = (4 - n)f'f'' - nff''' < 0$ , since  $f'(u) > 0$  and  $M(u_1) = 2[f'(u_1)]^2 > 0$ , so  $M(u) > 0$  for  $\beta \leq u < u_1$ . □

*Proof of theorem 3.1.* We first mention that the solution  $z$  of (2.4) can always be chosen as positive if  $n = 1$ , so the corresponding results in theorems 2.12, 2.14 and 2.16 (where  $n \geq 2$ ) are all true for  $n = 1$ .

We start by dividing the first quadrant of the  $(m, k)$ -plane into four parts using the lines  $m = k$  and  $k = 1$ . Define

$$\left. \begin{aligned} \text{I} &= \{(m, k) : m > k, k > 1\}, \\ \text{II} &= \{(m, k) : m > k, k < 1\}, \\ \text{III} &= \{(m, k) : m < k, k > 1\}, \\ \text{IV} &= \{(m, k) : m < k, k < 1\}. \end{aligned} \right\} \tag{3.2}$$

We will first discuss the exact multiplicity for  $(m, k)$  in these regions, and then discuss the cases for the border between them.

- (1) From the proof of lemma 3.2, we know that  $f'(0) = m - k$  and  $f''(0) = 2(k - 1)$ . First, if  $(m, k) \in \text{IV}$ , then  $f'(0) < 0$  and  $f''(u) < 0$  for all  $u > 0$ , and hence  $f(u) < 0$  for all  $u > 0$ , so that (1.4) has no positive solution. For  $(m, k) \in \text{III}$ ,  $f(u) < 0$  near  $u = 0$  and it is possible that  $f$  has one or two zeros in  $(0, \infty)$ . In fact, the zeros of  $f(u)$  are 0 and

$$u_{\pm} = \frac{1}{2}[(m - 1) \pm \sqrt{(m + 1)^2 - 4k}],$$

so that, in III,  $u_{\pm} > 0$  if and only if  $m > 1$ ,  $k > m$  and  $(m + 1)^2 - 4k > 0$ . So, if  $(m, k) \in \text{III}$  and  $(m + 1)^2 - 4k \leq 0$ , then  $f(u) \leq 0$  for all  $u > 0$  and (1.4) has no positive solution. Even when  $(m + 1)^2 - 4k > 0$  and  $(m, k) \in \text{III}$ , equation (3.1) is a necessary condition for the existence of positive solution, as follows from (2.21). Indeed,  $F(u_+) > F(u(0)) > 0$ . When  $(m + 1)^2 - 4k > 0$  and  $(m, k) \in \text{III}$ ,  $f(u) < 0$  for  $u \in (0, u_-)$  and  $f(u) > 0$  for  $u \in (u_-, u_+)$ . So, if there is a solution, then  $u(0) \in (u_-, u_+)$  and  $F(u_+) > 0$ .

- (2) If  $(m, k) \in \text{III}$ ,  $(m + 1)^2 - 4k > 0$  and (3.1) is satisfied, then all conditions in lemma 2.15 and theorem 2.16 are satisfied, so we can apply theorem 2.16 to prove part 3 of the theorem.
- (3) For  $(m, k) \in \text{I}$ ,  $f'(0) > 0$ ,  $f''(u) > 0$  for  $u \in (0, \sqrt[3]{k} - 1)$  and  $f''(u) < 0$  for  $u \in (\sqrt[3]{k} - 1, \infty)$ . From lemma 3.2,  $f$  also satisfies (1.3). So we can apply theorem 2.12.

- (4) For  $(m, k) \in \text{II}$ ,  $f'(0) > 0$ ,  $f''(u) < 0$  for all  $u > 0$ . Then the result in theorem 3.1 is well known in this case (see, for example, [19, theorem 6.2]). Also in this case, the bifurcation diagram is same as the one with  $k = 0$ , the classical logistic equation.
- (5) Finally, we handle the borderline cases. For  $k = 0$ , it is the classical logistic equation and the bifurcation curve is same as those for  $(m, k) \in \text{II}$ . For  $m = 0$ , it is obvious that  $f(u) < 0$  for all  $u > 0$  and thus there is no solution. For the border of II and IV,  $m = k$  and  $k < 1$ , then  $f'(0) = 0$  and  $f''(0) < 0$ , so  $f(u) < 0$  for all  $u > 0$  and there is no solution. For the border of I and II, the bifurcation curve is same as those for  $(m, k) \in \text{II}$ , except that at  $(\lambda, u) = (\lambda_1^0, 0)$ ,  $\lambda'(0) = 0$ , but  $\lambda''(0) > 0$ . (For a formula for  $\lambda''(0)$ , see lemma 2.5 or [21].) For the border of III and IV, it is obvious that  $f(u) < 0$  for all  $u > 0$  and hence there is no solution. For the border of I and III,  $f'(0) = 0$ ,  $f''(0) > 0$  and the conditions of theorem 2.14 are all satisfied, thus the solution curve is exactly C-shaped.

□

## Acknowledgments

P.K. was supported in part by the Taft Faculty Grant at the University of Cincinnati.

## References

- 1 Adimurti. Uniqueness of positive solutions of a quasilinear Dirichlet problem with exponential nonlinearity. *Proc. R. Soc. Edinb. A* **128** (1998), 895–906.
- 2 M. G. Crandall and P. H. Rabinowitz. Bifurcation, perturbation of simple eigenvalues and linearized stability. *Arch. Ration. Mech. Analysis* **52** (1973), 161–180.
- 3 D. G. De Figueiredo, P. L. Lions and R. D. Nussbaum. *A priori* estimates and existence of positive solutions of semilinear elliptic equations. *J. Math. Pures Appl.* **61** (1982), 41–63.
- 4 Y. Du and Y. Lou. Proof of a conjecture for the perturbed Gelfand equation from combustion theory (1997). (Preprint.)
- 5 Y. Du and Y. Lou. S-shaped global bifurcation curve and Hopf bifurcation of positive solutions to a predator–prey model. *J. Diff. Eqns* **144** (1998), 390–440.
- 6 B. Gidas, W.-M. Ni and L. Nirenberg. Symmetry and related properties via the maximum principle. *Commun. Math. Phys.* **68** (1979), 209–243.
- 7 C. S. Holling. Some characteristics of simple types of predation and parasitism. *Can. Entomol.* **91** (1959), 385–395.
- 8 D. Joseph and T. Lungren. Quasilinear Dirichlet problems driven by positive sources. *Arch. Ration. Mech. Analysis* **49** (1973), 241–269.
- 9 P. Korman. Solution curves for semilinear equations on a ball. *Proc. Am. Math. Soc.* **125** (1997), 1997–2005.
- 10 P. Korman. Curves of sign-changing solutions for semilinear equations. (Preprint.)
- 11 P. Korman, Y. Li and T. Ouyang. Exact multiplicity results for boundary value problems with nonlinearities generalising cubic. *Proc. R. Soc. Edinb. A* **126** (1996), 599–616.
- 12 P. Korman, Y. Li and T. Ouyang. An exact multiplicity result for a class of semilinear equations. *Commun. PDEs* **22** (1997), 661–684.
- 13 M. K. Kwong. On the Kolodner–Coffman method for the uniqueness problem of Emden–Fowler BVP. *Z. Angew. Math. Phys.* **41** (1990), 79–104.
- 14 M. K. Kwong and L. Q. Zhang. Uniqueness of the positive solution of  $\Delta u + f(u) = 0$  in an annulus. *Diff. Integ. Eqns* **4** (1991), 583–599.

- 15 C. S. Lin and W.-M. Ni. A counterexample to the nodal domain conjecture and a related semilinear equation. *Proc. Am. Math. Soc.* **102** (1988), 271–277.
- 16 P.-L. Lions. On the existence of positive solutions of semilinear elliptic equations. *SIAM Rev.* **24** (1982), 441–467.
- 17 W.-M. Ni and R. D. Nussbaum. Uniqueness and nonuniqueness for positive radial solutions of  $\Delta u + f(u, r) = 0$ . *Commun. Pure Appl. Math.* **38** (1985), 67–108.
- 18 T. Ouyang and J. Shi. Exact multiplicity of positive solutions for a class of semilinear problem. *J. Diff. Eqns* **146** (1998), 121–156.
- 19 T. Ouyang and J. Shi. Exact multiplicity of positive solutions for a class of semilinear problems. II. *J. Diff. Eqns* **158** (1999), 94–151.
- 20 R. Schaaf. *Global solution branches of two-point boundary value problems*. Lecture Notes in Mathematics, vol. 1458 (Springer, 1990).
- 21 J. Shi. Persistence and bifurcation of degenerate solutions. *J. Funct. Analysis* **169** (1999), 494–531.

(Issued 19 October 2001)