



Aurora kinase as a putative target to tick control

Bruno Moraes^{1,2}, Helga Gomes³, Luiz Saramago², Valdir Braz²,
Luís Fernando Parizi⁴, Gloria Braz⁵, Itabajara da Silva Vaz Jr^{4,6} ,
Carlos Logullo^{1,6}  and Jorge Moraes^{2,6}

Research Article

Cite this article: Moraes B, Gomes H, Saramago L, Braz V, Parizi LF, Braz G, da Silva Vaz Jr I, Logullo C, Moraes J (2024). Aurora kinase as a putative target to tick control. *Parasitology* **151**, 983–991. <https://doi.org/10.1017/S003118202400101X>

Received: 11 June 2024

Revised: 31 July 2024

Accepted: 6 August 2024

First published online: 15 November 2024

Keywords:

Aurora-kinase; BME26; CCT137690; embryonic cell; *Rhipicephalus microplus*; tick

Corresponding author:

Carlos Logullo;

Email: carlos.logullo@bioqmed.ufrj.br

¹Laboratório de Bioquímica de Artrópodes Hematófagos, Instituto de Bioquímica Médica Leopoldo de Meis, Universidade Federal do Rio de Janeiro, RJ, Brazil; ²Laboratório Integrado de Bioquímica Hatisaburo Masuda, NUPEM-Universidade Federal do Rio de Janeiro campus Macaé, Brazil; ³Laboratório de Tecido Conjuntivo, Hospital Universitário Clementino Fraga Filho and Instituto de Bioquímica Médica Leopoldo de Meis, Universidade Federal do Rio de Janeiro, RJ, Brazil; ⁴Centro de Biotecnologia and Faculdade de Veterinária, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil; ⁵Instituto de Química, Universidade Federal do Rio de Janeiro, RJ, Brazil and ⁶Instituto Nacional de Ciência e Tecnologia em Entomologia Molecular, Rio de Janeiro, RJ, Brazil

Abstract

Aurora kinases (AURK) play a central role in controlling cell cycle in a wide range of organisms. They belong to the family of serine-threonine kinase proteins. Their role in the cell cycle includes, among others, the entry into mitosis, maturation of the centrosome and formation of the mitotic spindle. In mammals, 3 isoforms have been described: A, B and C, which are distinguished mainly by their function throughout the cell cycle. Two aurora kinase coding sequences have been identified in the transcriptome of the cattle tick *Rhipicephalus microplus* (Rm-AURKA and Rm-AURKB) containing the aurora kinase-specific domain. For both isoforms, the highest number of AURK coding transcripts is found in ovaries. Based on deduced amino acid sequences, it was possible to identify non-conserved threonine residues which are essential to AURK functions in vertebrates and which are not present in *R. microplus* sequences. A pan AURK inhibitor (CCT137690) caused cell viability decline in the BME26 tick embryonic cell line. *In silico* docking assay showed an interaction between Aurora kinase and CCT137690 with exclusive interaction sites in Rm-AURKA. The characterization of exclusive regions of the enzyme will enable new studies aimed at promoting species-specific enzymatic inhibition in ectoparasites.

Introduction

Cell cycle is one of the most impressive controlled events in cell physiology. It governs cell growth, cell proliferation, DNA regulation and other major steps during mitosis (Matthews *et al.*, 2022). This process is driven by kinases, especially serine/threonine kinases (Arias and Hayward, 2006; Manning *et al.*, 2002). In most studied organisms, at least 2 different AURKs with distinct functions and cell localization have been found: Aurora A (AURKA) plays a central role in centrosome maturation and bipolar spindle assembly, while Aurora B (AURKB) is important in condensation, attachment to kinetochores and chromosome alignment (Carmena *et al.*, 2009; Wang *et al.*, 2014). Aurora C (AURKC) is present only in mammals, being essential for male fertility (Ounis *et al.*, 2015). Malfunction in AURK genes can lead to the formation of tumour cells (Fukuda *et al.*, 2005; Pérez-Fidalgo *et al.*, 2020).

The association between AURK dysregulation and cancer cell progression makes this protein family a potential oncogene. Several AURK inhibitors have been developed over the years, targeting AURKA and AURKB to block cell cycle progression and induce apoptosis, a useful strategy to control many types of tumours (Du *et al.*, 2021; Mou *et al.*, 2021). One of the most promising is CCT137690, which is a highly selective, orally bioavailable (imidazo[4,5-b] pyridine), with low IC₅₀ value against a wide range of tumour cell lines (Sogutlu *et al.*, 2021).

Aurora kinase was first discovered using *Drosophila melanogaster* as a model, showing its involvement in cell cycle progression (Glover, 1989). AURKA silencing in *D. melanogaster* led to a reduction in the length of astral microtubules in syncytial embryos, larval neuroblasts and cultured S2 cells (Giet *et al.*, 2002). In early development, AURKA is required for larval development, controlling proper timing through direct and indirect means. In larval tissues, AURKA is required for symmetric division rate and eventually development speed as was observed in central brain, wing disc and ring gland (Vaufrey *et al.*, 2018). Moreover, AURKA inactivation induces a reduction of ecdysteroids levels and a delay in pupariation as an indirect consequence of ring gland development deceleration (Vaufrey *et al.*, 2018). Overall, AURKA is considered to have an important role in arthropod development (Magnaghi-Jaulin *et al.*, 2019).

The study of this protein family is key for a better understanding of a range of cell cycle-related processes also in arthropod vectors, such as the tick *Rhipicephalus microplus*

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

(Carmena *et al.*, 2009). *Rhipicephalus microplus* tick is a cattle parasite in tropical and subtropical areas around the world, capable of spreading important veterinary diseases such as babesiosis and anaplasmosis (Some *et al.*, 2023). The parasitized cattle have a decreased commercial value, mainly due to the reduced milk production, weight loss and a lower quality of the leather. The reduction of this ectoparasite population is challenging, since a single female lays about 2000 eggs, which leads to a new infesting cycle (Senbill *et al.*, 2018). Infestation can result, directly or indirectly, in economic losses in the order of billions of dollars per year in Brazil (Grisi *et al.*, 2014). The application of non-selective anti-tick compounds may select resistant tick populations and contribute to environmental contamination of the environment (Obaid *et al.*, 2022; Waldman *et al.*, 2023a, 2023b). Hence the urgency to identify new biological targets such as enzymes, ion channels, receptors, to aid the development of new economic-viable alternative control methods (Graf *et al.*, 2004; Reck *et al.*, 2014; Mohs and Greig, 2017).

Compared to other arthropods, the knowledge related to tick physiology is scarce. To support the development of new control strategies, our research aims to supply information about new promising biological targets against ticks. Studies focused to uncover new drug target sites are necessary to help the control of *R. microplus* populations (Ozelame *et al.*, 2022; Maritz-Olivier *et al.*, 2023; Waldman *et al.*, 2023a, 2023b). In previous work, our group showed that cyclin-dependent kinases (CDKs) can be used to develop new strategies against arthropods. CDK inhibitor roscovitine decreased BME26 cell viability after 24 and 48 h of incubation and the vaccination using tick CDK as antigen reduced the amount of blood ingested and egg production by ticks (Gomes *et al.*, 2013, 2015). In the present work, we analysed the potential of the AURK protein family as novel physiological targets to control *R. microplus* infestations. We showed the effects of CCT137690 on cell growth and survival in an embryonic tick cell line (BME26) isolated from *R. microplus* embryos (Esteves *et al.*, 2008). The identification and characterization of new promising targets against this ectoparasite is one of the first steps to identify innovative control methods.

Methods and materials

BME26 cell maintenance

Cells were maintained following a previously described protocol (Esteves *et al.*, 2008). Briefly, adherent cells from 25 cm² confluent flasks were suspended into fresh complete medium (Munderloh and Kurtti, 2015) using a 22-gauge needle with a bent tip fitted to a plastic syringe. Cells were passaged every 2 weeks, and the medium replaced weekly. Culture density was determined with a Neubauer haemocytometer and cell viability was determined by the trypan blue (0.4%) exclusion method. Two weeks prior to use in assays, synchronized cells were prepared by seeding 1×10^7 cells into 5 mL of fresh complete medium (final volume), and grown at 34°C to ensure doubling (within 2 weeks), replacing the medium weekly.

Cell viability assay

BME26 cell suspension was seeded into 24-well plates at a density of 5×10^5 cells well⁻¹, to a final volume of 500 mL of complete medium and allowed to attach. After 24 h at 34°C, CCT137690 was added at the final concentrations indicated, and 0.1% dimethylsulfoxide was used in negative control wells. After 24 or 48 h of treatment, 50 mL of tetrazolium salt 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) prepared in serum-free medium (5 mg mL⁻¹) was added to each

well. After additional 2 h incubation, the media was completely discarded and 1 mL of acid-isopropyl alcohol (0.15% HCl in isopropyl alcohol) was added to dissolve the formazan crystals. The mixture was transferred to 1.5 mL tubes, spun at 6000×g for 15 min, and the clear supernatant collected in new tubes for absorbance measurement at 570 nm using quartz cuvettes in an UVmini-1240 UV-VIS spectrophotometer (Shimadzu, Japan).

Identification of AURK homologs from *R. microplus*

Protein sequences of AURK from *H. sapiens*, *Bos taurus*, *Mus musculus*, *Gallus gallus*, *Danio rerio*, *D. melanogaster*, *Anopheles gambiae* and *Caenorhabditis elegans* were downloaded from HomoloGene (<http://www.ncbi.nlm.nih.gov/homologene>). These proteins were further used as queries to conduct BLAST searches in the NCBI database (Altschul *et al.*, 1990). *Rhipicephalus microplus* Aurora kinase A protein sequence (GenBank: AHF48782.1) was found in NCBI and Aurora kinase B was assembled from annotation of comprehensive *R. microplus* transcriptome (Tirloni *et al.*, 2020).

Alignment and phylogenetic analyses

Classification of *R. microplus* AURK by similarity was performed solely by blast homology with AURKs from model organisms. A phylogenetic tree was then built using only AURKs found in *R. microplus* and their best sequence matches from Homologene Bank tool present in NCBI. The sequences from Homologene Bank were used to build a phylogenetic tree using neighbour-joining method in the MEGA software (Tamura *et al.*, 2021). The final tree was generated with 10 000 bootstraps.

Molecular modelling

Three-dimensional models of AURKs from *R. microplus* were constructed by comparative modelling using the SWISS-MODEL server (Waterhouse *et al.*, 2018) combining sequence, structural and functional information. The template recognition is based on profile–profile alignment guided by secondary structure and exposure predictions. The accurate template determination and sequence alignment algorithm enhances the reliability of the 3D structure. For the validation of the 3D model, the protein analysis tools available on the Structural Analysis and Verification Server (<http://nihserv-110er.mbi.ucla.edu/SAVES/>) and the visual inspections of the 3-dimensional models were made in the program PyMOL version 1.8x.

Electrostatic potential surface calculation and hydrophobic surface mapping

Poisson–Boltzmann electrostatic potential was calculated for the AURKA and AURKB predicted structures using APBS tool (Baker *et al.*, 2001) and the surface electrostatic potential map was visualized using PyMOL 1.8 version. To map the hydrophobic surface of AURKs, the hydrophobicity scale from Eisenberg was used as a reference to calculate the average hydrophobicity of each amino acid (Eisenberg *et al.*, 1984).

Structure conservation analysis

Multiple AURK sequences obtained from Homologene bank were aligned using clustaw Omega (Sievers and Higgins, 2014). ConSurf Server (Glaser *et al.*, 2003; Landau *et al.*, 2005) was used to determine the residue conservation level shown in the multiple alignment. Residues were considered conserved if assigned the maximum conservation grade (9).

Docking analysis

The docking analysis was performed using AutoDock 4.2 program and AutoDock Tools Version 1.5.4 (Morris *et al.*, 1998; Huey *et al.*, 2007). CCT137690 crystal structure was previously deposited in Protein Data Bank PDB ID: 2 × 6e (Bavetsias *et al.*, 2010). The structures were transferred to AutoDock 4.2 program (Morris *et al.*, 1998; Huey *et al.*, 2007) to create the ligand input file in the pdbqt format. All bond rotations and torsions for the ligand were automatically set in AutoDock Tools. The AURKA and AURKB from *R. microplus* were transferred to the Autodock 4.2 program and AutoDock Tools were used to prepare the proteins. All water molecules and the ligands were deleted, polar hydrogens were added and Gasteiger charges were calculated to create the pdbqt file for both proteins. The cubic box was made based on the 2 × 6e crystal, centred in the ligand position with 80 × 80 × 80 and calculated by Autogrid 4. Docking studies were performed using the empirical free energy function and the Lamarckian genetic algorithm applying a standard protocol; a total of 50 independent docking runs were carried out for each protein. Structures differing by less than 2 Å in positional root-mean-square deviation were clustered together and the selected complex for each ligand was that with the lowest binding energy. Redocking method was performed using PDB ID 2 × 6e crystal to validate the results. Two-dimensional interaction diagram was generated using Discovery Studio 3.5 version software Discovery Studio Client 3.5 version (Biovia, San Diego, CA, USA) and 3D analysis of the Protein-Ligand interaction was done using PyMOL 1.8 (The PyMOL Molecular Graphics System, Version 1.8 Schrodinger, LLC).

Statistical analysis

Experiments were performed in biological triplicate with technical triplicate. Graphs present the averages and respective standard deviations. The software GraphPad Prism 8.3 (www.graphpad.com).




Figure 1. Illustrative representation of the Rm-AURKA (A) and Rm-AURKB (B) proteins from *R. microplus* highlighting the STKc_Aurora domain. Aurora kinase characteristic conserved domain and activating threonine are highlighted. The sequences were identified based on the presence of a conserved threonine residue responsible for activating the loop in the same position as other well-studied Aurora kinase proteins (Walter *et al.*, 2000; Yasui *et al.*, 2004; Zorba *et al.*, 2014).




Figure 2. AURK orthologous proteins from *R. microplus* were identified in the available data obtained from a multi-tissue transcriptome (Tirloni *et al.*, 2020). Rm-AURKA and Rm-AURKB genes are transcribed in different organs.

com) was used to perform unpaired *t*-test or 2-way ANOVA followed by multiple comparisons, when applicable.

Results

Identification and phylogenetic analysis of putative AURKs in *R. microplus*

Two nucleotide sequences with similarity to AURK were obtained from a *R. microplus* transcriptome (Tirloni *et al.*, 2020) as described in the Methods section. The putative proteins were named Rm-AURKA and Rm-AURKB (Fig. 1). Also, the transcriptome analysis indicated the ovary as the organ with the highest number of transcripts (Fig. 2).

Both putative Rm-AURK showed a high similarity with related proteins in other species, including the conservation of key amino acids responsible for function specificity (Fig. 3 and Supplementary Fig. 1). Rm-AURKA protein contains 298 amino acids (with a molecular weight of 34.35 kDa and isoelectric point 8.99), whereas Rm-AURKB has 274 amino acids in the mature protein (31.76 kDa and isoelectric point 9.26). For comparison, the mature forms of mammalian AURKA and AURKB from *B. taurus*, the natural tick host, have 402 amino acids (molecular weight 45.46 kDa and isoelectric point 9.53) and 344 amino acids (molecular weight 39.40 kDa and isoelectric point 9.56), respectively. Rm-AURKA protein demonstrates 2 key amino acid substitutions when compared with vertebrates active site: Glu-124 and Met-200 in *R. microplus* are substituted for Gly-216 and Thr-292 in *Homo sapiens* (Fig. 3 and Supplementary Fig. 4). On the other hand, Rm-AURKB also presents changes in primary structure on the active site (Supplementary Fig. 1). A phylogenetic analysis of the AURK protein family members of selected organisms revealed that the AURKs from vertebrates are not in the same branch as the AURKs from ticks, which may indicate an evolutionary distance between ticks and vertebrates (Fig. 4).




Figure 3. Amino acid sequence alignment of AURKA from *Rhipicephalus microplus*, *Drosophila melanogaster*, *Ixodes scapularis*, *Bos taurus*, *Homo sapiens* and *Xenopus laevis*. Sequences were aligned using PRALINE multiple sequence alignment and were coloured according to a conservation rank. The black arrows represent non-conserved serine and threonine residues between the groups. Black dots represent non-conserved serine or threonine residues.




Figure 4. Phylogenetic tree containing various Aurora kinases from diverse organisms. An apparent distance can be observed in the origin of the gene between vertebrates (red branches) and invertebrates (other colours of branches). AURKA and AURKB from *R. microplus* are marked by a purple circle and a pink diamond, respectively. The phylogenetic tree was built using MEGA software (Tamura *et al.*, 2021) and the neighbour-joining method, with 10 000 bootstrap being used to generate the final tree.

Molecular modelling studies and comparative *Rm-AURKA* protein model

To study amino acid differences in AURKA proteins among species (Fig. 3), a conserved model was constructed using different AURKA sequences deposited in the GenBank to create a unique structure and identify different degrees of conservation throughout evolution. As expected, the active site region is highly conserved among species, but regions close to it have lower levels of conservation, suggesting they might be characteristic of each species (Fig. 5). *Rm-AURKA* sequence was used to generate a 3-dimensional protein model based on the *H. sapiens* AURKA structure (Fig. 6) (Bavetsias *et al.*, 2010). The overlay between *Rm-AURKA* and *B. taurus-AURKA* showed an exclusive loop present in tick protein (Fig. 6B).

Protein hydrophobic surface is essential for folding, stability and the formation of compact nucleus. These interactions are also essential for anchoring ligands and some protein-protein interactions (Cherry and Fidantsef, 2003; Almeida *et al.*, 2021).

The hydrophobic surface of *Rm-AURKA* model showed a characteristic nucleus with hydrophobic amino acids (Fig. 6C1), as found in *B. taurus* (Fig. 6C3). However, in other portions, *Rm-AURKA* protein (Fig. 6C2) has a pattern that differs from that seen in *B. taurus* (Fig. 6C4). The electrostatic surfaces of *Rm-AURKA* (Fig. 6D) and AURKB (Supplementary Fig. 2D) generally show a larger predominance of electronegative residues compared with other animals, including *B. taurus* (Figs 6D3 and D4) (Supplementary Figs 2D3 and D4). These regions with greater differences in electrostatic properties can be explored as regions specific to the tick protein.

Molecular docking and effect on cell viability

To investigate the interactions between the specific pan-aurora kinase inhibitor CCT137690 and *Rm-AURKA*, a molecular docking experiment was performed to predict possible interactions between the protein and the ligand. Top-scoring docking result




Figure 5. Posterior (A) and anterior view (B) of Aurora kinases conservation ranking. Rm-AURKA was used as a query to construct this 3D model as described in methods. Representation in spheres, using ConSurf conservation ranking.




Figure 6. Structural comparison and physicochemical properties of *B. taurus* and *R. microplus* AURKA models. (A) Structure of the comparative model of Rm-AURKA showing in detail the putative phosphorylating residue Thr195. The structure was based on *H. sapiens* AURKA deposited in PDB (ID 2 × 6e). (B) Overlaps between *R. microplus* AURKA (red) and *B. taurus* AURKA (black). Dashed circle highlights structural differences between *R. microplus* and *B. taurus* proteins. The 3-dimensional models were constructed using the Swiss-Model server (Waterhouse *et al.*, 2018) and their energies were minimized using the SAVES platform (Colovos and Yeates, 1993). The *B. taurus* sequence was obtained from GenBank Accession: (NP_001033117.1) (Benson *et al.*, 2013). (C) Electrostatic profiles of both protein models, with 180° rotations shown on the right. The red areas represent negative charges, while the blue areas indicate positive charges. (D) Hydrophobicity distribution in the protein models, with 180° rotations also presented on the right. Red regions represent increased hydrophobicity. All figures were generated using the PyMol software (DeLano, 2002).

is similar to the one found in the AURKA crystal deposited in PDB data bank (Fig. 7A). The ligand interacts with Rm-AURKA amino acids in the active site with a theoretical $K_i = 1.4073109 \times 10^{-8}$, and forms a hydrogen bond with a distance of 3.1 Å (Fig. 7B). For Rm-AURKB protein, the analysis showed a theoretical $K_i = 1.6664252 \times 10^{-8}$, and a hydrogen bond with a distance of 3.7 Å (Supplementary Fig. 3B). As observed in studies with cancer cells, AURKA protein interacts more strongly with CCT137690 than AURKB, which is also the case for the tick proteins (Bavetsias *et al.*, 2010).

To assess the effect of AURK inhibition on cell viability, tick embryonic BME26 cells were incubated with different

concentrations of CCT137690 for 24 or 48 h. At the lowest tested concentration, cell viability decreased to 60% after 24 h incubation, and to 40% after 48 h (Fig. 8). Compared with results in HeLa and HCT116 cells (Bavetsias *et al.*, 2010; Faisal *et al.*, 2011), BME26 cells showed a higher GI_{50} compared to cancer cells models.

Discussion

In this paper, we characterize the AURK proteins in *R. microplus* and put them forward as putative targets to develop new tick control methods. The reproduction strategy adopted by many




Figure 7. Effect of CCT137690 AURK inhibitor on BME26 cells after 24 or 48 h, assessed by cell viability assay (MTT). MTT reaction was measured by spectrophotometry at 570 nm. Graph represents 3 independent experiments in triplicate (1-way ANOVA, $P < 0.05$).

arthropods, which can lay thousands of viable eggs, highlights this process as an attractive target for investigating new methods of population control (Smaghe *et al.*, 2019). There is a considerable number of insects and other arthropods which show high fertility, but in tick this can be even more expressive, with 1 female being able to lay more than thousands of eggs (Ma *et al.*, 2016). Previous results by our group using as a model an embryonic cell line from the hard tick *R. microplus* (BME26) have shown a susceptibility to roscovitine, a CDK inhibitor. Similar to Rm-CDK, Rm-AURKA and Rm-AURKB show a high number of transcripts in certain tick organs, such as salivary gland and ovary (Fabres *et al.*, 2010; Gomes *et al.*, 2013).

Aurora kinases transcripts were observed at high numbers in the ovary (Fig. 2), an organ marked by intense cell division, and that participates in the oogenesis process in arthropods (Nguyen and Schindler, 2017). These observations suggest a possible role of AURK in controlling cell cycle in this organ. The fact that the cell cycle is a well-studied process across a wide range of organisms makes it possible to draw parallels and make comparisons between different species (von der Dunk *et al.*, 2022). Nevertheless, despite being extensively studied in vertebrate and invertebrate models, studies on cell cycle control in invertebrate disease vectors have apparently been to date largely

neglected (Lorenzo *et al.*, 2014; Sullivan, 2016; Valenzuela and Aksoy, 2018).

Our main study hypothesis is based on structural differences among AURKs, particularly tick vs bovine proteins. Rm-AURK amino acid sequence (Fig. 3) showed 78% of similarity with the *B. taurus* model, mainly within the active site but interestingly not in other essential protein regions (Fig. 5). Additional major changes can be identified in vertebrates. For example, *H. sapiens* and *Xenopus laevis* share Hs-Ser342/Xl-Ser349 (respectively), which is important for interaction with PAK kinase (Supplementary Fig. 5) (Pascreau *et al.*, 2008; Korobeynikov *et al.*, 2019), while *R. microplus* sequences have Rm-Arg249 in the same position, with different physicochemical characteristics that may impact other divergent protein regions (Fig. 3). Another important residue in Hs-AURKA is Thr287, which is part of the activation loop, but is absent in tick sequences (Supplementary File 6) (Rowan *et al.*, 2013). Putative AURKB-like protein also showed high similarity with the bovine model in the active site, while presenting important differences between ticks and vertebrates (Supplementary Fig. 1). The KEN motif is highly conserved in vertebrate AURK sequences, being responsible for protein degradation *via* ubiquitination mediated by the anaphase-promoting complex (APC) (Nguyen *et al.*, 2005). Interestingly, the KEN motif was not found in putative Rm-AURKB (Supplementary Fig. 1), suggesting an alternative degradation pathway different from that of vertebrate organisms. It is possible that ticks lack a specialized gene machinery for degradation of AURKB, as the interaction with the Cdc20 subunit of the APC is necessary for ubiquitination of AURKB (Nguyen *et al.*, 2005).

These specific differences between ticks and vertebrates can be inferred by phylogenetic analysis (Fig. 4). The obtained results indicate a separation in the vertebrate and invertebrate groups which can be divided into hexapods and arachnids. This phylogenetic distance suggests that throughout evolution, these genes were possibly acquiring new mutations that supported the differences between ticks and vertebrates. Nevertheless, central regions for the protein to function such as the active site and its shape are maintained, while N and C terminal regions have a greater degree of variability. Several portions of the protein have remained conserved (Fig. 5); regions close to the protein hydrophobic hinge, where the active site is located, appear to be more conserved,




Figure 8. Interaction between CCT137690 and Rm-AURKA. (A) Representation of the AURKA comparative model in spheres, highlighting the amino acids belonging to the active site of the enzyme (blue). (B) Top-scoring pose obtained by docking of CCT137690 with Rm-AURKA comparative model. Hydrogen atoms have been omitted for improved view. Hydrogen bonds are depicted as yellow dashed lines. Docking was performed using AUTODOCK Vina 4.2 program and the model was visualized in PyMOL 1.8.

showing point mutations. The superposition of *R. microplus* and *B. taurus*-predicted structures showed exclusive regions in the tick protein, which are potentially attractive targets for drug design (Fig. 7; Supplementary Fig. 2B). In addition, the electrostatic and hydrophobic surfaces (Fig. 6D and C, respectively) also exhibit interesting different areas with distinct patterns between tick and bovine proteins, which may suggest different interactions with substrate and ligands. These distinct physicochemical characteristics may help in identifying new tick-specific protein ligands.

Molecular docking demonstrated that the anchorage site of the compound CCT137690 is similar to that observed in the *H. sapiens* protein crystal previously deposited in the Protein Data Bank (PDB ID: 2 × 6e) (Bavetsias *et al.*, 2010). Rm-AURKA-like protein has an alanine residue (ALA-120) that interacts with the compound *via* hydrogen bond at a shorter distance than Rm-AURKB; the compound is capable to interact differently comparing tick and mammal models. The differences in interactions between the proteins can help to explain the theoretical Ki differences observed between tick and cancer cells (Fig. 7B) (Bavetsias *et al.*, 2010). The GI⁵⁰ of this compound in cervical carcinoma cells in ovarian cancer cells (A2780) is 0.35 μM (Bavetsias *et al.*, 2010). The BME26 cell line appears to be less sensitive to CCT137690, with a GI⁵⁰ of 6.53 μM indicating that, although the cells and the AURKs present there have different characteristics, the inhibitor still affects cell viability. The putative mechanism involves inhibition of AURK enzyme activity and cell cycle progression, highlighting these components as possible drug targets to control populations of disease vectors. Limited knowledge about AURKs in arthropods can hinder the identification of such novel targets. While most studies to date have focused on classical cell biology involving this enzyme, here we propose a new approach to advance those efforts. This study is aimed in the characterization of these 2 proteins; more studies are needed to indicate the better approach to develop new control methods based on Rm-AURKs. In light of our present findings, the use of AURK inhibitors for arthropod control merits deeper investigation to aid the development of new efficient strategies.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S003118202400101X>.

Data availability statement. The authors confirm that the data supporting the findings of this study are available within the article and/or its supplementary materials.

Author contributions. Conceived and designed the experiments: B. M., H. G. Performed the experiments: B. M., H. G., L. S., V. B., L. F. P. Analysed the data: B. M., H. G., L. S., V. B. Contributed reagents/materials/analysis tools: L. F. P., G. B., I. S. V., C. L., J. M. Wrote the paper: B. M., H. G. Supervision, revision and final approval of the manuscript: B. M., H. G., L. S., V. B., L. F. P., G. B., I. S. V., C. L., J. M. Conducted statistical analysis: B. M. Supervised statistical analysis: H. G., C. L., J. M. All authors read and approved the final version of the manuscript.

Financial support. This work was supported by grants from CNPq-Instituto Nacional de Ciência e Tecnologia, INCT-Entomologia Molecular and FAPERJ (Brazil).

Competing interests. None.

Ethical standards. Not applicable.

References

- Almeida FCL, Sanches K, Pinheiro-Aguiar R, Almeida VS and Caruso IP (2021) Protein surface interactions-theoretical and experimental studies. *Frontiers in Molecular Biosciences* **8**, 706002.
- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.

- Arias AM and Hayward P (2006) Filtering transcriptional noise during development: concepts and mechanisms. *Nature Reviews Genetics* **7**, 34–44.
- Baker NA, Sept D, Joseph S, Holst MJ and McCammon JA (2001) Electrostatics of nanosystems: application to microtubules and the ribosome. *Proceedings of the National Academy of Sciences* **98**, 10037–10041.
- Bavetsias V, Large JM, Sun C, Boulouc N, Kosmopoulou M, Matteucci M, Wilsher NE, Martins V, Reynisson J, Atrash B, Faisal A, Urban F, Valenti M, de Haven Brandon A, Box G, Raynaud FI, Workman P, Eccles SA, Bayliss R, Blagg J, Linardopoulos S and McDonald E (2010) Imidazo[4,5-b]pyridine derivatives as inhibitors of Aurora kinases: lead optimization studies toward the identification of an orally bioavailable pre-clinical development candidate. *Journal of Medicinal Chemistry* **53**(14), 5213–5228. doi: 10.1021/jm100262j
- Carmena M, Ruchaud S and Earnshaw WC (2009) Making the Auroras glow: regulation of Aurora A and B kinase function by interacting proteins. *Current Opinion in Cell Biology* **21**, 796–805.
- Cherry JR and Fidantsef AL (2003) Directed evolution of industrial enzymes: an update. *Current Opinion in Biotechnology* **14**, 438–443.
- Colovos C and Yeates TO (1993) Verification of protein structures: patterns of nonbonded atomic interactions. *Protein Science* **2**, 1511–1519.
- DeLano WL (2002) Pymol: an open-source molecular graphics tool. *CCP4 Newsletter on Protein Crystallography* **40**, 82–92.
- Du R, Huang C, Liu K, Li X and Dong Z (2021) Targeting AURKA in cancer: molecular mechanisms and opportunities for cancer therapy. *Molecular Cancer* **20**, 15.
- Eisenberg D, Schwarz E, Komaromy M and Wall R (1984) Analysis of membrane and surface protein sequences with the hydrophobic moment plot. *Journal of Molecular Biology* **179**, 125–142.
- Esteves E, Lara FA, Lorenzini DM, Costa GHN, Fukuzawa AH, Pressinotti LN, Silva JRM, Ferro JA, Kurtti TJ, Munderloh UG and Daffre S (2008) Cellular and molecular characterization of an embryonic cell line (BME26) from the tick *Rhipicephalus (Boophilus) microplus*. *Insect Biochemistry and Molecular Biology* **38**, 568–580.
- Fabres A, De Andrade CP, Guizzo M, Sorgine MH, Paiva-Silva Gde O, Masuda A, Vaz Ida Jr. S and Logullo C (2010) Effect of GSK-3 activity, enzymatic inhibition and gene silencing by RNAi on tick oviposition and egg hatching. *Parasitology* **137**, 1537–1546.
- Faisal A, Vaughan L, Bavetsias V, Sun C, Atrash B, Avery S, Jamin Y, Robinson SP, Workman P, Blagg J, Raynaud FI, Eccles SA, Chesler L and Linardopoulos S (2011) The aurora kinase inhibitor CCT137690 downregulates MYCN and sensitizes MYCN-amplified neuroblastoma in vivo. *Molecular Cancer Therapeutics* **10**, 2115–2123.
- Fukuda T, Mishina Y, Walker MP, Diaugustine RP and Carolina N (2005) Conditional transgenic system for mouse Aurora A kinase: degradation by the ubiquitin proteasome pathway controls the level of the transgenic protein. *Molecular and Cellular Biology* **25**, 5270–5281. doi: 10.1128/MCB.25.12.5270
- Giet R, McLean D, Descamps S, Lee MJ, Raff JW, Prigent C and Glover DM (2002) Drosophila Aurora A kinase is required to localize D-TACC to centrosomes and to regulate astral microtubules. *Journal of Cell Biology* **156**, 437–451.
- Glaser F, Pupko T, Paz I, Bell RE, Bechor-Shental D, Martz E and Ben-Tal N (2003) ConSurf: identification of functional regions in proteins by surface-mapping of phylogenetic information. *Bioinformatics* **19**, 163–164.
- Glover DM (1989) Mitosis in Drosophila. *Journal of Cell Science* **92**(Pt 2), 137–146.
- Gomes H, Romeiro NC, Braz GRC, de Oliveira EAG, Rodrigues C, da Fonseca RN, Githaka N, Isezaki M, Konnai S, Ohashi K, da Silva Vaz I, Logullo C and Moraes J (2013) Identification and structural-functional analysis of cyclin-dependent kinases of the cattle tick *Rhipicephalus (Boophilus) microplus*. *PLoS ONE* **8**, 1–12.
- Gomes H, Moraes J, Githaka N, Martins R, Isezaki M, Vaz Ida Jr. S, Logullo C, Konnai S and Ohashi K (2015) Vaccination with cyclin-dependent kinase tick antigen confers protection against Ixodes infestation. *Veterinary Parasitology* **211**, 266–273.
- Graf JE, Gogolewski R, Leach-Bing N, Sabatini GA, Molento MB, Bordin EL, ... Arantes GJ (2004) Tick control: an industry point of view. *Parasitology* **129**(Suppl.), S427–S442.
- Grisi L, Leite RC, Martins JR, Barros AT, Andreotti R, Caçado PH, León AA, Pereira JB and Villela HS (2014) Reassessment of the potential economic impact of cattle parasites in Brazil. *Revista Brasileira de Parasitologia Veterinária* **23**, 150–156.

- Huey R, Morris GM, Olson AJ and Goodsell DS (2007) A semiempirical free energy force field with charge-based desolvation. *Journal of Computational Chemistry* **28**, 1145–1152.
- Korobeynikov V, Borakove M, Feng Y, Wuest WM, Koval AB, Nikonova AS, Serebriiskii I, Chernoff J, Borges VF, Golemis EA and Shagisultanova E (2019) Combined inhibition of Aurora A and p21-activated kinase 1 as a new treatment strategy in breast cancer. *Breast Cancer Research and Treatment* **177**, 369–382.
- Landau M, Mayrose I, Rosenberg Y, Glaser F, Martz E, Pupko T and Ben-Tal N (2005) ConSurf 2005: the projection of evolutionary conservation scores of residues on protein structures. *Nucleic Acids Research* **33**(Suppl. 2), 299–302.
- Lorenzo M, Vidal D and Zarbin P (2014) Control of neglected disease insect vectors: future prospects for the use of tools based on behavior manipulation-interference. *Journal of the Brazilian Chemical Society* **25**. doi: 10.5935/0103-5053.20140202
- Ma M, Chen Z, Liu A, Ren Q, Liu J, Liu Z, Li Y, Yin H, Guan G and Luo J (2016) Biological parameters of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae) fed on rabbits, sheep, and cattle. *Korean Journal of Parasitology* **54**, 301–305.
- Magnaghi-Jaulin L, Eot-Houllier G, Gallaud E and Giet R (2019) Aurora A protein kinase: to the centrosome and beyond. *Biomolecules* **9**, 28.
- Manning G, Whyte DB, Martinez R, Hunter T and Sudarsanam S (2002) The protein kinase complement of the human genome. *Science* **298**(5600), 1912–1934. doi: 10.1126/science.1075762
- Maritz-Olivier C, Ferreira M, Olivier NA, Crafford J and Stutzer C (2023) Mining gene expression data for rational identification of novel drug targets and vaccine candidates against the cattle tick, *Rhipicephalus microplus*. *Experimental & Applied Acarology* **91**, 291–317.
- Matthews HK, Bertoli C and de Bruin RAM (2022) Cell cycle control in cancer. *Nature Reviews Molecular Cell Biology* **23**, 74–88.
- Mohs RC and Greig NH (2017) Drug discovery and development: role of basic biological research. *Alzheimer's and Dementia* **3**, 651–657.
- Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK and Olson AJ (1998) Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function. *Journal of Computational Chemistry* **19**, 1639–1662.
- Mou PK, Yang EJ, Shi C, et al. (2021) Aurora kinase A, a synthetic lethal target for precision cancer medicine. *Experimental and Molecular Medicine* **53**, 835–847.
- Munderloh UG and Kurtti TJ (2015) Formulation of medium for tick cell culture. *Experimental & Applied Acarology* **7**, 219–229.
- Nguyen AL and Schindler K (2017) Specialize and divide (twice): functions of three Aurora kinase homologs in mammalian oocyte meiotic maturation. *Trends in Genetics* **33**, 349–363.
- Nguyen HG, Chinnappan D, Urano T and Ravid K (2005) Mechanism of Aurora-B degradation and its dependency on intact KEN and A-boxes: identification of an aneuploidy-promoting property. *Molecular and Cellular Biology* **25**, 4977–4992.
- Obaid MK, Islam N, Alouffi A, Khan AZ, da Silva Vaz Jr. I, Tanaka T and Ali A (2022) Acaricides resistance in ticks: selection, diagnosis, mechanisms, and mitigation. *Frontiers in Cellular and Infection Microbiology* **12**, 941831.
- Ounis L, Zoghmar A, Coutton C, Rouabah L, Hachemi M, Martinez D, Martinez G, Bellil I, Khelifi D, Arnoult C, Fauré J, Benbouhedja S, Rouabah A and Ray PF (2015) Mutations of the aurora kinase C gene causing macrozoospermia are the most frequent genetic cause of male infertility in Algerian men. *Asian Journal of Andrology* **17**(1), 68–73.
- Ozelame KPC, Mattia MMC, e Silva LAD, Randall LM, Corvo I, Saporiti T, Seixas A, da Silva Vaz Jr. I and Alvarez G (2022) Novel tick glutathione transferase inhibitors as promising acaricidal compounds. *Ticks and Tick-Borne Diseases* **13**, 101970.
- Pascreau G, Delcros JG, Morin N, Prigent C and Arlot-Bonnemains Y (2008) Aurora-A kinase Ser349 phosphorylation is required during *Xenopus laevis* oocyte maturation. *Developmental Biology* **317**, 523–530.
- Pérez-Fidalgo JA, Gambardella V, Pineda B, Burgues O, Piñero O and Cervantes A (2020) Aurora kinases in ovarian cancer. *ESMO Open* **5**, e000718.
- Reck J, Klafke GM, Webster A, Dall'Agnol B, Scheffer R, Souza UA, Corassini VB, Vargas R, dos Santos JS and Martins JR (2014) First report of fluazuron resistance in *Rhipicephalus microplus*: a field tick population resistant to six classes of acaricides. *Veterinary Parasitology* **201**, 128–136.
- Rowan FC, Richards M, Bibby RA, Thompson A, Bayliss R and Blagg J (2013) Insights into Aurora-A kinase activation using unnatural amino acids incorporated by chemical modification. *ACS Chemical Biology* **8**, 2184–2191. doi: 10.1021/cb400425t
- Senbill H, Hazarika LK, Baruah A, Borah DK, Bhattacharyya B and Rahman S (2018) Life cycle of the southern cattle tick, *Rhipicephalus (Boophilus) microplus* Canestrini 1888 (Acari: Ixodidae) under laboratory conditions. *Systematic and Applied Acarology* **23**, 1169–1179.
- Sievers F and Higgins DG (2014) *Clustal Omega, Accurate Alignment of Very Large Numbers of Sequences*, Vol. 1079. Totowa, NJ: Humana Press, pp. 105–116.
- Smagge G, Zotti M and Retnakaran A (2019) Targeting female reproduction in insects with biorational insecticides for pest management: a critical review with suggestions for future research. *Current Opinion in Insect Science* **31**, 65–69.
- Sogutlu F, Kayabasi C, Yelken BO, Asik A, Gasimli R, Kipcak S, Susluer SY, Avci CB and Gunduz C (2021) The evaluation of effect of Aurora kinase inhibitor CCT137690 in melanoma and melanoma cancer stem cell. *Anti-Cancer Agents in Medicinal Chemistry* **21**, 1564–1574.
- Some MV, Biguezoton AS, Githaka N, Adakal H, Dayo GK, Belem A, Zoungrana S, Stachurski F and Chevillon C (2023) The potential of *Rhipicephalus microplus* as a vector of *Ehrlichia ruminantium* in West Africa. *Ticks and Tick-Borne Diseases* **14**, 102117.
- Sullivan W (2016) The untapped cell biology of neglected tropical diseases. *Molecular Biology of the Cell* **27**, 739–743.
- Tamura K, Stecher G and Kumar S (2021) MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* **38**, 3022–3027.
- Tirloni L, Braz G, Nunes RD, Gandara ACP, Vieira LR, Assumpcao TC, Sabadin GA, da Silva RM, Guizzo MG, Machado JA, Costa EP, Santos D, Gomes HF, Moraes J, Dos Santos Mota MB, Mesquita RD, de Souza Leite M, Alvarenga PH, Lara FA, Seixas A, da Fonseca RN, Fogaça AC, Logullo C, Tanaka AS, Daffre S, Oliveira PL, da Silva Vaz Jr. I and Ribeiro JMC (2020) A physiologic overview of the organ-specific transcriptome of the cattle tick *Rhipicephalus microplus*. *Scientific Reports* **10**, 18296.
- Valenzuela JG and Aksoy S (2018) Impact of vector biology research on old and emerging neglected tropical diseases. *PLoS Neglected Tropical Diseases* **12**, e0006365.
- Vaufrey L, Balducci C, Lafont R, Prigent C and Le Bras S (2018) Size matters! Aurora A controls *Drosophila* larval development. *Developmental Biology* **440**, 88–98.
- von der Dunk SHA, Snel B and Hogeweg P (2022) Evolution of complex regulation for cell-cycle control. *Genome Biology and Evolution* **14**, evac056.
- Waldman J, Klafke GM, Tirloni L, Logullo C and da Silva Vaz Jr. I (2023a) Putative target sites in synganglion for novel ixodid tick control strategies. *Ticks and Tick-Borne Diseases* **14**, 102123.
- Waldman J, Marcondes Klafke G and da Silva Vaz Júnior I (2023b) Mechanisms of Acaricide resistance in ticks. *Acta Scientiae Veterinariae* **51**. doi: 10.22456/1679-9216.128913
- Walter AO, Seghezzi W, Korver W, Sheung J and Lees E (2000) The mitotic serine/threonine kinase Aurora2/AIK is regulated by phosphorylation and degradation. *Oncogene* **19**, 4906–4916.
- Wang G, Jiang Q and Zhang C (2014) The role of mitotic kinases in coupling the centrosome cycle with the assembly of the mitotic spindle. *Journal of Cell Science* **127**(Pt 19), 4111–4122. doi: 10.1242/jcs.151753
- Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, De Beer TAP, Rempfer C, Bordoli L, Lepore R and Schwede T (2018) SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Research* **46**, W296–W303.
- Yasui Y, Urano T, Kawajiri A, Nagata KI, Tatsuka M, Saya H, Furukawa K, Takahashi T, Izawa I and Inagaki M (2004) Autophosphorylation of a newly identified site of Aurora-B is indispensable for cytokinesis. *Journal of Biological Chemistry* **279**, 12997–13003.
- Zorba A, Buosi V, Kutter S, Kern N, Pontiggia F, Cho Y-J and Kern D (2014) Molecular mechanism of Aurora A kinase autophosphorylation and its allosteric activation by TPX2. *ELife* **3**, 1–24.