


VARIABILITY OF RADIOCARBON RESERVOIR AGE EFFECTS IN LAKES AND RIVERS IN ANATOLIA AND LESSER CAUCASUS

Michel Fontugne^{1,2} • Christine Hatté^{2,3*}  • Nadine Tisnérat-Laborde² • Vincent Ollivier¹ • Catherine Kuzucuoglu⁴

¹CNRS, Aix Marseille University, Ministère de la Culture, LAMPEA, 13097 Aix-en-Provence, France

²Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, UMR 8212, CEA CNRS UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

³Institute of Physics, Silesian University of Technology, 44-100 Gliwice, Poland

⁴Paris 1 and Paris 12 Universities and CNRS, Laboratory of Physical Geography (UMR 8591), 2 rue Henri Dunant, 94320 Thiais, France

ABSTRACT. Multiproxy sedimentary sequence analysis constitutes the basis for reconstructions of past paleoenvironments and climate evolution. These sequences are, for the most part, obtained by coring in lakes, maars or crater lakes whose waters can record volcanic activity or karstic contributions, especially in Eastern Anatolia and the Lesser Caucasus. The reservoir age effect in these geological contexts leads to an apparent aging of the radiocarbon ages which also affects the plants and animals developing in or near these waters and consequently the population consuming them. We present here some results obtained from modern samples taken from Mediterranean, central and eastern Anatolian lakes, from the Van and Sevan lakes and along the Kura River and its tributaries from the Lesser Caucasus. The effect of volcanic CO₂ outgassing in the vicinity of maar crater lakes is also discussed.

KEYWORDS: Anatolia, Caucasus, karstic area, reservoir age, volcanic area.

INTRODUCTION

For about forty years, the estimation of reservoir ages to correct the aging of radiocarbon dates has been a major concern of archaeologists and geologists. This reservoir effect results, during the synthesis of the dated organic sample, from the use of carbon whose ¹⁴C activity is different from that of contemporary atmospheric CO₂. It is defined as:

$$R(^{14}\text{C reservoir ages}) = ^{14}\text{C age of sample} - ^{14}\text{C age of contemporaneous atmospheric CO}_2$$

and is expressed in ¹⁴C years (Mangerud 1972).

This reservoir effect in non-marine water-bodies is known to have multiple origins: (1) the hard water effect which results mainly from the dissolution of fossil carbonate rocks; (2) degassing of volcanic CO₂ in the atmosphere or in lakes; or even (3) confinement (cessation of exchanges with the atmosphere). These inputs lead to an aging of lacustrine or underground waters, inducing a reservoir age effect and biasing the radiocarbon dates of the palaeoclimatological record from such environments (Deevey et al. 1954; Deevey and Stuiver 1964). As the original source of carbon for all the aquatic trophic chain, the depleted ¹⁴C signature of Dissolved Inorganic Carbon (DIC) is reflected in all the halieutic resources of these lakes. This reservoir effect then propagates in terrestrial animals and humans consuming these food resources, and is reflected in the skeleton of the lake population (Oana and Deevey 1960; Philippsen and Heinemeier 2013, among others). In Asia Minor and the Caucasus, these three aging processes are frequent and are sometimes cumulative.

*Corresponding author. Email: christine.hatte@lsce.ipsl.fr

While the inland seas (Black and Caspian seas) (see among others Jones and Gagnon 1994; Kuzmin et al. 2007; Fontugne et al. 2009) and the northern Greater Caucasus in Russia (Shishlina et al. 2007; Higham et al. 2010) have already been partially studied, practically no age reservoir data have been published for the Lesser Caucasus and Anatolia. Here, we present age reservoir results obtained between 1999 and 2014. The spatial resolution of the results on the regional to local scales is somewhat incomplete, mainly because some of these regions (e.g. Armenia, Georgia, Azerbaijan, Eastern Taurus, etc.) have been or are still, depending on the period, inaccessible due to recurrent armed conflicts.

MATERIAL AND METHODS

The samples were collected between 1999 and 2014 during different scientific projects in Turkey and the Caucasus. The location, the nature of the samples and their date of collection are reported in Table 1 and Figure 1.

Chemical Treatment and Conversion to CO₂

Water Samples

After collection, surface water samples were immediately poisoned with mercury chloride and stored hermetically in 250 mL glass bottles until laboratory analysis was performed according to the methods described by Duplessy (1972). In the laboratory, water samples between 50 and 100 mL were introduced in a vacuum line, acidified with phosphoric acid, and bubbled with helium gas in order to optimize the extraction of the total dissolved carbon dioxide (Bard et al. 1988). The evolved CO₂ was purified, trapped and quantified. An aliquot of CO₂ gas was also sampled to measure the stable carbon isotopic composition ($\delta^{13}\text{C}$) with a Fisons-OPTIMA mass spectrometer. The precision was better than 0.05‰.

Organic Samples

Fish samples were bought from fishermen fishing by the river. Fish bone and flesh were dried and stored at -18°C . Fish were collected as part of a paleodietary study using stable isotopy. Collagen was extracted according to the procedures in use in this community (Dufour et al. 1999), i.e. including a lipid elimination step (solvent extraction (Bligh and Dyer 1959); no chemical treatment was carried out on fish flesh. Reed and acacia leaves were dried as soon as possible after collection and were processed according to the standard acid-alkali-acid (AAA) treatment, i.e. 1M HCl, 0.1M then 1M NaOH, and 0.1M and 1M HCl. All treatments were performed at room temperature either in an ultrasonic bath or under agitation. Rinsing with ultrapure water followed each step.

About 1 mg of clean organic sample was then sealed in a quartz tube under vacuum with an excess of copper oxide and silver wire. The tubes were placed in an oven at 840°C for 5 hours to transform the organic matter into CO₂.

CO₂ Reduction and Measurements

Evolved CO₂ was either sent to the Gif-sur-Yvette Accelerator Mass Spectrometry (AMS) ¹⁴C laboratory (samples numbered GifA, Arnold et al. 1989) that operated a Tandemtron at that time or to the French national AMS facility in Saclay (LMC14, samples numbered SacA-, Cottreau et al. 2007) that operates a NEC AMS. CO₂ was thus graphitized according to protocols slightly modified from Arnold et al. (1989) and measured on AMS. See Dumoulin

Table 1 List of samples used to constrain reservoir age variability. The first three columns give the sample description, collection year and location. The last four columns concern laboratory handling: i – the fraction that was extracted from the sample for ^{14}C analysis, ii – lab identification for measurement and chemical treatment (if only one number, identification is for both chemistry and measurement); please note that the numbers of samples processed in 1999 by the Tandetron team are lost and cannot be recovered because the database crashed, iii – the ^{14}C age, and iv – the R that is calculated deriving from the ^{14}C atmospheric data of the collection year as provided by Hua et al. (2022) for the NH2 zone.

Sample identification							^{14}C data			
Sample N°	Location	Latitude	Longitude	Sample origin	Collection year	Type of sample	Analyzed fraction	Lab identification	Conventional age ± 1 sigma (BP)	R ** (^{14}C yr)
Antalya region										
1	Antalya (DSI PETROL)	36°55'17"N	30°40'35"E	Water table: (well -80m) (water pumping station)	2006	water	DIC*	SacA 5790	8250 \pm 40	8730 \pm 45
2	Öküzini marshes	37°06'02"N	30°35'04"E	Kırkgözüler (spring-fed lake north of ÖKüzini)	2006	water	DIC	SacA 5791	15810 \pm 60	16290 \pm 65
3	Öküzini marshes	37°04'56"N	30°34'50"E	Karstic source springing in the Öküzini marsh	2006	water	DIC	SacA 5792	17150 \pm 80	17630 \pm 80
4	Öküzini marshes	37°06'36"N	30°34'50"E	Pınarbaşı (water table in marsh at Plain center)	2006	water	DIC	SacA 5793	13360 \pm 50	13840 \pm 55
Central Anatolia region										
5	Acıgözü	37°42'44"N	33°40'21"E	Crater Lake (saline)	1999	water	DIC	GifA (1999)	12020 \pm 120	12770 \pm 120
6	Meke gölü	37°41'24.3"N	33°38'24"E	Crater lake (saline)	1999	water	DIC	GifA (1999)	2150 \pm 70	2895 \pm 75
7	Karapınar	~38°31'31"N	~34°32'53.7"E	Water table at Mr Aydınbelge's home (karstic)	1999	water	DIC	GifA (1999)	13770 \pm 130	14520 \pm 130
8a	Karacaören maar, near Eski Acıgol (1330 m)	38°32'18"N	34°35'28"E	Karacaören (Hot springs) (-80m) near crater lake	1999	water	DIC	GifA (1999)	\geq 43500	\geq 43500
8b	Eski Acıgol (1270m)	38°33'01"N	34°32'41"E	see Roberts et al. The Holocene (2001)	1995	modern soil	bulk OM	GrN-23467	1360 \pm 30	2265 \pm 35
8c	Eski Acıgol (1270m)			see Roberts et al. The Holocene (2001)		Sediment cores	bulk OM	Ra/Th and U/Th ages		3100 \pm 35
9	Nar Gölü (1363m)	38°20'16.0"N	34°27'20.8"E	Few meters from the shore of the crater lake	2002	reed	bulk OM	GifA17090/ Gif-11796	1525 \pm 30	2150 \pm 35

(Continued)

Table 1 (*Continued*)

Sample identification							¹⁴ C data				
Sample N°	Location	Latitude	Longitude	Sample origin	Collection year	Type of sample	Analyzed fraction	Lab identification	Conventional age ± 1 sigma (BP)	R ** (¹⁴ C yr)	
10	Nar Gölü (1363m)	38°20'12.5"N	34°27'12.1"E	100 meters from the shore of the crater lake	2002	acacia leaves	bulk OM	SacA44568/ Gif-11797	pMC 102 ± 0.3	400 ± 35	
11	Nar Gölü (1363m)	38°20'17.5"N	34°27'23.0"E	Crater lake	1999	water	DIC	GifA (1999)	26100 ± 390	26850 ± 390	
12	Obruks Plateau	37°35'26.4"N	33°35'50.1"E	Source (Water table in karst)	1999	water	DIC	GifA (1999)	4420 ± 80	5170 ± 80	
13	Çora maar (near Erciyes dağ)	38°39'36"N	35°17'59"E	Core Interstitial surface Water in maar (vole)	2006	water	DIC	SacA 5794	1420 ± 30	1900 ± 35	
Lake Van											
14	South of the lake (Taurus)	38°26'N	42°41'E	Collected in the karstic rejection (white plume)	2009	water	DIC	SacA 15095/ Gif-12407	4200 ± 35	4600 ± 40	
15	South of the lake (Taurus)			Collected outside the karstic rejection	2009	water	DIC	SacA 15096/ Gif-12408	1935 ± 35	2335 ± 40	
16	South of the lake (Taurus)			Collected outside the karstic rejection	2009	water	DIC	SacA 15097/ Gif-12409	1845 ± 30	2245 ± 35	
	Center of the lake	38°36'52"N	42°53'04"E	Makaroğlu et al. Quat Int. (2016)		sediment	bulk OM			2500-2800	
Armenian Lakes											
17	Lac Yerevan (908m)	40°09'38"N	44°28'43"E	Artificial lake near Erevan city	2014	farm trout bone	collagen	SacA41910/ Gif-13147	20 ± 30	230 ± 35	
18	Lac Sevan (Alt 1895m)	~40°34'N	~45°00'E	Lake	2014	wild trout bone	collagen	SacA41912/ Gif-13149	730 ± 30	940 ± 35	
19	Lac Sevan (Alt 1895m)	~40°34'N	~45°00'E	Lake	2014	carp bone	collagen	SacA41913/ Gif-13150	665 ± 30	875 ± 35	
20	Lac Sevan (Alt 1895m)	40°34'01"N	45°00'28"E	Lake	2014	water	DIC	SacA36821/ GifA-14010/Gif-13072	950 ± 30	1160 ± 35	
Kura Valley (Georgia)											
21	Tbilissi - Kura river	41°43'35"N	44°46'53"E	Kura River Near Tbilissi	2014	Laebobarbus (<i>Varicorhinus Sapoeta</i>) bone	collagen	SacA41911/ Gif-13148	pMC 102 ± 0,28	0 ± 35	

Table 1 (Continued)

Sample identification							¹⁴ C data			
Sample N°	Location	Latitude	Longitude	Sample origin	Collection year	Type of sample	Analyzed fraction	Lab identification	Conventional age ± 1 sigma (BP)	R ** (¹⁴ C yr)
Kura valley (Azerbaijan)										
22	Confluence with Agstafa river	41°14'29.2"N	45°26'12.5"E	Geologic bedrock (limestone)	2013	fish flesh	bulk OM	SacA 38572/ Gif-13109	1065 ± 30	1305 ± 35
23	Confluence with Agstafa river	–	–	Geologic bedrock (limestone)	2013	fish flesh	bulk OM	SacA 38573/ Gif-13110	900 ± 30	1140 ± 35
24	Bridge near confluence Tovuz river	41°03'51"N	45°47'41" E	Geologic bedrock (limestone)	2014	fish flesh	bulk OM	SacA40568/ Gif-13127	910 ± 30	1120 ± 35
25	Upper stream of Zeyem river	40°38'19" N	45°39'37" E	Geologic bedrock (volcanic)	2014	fish flesh	bulk OM	SacA40569/ Gif-13128	150 ± 30	360 ± 35
Mil Plain										
26	(near Agiabedi)	39°53'35.7"N	47°25'24.9"E	Irrigation channel from Aras River Geologic bedrock (Caspian Sea and Araxe River sediments)	2013	fish flesh	bulk OM	SacA 38574/ Gif-13111	205 ± 30	445 ± 35

*DIC dissolved inorganic carbon,

**R calculated using Hua et al. (2022) to know the atmospheric ¹⁴C content of the collection year




Figure 1 Location and geological context of the samples. 1a – Map of the Turkish region that shows the location of samples 1 to 16. 1b – Map of the Kura River and tributaries, lesser Caucasus and Eastern Anatolia (adapted from Ollivier et al. 2016, 2018). The red stars point to the location of the analyzed samples listed in Table 1. 1c – Geological map and 1d – profile of the Lesser Caucasus in the studied area (adapted from Ollivier et al. 2016 and Sosson et al. 2010). The dashed red line in 1c shows the AB section of the profile shown in 1d.

et al. (2017) and Moreau et al. (2020) for updated details at LMC14; Hatté et al. (2023), Tisnérat-Laborde et al. (2015) and Thil et al. (submitted) for updated details at LSCE.

Results are expressed as conventional ^{14}C age, rounded up as recommended by Stuiver and Polach (1977). The atmospheric ^{14}C age values were calculated using “Atmospheric ^{14}C age = $-8033 \cdot \ln(F^{14}\text{C})$ ”. All samples belong to NH zone 2 defined in Hua et al. (2022 and suppl. mat.). The average value of $F^{14}\text{C}$ of the year of collection was used for calculation. Reservoir age was calculated thanks to the equation previously reported. All results are reported in Table 1.

RESULTS AND DISCUSSION

First of all, it is worth pointing out that not all reservoir ages have the same meaning. Thus, those obtained on organic samples give a more integrated average value while the samples of dissolved bicarbonate (DIC) are more or less punctual, reflecting a degree of re-equilibration of the mineral carbon dissolved in these waters with the atmosphere. Furthermore, the reservoir age may be underestimated due to the variable input of bomb ^{14}C stored in soils and peats in lakes and rivers (Marchenko et al. 2021). Alternatively, it may be overestimated due to artificially high atmospheric values that are not yet in equilibrium with the old carbon reservoir (although the most recent samples (e.g. after 2000 AD) will be minimally affected).

The Antalya Region (Turkey)

The results are reported in Table 1. The Öküzini marshes extend at the edge of the Antalya Plain travertines, at the foot of the Western Taurus thick limestone massif (Kuzucuoğlu *et al.* 2001). Samples 2 to 4 taken near the springs feeding the marsh show reservoir ages varying between 13,800 and 17,600 ¹⁴C years. Further south, water from a well (sample 1) coming from the aquifer delivers a reservoir age of 8730 ± 45 ¹⁴C years which records the partial re-equilibration of these waters with the atmosphere.

The Central Anatolia Region (Turkey)

In a limestone and volcanic context, the waters of the aquifer also have variable reservoir ages: from 5170 ± 80 ¹⁴C years for sample 12, from the Obruks Plateau (doline lake in the limestone context north of Karapinar, Kuzucuoğlu 2019), to 14,520 ± 130 ¹⁴C years for a deep aquifer well at Karapinar (sample 7).

On the Anatolian Plateau, many volcanic edifices are associated with crater lakes, three of which were sampled (5, 6, 11). The surface water reservoir ages are extremely variable: 2895 ± 75 years for Meke Gölü (sample 6, basaltic maar, south of Karapinar), 12,770 ± 120 ¹⁴C years for Acigölü (sample 5, basaltic maar, east of Karapinar, most probably Holocene) and 26,850 ± 390 ¹⁴C years for Nar Gölü (sample 11, basaltic maar, north of the Göllüdağ massif). The high ages in Karapinar and Göllüdağ regions in fact record the CO₂ degassing of a volcanic edifice in the lake waters. Such outgassing is usually recorded by plants growing nearby (e.g. Pasquier-Cardin *et al.* (1999)). In the Nar Gölü crater which extends to the bottom of a basin bordered by the walls of the crater, about a hundred meters high, this degassing also marks the vegetation of the banks with reservoir ages decreasing rapidly from 2150 ± 35 ¹⁴C years for the reeds a few meters from the shore of the lake (sample 9) to 400 ± 35 ¹⁴C years for the leaves of the acacia 100 m further away (sample 10).

Further east, near the Erciyes volcano, the interstitial waters of the Çora maar (basaltic) give an age of 1900 ± 35 ¹⁴C years (Sample 13) used by Gauthier *et al.* (2014) for pollen sequence chronology.

To the east of the Eski Acigöl crater (rhyolitic maar, northern Cappadocia, Kuzucuoğlu *et al.* 1998), the Karacaören hot springs (sample 8a) have waters devoid of ¹⁴C. By their degassing and their contribution to the swamp in the center of the crater, the hot springs are at the origin of reservoir effects. Roberts *et al.* (2001) mentioned a modern soil age of 1630 BP, meaning a reservoir age of 2265 ± 35 ¹⁴C years (sample 8b). In the same study, Roberts *et al.* calculated, from Ra/Th and U/Th dating along sediment cores taken from the swamp, a reservoir age of 3100 ± 35 ¹⁴C years that has remained fairly constant for the last 16 millennia (sample 8c). The reservoir age difference between the 3100 ¹⁴C years offset for the sediment core and that of 2265 ¹⁴C years for soil may be due to a variable input of bomb ¹⁴C, stored in soils and peats, into lakes and rivers in agreement with the observations by Marchenko *et al.* (2021) in Western Siberia.

The Eastern Anatolia Region (Turkey)

Lake Van is bordered to the north by the volcanic complexes of Nemrut Dag and Süphan Dag and to the south by the limestone massif of the Eastern Taurus mountains. Lake Van is a large soda lake, with a hyper-alkalinity of the water explained by discharges of strongly mineral

deposits related to volcanic and hydrothermal activity, to which are added the carbonated contributions by karst inputs from the Taurus Mountains and rivers (Kuzucuoglu *et al.* 2010). Three samples were taken from the south of the lake: sample 14, which comes from a karst water discharge materialized in the lake by a whitish plume resulting from carbonate precipitation, has a reservoir age of 4600 ± 40 ^{14}C years; samples 15 and 16 taken outside this karstic discharge give similar results, 2335 ± 40 and 2245 ± 35 ^{14}C years, respectively. These intermittent karstic discharges linked to episodes of rain on the Taurus mountains are likely to increase the hard water effect but their influence remains limited to small areas to the south of the lake.

Previous work carried out on sedimentary cores and dated from varve counts made it possible to calculate reservoir ages. Lemcke (1996) and Kempe *et al.* (2002) working on cores retrieved in the center of the deepest depression of the lake (western part), estimated reservoir ages ranging from 2600 years in the surface sediment to about 4700 years in the Late Glacial. From cores extracted in the north of the same depression and at a lower depth, Makaroglu *et al.* (2016) calculated an average reservoir age of between 2500 and 2800 years for sediments collected from the central-southern part of the lake (lower than the previous one). Our measurements are in good agreement with these studies and thus reinforce the confidence in the estimates obtained by the varve counts. Unfortunately, the eastern part of the lake, where the rivers allow the development of greater biological activity at their mouth, has not yet been documented.

The Armenian Lakes

In Armenia, Lake Sevan, despite its resemblance to the volcanic environment of Lake Van, is not a sodic lake, thus allowing the development of a flourishing aquatic life. The water sample (20) has an age of 1160 ± 35 ^{14}C years which is almost entirely recorded in the flesh of a carp (Sample 19: 875 ± 35 ^{14}C years) and a trout (Sample 18, 940 ± 35 ^{14}C years) living in the western part of the lake. On the other hand, near Yerevan city, the artificial lake devoted to trout farming does not seem to be subject to a notable volcanic influence, as indicated by the age of sample 17 (trout flesh, 230 ± 35 ^{14}C years).

The Kura Valley (Georgia and Azerbaijan) and the Plain of Mil (Azerbaijan)

This variability of reservoir ages is also noted in Georgia and Azerbaijan for the Kura River and its tributaries (Samples 21 to 26). In a volcanic context, fish from the Kura in Tbilisi and that from Zeyem Caye indicate fairly low reservoir ages of 0 ± 35 ^{14}C years and 360 ± 35 ^{14}C years, respectively. Conversely, fish sampled in the Kura River upstream from the mouth of the Zeyem caye show significantly higher ages: 1305 ± 35 , 1140 ± 35 and 1120 ± 35 ^{14}C years. These ages result from the exchanges of the Kura River with the water table in the Cretaceous limestones which constitute the bed of the Kura in this region (Figure 1b–d). Such an effect has been described in detail by Coularis *et al.* (2016) for the Loire and its tributaries. In the plain of Mil, on a substrate consisting of sediment from the Caspian Sea and the Araxes River, fish from an irrigation canal give an intermediate age of 445 ± 35 ^{14}C years, in good agreement with previous observations.

CONCLUSION

The aging of the radiocarbon ages of the waters of the Anatolian lakes or the Lesser Caucasus is a general phenomenon resulting from volcanic activity (Central and Eastern Anatolia) and/

or from karstic discharges from the Taurus mountains (Öküzini in Mediterranean Anatolia, Central Anatolian volcanic province, Lake Van region) and from the Tertiary limestone basement in Central Anatolia. In the crater lake where it has been measured (Nar Gölü), the vegetation in the immediate surroundings records the degassing of the waters. These effects of degassing are observed in all the sites studied but often concern only restricted areas, located a few tens or hundreds of meters from the emitting source. The low geographical impact was also noted for the karstic discharges south of Lake Van. Such reservoir ages of lake water are almost all recorded by the aquatic fauna as evidenced by the results of Lake Van and Lake Sevan. For the rivers of the Caucasus, the reservoir ages are dependent on the geological substrate: ages on the limestone formations are high compared to those on the granite or volcanic formations, confirming the observations of Coularis *et al.* (2016).

This study shows, through the multiplicity of situations, a great variability in reservoir ages resulting, among other things, from the distance to the source of dead carbon and the level of re-equilibration of the latter with atmospheric CO₂.

ACKNOWLEDGMENTS

Thanks are due to E. Herrscher for fish collagen extraction samples, J-P. Dumoulin and L. Beck for Saclay AMS facilities, to B. Lyonnet, B. Helwing and S. Hansen, Kura Projects Managers, for their help in the field. This study was funded by the bilateral French (CNRS) / Turkish (TUBITAK) cooperation agreement (“Volcanism in Anatolia, ANOVAN”), the European Project ARIDUSEUROMED (ENV4-CT95-0062) and the French/German joint projects ANR-DFG (“Ancient Kura and Kura in Motion”). Fish were collected during the ORIMIL Project which aimed to reconstruct Neolithic and Bronze Age diets using stable isotopes. This research also benefited from the CEA measurements quota on the ARTEMIS program for ¹⁴C measurements.

REFERENCES

- Arnold M, Bard E, Maurice P, Valladas H, Duplessy J-C. 1989. ¹⁴C Dating with the Gif-sur-Yvette Tandemron Accelerator: Status Report and Study of Isotopic Fractionation in the Sputter Ion Source. *Radiocarbon* 31(3):284–291. doi: [10.1017/S0033822200011814](https://doi.org/10.1017/S0033822200011814)
- Bard E, Arnold M, Ostlund HG, Maurice P, Monfray P, Duplessy JC. 1988. Penetration of bomb radiocarbon in the tropical Indian Ocean measured by means of accelerator mass spectrometry. *Earth and Planetary Sciences Letters* 87:379–389. doi: [10.1016/0012-821X\(88\)90002-7](https://doi.org/10.1016/0012-821X(88)90002-7)
- Bligh EG, Dyer WJ. 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, 37:911–917. doi: [10.1139/o59-099](https://doi.org/10.1139/o59-099)
- Cottéreau E, Arnold M, Moreau C, Baqué D, Bavay D, Caffy I, Comby C, Dumoulin J-P, Hain S, Perron M, Salomon J, Setti V, 2007. ARTEMIS, the new ¹⁴C AMS at LMC14 in Saclay, France. *Radiocarbon* 49(2):291–299. doi: [10.1017/S0033822200042211](https://doi.org/10.1017/S0033822200042211)
- Coularis C, Tisnérat-Laborde N, Pastor L, Siclet F, Fontugne M. 2016. Temporal and spatial variations of freshwater reservoir ages in the Loire River watershed. *Radiocarbon* 58:549–563. doi: [10.1017/RDC.2016.36](https://doi.org/10.1017/RDC.2016.36)
- Deevey ES, Stuiver M. 1964. Distribution of natural isotopes of carbon in Linsley Pond and other New England lake. *Limnol. & Oceanogr.* 1: 1–11. doi: [10.4319/lo.1964.9.1.0001](https://doi.org/10.4319/lo.1964.9.1.0001)
- Deevey ES, Gross MS, Hutchinson GE, Kraybill HL. 1954. The Natural C Contents of Materials from Hard-Water Lakes. *Proc. Natl. Acad. Sci. U.S.A.* 40:285–288. doi: [10.1073/pnas.40.5.285](https://doi.org/10.1073/pnas.40.5.285)
- Dufour E, Boscherens H, Mariotti A. 1999. Palaeodietary implications of the isotopic variability in Eurasian Lacustrine Fish. *J. Archeol. Sci.*, 26:617–627. doi: [10.1006/jasc.1998.0379](https://doi.org/10.1006/jasc.1998.0379)
- Dumoulin JP, Comby-Zerbino C, Delqué-Količ E, Moreau C, Caffy I, Hain S, Perron M, Thellier B, Setti V, Berthier B, Beck L. 2017. Status report on sample preparation protocols developed at the LMC14 Laboratory, Saclay, France: from

- sample collection to ^{14}C AMS measurement. *Radiocarbon* 59:713–726. doi: [10.1017/RDC.2016.116](https://doi.org/10.1017/RDC.2016.116)
- Duplessy JC. 1972. La géochimie des isotopes stables du carbone dans la mer. Thèse Univ. Paris 6. 196 p.
- Fontugne M, Guichard F, Bentalab I, Strehle C, Lericolais G. 2009. Variations in ^{14}C reservoir ages in Black Sea waters and sedimentary organic carbon during the anoxic periods: influence of photosynthetic versus chemoautotrophic production. *Radiocarbon* 51:969–976. doi: [10.1017/S0033822200034044](https://doi.org/10.1017/S0033822200034044)
- Gauthier A, Mouralis D, Kuzucuoglu C, Fontugne M, Evren Atakay E, Evcimen Ö. 2014. Changements environnementaux en Anatolie centrale depuis la fin du LGM : la séquence du maar de Çora (Erciyes). International Congress AFEQ - CNF INQUA Q9 Lyon March 2014, Book of Abstracts.
- Hatté C, Arnold M, Dapoigny A, Daux V, Delibrias G, Du Boisgheueuc D, Fontugne M, Gauthier C, Guillier M-T, Jacob J, Jaudon M, Kaltnecker E, Labeyrie J, Noury C, Paterne M, Pierre M, Phouybandhyt B, Poupeau J-J, Tannau J-F, Thil F, Tisnéat-Laborde N, Valladas H. 2023. Radiocarbon dating on ECHOMICADAS, LSCE, Gif-sur-Yvette, France: new and updated chemical procedures. *Radiocarbon*. doi: [10.1017/RDC.2023.46](https://doi.org/10.1017/RDC.2023.46)
- Higham T, Warren R, Belinskij A, Härke H, Wood R. 2010. Radiocarbon dating, stable isotope analysis, and diet-derived offsets in ^{14}C ages from the Klin-Yar site, Russian North Caucasus. *Radiocarbon* 52(2–3):653–670. doi: [10.1017/S0033822200045689](https://doi.org/10.1017/S0033822200045689)
- Hua Q, Turnbull JC, Santos GM, Rakowski AZ, Ancapichún S, De Pol-Holz R, Hammer S, Lehman SJ, Levin I, Miller JB, Palmer JG, Turney CSM. 2022. Atmospheric Radiocarbon for The Period 1950–2019. *Radiocarbon* 68: 723–745. doi: [10.1017/rdc.2021.95](https://doi.org/10.1017/rdc.2021.95)
- Jones GA, Gagnon AR. 1994. Radiocarbon chronology of Black Sea sediments: Deep-Sea Research 41: 531–557. doi: [10.1016/0967-0637\(94\)90094-9](https://doi.org/10.1016/0967-0637(94)90094-9)
- Kempe S, Landmann G, Müller G. 2002. A floating varve chronology from the Last Glacial Maximum terrace of Lake Van/Turkey. *Zeitschrift für Geomorphologie, Supp. Iss.*, 126:97–114.
- Kuzmin YV, Nevesskaya LA, Krivonogov SK, Burr GS. 2007. Apparent ^{14}C ages of the ‘pre-bomb’ shells and correction values (R , ΔR) for Caspian and Aral Seas (Central Asia). *Nucl. Instrum. Methods Phys. Res. Sect. B* 259:463–466. doi: [10.1016/j.nimb.2007.01.187](https://doi.org/10.1016/j.nimb.2007.01.187)
- Kuzucuoglu C. 2019. Geomorphological landscapes and Pleistocene archives in the Konya Plain. In: Kuzucuoglu C, Çiner A, Kazancı N, editors. *Landscapes and landforms of Turkey*. Berlin: Springer. 353–358. doi: [10.1007/978-3-030-03515-0_17](https://doi.org/10.1007/978-3-030-03515-0_17)
- Kuzucuoglu C, Emery-Barbier A, Fontugne M, Kunesh S. 2001. The Öküzini marshes: a new Upper Pleistocene record on the Anatolian Mediterranean Coast. In: Yalçinkaya I, Otte M, Kozłowski J, Bar-Yosef O, editors. *Öküzini: final Paleolithic evolution in southwest Anatolia*. ERAUL 96. Liège. p. 79–82: 85–90.
- Kuzucuoglu C, Pastre J-F, Black S, Ercan T, Fontugne M, Guillou H, Hatté C, Karabiyikoglu M, Orth P, Türkecan A. 1998. Identification and dating of tephra layers from Quaternary sedimentary sequences of inner Anatolia. *Journal of Volc. Geotherm. Research* 85:153–172. doi: [10.1016/S0377-0273\(98\)00054-7](https://doi.org/10.1016/S0377-0273(98)00054-7)
- Kuzucuoglu C, Christol A, Mouralis D, Doğu AF, Akköprü E, Fort M, Brunstein D, Zorer H, Fontugne M, Scaillet S, Reyss JL, Guillou H, Karabiyikoglu M. 2010. Formation of the Upper Pleistocene terraces of Lake Van (Turkey). *Journal of Quaternary Science* 25(7):1124–1137. doi: [10.1002/jqs.1431](https://doi.org/10.1002/jqs.1431)
- Lemcke G. 1996. Paläoklimarekonstruktion am Van See (Ostanatolien, Türkei) [PhD dissertation]. Göttingen Univ. p. 195.
- Makaroglu O, Çagatay MN, Naci Orbay N, Pesonen LJ. 2016. The radiocarbon reservoir age of Lake Van, eastern Turkey. *Quat. Int.* 408:113–122. doi: [10.1016/j.quaint.2015.11.008](https://doi.org/10.1016/j.quaint.2015.11.008)
- Marchenko ZV, Svyatko SV, Grishin AE. 2021. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ isotope analysis of modern freshwater fish in the south of Western Siberia and its potential for palaeoreconstructions. *Quaternary International*, 598, 97–109. doi: [10.1016/j.quaint.2021.06.006](https://doi.org/10.1016/j.quaint.2021.06.006)
- Mangerud J. 1972. Radiocarbon dating of marine shells, including a discussion of apparent age of recent shells from Norway. *Boreas* 1:143–172. doi: [10.1111/j.1502-3885.1972.tb00147.x](https://doi.org/10.1111/j.1502-3885.1972.tb00147.x)
- Moreau C, Messenger C, Berthier B, Hain S, Thellier B, Dumoulin J-P, Caffy I, Sieudat M, Delqué-Količ E, Mussard S, Perron M, Setti V, Beck L. 2020. ARTEMIS, the ^{14}C AMS Facility of the LMC14 National Laboratory: a status report on quality control and microsample procedures. *Radiocarbon*, 1–16. doi: [10.1017/rdc.2020.73](https://doi.org/10.1017/rdc.2020.73)
- Oana S, Deevey ES. 1960. Carbon 13 in lake water and its possible bearing on paleolimnology. *Am.J.Sci.*, 258-A:253–272.
- Ollivier V, Fontugne M, Lyonnet B, Chataigner C. 2016. Base level changes, river avulsions and Holocene human settlement dynamics in the Caspian Sea area (middle Kura valley, South Caucasus). *Quaternary International* 395, 79–94. doi: [10.1016/j.quaint.2015.03.017](https://doi.org/10.1016/j.quaint.2015.03.017)
- Ollivier V, Fontugne M, Hamon C, Decaix A, Hatté C, Jalabadze M. 2018. Neolithic water

- management and flooding in the Lesser Caucasus (Georgia). *Quaternary Science Reviews* 197: 267–287. doi: [10.1016/j.quascirev.2018.08.016](https://doi.org/10.1016/j.quascirev.2018.08.016)
- Ollivier V, Fontugne M, Lyonnet B. 2015. Geomorphic response and ^{14}C chronology of base-level changes induced by Late Quaternary Caspian Sea mobility (middle Kura Valley, Azerbaijan). *Geomorphology* 230, 109–124. doi: [10.1016/j.geomorph.2014.11.010](https://doi.org/10.1016/j.geomorph.2014.11.010)
- Pasquier-Cardin, Allard P, Ferreira T, Hatté C, Couthino R, Fontugne M, Jaudon M. 1999. Magma-derived CO_2 emissions recorded in ^{14}C and ^{13}C of plants growing in Furnas caldera, Azores. *Journal of Volcanology and Geothermal Research* 92, 195–207. doi: [10.1016/S0377-0273\(99\)00076-1](https://doi.org/10.1016/S0377-0273(99)00076-1)
- Philippsen B, Heinemeier J. 2013. Freshwater reservoir effect variability in Northern Germany. *Radiocarbon* 55:1085–1101. doi: [10.1017/S0033822200048001](https://doi.org/10.1017/S0033822200048001)
- Roberts N, Reed JM, Leng MJ, Kuzucuoglu C, Fontugne M, Bertaux J, Woldring JH, Bottema S, Black S, Hunt SE, Karabiyikoglu M. 2001. The tempo of Holocene climatic change in the eastern Mediterranean region: new high-resolution crater-lake sediment data from central Turkey. *The Holocene* 11:721–736. doi: [10.1191/09596830195744](https://doi.org/10.1191/09596830195744)
- Shishlina NI, van der Plicht J, Hedges REM, Zazovskaya EP, Sevastyanov VS, Chichagova OA. 2007. The catacomb cultures of the North-West Caspian steppe: ^{14}C chronology, reservoir effect, and paleodiet. *Radiocarbon* 49(2): 713–726. doi: [10.1017/S0033822200042600](https://doi.org/10.1017/S0033822200042600)
- Sosson M, Rolland Y, Muller C, Danelian T, Melkonyan R, Kekelia S, Adamia S, Babazadeh V, Kangarli T, Avagyan A, Galoyan G, Mosar J. 2010. Subductions, obduction and collision in the Lesser Caucasus (Armenia, Azerbaijan, Georgia), new insights. In: Sosson M, Kaymakci N, Stephenson R, Bergerat F, Starostenko V, editors. *Sedimentary basin tectonics from the Black Sea and Caucasus to the Arabian Platform*. Geological Society of London, Special Publication 340:329–352.
- Stuiver M, Polach HA. 1977. Discussion reporting ^{14}C data. *Radiocarbon* 19:355–363. doi: [10.1017/S0033822200003672](https://doi.org/10.1017/S0033822200003672)
- Thil F, Tisnérat-Laborde N, Hatté C, Noury C, Paterne M, Phouybandhyt B, submitted. Microsample analysis with ECHoMICADAS facility: current status. *Radiocarbon*.
- Tisnérat-Laborde N, Thil F, Synal H.-A, Cersoy S, Hatté C, Gauthier C, Massault M, Michelot J.-L, Noret A, Siani G, Tombret O, Vigne J.-D, Zazzo A. 2015. ECHoMICADAS: A new compact AMS system to measuring ^{14}C for Environment, Climate and Human Sciences, 22nd International Radiocarbon Conference, Dakar, Senegal, 16–20 November 2015. PHYS-O.05.