

Gender differences in the structure of risk for alcohol use disorder in adolescence and young adulthood

K. T. Foster^{1*}, B. M. Hicks², W. G. Iacono³ and M. McGue³

¹Department of Psychology, University of Michigan, Ann Arbor, MI, USA

²Department of Psychiatry, University of Michigan, Ann Arbor, MI, USA

³Department of Psychology, University of Minnesota, Minneapolis, MN, USA

Background. Gender differences in the prevalence of alcohol use disorder (AUD) have motivated the separate study of its risk factors and consequences in men and women. However, leveraging gender as a third variable to help account for the association between risk factors and consequences for AUD could elucidate etiological mechanisms and clinical outcomes.

Method. Using data from a large, community sample followed longitudinally from 17 to 29 years of age, we tested for gender differences in psychosocial risk factors and consequences in adolescence and adulthood after controlling for gender differences in the base rates of AUD and psychosocial factors. Psychosocial factors included alcohol use, other drug use, externalizing and internalizing symptoms, deviant peer affiliation, family adversity, academic problems, attitudes and use of substances by a romantic partner, and adult socio-economic status.

Results. At both ages of 17 and 29 years, mean levels of psychosocial risks and consequences were higher in men and those with AUD. However, the amount of risk exposure in adolescence was more predictive of AUD in women than men. By adulthood, AUD consequences were larger in women than men and internalizing risk had a stronger relationship with AUD in women at both ages.

Conclusions. Despite higher mean levels of risk exposure in men overall, AUD appears to be a more severe disorder in women characterized by higher levels of adolescent risk factors and a greater magnitude of the AUD consequences among women than men. Furthermore, internalizing symptoms appear to be a gender-specific risk factor for AUD in women.

Received 29 July 2014; Revised 5 May 2015; Accepted 6 May 2015; First published online 29 June 2015

Key words: Adolescent risk factors, adult consequences, alcohol use disorder, gender differences.

Introduction

Relative to women, men consume alcohol more frequently and in greater quantities, and so have higher rates of alcohol use disorder (AUD) [Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV) abuse, 24.6%; dependence, 17.4%] than women (abuse, 11.5%; dependence, 8.0%) (Keyes *et al.* 2008). These differences have encouraged the separate study of risk exposure and outcomes in men and women; however, leveraging gender as a third variable to help account for the association between psychosocial factors and AUD may advance understanding of etiological mechanisms and clinical outcomes (Rutter *et al.* 2003).

Two types of gender effects – mean-level and structural – help to organize our understanding of the links among gender, AUD and important psychosocial variables. Mean-level gender effects (i.e. the main effect of gender) refer to the absolute amount or severity of risk exposure experienced by men and women with AUD (e.g. a risk factor may occur at a higher rate in men than women). Another important source of gender differences are structural effects or the strength of the association between a risk factor and AUD within each gender. In particular, gender differences in both AUD and a psychosocial variable can obscure gender's moderating effects (i.e. interaction effects). For instance, a psychosocial variable could have a stronger association with AUD in one gender than the other, irrespective of mean-level gender differences for that variable. Risk factors with strong structural effects on AUD may be more potent in one gender and, consequently, require a lower mean level of exposure to

* Address for correspondence: K. T. Foster, University of Michigan, East Hall, 530 Church Street, Ann Arbor, MI 48109, USA.
(Email: ktfooster@umich.edu)

produce AUD. By controlling for the mean-level effect of gender, the strength of the association between AUD and each psychosocial variable (i.e. interaction effects) can be estimated directly.

Delineating the mean-level and structural effects of gender for multiple psychosocial variables can increase insight regarding the accumulation of impairment across different domains that may comprise gender-specific pathways for AUD. Further, evaluating mean-level and structural effects at key developmental periods for AUD will help identify patterns of psychosocial impairment that contribute to the onset and persistence of AUD. Specifically, late adolescence (when early-onset AUD cases emerge) and young adulthood (when drinking reduces and serious consequences accumulate if AUD persists) are particularly informative periods to examine how gender differences in early risk factors and consequences of AUD underlie gender difference in its prevalence.

Common risk factors and outcomes for AUD

AUD represents the end point in a long history of biological, psychosocial and environmental risk factors interacting and accumulating over the course of development (Caspi et al. 1995, 1996; Masse & Tremblay, 1997; Blazei et al. 2006; Wong et al. 2006; Zucker, 2006). Importantly, these variables may exhibit gender differences in their mean-level and structural associations with AUD. For example, behavioral disinhibition – a heritable cluster of disinhibited personality traits and externalizing disorders (Moffitt et al. 2001; Rutledge & Sher, 2001; Krueger et al. 2002; Slutske et al. 2002; Kendler et al. 2003) – increases the odds of early-onset AUD and other problem behaviors (e.g. drug use, delinquency, precocious sexual behavior; Hawkins et al. 1992; Iacono et al. 2008). A cycle of coercive parent-child interactions, conflict with socializing agents (e.g. teachers and prosocial peers) and affiliation with deviant peers (Dishion et al. 1991; Tangney et al. 1996; Patterson & Yoerger, 1997, 1999; Granic & Patterson, 2006) also contributes to AUD and related adult impairment (e.g. unemployment, romantic partnership problems and life satisfaction; Cranford et al. 2011). Other non-specific risk factors for AUD include internalizing disorders and exposure to traumatic life events like physical and sexual abuse and assault (Cutler & Nolen-Hoeksema, 1991; Widom et al. 1995; Wilsnack et al. 1997; Kilpatrick et al. 2000; Nolen-Hoeksema & Hilt, 2006). While these patterns of risk and consequences are generally associated with AUD, there may be gender differences in their mean level and the strength of their association with AUD (i.e. structural effects). Examining the nature of gender

differences in risk factors and consequences can help explain the prevalence, etiological course and relative severity of AUD.

Gender differences in mean levels of risk factors for AUD

The greater prevalence of AUD in men suggests that men experience higher mean levels of risk exposure than women (i.e. between-gender differences in those with AUD). To express AUD, then, a woman must be more deviant relative to the norm for her gender (i.e. within-gender mean-level effects) than a man. Therefore, elucidating gender differences in the risks and consequences for AUD requires comparing men and women with alcohol use problems with those of the same gender that do not (e.g. AUD women *versus* non-AUD women). However, risk factors are often studied individually using between-gender comparisons of mean-level effects, making it difficult to discern their relative contributions to the development of AUD in men and women (Labouvie & McGee, 1986; Waldeck & Miller, 1997; Moffitt et al. 2001; Petry et al. 2002). For example, though mean levels of behavioral disinhibition and sexual trauma vary significantly by gender, both are equally predictive of AUD in boys and girls (Stein et al. 1988; Cutler & Nolen-Hoeksema, 1991; Moffitt et al. 2001; Iacono et al. 2008). Studies conducting between-gender comparisons of a single factor are not well equipped to test the etiological importance of these factors as gender-specific pathways to AUD. Within-gender comparisons of mean levels of risk between those with and without AUD estimate the importance of a risk factor on AUD separately for men and women. These comparisons are vital for identifying which risk factors are more predictive of AUD in men relative to women.

Gender differences in structural associations between AUD risk factors and consequences

Another possible cause of gender differences in AUD is that psychosocial factors may have different structural associations with AUD in men and women. Notably, the association (e.g. correlation) between AUD and several psychosocial factors differs across gender. For example, alcohol's rewarding effects have been linked with AUD in men (Schuckit, 1994; Wilhelmsen et al. 2003) while, the lower threshold for alcohol-related impairment and toxicity in women has been conceptualized as a deterrent of heavy drinking (Klassen & Wilsnack, 1986; Niaura et al. 1987; Nixon, 1994; Blume & Russell, 2001). Also, internalizing disorders may play a more prominent role in the development of AUD among women (Nolen-Hoeksema, 2004). For example, even after controlling for higher rates of

depression in women relative to men, depressive symptoms have been prospectively associated with AUD in women (Kendler *et al.* 1997; Brady & Randall, 1999; Sannibale & Hall, 2001). Finally, even among those who desist by young adulthood, a greater proportion of women than men exhibit enduring consequences of AUD, including prolonged polysubstance abuse, psychiatric problems and poor psychosocial adjustment (Hicks *et al.* 2010a, b; Foster *et al.* 2014). As such, AUD may be less prevalent in women because it is a more extreme form of psychopathology, requiring a greater loading of risk before the disorder is expressed. These findings suggest then that gender-specific associations between AUD and its risk factors may contribute to gender differences in the prevalence of AUD.

Leveraging gender differences to study the etiology of AUD

Research on AUD has often been constrained within gender under the assumption that the link between AUD and its risks and consequences differs by gender. Consequently, few studies have compared the relative effects of multiple risk factors and outcomes for AUD in men and women in the same study. Without such tests, it is unclear if psychosocial risks and outcomes are simply more prevalent in one gender, or if gender-specific influences increase their association with AUD in one gender more than the other. To examine the potential moderating role of gender, we directly compared the effects of several well-replicated risk factors for and outcomes of AUD in a large, community sample of men and women at 17 and 29 years of age. Specifically, separately for men and women, we first estimated the odds of developing AUD by 29 years of age given the mean level of risk evident for each risk factor at 17 years of age after controlling for the average amount of exposure to the risk factor within each gender. We then estimated the association between AUD and several psychosocial outcomes at 29 years of age, after adjusting for gender differences in the base rates of AUD and the psychosocial outcome. We hypothesized that mean-level but not structural gender effects would be present across risk factors. One exception, however, was internalizing disorders for which we predicted a stronger association with AUD in women relative to men.

Method

Sample

Participants were male ($n=578$) and female ($n=674$) twins of the Minnesota Twin Family Study (MTFS), a prospective, community-based study designed to investigate the etiology of substance use disorders (for

extensive details on study design, see Iacono *et al.* 1999). Twin pairs born between the years of 1972 and 1979 were recruited from Minnesota public birth records at 17 years of age. Of the 90% of families located, 83% completed the in-person laboratory assessment at the University of Minnesota. Nearly all participants were of European American ancestry (96%) and were similar to non-participating families in parental occupation, education and history of mental health treatment.

Assessment

At the age 17 years assessment, multiple informants (twins, parents and teachers) provided information on alcohol and other substance use along with psychiatric, psychosocial and environmental functioning. Follow-up assessments occurred every 3–5 years at the target ages of 20 years old ($n=1110$, 89% retention rate, 83% of men and 93% of women), 24 years old ($n=1159$, 92% retention rate, 94% of men and 91% of women) and 29 years old ($n=1164$, 93% retention rate, 91% of men and 94% of women). The current report focused on risk factors and domains of psychosocial functioning at the ages of 17 and 29 years (Hicks *et al.* 2009, 2010a, b) to assess both risk and outcomes for lifetime AUD by the age of 29 years. More comprehensive descriptions of the measures are provided elsewhere (Hicks *et al.* 2009, 2010a, b).

AUD diagnosis

Trained staff administered the Substance Abuse Module (Robins *et al.* 1987) of the Composite International Diagnostic Interview (Robins *et al.* 1988) to determine lifetime AUD status at the age of 17 years. Subsequent evaluations measured AUD symptoms since the last assessment. Consistent with DSM-5, AUD was defined as two or more symptoms of alcohol abuse or dependence for at least one assessment by the age of 29 years (men, $n=316$; women, $n=155$). Multiple studies using the MTFS sample have demonstrated the validity of this approach (Elkins *et al.* 2004, 2006, 2007; McGue & Iacono, 2005). For each gender, an AUD group was compared with a non-AUD group (i.e. no more than one AUD symptom at any assessment; men, $n=226$; women, $n=449$) on the risk factors and outcomes.

Measures of risk and impairment

Prior studies using the MTFS sample have linked several measures of risk and consequences with AUD in both genders (Hicks *et al.* 2010a, b; Foster *et al.* 2014). Using principal components analysis and theoretical considerations, we combined variables into composites (i.e. mean z-score across constituent variables) to assess

critical domains of AUD risk exposure and impairment at the ages of 17 and 29 years. Whenever possible, the same measures were used to assess each factor across different assessment periods. However, certain domains were age-specific including family adversity and academic problems in adolescence (age 17 years only) and romantic partner relationships and socioeconomic status in adulthood (age 29 years only).

Alcohol, nicotine and illicit substance use

Alcohol use was assessed using past-year average quantity and the maximum number of drinks consumed in 24 h. Nicotine and illicit drug use was estimated using DSM-IV symptoms of nicotine dependence and abuse/dependence for illicit drugs, along with quantity and frequency of use and the number of drug classes tried. Substances assessed included nicotine, amphetamine, cannabis, cocaine, hallucinogen, inhalant, opioid, phencyclidine (PCP) and sedatives. The illicit drug class with the greatest number of reported symptoms was used for each participant's drug abuse/dependence variable.

Externalizing symptoms

At the age of 17 years, symptoms of adult antisocial behavior were evaluated using a structured interview similar to the Structured Clinical Interview for DSM-IV Axis II (SCID-II) module for antisocial personality disorder. Personality traits of disinhibition were assessed using the behavioral constraint (i.e. inclination toward planning, traditional social values, and caution) factor of the Multidimensional Personality Questionnaire (MPQ; Tellegen & Waller, 2008). At the age of 29 years, symptoms of adult antisocial behavior over the past 6 years were evaluated in conjunction with behavioral constraint.

Internalizing distress

Internalizing distress was assessed using lifetime symptoms of major depressive disorder, negative emotionality and significant mental health problems (i.e. prior suicide attempts, mental health treatment, or psychiatric hospitalization). Symptoms of major depression were assessed using the Structured Clinical Interview for DSM-III-R. Trait negative emotionality (i.e. propensity toward breakdown under stress and a suspicious, aggressive interpersonal style) was assessed using the MPQ. Mental health problems were assessed using the Lifetime Events Interview (Bemmels et al. 2008). At the age of 29 years, the same variables were used to estimate internalizing distress (i.e. major depression symptoms, mental health problems over the past 6 years, and negative emotionality scores).

Deviant peer group affiliation

Adolescent peer groups were assessed for antisocial ($\alpha=0.82$; e.g. my friends enjoy getting drunk, get into fights, can't seem to hold a job) and prosocial behaviors ($\alpha=0.60$; e.g. my friends work hard, do volunteer work, have a regular job) using a teacher rating form (five items each; Walden et al. 2004). At the age of 29 years, participants reported antisocial (coded positive) and prosocial (coded negative) qualities of their own peer group (27-item questionnaire).

Family adversity

At the age of 17 years, family adversity was indexed by socio-economic status for the family of origin, quality of the parent-child relationship, and parental externalizing disorder symptoms. Socio-economic status was defined as the mean z-score for each parent's years of education, occupational status (Hollingshead index) and annual income. The Parent Environment Questionnaire (PEQ; Elkins et al. 1997) measured quality of the parent-child relationship from each parent and adolescent (mean z-score of the three informant ratings for the first principal component of the PEQ scales; Hicks et al. 2009). Parental externalizing disorders were indexed using the symptoms of antisocial personality disorder and alcohol, nicotine and drug abuse/dependence.

Academic problems

At the age of 17 years, difficulties in school were evaluated using the Academic History Questionnaire (Johnson et al. 2006) that queried mother and child for cumulative grade-point average and positive engagement with academics (seven items; $\alpha=0.83$).

Adult romantic partner drug use

Participants in a current romantic relationship (i.e. married, cohabiting, or consistently dating the same person for 3 months or more) at the age of 29 years reported their partner's past-year drinking patterns including the frequency, quantity and proportion of intoxicating drinking episodes and attitudes toward substance use (e.g. 'my spouse/partner would be upset if he knew I was smoking'; 'my spouse/partner would purchase alcohol if I asked him to'; 'my spouse's/partner's friends use marijuana') using an 11-item scale ($\alpha=0.84$).

Adult socio-economic status

Measures of educational attainment, a Hollingshead rating of current occupational status, and annual income all reported in the Life Events Interview and

the Social Adjustment Inventory were used to create a composite for socio-economic status.

Statistical analysis

A series of hierarchical linear models was fit to estimate the associations between AUD, gender and the risk factors assessed at the age of 17 years. Generalized estimating equations were used to model the associations between adolescent risk exposure and the odds of developing AUD by the age of 29 years in men and women using the following model:

$$\text{AUD}_{ij} = \gamma_{00} + \gamma_{10} \times \text{GENDER}_{ij} + \gamma_{20} \times \text{RISK}_{ij} + \gamma_{30} \times \text{GENDER} \times \text{RISK}_{ij} + \mu_{0j}.$$

Data for this model were mean-centered within each gender to facilitate interpretation as follows. The main effect of gender (γ_{10}) was the increase in the odds of AUD by the age of 29 years given gender status and an average level of risk exposure for that gender (e.g. increase in odds for men compared with women, given average levels of risk within men and women). The main effect for the risk factor (γ_{20}) was estimated as the increase in the odds of AUD by the age of 29 years given a 1 s.d. increase in risk exposure. The gender \times risk interaction term (γ_{30}) tested whether the association between AUD and the risk factor was moderated by gender. Models were fit using the Bernoulli option in HLM 7.0 (SSI Scientific Software, USA) specifically designed to predict binary outcomes (i.e. AUD or non-AUD by the age of 29 years in this case). Data were nested within families (γ_{00}) to adjust for the non-independence of the twin data and any non-normal distributions for the risk factor variables. A residual term was also included (μ_{0j}) to account for variation in the outcome not accounted for by the predictor variables.

For the age 29 years assessment, we estimated the associations between lifetime AUD, gender and several psychosocial consequences using the following model:

$$\text{OUTCOME}_{ij} = \gamma_{00} + \gamma_{10} \times \text{GENDER}_{ij} + \gamma_{20} \times \text{AUD}_{ij} + \gamma_{30} \times \text{GENDER} \times \text{AUD}_{ij} + \mu_{0j}.$$

This model included the main effects of gender (γ_{10}) and AUD status (γ_{20}) and the gender \times AUD interaction (γ_{30}) in the prediction of each psychosocial outcome at the age of 29 years. Parameters were adjusted for other variables in the model so that the effect of gender on the outcome was adjusted for AUD *versus* non-AUD group differences in outcome, while the effect of AUD was adjusted for gender differences on the outcome. The gender \times AUD interaction term tested whether gender moderated the association between AUD and the adult outcome. Models for outcome

variables at the age of 29 years were fit using the cluster option and the MLR estimator in Mplus 5.0 (<https://www.statmodel.com/>) that is appropriate for continuous outcomes (i.e. degree of the risk outcome). All standard errors and p values were adjusted for the non-independence of the family-level data (i.e. nested by γ_{00}) and any non-normal distributions for risk factor variables. A residual term was also included (μ_{0j}) to account for variance in the outcome independent of gender, AUD and their interaction.

Results

Over a third of the sample ($n=471$, 37.6%) reported two or more symptoms of AUD at one or more assessments by the age of 29 years. In our sample, lifetime AUD was more prevalent among men than women [odds ratio (OR) 2.37, 95% confidence interval (CI) 1.90–2.90].

Risk exposure at the age of 17 years and lifetime AUD outcomes by gender

Results for the main effects of gender, risk factor, and the gender \times risk factor interaction terms at the age of 17 years are reported in [Table 1](#). The average level of risk exposure common to boys at the age of 17 years significantly increased the odds of developing AUD relative to the average level of risk exposure common to girls at the age of 17 years. The significant main effects of risk exposure at the age of 17 years indicated that higher levels of risk increased the odds of developing AUD by the age of 29 years. Within both men and women, a 1 s.d. increase in alcohol use, other drug use, externalizing problems, deviant peers, family adversity and academic problems increased the odds of AUD. Internalizing distress at the age of 17 years was associated with increased odds of AUD in women (OR 1.55, 95% CI 1.30–1.85, $p<0.001$) but not men (OR 1.23, 95% CI 0.96–1.57, $p=0.098$), suggesting a gender-specific risk factor for AUD. Finally, the association between each risk exposure and AUD was stronger for women except for family adversity, suggesting that similar increases in risk for both genders are linked with more severe consequences in women compared with men (see [Fig. 1](#)). For instance, a 1 s.d. increase in drug use was associated with a greater increase in the odds of AUD in women (OR 2.59, 95% CI 2.02–3.32) compared with men (OR 1.78, 95% CI 1.36–2.32). Consequently, we detected gender \times risk factor interactions for alcohol use, other substance use, deviant peers and academic problems, such that greater risk exposure on these variables had a significantly stronger association with AUD in women relative to men.

Table 1. T-score means, standard deviations, Cohen's *d* and results for the generalized estimating equation using risk factors at age 17 years and gender to predict the odds of developing AUD by the age of 29 years

Predictors at age 17 years	Gender difference		AUD status by age 29 years			Risk effect at age 17 years ^d		Gender effect (men <i>v.</i> women) ^e	Risk × gender ^f
	Overall: mean (s.d.) ^a	<i>d</i> ^b	Control: mean (s.d.) ^a	AUD: mean (s.d.) ^a	<i>d</i> ^c	β (s.e.)	OR (95% CI)	β (s.e.)	<i>p</i>
Alcohol use									
Total	50.0 (10.0)		46.4 (6.2)	56.1 (12.0)	1.02				
Women	48.6 (7.9)		46.4 (5.6)	55.1 (9.9)	1.09	1.46 (0.15)	4.31 (3.20–5.80)***		
Men	52.3 (12.0)	0.36	46.3 (7.2)	56.5 (12.9)	0.98	1.02 (0.12)	2.79 (2.19–3.54)***	1.78 (0.16)***	0.025*
Other drug use									
Total	50.0 (10.0)		47.6 (6.3)	53.5 (13.0)	0.58				
Women	50.1 (9.8)		47.7 (5.7)	56.9 (14.9)	0.82	0.95 (0.12)	2.59 (2.02–3.32)***		
Men	50.0 (10.3)	−0.01	47.4 (7.3)	51.9 (11.6)	0.46	0.57 (0.13)	1.78 (1.36–2.32)***	1.51 (0.15)***	0.043*
Externalizing									
Total	50.0 (10.0)		47.2 (7.3)	54.7 (11.7)	0.77				
Women	48.0 (9.1)		46.1 (7.0)	53.5 (12.0)	0.76	0.90 (0.12)	2.46 (1.95–3.10)***		
Men	52.8 (10.5)	0.49***	49.3 (7.6)	55.2 (11.6)	0.61	0.63 (0.11)	1.89 (1.51–2.35)***	1.56 (0.15)***	0.112
Internalizing									
Total	50.0 (10.0)		48.8 (9.3)	51.5 (10.2)	0.28				
Women	50.0 (11.1)		48.9 (10.0)	54.7 (13.1)	0.51	0.43 (0.08)	1.55 (1.30–1.85)***		
Men	49.5 (8.0)	0.09	48.7 (7.7)	50.0 (8.1)	0.16	0.20 (0.12)	1.23 (0.96–1.57)	1.45 (0.15)***	0.129
Deviant peers									
Total	50.0 (10.0)		48.1 (8.5)	52.5 (11.0)	0.45				
Women	49.8 (9.3)		48.2 (8.4)	54.8 (10.2)	0.71	0.74 (0.12)	2.11 (1.65–2.69)***		
Men	50.0 (10.4)	0.02	48.0 (8.7)	51.5 (11.2)	0.35	0.34 (0.10)	1.41 (1.14–1.73)**	1.55 (0.16)***	0.014*
Family adversity									
Total	50.0 (10.0)		48.6 (9.6)	51.6 (9.9)	0.32				
Women	50.0 (11.0)		48.8 (10.6)	53.3 (11.6)	0.40	0.37 (0.09)	1.46 (1.21–1.76)***		
Men	49.7 (8.4)	−0.02	48.1 (7.4)	50.9 (8.9)	0.35	0.44 (0.12)	1.56 (1.21–2.01)**	1.45 (0.15)***	0.687
Academic problems									
Total	50.0 (10.0)		47.8 (9.0)	52.9 (10.5)	0.53				
Women	48.6 (9.7)		46.8 (8.9)	53.6 (10.1)	0.71	0.71 (0.10)	2.03 (1.65–2.52)***		
Men	51.4 (10.0)	0.28	49.6 (8.8)	52.6 (10.7)	0.30	0.30 (0.10)	1.35 (1.11–1.65)**	1.47 (0.15)***	0.006**

AUD, Alcohol use disorder; s.d., standard deviation; s.e., standard error; OR, odds ratio; CI, confidence interval.

^a T-score means arranged by gender and AUD status (women: *n* = 155 AUD, 449 control; men: *n* = 316 AUD, 226 non-AUD) reflect mean level of risk exposure for each group at the age of 17 years.

^{b,c} Cohen's *d* effect sizes estimate the magnitude of the difference in the average level of risk between males and females overall (i.e. all men *versus* all women; *d*^b) and the change in risk exposure coinciding with AUD in each gender (e.g. control women *versus* AUD women; *d*^c).

^d The risk effect for each gender estimates the increased odds of developing AUD by the age of 29 years as a result of a 1 s.d. increase in risk exposure for a person of that gender (i.e. potency of the risk).

^e The gender effect estimates the increase in odds of developing AUD by the age of 29 years in men compared with women given an average level of risk exposure within each gender. A positive β value denotes higher odds in men compared with women.

^f The risk × gender interaction effect was tested if gender moderated the relationship between the risk factor and the AUD outcome.

* *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001.

Consequences of AUD at the age of 29 years

Results for the main effects for gender, AUD, and the gender × AUD interactions for the age 29 years outcomes are reported in Table 2. Lifetime AUD predicted

greater alcohol consumption, nicotine and illicit drug use, internalizing distress, externalizing problems, deviant peer affiliation and substance use by a romantic partner at the age of 29 years. Men exhibited




Fig. 1. Odds of developing alcohol use disorder by the age of 29 years for each 1 S.D. increase from the average level of risk exposure for a person of that gender at the age of 17 years. OR, Odds ratio; CI, confidence interval.

significantly greater alcohol use, other substance use, externalizing symptoms, deviant peer affiliation and socio-economic status. Women reported greater partner substance use. Although mean-level comparisons at the age of 29 years suggest that men with AUD were more impaired than women with AUD, the difference in psychosocial outcome between non-AUD and AUD groups was larger among women than men for alcohol use, drug use, internalizing, deviant peers and romantic partner drug use. That is, AUD coincided with greater overall decrements in functioning among women than men compared with those of the same gender without the disorder (see Fig. 2). For example, the effect size of AUD on other drug use was larger in women ($d=1.00$) relative to men ($d=0.65$). We also detected a gender \times AUD interaction for internalizing distress, such that the differences between AUD and non-AUD groups were greater among women than men.

Discussion

The higher prevalence of AUD in men relative to women suggests that mean levels of risk exposure for AUD are either greater in men or that certain risk factors have differential effects across gender. To test the moderating role of gender, we estimated the strength of the association between AUD and several established risk factors and negative outcomes after adjusting for gender differences in their prevalence. Our results confirmed the hypothesis that women with AUD have a greater loading of risk at 17 years old

and that AUD increases mean levels of psychosocial impairment in young adulthood for both men and women, but that internalizing distress has a stronger structural relationship with AUD for women than men.

Greater exposure to each risk factor was associated with increased odds of AUD by 29 years old. Further, men tended to have higher mean levels of risk exposure that contributed to a higher prevalence of AUD in men relative to women. Despite the higher level of absolute risk in men, AUD in women was associated with an especially high level of risk exposure during adolescence relative to their gender norm and a higher level of risk exposure was necessary for women to exhibit AUD relative to men. Gender variation in the psychosocial consequences of AUD during young adulthood followed a similar pattern. That is, the magnitude of the difference between AUD and non-AUD impairment levels at 29 years old (i.e. effect size) was larger in women than men for most variables. Compared with their gender norm, women with AUD tended to experience both higher risk exposure in adolescence and more negative outcomes in young adulthood relative to men with AUD. Consequently, AUD appears to be a more severe form of psychopathology in women, with a risk structure that is present early in development (i.e. at least by adolescence).

We detected several interactions between gender and adolescent risk factors, such that increases in alcohol use, other substance use, deviant peer relationships and academic problems increased the odds of

Table 2. T-score means, standard deviations, Cohen's *d* and β values when AUD status and gender predict age 29 years psychosocial functioning

Criterion at age 29 years	Overall: mean (s.d.) ^a	AUD status by age 29 years		AUD effect ^d		Gender effect ^e		AUD × gender ^f
		Control: mean (s.d.) ^a	AUD: mean (s.d.) ^a	<i>d</i> ^b	β	<i>d</i> ^c	β	β
Alcohol use								
Total	50.0 (10.0)	46.0 (6.9)	56.5 (10.7)	1.17***	0.43***	0.84***	0.24***	0.01
Women	46.6 (7.7)	44.4 (5.4)	53.2 (9.7)	1.11***				
Men	54.5 (10.8)	49.2 (8.5)	58.1 (10.8)	0.92***				
Other drug use								
Total	50.0 (10.0)	46.9 (6.8)	54.7 (12.0)	0.82***	0.40***	0.37*	0.08*	-0.05
Women	48.5 (8.0)	46.4 (5.8)	54.8 (10.0)	1.00***				
Men	51.7 (11.6)	47.8 (8.9)	54.7 (12.8)	0.65***				
Externalizing								
Total	50.0 (10.0)	46.8 (8.1)	55.0 (10.6)	0.87***	0.29***	0.75***	0.22***	0.06
Women	46.7 (8.4)	45.3 (7.4)	51.1 (9.5)	0.69***				
Men	53.8 (10.3)	49.7 (8.6)	56.9 (10.5)	0.74***				
Internalizing								
Total	50.0 (10.0)	48.9 (8.9)	51.4 (11.0)	0.26***	0.27***	-0.08	-0.02	-0.18**
Women	50.3 (10.6)	49.0 (9.3)	54.3 (13.3)	0.46***				
Men	49.5 (8.9)	48.7 (8.2)	50.1 (9.3)	0.16				
Deviant peers								
Total	50.0 (10.0)	47.0 (8.8)	54.8 (9.7)	0.84***	0.32***	0.60***	0.17***	0.01
Women	47.5 (9.1)	45.9 (8.4)	52.4 (9.4)	0.73***				
Men	53.2 (10.1)	49.3 (9.3)	56.0 (9.7)	0.70***				
Romantic partner drug use								
Total	50.0 (10.0)	48.2 (9.4)	52.9 (10.1)	0.48***	0.38***	-0.36***	-0.25***	-0.09
Women	51.5 (10.1)	49.3 (9.5)	57.5 (10.0)	0.80***				
Men	48.1 (9.3)	44.8 (8.5)	50.5 (9.2)	0.64***				
Socio-economic status								
Total	50.0 (10.0)	50.2 (10.3)	49.7 (9.3)	-0.05	-0.05	0.20**	0.14**	-0.03
Women	49.1 (10.1)	49.3 (10.2)	48.4 (9.8)	-0.09				
Men	51.0 (9.6)	52.0 (10.3)	50.3 (8.9)	-0.16*				

AUD, Alcohol use disorder; s.d., standard deviation.

^a T-score means arranged by gender and AUD status (women: $n = 155$ AUD, 449 control; men: $n = 316$ AUD, 226 non-AUD) reflect mean level of consequences for each group at the age of 29 years.

^{b,c} Cohen's *d* effect sizes estimate the magnitude of the change in consequence factor coinciding with AUD in each gender (i.e. control women *versus* AUD women; d^b) and the difference in each mean-level consequence between men and women at the age of 29 years (i.e. all men *versus* all women; d^c).

^d The AUD effect estimates the level of consequences at the age of 29 years associated with lifetime AUD status compared with control status.

^e The gender effect estimates the difference in consequences at the age of 29 years for men compared with women.

^f The AUD × gender interaction effect tests if gender moderates the relationship between AUD and consequences at the age of 29 years.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

developing AUD more dramatically in women compared with men. Notably, these risks are not necessarily associated with concurrent AUD, as our AUD groups were derived using lifetime diagnoses by the age of 29 years. That is, the risk structure for AUD appears to emerge by adolescence, irrespective of the onset and chronicity of alcohol problems. The higher levels of adolescent risk exposure and young adult

consequences associated with AUD in women provide further evidence that it is a more severe and debilitating disorder in women than men. While psychosocial problems in young adulthood may be consequences of AUD they may, alternatively, also reflect the persistence of the high loading of risk present in adolescence for women with AUD. Studies aiming to understand the etiology of AUD in women would benefit from




Fig. 2. Cohen's d effect sizes for the main effect of alcohol use disorder within each gender at the age of 29 years. SES, Socio-economic status.

examining risk structure at even earlier ages to track how risk exposure relates to the onset and persistence of AUD and psychosocial problems.

Consistent with previous reports (Nolen-Hoeksema, 2004; Foster *et al.* 2014), internalizing distress exhibited a unique structural relationship with AUD in women compared with men, suggesting it may be a gender-specific risk factor for AUD. In women, increases in internalizing distress significantly increased the odds of AUD during adolescence and also had a significant relationship with AUD in young adulthood. In contrast, AUD had a near-zero association with internalizing distress in men at both ages, suggesting that it is neither a risk for nor a consequence of AUD in men. While previous literature has documented that women develop internalizing symptoms at a higher rate than men, our results suggest that, even after controlling for gender differences in their prevalence, internalizing symptoms probably play a role in the development of AUD in women but not men. The early emergence of internalizing symptoms in girls may potentiate alcohol use problems later in life through a developmental cascade. For example, symptoms of depression and anxiety that are more common in girls than boys during adolescence may be commonly associated with difficulties in school, work and peer relationships during puberty and catalyse alcohol use problems as a method of coping with negative emotions. As a result, alcohol use may exacerbate internalizing distress indirectly through its negative influence on psychosocial development.

The temporal relationship between internalizing distress and alcohol problems, however, remains unclear. Another possibility is that girls may engage in early and heavy use of alcohol independent of internalizing distress. Heavy alcohol use by adolescent girls has been shown to impair neurocognitive functioning

(Squeglia *et al.* 2010, 2011) and may increase isolation, disrupt social relationships and hinder academic engagement. A lack of stability in social support and academic success may substantially diminish girls' self-esteem and efficacy for coping with negative emotions in adaptive and prosocial ways (Lopez & DuBois, 2005). Subsequently, internalizing symptoms may emerge during young adulthood. Directly testing the temporal relationship between internalizing distress and alcohol problems using longitudinal methods will be vital for explicating this aspect of women's vulnerability to AUDs.

Overall, we provided evidence that AUD is a more severe disorder in women and that internalizing distress may play a gender-specific role in AUD symptoms among women. As only a few studies of gender differences have directly compared men and women in the association between multiple risk variables and the development of AUD at multiple time points, this research represents an important advancement of current research. However, these findings are limited in a number of ways. First, the same associations between each risk factor and AUD may not apply to more diverse samples of men and women. Replication among a more racially and ethnically diverse sample is needed to determine the generalizability of our findings. Second, the associations between risk factors and AUD may be better explained by a third variable that also varies by gender. Third, the use of multiple comparisons is not ideal but allowed for the comparison of the relative contributions of a number of risks to identify candidates for causal pathways that should be validated through future replication of this work and other investigations of individual risk factors. Finally, our analyses do not address the co-development between each risk exposure and AUD. Future work in these areas will be important

for determining the etiological role of these for AUD and its clinical features (i.e. onset and course).

Acknowledgements

This research was supported by National Institute on Drug Abuse Awards R37 DA005147 and R01 DA034606 and National Institute on Alcohol Abuse and Alcoholism Award R01AA 009367. K.T.F. was supported by National Institute on Alcohol Abuse and Alcoholism F31 Fellowship Award AA 023121. B.M.H. was supported by National Institute on Drug Abuse Award K01 DA 025868. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Declaration of Interest

None.

References

- Bemmels HR, Burt SA, Legrand LN, Iacono WG, McGue M** (2008). The heritability of life events: an adolescent twin and adoption study. *Twin Research and Human Genetics* **11**, 257–265.
- Blazei RW, Iacono WG, Krueger RF** (2006). Intergenerational transmission of antisocial behavior: how do kids become antisocial adults? *Applied and Preventive Psychology* **11**, 230–253.
- Blume SB, Russell M** (2001). Alcohol and substance abuse in obstetrics and gynecology practice. In *Psychological Aspects of Women's Health Care: The Interface Between Psychiatry and Obstetrics and Gynecology*, 2nd edn. (ed. N. L. Stotland and D. E. Steward), pp. 421–439. American Psychiatric Press: Washington, DC.
- Brady KT, Randall CL** (1999). Gender differences in substance use disorders. *Psychiatric Clinics of North America* **22**, 241.
- Caspi A, Henry B, McGee R, Moffitt T, Silva P** (1995). Temperamental origins of child and adolescent behavior problems from age 3 to age 15. *Child Development* **66**, 55–68.
- Caspi A, Moffitt TE, Newman DL, Silva PA** (1996). Behavioral observations at age 3 years predict adult psychiatric disorders: longitudinal evidence from a birth cohort. *Archives of General Psychiatry* **53**, 1033–1039.
- Cranford JA, Floyd FJ, Schulenberg JE, Zucker RA** (2011). Husbands' and wives' alcohol use disorders and marital interactions as longitudinal predictors of marital adjustment. *Journal of Abnormal Psychology* **120**, 210–222.
- Cutler SE, Nolen-Hoeksema S** (1991). Accounting for sex differences in depression through female victimization: childhood sexual abuse. *Sex Roles* **24**, 425–438.
- Dishion TJ, Patterson GR, Stoolmiller M, Skinner ML** (1991). Family, school, and behavioral antecedents to early adolescent involvement with antisocial peers. *Developmental Psychology* **27**, 172–180.
- Elkins I, King S, McGue M, Iacono W** (2006). Personality traits and the development of nicotine, alcohol, and illicit drug disorders: prospective links from adolescence to young adulthood. *Journal of Abnormal Psychology* **115**, 26–39.
- Elkins IJ, McGue M, Iacono WG** (2007). Prospective effects of attention-deficit/hyperactivity disorder, conduct disorder, and sex on adolescent substance use and abuse. *Archives of General Psychiatry* **64**, 1145–1152.
- Elkins IJ, McGue M, Iacono WG, Tellegen A** (1997). Genetic and environmental influence on parent–son relationships: evidence for increasing genetic influence during adolescence. *Developmental Psychology* **33**, 351–363.
- Elkins I, McGue M, Malone S, Iacono W** (2004). The effect of parental alcohol and drug disorders on adolescent personality. *American Journal of Psychiatry* **161**, 670–676.
- Foster KT, Hicks BM, Iacono WG, McGue M** (2014). Alcohol use disorder in women: risks and consequences of an adolescent onset and persistent course. *Psychology of Addictive Behaviors* **28**, 322–335.
- Granic I, Patterson GR** (2006). Toward a comprehensive model of antisocial development: a dynamic systems approach. *Psychological Review* **113**, 101–131.
- Hawkins JJD, Catalano RF, Miller JY** (1992). Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention. *Psychological Bulletin* **112**, 64–105.
- Hicks BM, DiRago AC, Iacono WG, McGue M** (2009). Gene–environment interplay in internalizing disorders: consistent findings across six environmental risk factors. *Journal of Child Psychology and Psychiatry* **50**, 1309–1317.
- Hicks BM, Durbin CE, Blonigen DM, Iacono WG, McGue M, Zucker R** (2010a). Alcohol dependence and personality change in young adulthood: effects of an adolescent onset, persistence, and desistence. *Alcoholism-Clinical and Experimental Research* **34**, 242A–242A.
- Hicks BM, Iacono WG, McGue M** (2010b). Consequences of an adolescent onset and persistent course of alcohol dependence in men: adolescent risk factors and adult outcomes. *Alcoholism-Clinical and Experimental Research* **34**, 819–833.
- Iacono WG, Carlson SR, Taylor J, Elkins IJ, McGue M** (1999). Behavioral disinhibition and the development of substance-use disorders: findings from the Minnesota Twin Family Study. *Development and Psychopathology* **11**, 869–900.
- Iacono WG, Malone SM, McGue M** (2008). Behavioral disinhibition and the development of early-onset addiction: common and specific influences. *Annual Review of Clinical Psychology* **4**, 12.1–12.24.
- Johnson W, McGue M, Iacono WG** (2006). Genetic and environmental influences on academic achievement trajectories during adolescence. *Developmental Psychology* **42**, 514–532.

- Kendler KS, Prescott C, Myers J, Neale M** (2003). The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Archives of General Psychiatry* **60**, 929–937.
- Kendler KS, Walters EE, Kessler RC** (1997). The prediction of length of major depression episodes: results from an epidemiological sample of female twins. *Psychological Medicine* **27**, 107–117.
- Keyes KM, Grant BF, Hasin DS** (2008). Evidence for a closing gender gap in alcohol use, abuse, and dependence in the United States population. *Drug and Alcohol Dependence* **93**, 21–29.
- Kilpatrick D, Acierno R, Saunders B, Resnick H, Best C, Schnurr P** (2000). Risk factors for adolescent substance abuse and dependence: data from a national sample. *Journal of Consulting and Clinical Psychology* **68**, 19–30.
- Klassen A, Wilsnack S** (1986). Sexual experience and drinking among women in a United-States national survey. *Archives of Sexual Behavior* **15**, 363–392.
- Krueger R, Hicks B, Patrick C, Carlson S, Iacono W, McGue M** (2002). Etiologic connections among substance dependence, antisocial behavior, and personality: modeling the externalizing spectrum. *Journal of Abnormal Psychology* **111**, 411–424.
- Labouvie E, McGee C** (1986). Relation of personality to alcohol and drug-use in adolescence. *Journal of Consulting and Clinical Psychology* **54**, 289–293.
- Lopez C, DuBois D** (2005). Peer victimization and rejection: investigation of an integrative model of effects on emotional, behavioral, and academic adjustment in early adolescence. *Journal of Clinical Child and Adolescent Psychology* **34**, 25–36.
- Masse L, Tremblay R** (1997). Behavior of boys in kindergarten and the onset of substance use during adolescence. *Archives of General Psychiatry* **54**, 62–68.
- McGue M, Iacono W** (2005). The association of early adolescent problem behavior with adult psychopathology. *American Journal of Psychiatry* **162**, 1118–1124.
- Moffitt TE, Caspi A, Rutter M, Silva PA** (2001). *Sex Differences in Antisocial Behaviour: Conduct Disorder, Delinquency, and Violence in the Dunedin Longitudinal Study*. Cambridge University Press: Cambridge.
- Niaura R, Nathan P, Frankenstein W, Shapiro A, Brick J** (1987). Gender differences in acute psychomotor, cognitive, and pharmacokinetic response to alcohol. *Addictive Behaviors* **12**, 345–356.
- Nixon S** (1994). Cognitive deficits in alcoholic women. *Alcohol Health and Research World* **18**, 228–232.
- Nolen-Hoeksema S** (2004). Gender differences in risk factors and consequences for alcohol use and problems. *Clinical Psychology Review* **24**, 981–1010.
- Nolen-Hoeksema S, Hilt L** (2006). Possible contributors to the gender differences in alcohol use and problems. *Journal of General Psychology* **133**, 357–374.
- Patterson GR, Yoerger K** (1997). A developmental model for late-onset delinquency. In *Motivation and Delinquency* (ed. D. W. Osgood), pp. 119–177. University of Nebraska Press: Lincoln, NE.
- Patterson GR, Yoerger K** (1999). Intraindividual growth in covert antisocial behaviour: a necessary precursor to chronic juvenile and adult arrests? *Criminal Behaviour and Mental Health* **9**, 24–38.
- Petry N, Kirby K, Kranzler H** (2002). Effects of gender and family history of alcohol dependence on a behavioral task of impulsivity in healthy subjects. *Journal of Studies on Alcohol* **63**, 83–90.
- Robins LM, Baber T, Cottler LB** (1987). *Composite International Diagnostic Interview: Expanded Substance Abuse Module*. L. M. Robins: St Louis, MO.
- Robins LM, Wing J, Wittchen H, Weler J, Babor T, Burke J, Farmer A, Jablenski A, Pickens R, Regier D, Sartorius N, Towle L** (1988). The Composite International Diagnostic Interview: an epidemiologic instrument suitable for use in conjunction with different diagnostic systems and in different cultures. *Archives of General Psychiatry* **45**, 1069–1077.
- Rutledge P, Sher K** (2001). Heavy drinking from the freshman year into early young adulthood: the roles of stress, tension-reduction drinking motives, gender and personality. *Journal of Studies on Alcohol* **62**, 457–466.
- Rutter M, Caspi A, Moffitt T** (2003). Using sex differences in psychopathology to study causal mechanisms: unifying issues and research strategies. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **44**, 1092–1115.
- Sannibale C, Hall W** (2001). Gender-related symptoms and correlates of alcohol dependence among men and women with a lifetime diagnosis of alcohol use disorders. *Drug and Alcohol Review* **20**, 369–383.
- Schuckit MA** (1994). Low-level of response to alcohol as a predictor of future alcoholism. *American Journal of Psychiatry* **151**, 184–189.
- Slutske WS, Heath AC, Madden PA, Bucholz KK, Statham DJ, Martin NG** (2002). Personality and the genetic risk for alcohol dependence. *Journal of Abnormal Psychology* **111**, 124–133.
- Squeglia LM, Schweinsburg AD, Pulido C, Tapert SF** (2011). Adolescent binge drinking linked to abnormal spatial working memory brain activation: differential gender effects. *Alcoholism-Clinical and Experimental Research* **35**, 1831–1841.
- Squeglia LM, Spadoni AD, Infante MA, Myers MG, Tapert SF** (2010). Initiating moderate to heavy alcohol use predicts changes in neuropsychological functioning for adolescent girls and boys. *Psychology of Addictive Behaviors* **23**, 715–722.
- Stein JA, Golding JM, Siegel JM, Burnam MA, Sorenson SB** (1988). Long-term psychological sequelae of child sexual abuse: the Los Angeles Epidemiologic Catchment Area study. In *Lasting Effects of Child Sexual Abuse* (ed. G. E. Wyatt and G. J. Powell), pp. 135–154. Sage: Thousand Oaks, CA.
- Tangney JP, Miller RS, Flicker L, Barlow DH** (1996). Are shame, guilt, and embarrassment distinct emotions? *Journal of Personality and Social Psychology* **70**, 1256–1269.
- Tellegen A, Waller NG** (2008). Exploring personality through test construction: development of the Multidimensional Personality Questionnaire. In *Handbook of Personality Theory*

- and Testing: Vol. II. Personality Measurement and Assessment (ed. G. J. Boyle, B. Matthews and D. H. Saklofske), pp. 261–292. Sage: Thousand Oaks, CA.
- Waldeck T, Miller L** (1997). Gender and impulsivity differences in licit substance use. *Journal of Substance Abuse* **9**, 269–275.
- Walden B, McGue M, Iacono WG, Burt SA, Elkins I** (2004). Identifying shared environmental contributions to early substance use: the respective roles of peers and parents. *Journal of Abnormal Psychology* **113**, 440–450.
- Widom C, Ireland T, Glynn P** (1995). Alcohol-abuse in abused and neglected children followed-up: are they at increased risk? *Journal of Studies on Alcohol* **56**, 207–217.
- Wilhelmsen K, Schuckit M, Smith T, Lee J, Segall S, Feiler H, Kalmijn J** (2003). The search for genes related to a low-level response to alcohol determined by alcohol challenges. *Alcoholism-Clinical and Experimental Research* **27**, 1041–1047.
- Wilsnack S, Vogeltanz N, Klassen A, Harris T** (1997). Childhood sexual abuse and women's substance abuse: national survey findings. *Journal of Studies on Alcohol* **58**, 264–271.
- Wong MM, Nigg JT, Zucker RA, Puttler LI, Fitzgerald HE, Jester JM, Glass JM, Adams K** (2006). Behavioral control and resiliency in the onset of alcohol and illicit drug use: a prospective study from preschool to adolescence. *Child Development* **77**, 1016–1033.
- Zucker RA** (2006). Alcohol use and the alcohol use disorders: a developmental-biopsychosocial systems formulation covering the life course. In *Developmental Psychopathology: Vol. 3. Risk, Disorder and Adaptation*, 2nd edn. (ed. D. Cicchetti and D. J. Cohen), pp. 620–656. Wiley: New York.