

# An artificial pool experiment in Antarctic sea ice: effects of sea ice melting on physical and biogeochemical components of pool water

DAIKI NOMURA<sup>1,2,3</sup>, DAISUKE SIMIZU<sup>4</sup>, SUCHANA CHAVANICH<sup>5</sup>, HIDEO SHINAGAWA<sup>6</sup> and MITSUO FUKUCHI<sup>1</sup>

<sup>1</sup>National Institute of Polar Research, 10-3 Midori-cho, Tachikawa, Tokyo 190-8501, Japan

<sup>2</sup>Japan Society for the Promotion of Science (JSPS), 6 Ichiban-cho, Chiyoda, Tokyo 102-8471, Japan

<sup>3</sup>Current address: Norwegian Polar Institute, Fram Centre, NO-9296 Tromsø, Norway

<sup>4</sup>Institute of Low Temperature Science, Hokkaido University, Kita-19, Nishi-8, Kita-ku, Sapporo, Hokkaido 060-0819, Japan

<sup>5</sup>Reef Biology Research Group, Department of Marine Science, Faculty of Science, Chulalongkorn University, 254 Phayathai Road, Pathumwan, Bangkok 10330, Thailand

<sup>6</sup>Shimoda Marine Research Center, University of Tsukuba, 5-10-1 Shimoda, Shizuoka 415-0025, Japan  
daiki.nomura@npolar.no

**Abstract:** We performed an artificial pool experiment in the Antarctic multi-year land-fast ice to examine and simulate the effect of sea ice melting on physical and biogeochemical components of the sea ice field. The input of snow and ice meltwater resulted in warmer, low salinity water at the surface of the pool and probably stratification of the less dense water. Current speed measurements also pointed to water stratification within the pool. Rapid phytoplankton growth in the pool resulted in drastic decreases in concentrations of dissolved inorganic carbon and nutrients ( $\text{NO}_3^-$  and  $\text{Si}(\text{OH})_4$ ) in the surface waters of the pool, particularly depleted for  $\text{NO}_3^-$ . There was high correlation between variations of dissolved inorganic carbon and nutrient concentrations, but the apparent uptake ratios of these components deviated from that generally applied to marine phytoplankton. The sequence of changes in the physical and biogeochemical components of the pool water suggests that the onset of rapid phytoplankton growth was closely related to the water stratification, which provided stable conditions for phytoplankton bloom even though the supply of nutrients from under-ice water would have declined.

Received 18 August 2011, accepted 26 February 2012, first published online 16 May 2012

**Key words:** dissolved inorganic carbon, nutrients, phytoplankton bloom, Southern Ocean, water stratification


## Introduction

Ongoing climate change will lead to increased sea ice melting and ice-free summer in the Arctic Ocean within the next few decades (e.g. Stroeve *et al.* 2007). As the ice melt season becomes longer, the large quantities of ice melt freshwater supplied to the ocean surface will influence the heat budget, water temperature, ice cover, and water stratification by reducing vertical mixing in the water column (e.g. Steele *et al.* 2010). In addition to changing the physical structure of the atmosphere-ocean system, these effects also influence biogeochemical processes and ecosystems through changes in vertical and horizontal transport of chemical substances and biological components, and in food web structure, thereby altering the processes affecting gas flux between the ocean and atmosphere (e.g. Arrigo *et al.* 1999, 2008, Bates & Mathis 2009). Arrigo *et al.* (1999) found that the phytoplankton community structure in the Ross Sea is related to the water stratification. Water stratification could favour the growth of diatoms, which would reduce nutrient

utilization efficiency and rates of the production in nutrient limited waters. Therefore, they indicated that the capacity of the biological community to draw down the atmospheric  $\text{CO}_2$  and transport it to the deep ocean could decrease with the increase of the water stratification (Arrigo *et al.* 1999).

There are clear relationships in spring and summer between the physical regime and biological productivity at the ice-edge at the seasonal timescale (Smith & Nelson 1985, Niebauer *et al.* 1990, Pérez *et al.* 1994, Jin *et al.* 2007). These studies showed that water stratification due to the supply of freshwater to the surface advances the ice-edge bloom that is seeded from ice algae in the meltwater supply. The nutrients in the surface water are eventually depleted through consumption by phytoplankton. Similar depletions are also found in the ice-gap layers of sea ice (Kattner *et al.* 2004, Papadimitriou *et al.* 2009), surface ponds (Kennedy *et al.* 2002), crack pools between the sea ice (Gleitz *et al.* 1996), and under-ice water (e.g. McMinn *et al.* 2000).

In this study, we conducted an artificial pool experiment in the Antarctic multi-year land-fast ice to examine and simulate the effect of sea ice melting on physical and



**Fig. 1.** Schematic illustration of the artificial pool used in this study showing the placement of temperature (T) and salinity (S) sensors and current meters.

biogeochemical components in the sea ice field. Our results provide a detailed description of the mechanisms and relationships between water stratification, biological productivity, and related biogeochemical components in the open water within the sea ice.

## Materials and methods

### Experimental setting

We constructed an artificial pool in the multi-year land-fast ice in Lützow-Holm Bay, East Antarctica (68°59'54"S, 39°37'03"E) from 22–24 December 2009 (Fig. 1). First, snow on the sea ice was removed with a snow shovel, and holes were drilled along the outline of the pool (1.5 m x 1.5 m) with a Jiffy ice drill (Model 30, Feldmann Engineering & Manufacturing Co. Inc, WI, USA). The sea ice was then crushed with a long chisel and removed with ice tongs and a net. The sea ice was 2.2 m thick. The depth of the snow on the ice surrounding the pool decreased from 0.4–0.1 m during the study period.


Three types of sensors were attached to a line supported by a tripod and passing through the water column at the centre of the pool. The sensors were used for long-term measurements of temperature, salinity and current speed (24 December 2009–17 January 2010). The temperatures at 1.0 m above the pool water surface and at water depths of 0.01, 0.05, and 0.10 m were measured by thermistor probe (TR-5106, T&D Corp, Nagano, Japan), and temperature data were stored in a data logger (RTR-52A, T&D Corp). The temperature and salinity at water depths of 0.3, 2.0 and 3.5 m were measured by memory-type conductivity-temperature (CT) sensor (Compact CT, JFE Advantech Co. Ltd, Hyogo, Japan). During 8–9 January 2010, the thermistor probe and the CT sensor were removed for a battery change and data download and then reinstalled. Current speeds at water depths of 0.6 and 3.0 m were measured by memory-type current meter (Compact EM, JFE Advantech Co. Ltd), which also had a temperature sensor. During 1–9 January 2010 the current meter could not record data because of a program error related to the transition from 2009 to 2010.

### Pool water sampling

Samples of the pool water were collected at depths of 0.01, 0.05, 0.3, 1.5 and 3.5 m on 24 and 26 December 2009 and on 3, 5, 11, and 15 January 2010 by using a Teflon tube and a diaphragm pump (EWP-01, As One Corporation, Osaka, Japan). Water was collected into a 12 ml glass screw-cap vial (Nichiden-Rika Glass Co. Ltd, Kobe, Japan) for measurement of salinity, a 120 ml amber glass vial (Maruemu Co. Ltd, Osaka, Japan) for measurement of dissolved inorganic carbon (DIC) concentration, a 10 ml polyethylene screw-cap vial (Eiken Chemical Co. Ltd, Tokyo, Japan) for measurement of inorganic nutrient concentrations, and into a 500 ml Nalgene polycarbonate bottle (Thermo Fisher Scientific Inc, Waltham MA, USA) for measurement of chlorophyll *a* (chl *a*) concentration. To stop biological activity 200  $\mu$ l of a saturated mercuric chloride solution was added to the DIC samples. Samples for measurement of salinity and DIC were stored in a refrigerator (+4°C), and those for measurement of nutrient concentrations in a freezer (-30°C) until further analysis. Samples for chl *a* measurement were filtered through 25 mm Whatman GF/F filters immediately after returning to the laboratory near the sampling station. Chlorophyll pigments on the filters were extracted in dimethylformamide (Suzuki & Ishimaru 1990) for 24 h at *c.* -80°C.


### Sample analysis

Salinity was measured with a salinity analyser (SAT-210, Toa Electronics Ltd, Tokyo, Japan). The salinity analyser was calibrated with standard seawater from the International Association for the Physical Science of the




**Fig. 2.** Temperature time series data. **a.** Air temperature, and **b.** depth profiles of pool water temperature obtained by moored sensors. Colour scale and contour lines indicate the temperature in °C. The contour plot was created using Ocean Data View software version 3.4.3.

Ocean (P series; Ocean Scientific International Ltd, Hampshire, UK). The standard deviation for salinities of 15 subsamples taken from a reference water sample ( $S = 10.00$ ) was 0.03.



**Fig. 3.** Time-series data recorded by moored sensors. **a.** Salinity, and **b.** current speed.



**Fig. 4.** Time-series of pool water biogeochemical components. **a.** Salinity, **b.** dissolved inorganic carbon (DIC), **c.**  $\text{NO}_3^-$ , **d.**  $\text{Si(OH)}_4$ , and **e.** chl *a*. The contour plots were created using Ocean Data View software version 3.4.3.

Dissolved inorganic carbon was determined by coulometry (Johnson *et al.* 1985). The DIC measurements were calibrated by using reference materials for seawater (Batch AG; KANSO Technos Co. Ltd, Osaka, Japan) traceable to the Certified Reference Material distributed by A.G. Dickson (Scripps Institution of Oceanography, La Jolla, CA, USA). The standard deviation for DICs of 20 subsamples from the reference material for seawater ( $\text{DIC} = 2035.4 \mu\text{mol kg}^{-1}$ ) was  $1.4 \mu\text{mol kg}^{-1}$ .

Nutrient concentrations were determined by using an auto-analyser system according to the Joint Global Ocean Flux Study (JGOFS) spectrophotometric method (Knap *et al.* 1996). Different systems were used for nitrate ( $\text{NO}_3^-$ ) (AACS III, Bran+Luebbe, Tokyo, Japan) and silicic acid ( $\text{Si(OH)}_4$ ) (Quattro-Marine 5ch, SEAL Analytical Ltd, Hampshire, UK). The analysers were calibrated with standard reference materials for nutrient analysis (Lots AW and BG; KANSO Technos Co. Ltd). The standard

**Table I.** Summary of all pool water data for each depth. The prefix “n-” indicates data normalised to a salinity of 34.6. n-DIC = normalized dissolved inorganic carbon, nd = no data.

Depth (m)	Date	Salinity	DIC ( $\mu\text{mol kg}^{-1}$ )	$\text{NO}_3^-$ ( $\mu\text{mol kg}^{-1}$ )	$\text{Si(OH)}_4$ ( $\mu\text{mol kg}^{-1}$ )	Chl <i>a</i> ( $\mu\text{g l}^{-1}$ )	n-DIC ( $\mu\text{mol kg}^{-1}$ )	n- $\text{NO}_3^-$ ( $\mu\text{mol kg}^{-1}$ )	n- $\text{Si(OH)}_4$ ( $\mu\text{mol kg}^{-1}$ )
0.01	24 December 2009	32.7	2029.2	25.2	55.4	0.0	2148.9	26.7	58.6
	26 December 2009	23.6	1461.8	13.1	39.3	0.1	2144.6	19.2	57.7
	3 January 2010	21.1	1214.5	1.3	12.9	0.4	1993.6	2.2	21.2
	5 January 2010	14.3	877.4	1.6	10.3	0.1	2125.2	4.0	24.9
	11 January 2010	5.1	281.4	0.2	2.9	0.1	1920.0	1.1	19.6
	15 January 2010	5.5	259.1	0.0	2.7	1.1	1639.9	0.0	16.9
0.05	24 December 2009	32.7	2045.6	24.6	57.2	0.0	2161.8	26.0	60.5
	26 December 2009	32.1	1963.5	19.2	55.2	0.0	2114.0	20.7	59.4
	3 January 2010	26.6	1513.1	4.4	23.4	0.3	1970.5	5.8	30.4
	5 January 2010	18.6	1041.9	1.7	11.1	0.2	1938.5	3.1	20.6
	11 January 2010	6.9	345.1	0.2	2.9	0.2	1730.0	0.8	14.5
	15 January 2010	9.2	389.5	0.1	2.5	3.4	1463.2	0.4	9.3
0.3	24 December 2009	32.8	2043.9	25.0	57.5	0.1	2155.4	26.3	60.6
	26 December 2009	32.7	1997.2	20.1	54.9	0.0	2112.8	21.3	58.1
	3 January 2010	30.8	1817.2	12.6	45.0	0.2	2038.8	14.1	50.5
	5 January 2010	29.4	1404.8	6.5	22.8	0.3	1656.1	7.7	26.9
	11 January 2010	13.7	654.8	1.2	5.6	0.3	1653.7	3.1	14.1
	15 January 2010	12.6	534.6	0.5	2.8	3.8	1468.4	1.4	7.7
1.5	24 December 2009	34.6	2159.3	26.2	63.7	0.1	2161.3	26.2	63.8
	26 December 2009	34.1	nd	24.6	58.7	0.0	nd	24.9	59.6
	3 January 2010	33.0	2000.6	18.7	56.5	0.3	2096.5	19.6	59.3
	5 January 2010	25.3	1642.5	9.6	28.9	0.1	2248.3	13.1	39.6
	11 January 2010	21.5	1043.6	1.8	9.5	0.2	1679.6	3.0	15.3
	15 January 2010	23.8	1129.9	1.3	8.2	0.5	1639.4	1.9	11.9
3.5	24 December 2009	34.6	2145.5	25.7	62.1	0.1	2146.4	25.7	62.2
	26 December 2009	34.6	2171.1	27.6	62.4	0.1	2170.9	27.6	62.4
	3 January 2010	34.5	2149.0	24.4	67.2	0.1	2155.3	24.4	67.4
	5 January 2010	34.6	2168.9	27.7	66.5	0.0	2172.0	27.7	66.6
	11 January 2010	34.5	2149.1	26.4	67.6	0.1	2153.2	26.4	67.7
	15 January 2010	34.5	2098.2	26.6	66.5	0.0	2102.2	26.7	66.6

deviation for nutrient concentrations in 20 subsamples from a reference seawater sample ( $10.0 \mu\text{mol kg}^{-1} \text{NO}_3^-$ ,  $120.1 \mu\text{mol kg}^{-1} \text{Si(OH)}_4$ ) was  $0.3 \mu\text{mol kg}^{-1}$  for  $\text{NO}_3^-$  and  $1.1 \mu\text{mol kg}^{-1}$  for  $\text{Si(OH)}_4$ .

Chlorophyll *a* concentrations were determined using a fluorometer (Model 10AU, Turner Designs Inc, Sunnyvale CA, USA) following methods described by Parsons *et al.* (1984). The fluorometer was calibrated before measurements by using calibration standards ( $0.28\text{--}282.3 \mu\text{g l}^{-1}$  chl *a*) made by stepwise dilution of a liquid chl *a* standard (Wako Pure Chemical Industries Ltd, Osaka, Japan) with *N,N*-dimethylformamide.

## Results

### Temperatures of air and pool water

Mean air temperatures throughout the study period were close to  $0^\circ\text{C}$  although there were daily variations (mean  $\pm$  SD,  $-0.3 \pm 2.2^\circ\text{C}$ ) (Fig. 2a). Water temperatures in the upper part of the pool (shallower than about 1.2 m) were below zero ( $-1.0 \pm 0.6^\circ\text{C}$ ) from the start of the study until 8 January 2010, and then increased to positive values

( $+0.9 \pm 1.3^\circ\text{C}$ ) throughout the rest of the study period (Fig. 2b). Water temperatures towards the bottom part of the pool (deeper than about 1.2 m) were below zero ( $-1.6 \pm 0.1^\circ\text{C}$ ) throughout the study period (Fig. 2b).


### Salinity and current speed

Salinity of the pool water at 0.3 m depth decreased drastically from a range of 30.8–33.6 on 24–31 December 2009 to 8.6–13.7 on 10–17 January 2010 (Fig. 3a). In contrast, salinity of the pool water at depths of 2.0 and 3.5 m was almost constant at  $33.7 \pm 0.3$ , except for 3 January and 8–12 January 2010. The salinity at 2.0 m decreased to 30.7 on 3 January and to 28.5 on 10 January 2010.

Current speed at 0.6 m depth was almost constant at  $0.7 \pm 0.1 \text{ cm s}^{-1}$  during the study period (Fig. 3b). In contrast, the current speed at 3.0 m varied from  $0.5\text{--}9.6 \text{ cm s}^{-1}$  during the study period with a mean of  $3.6 \text{ cm s}^{-1}$ .

### Salinity, DIC, nutrients and chl *a*

The results from analyses of bottle samples were used to create time series depth profiles of salinity, DIC,  $\text{NO}_3^-$ ,




**Fig. 5.** Time-series of pool water biogeochemical components normalized (“n-” prefix) to a salinity of 34.6. **a.** Normalized dissolved inorganic carbon (n-DIC), **b.**  $n\text{-NO}_3^-$ , and **c.**  $n\text{-Si(OH)}_4$ . The contour plots were created using Ocean Data View software version 3.4.3.

$\text{Si(OH)}_4$  and chl *a* concentrations (Fig. 4). Detailed values for each parameter are presented in Table I. Pool water salinity was almost constant vertically on 24 and 26 December 2009 ( $32.4 \pm 3.3$ ). By 15 January 2010 the salinity at the top of the pool (0.01 m) had decreased drastically to 5.5. Throughout the study period, the salinity at 3.5 m remained constant ( $34.6 \pm 0.0$ ) (Fig. 4a, Table I). The salinity variations detected in bottle samples were consistent with those recorded by the moored sensors (Fig. 3a).

Variations in DIC,  $\text{NO}_3^-$  and  $\text{Si(OH)}_4$  concentrations were similar to those of salinity, with drastic decreases in pool surface water (0.01 m) from  $> 2000 \mu\text{mol kg}^{-1}$  to  $259.1 \mu\text{mol kg}^{-1}$  for DIC, from  $> 25 \mu\text{mol kg}^{-1}$  to  $0.0 \mu\text{mol kg}^{-1}$  for  $\text{NO}_3^-$ , and from  $> 55 \mu\text{mol kg}^{-1}$  to  $2.7 \mu\text{mol kg}^{-1}$  for  $\text{Si(OH)}_4$  over the course of the study period (Fig. 4b–d, Table I). Concentrations at 3.5 m remained almost constant through the study period:  $2147.0 \pm 26.3 \mu\text{mol kg}^{-1}$  for DIC,  $26.4 \pm 1.2 \mu\text{mol kg}^{-1}$  for  $\text{NO}_3^-$ , and  $65.4 \pm 2.4 \mu\text{mol kg}^{-1}$  for  $\text{Si(OH)}_4$ .


The chl *a* concentrations were extremely low at all depths on 24 and 26 December 2009 ( $0.1 \pm 0.0 \mu\text{g l}^{-1}$ ). Chlorophyll *a* concentrations then increased drastically to  $3.4 \mu\text{g l}^{-1}$  at 0.05 m depth and to  $3.8 \mu\text{g l}^{-1}$  at 0.3 m depth on 15 January 2010 (Fig. 4e, Table I). These trends in chl *a* concentrations are opposite to those of salinity, DIC,  $\text{NO}_3^-$  and  $\text{Si(OH)}_4$  (Fig. 4). The colour of the pool water visibly changed from clear to green between the samplings on 5 and 11 January 2010.



**Fig. 6.** Relationships between salinity-normalized dissolved inorganic carbon (n-DIC) and nutrients in pool water and salinity. **a.** n-DIC vs salinity, **b.**  $n\text{-NO}_3^-$  vs salinity, and **c.**  $n\text{-Si(OH)}_4$  vs salinity. The “n-” indicates normalization to a salinity of 34.6. Solid and dashed lines indicate the mean and range, respectively, during the study period of the values in the pool at 3.5 m depth.

#### *Salinity-normalized DIC and nutrients*

To assess effects of dilution or concentration processes on the biogeochemical components of the pool water, we normalized each component to a salinity of 34.6, which was



**Fig. 7.** Relationships between salinity-normalized biogeochemical components in pool water. **a.** Normalized dissolved inorganic carbon (n-DIC) vs n-NO<sub>3</sub><sup>-</sup>, **b.** n-DIC vs n-Si(OH)<sub>4</sub>, and **c.** n-NO<sub>3</sub><sup>-</sup> vs n-Si(OH)<sub>4</sub>. The “n-” indicates normalization to a salinity of 34.6. The dashed lines indicate the Redfield-Brzezinski nutrient ratio.

the mean salinity of the pool water at a depth of 3.5 m during the sampling period (Table I). The trends in concentrations of normalized DIC (n-DIC) and nutrients (n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub>) (Fig. 5) were similar to those of DIC, NO<sub>3</sub><sup>-</sup> and Si(OH)<sub>4</sub> (Fig. 4), which decreased drastically in the pool surface water (shallower than 1.5 m): n-DIC decreased from > 2100 μmol kg<sup>-1</sup> to < 1500 μmol kg<sup>-1</sup>, n-NO<sub>3</sub><sup>-</sup> decreased from > 20 μmol kg<sup>-1</sup> to almost zero and n-Si(OH)<sub>4</sub> decreased from > 60 μmol kg<sup>-1</sup> to < 10 μmol kg<sup>-1</sup> over the study period (Fig. 5, Table I). Throughout the study period, concentrations of n-DIC, n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub> were almost constant at 3.5 m depth (Fig. 5, Table I).

We determined the relationships between the concentration of each component and the pool water salinity (Fig. 6). Generally, n-DIC, n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub> concentrations were lower than the mean value of the pool water at the depth of 3.5 m, particularly at lower salinities (Fig. 6). Lowest n-DIC, n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub> concentrations correspond to decreases from the mean pool water values at 3.5 m of 686.8 μmol kg<sup>-1</sup> for n-DIC, 26.4 μmol kg<sup>-1</sup> for n-NO<sub>3</sub><sup>-</sup>, and 57.8 μmol kg<sup>-1</sup> for n-Si(OH)<sub>4</sub>, or 31.9%, 100.0% and 88.2%, respectively.

We also compared the relationships between each pair of components in the pool water (Fig. 7). There were high degrees of correlation for each relationship: n-NO<sub>3</sub><sup>-</sup> vs n-DIC,  $r^2 = 0.63$ ,  $P < 0.001$ ; n-Si(OH)<sub>4</sub> vs n-DIC,  $r^2 = 0.73$ ,  $P < 0.001$ ; and n-Si(OH)<sub>4</sub> vs n-NO<sub>3</sub><sup>-</sup>,  $r^2 = 0.95$ ,

$P < 0.001$ . However, for n-NO<sub>3</sub><sup>-</sup> vs n-DIC and n-Si(OH)<sub>4</sub> vs n-DIC, the data plotted for the final stage of the study period (11 and 15 January 2010) deviated widely from the line of Redfield-Brzezinski nutrient ratio for marine phytoplankton (C:N:Si:P = 105:16:15:1; Brzezinski 1985) at the low n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub> (Fig. 7a & b). For n-Si(OH)<sub>4</sub> vs n-NO<sub>3</sub><sup>-</sup>, most of the data plotted deviated from the line of Redfield-Brzezinski ratio (Fig. 7c).

## Discussion and conclusions

The abrupt increase of temperature and decrease of salinity observed during the study period, particularly in the upper part of the pool (Figs 2b, 3a & 4a) are attributed to the input of meltwater from the snow around the pool and the melting of sea ice because of the input of solar radiation to the pool water. During the study period, the depth of snow surrounding the pool decreased drastically, suggesting that the snow meltwater was supplied to the pool mainly from the bottom of the snow layer.

The influx of higher temperature and lower salinity water induced stratification, with the less dense water in the surface layer of the pool (Figs 2b, 3a & 4a). The water stratification was enhanced by the greatly reduced current speeds at the surface of the pool. The current speed at the top of the pool (mean 0.7 cm s<sup>-1</sup> at 0.6 m depth) was much lower than that in the under-ice water below the pool (mean 3.6 cm s<sup>-1</sup> at 3.0 m depth) (Fig. 3b).

Shirasawa & Leppäranta (2003) measured similar current speeds ( $2\text{--}3\text{ cm s}^{-1}$ ) under the ice in a semi-enclosed lagoon in the southern part of the Okhotsk Sea. They indicated that the ice cover reduced the wind driven mixing at the water surface. Although the air temperature occasionally fell below  $-5^{\circ}\text{C}$  during the second half of the study period (Fig. 2a), the surface layer of the pool remained at temperatures that were relatively warmer (Fig. 2b). These results suggest that no vertical mixing was induced in the pool through cooling of the pool surface water.

Concentrations of the biogeochemical components (DIC,  $\text{NO}_3^-$  and  $\text{Si(OH)}_4$ ) decreased drastically in the top layer of the pool, tracking the variation of the salinity (Fig. 4, Table I). At first glance, these results suggest that these biogeochemical components were diluted by the freshwater supply from the outside of the pool. However, even after these components were normalized to salinity, we observed decreases in n-DIC, n- $\text{NO}_3^-$  and n- $\text{Si(OH)}_4$  with salinity and time in the surface layer of the pool (Figs 5 & 6). In contrast to the variations of biogeochemical components, chl *a* concentrations increased in the surface layer with time (Fig. 4e). These results clearly suggest that the DIC and nutrients were consumed during photosynthesis by the phytoplankton at the surface of the pool.

A drastic increase in the maximum chl *a* concentration was observed at the end of the experiment (15 January 2010, Fig. 4e), indicating that it took about three weeks for a phytoplankton bloom to occur in the pool even though the pool water was exposed to sunlight immediately after the onset of the experiment (24 December 2009). These results suggest that the timing of the phytoplankton bloom in the pool was affected by the degree of water column stratification in the pool. This is a situation similar to those at the ice-edge during the ice melting season. Ice-edge blooms are closely related to the water stratification, and the blooms are seeded from ice algae, which are released from sea ice with ice meltwater (Smith & Nelson 1985, Niebauer *et al.* 1990, Pérez *et al.* 1994, Jin *et al.* 2007).

Despite the favourable conditions provided for phytoplankton blooms by the water stratification, the stratification eventually resulted in nutrient depletion, particularly of  $\text{NO}_3^-$ , in the upper layer of the pool where salinity was low (Fig. 6b, Table I). The supply of nutrients from under the pool declined because of the water stratification in the pool, and the bloom presumably would have ended after our experiment. Therefore, it appears that phytoplankton succession in surface waters of semi-enclosed marine systems is strongly affected by the balance between the water stratification and nutrient supply from the under-ice water. Nutrient depletion has been observed at the ice-edge (e.g. Niebauer *et al.* 1990) and in the relatively closed sea ice systems in the ice-gap layer at the top of sea ice (e.g. Kattner *et al.* 2004, Papadimitriou *et al.* 2009), surface ponds (Kennedy *et al.* 2002) and crack pools between the sea ice (Gleitz *et al.* 1996).

Relationships between n-DIC and n- $\text{NO}_3^-$  (Fig. 7a) and n- $\text{Si(OH)}_4$  (Fig. 7b) in this study indicated that the data plotted for the final stage of the study period (11 and 15 January 2010) were below the line of Redfield-Brzezinski ratio at the low n- $\text{NO}_3^-$  and n- $\text{Si(OH)}_4$  (Fig. 7a & b). This suggests that the DIC consumption relative to  $\text{NO}_3^-$  and  $\text{Si(OH)}_4$  was higher than that expected for general phytoplankton growth. In previous studies, over-consumption of carbon was found in blooms in a mesocosm (Engel *et al.* 2002), stratified waters (Arrigo *et al.* 1999) and under various sea ice conditions (Gleitz *et al.* 1996, Günther *et al.* 1999, Papadimitriou *et al.* 2009). These carbon over-consumptions were attributed to the production of the carbohydrates and lipids (Fahl & Kattner 1993, Gleitz *et al.* 1996, Schartau *et al.* 2007, Papadimitriou *et al.* 2009) and to the substantial utilization by phytoplankton of ammonium and urea in addition to nitrate (e.g. Papadimitriou *et al.* 2009). In this study, there were no data collected that would explain the main reasons for the over-consumption of carbon.

There are other factors that might help explain the deviations from the Redfield-Brzezinski ratio observed in this study (Fig. 7a & b). One factor is the non-conservative nature of DIC in the pool water during the study period. Gas exchange between pool water and the atmosphere would change the DIC concentration in the pool water. The blooming of phytoplankton during the study period (Fig. 4e) suggests that the partial pressure of  $\text{CO}_2$  ( $p\text{CO}_2$ ) in the pool water became lower than that of the atmosphere. Therefore, atmospheric  $\text{CO}_2$  would have been absorbed by the pool water, increasing the DIC concentration. In addition, calcium carbonate ( $\text{CaCO}_3$ ) precipitation or dissolution would change the DIC concentration in the pool water. In the low temperature, high salinity conditions of sea ice,  $\text{CaCO}_3$  crystals in the form of ikaite ( $\text{CaCO}_3 \cdot 6\text{H}_2\text{O}$ ) have been found in the Arctic and Antarctic (Dieckmann *et al.* 2008, 2010). The formation of this  $\text{CaCO}_3$  releases  $\text{CO}_2$  into the sea ice, whereas the dissolution of this  $\text{CaCO}_3$  consumes  $\text{CO}_2$  and decreases the  $p\text{CO}_2$  in water (e.g. Papadimitriou *et al.* 2004). This latter process usually occurs during the ice melting season. Therefore, in this study, gas exchange and  $\text{CaCO}_3$  dissolution would have tended to increase the DIC concentration in the pool water. Although these processes could potentially alter the C:N:Si ratios away from the Redfield-Brzezinski ratio (Fig. 7a & b), processes other than gas exchange and carbonate dissolution were responsible for carbon over-consumption in this study.

It is said that the phytoplankton community structure in the Ross Sea is related to the water stratification (Arrigo *et al.* 1999). Water stratification could favour the growth of diatoms, which would cause the over-consumption of carbon. In addition, the reduction of nutrient utilization efficiency and rates of the production occurred in the diatoms dominated waters (Arrigo *et al.* 1999). Therefore,

they indicated that the capacity of the biological community to draw down the atmospheric CO<sub>2</sub> and transport it to the deep ocean could decrease with the increase of the water stratification (Arrigo *et al.* 1999).

In order to estimate the rate of production by the phytoplankton in the pool, changes in depth-integrated n-DIC were evaluated during the study period. The depth-integrated n-DIC decreased with time, and the result indicated that the production rate was 0.84 g C m<sup>-2</sup> day<sup>-1</sup>. Same order of the production rate (0.93 g C m<sup>-2</sup> day<sup>-1</sup>) was observed in the Ross Sea when the phytoplankton community was dominated by diatoms (Arrigo *et al.* 1999). This result and the over-consumption of carbon found in this study suggest that diatoms were dominated within the stratified pool water.

Most of the data plotted deviated from the line of Redfield-Brzezinski ratio for n-Si(OH)<sub>4</sub> vs n-NO<sub>3</sub> (Fig. 7c), but n-Si(OH)<sub>4</sub> and n-NO<sub>3</sub> were highly correlated ( $r^2 = 0.95$ ,  $P < 0.001$ ). The slope of the regression line for the relationship between n-NO<sub>3</sub><sup>-</sup> and n-Si(OH)<sub>4</sub> was 0.5. This value was smaller than the Redfield-Brzezinski ratio (N:Si = 1.1). It is said that the N:Si ratio varies considerably, particularly in polar environments (Jennings *et al.* 1984, Arrigo & Sullivan 1994). Values of 0.4 have been reported for the Southern Ocean (Jennings *et al.* 1984) and for sea ice algae growing in McMurdo Sound, Antarctica, where a value of 0.4 was used for modelling microalgal growth (Arrigo & Sullivan 1994). These values are in good agreement with the slope of 0.5 obtained in this study. However, the active uptake of Si by diatoms in sea ice environments is governed by the many factors that control growth, including temperature, light conditions and the nutrient composition of the environment (e.g. Thomas & Papadimitriou 2003).

Based on an artificial pool experiment in the Antarctic multi-year land-fast ice, we could examine and simulate the effect of sea ice melting on physical and biogeochemical components. Our results provide a detailed description of the mechanisms and relationships between water stratification, biological productivity, and related biogeochemical components in the open water within the sea ice and ice-edge regions in the polar seas.

### Acknowledgements

We express heartfelt thanks to Dr S. Kudoh for his help in making the artificial pool. We thank Dr Y.W. Watanabe for providing the instrument for DIC analysis, and to Prof J. Kanda, Ms S. Yasui and Ms C. Oouchida for nutrient analyses. We also thank Dr P. Assmy, Dr S. Koga, Mr H. Shimoda, Ms S. Youkongkaew and all of the members of the 51st Japanese Antarctic Research Expedition (JARE-51) for their comment and support in the field. We also thank National Research University Project of CHE Fund (CC1043A). This paper was supported by funds from the

National Institute of Polar Research for English proofreading. The constructive comments of the reviewers are also gratefully acknowledged.

### References

- ARRIGO, K.R. & SULLIVAN, C.W. 1994. A high-resolution bio-optical model of microalgal growth: tests using sea ice algal community time series data. *Limnology and Oceanography*, **39**, 609–631.
- ARRIGO, K.R., ROBINSON, D.H., WORTHEN, D.L., DUNBAR, R.B., DITULLIO, G.R., VANWOERT, M. & LIZOTTE, M.P. 1999. Phytoplankton community structure and the drawdown of nutrients and CO<sub>2</sub> in the Southern Ocean. *Science*, **283**, 365–367.
- ARRIGO, K.R., VAN DIJKEN, G.L. & PABI, S. 2008. The impact of a shrinking Arctic ice cover on marine primary production. *Geophysical Research Letters*, 10.1029/2008GL035028.
- BATES, N.R. & MATHIS, J.T. 2009. The Arctic Ocean marine carbon cycle: evaluation of air-sea CO<sub>2</sub> exchanges, ocean acidification impacts and potential feedbacks. *Biogeosciences*, **6**, 2433–2459.
- BRZEZINSKI, M.A. 1985. The Si:C:N ratio of marine diatoms: interspecific variability and the effect of some environmental variables. *Journal of Phycology*, **21**, 347–357.
- DIECKMANN, G.S., NEHRKE, G., PAPADIMITRIOU, S., GÖTTLICHER, J., STEININGER, R., KENNEDY, H., WOLF-GLADROW, D. & THOMAS, D.N. 2008. Calcium carbonate as ikaite crystals in Antarctic sea ice. *Geophysical Research Letters*, 10.1029/2008GL033540.
- DIECKMANN, G.S., NEHRKE, G., UHLIG, C., GÖTTLICHER, J., GERLAND, S., GRANSKOG, M.A. & THOMAS, D.N. 2010. Brief Communication: Ikaite (CaCO<sub>3</sub>·6H<sub>2</sub>O) discovered in Arctic sea ice. *The Cryosphere*, **4**, 227–230.
- ENGEL, A., GOLDTHWAIT, S., PASSOW, U. & ALLDREDGE, A.L. 2002. Temporal decoupling of carbon and nitrogen dynamics in a mesocosm diatom bloom. *Limnology and Oceanography*, **47**, 753–761.
- FAHL, K. & KATTNER, G. 1993. Lipid content and fatty acid composition of algal communities in sea ice and water from the Weddell Sea (Antarctica). *Polar Biology*, **13**, 405–409.
- GLEITZ, M., GROSSMANN, S., SCHAREK, R. & SMETACEK, V. 1996. Ecology of diatom and bacterial assemblages in water associated with melting summer sea ice in the Weddell Sea, Antarctica. *Antarctic Science*, **8**, 135–146.
- GÜNTHER, S., GLEITZ, M. & DIECKMANN, G.S. 1999. Biogeochemistry of Antarctic sea ice: a case study on platelet ice layers at Drescher Inlet, Weddell Sea. *Marine Ecology Progress Series*, **177**, 1–13.
- JENNINGS, J.C., GORDON, L.I. & NELSON, D.M. 1984. Nutrient depletion indicates high productivity in the Weddell Sea. *Nature*, **309**, 51–54.
- JIN, M.B., DEAL, C., WANG, J., ALEXANDER, V., GRADINGER, R., SAITOH, S., IIDA, T., WAN, Z.W. & STABENO, P. 2007. Ice-associated phytoplankton blooms in the southeastern Bering Sea. *Geophysical Research Letters*, 10.1029/2006GL028849.
- JOHNSON, K.M., KING, A.E. & SIEBURTH, G.M. 1985. Coulometric TCO<sub>2</sub> analyses for marine studies: an introduction. *Marine Chemistry*, **16**, 61–82.
- KATTNER, G., THOMAS, D.N., HAAS, C., KENNEDY, H. & DIECKMANN, G.S. 2004. Surface ice and gap layers in Antarctic sea ice: highly productive habitats. *Marine Ecology Progress Series*, **277**, 1–12.
- KENNEDY, H., THOMAS, D.N., KATTNER, G., HAAS, C. & DIECKMANN, G.S. 2002. Particulate organic matter in Antarctic summer sea ice: concentration and stable isotopic composition. *Marine Ecology Progress Series*, **238**, 1–13.
- KNAP, A., MICHAELS, A., CLOSE, A., DUCKLOW, H. & DICKSON, A., eds. 1996. *Protocols for the Joint Global Ocean Flux Study (JGOFS) core measurements. JGOFS Report*, **19**. Paris: UNESCO, 170 pp.
- MCMINN, A., BLEAKLEY, N., ROBERTS, D. & TRENNERY, L. 2000. Effect of permanent sea ice cover and different nutrient regimes on the phytoplankton succession of fjords of Vestfold Hills Oasis, eastern Antarctica. *Journal of Plankton Research*, **22**, 287–303.



- NIEBAUER, H.J., ALEXANDER, V. & HENRICH, S. 1990. Physical and biological oceanographic interaction in the spring bloom at the Bering Sea marginal ice-edge zone. *Journal of Geophysical Research*, **95**, 22 229–22 242.
- PAPADIMITRIOU, S., KENNEDY, H., KATTNER, G., DIECKMANN, G.S. & THOMAS, D.N. 2004. Experimental evidence for carbonate precipitation and CO<sub>2</sub> degassing during sea ice formation. *Geochimica et Cosmochimica Acta*, **68**, 1749–1761.
- PAPADIMITRIOU, S., THOMAS, D.N., KENNEDY, H., KUOSA, H. & DIECKMANN, G.S. 2009. Inorganic carbon removal and isotopic enrichment in Antarctic sea ice gap layers during early austral summer. *Marine Ecology Progress Series*, **386**, 15–27.
- PARSONS, T.R., TAKAHASHI, M. & HARGRAVE, B. 1984. *Biological oceanographic processes*, 3rd ed. Oxford: Pergamon Press, 330 pp.
- PÉREZ, F.F., FIGUEIRAS, F.G. & RÍOS, A.F. 1994. Nutrient depletion and particulate matter near the ice-edge in the Weddell Sea. *Marine Ecology Progress Series*, **112**, 143–153.
- SCHARTAU, M., ENGEL, A., SCHRÖTER, J., THOMAS, S., VÖLKER, C. & WOLF-GLADROW, D. 2007. Modelling carbon overconsumption and the formation of extracellular particulate organic carbon. *Biogeosciences*, **4**, 433–454.
- SHIRASAWA, K. & LEPPÄRANTA, M. 2003. Hydrometeorological and sea ice conditions at Saroma-ko lagoon, Hokkaido, Japan. *Report Series in Geophysics*, **46**. Helsinki: University of Helsinki, Division of Geophysics, 161–168.
- SMITH, W.O. & NELSON, D.M. 1985. Phytoplankton bloom produced by a receding ice-edge in the Ross Sea: spatial coherence with the density field. *Science*, **227**, 163–166.
- STEELE, M., ZHANG, J. & ERMOLD, W. 2010. Mechanisms of summertime upper Arctic Ocean warming and the effect on sea ice melt. *Journal of Geophysical Research*, 10.1029/2009JC005849.
- STROEVE, J., HOLLAND, M.M., MEIER, W., SCAMBOS, T. & SERREZE, M. 2007. Arctic sea ice decline: faster than forecast. *Geophysical Research Letters*, 10.1029/2007GL029703.
- SUZUKI, R. & ISHIMARU, T. 1990. An improved method for the determination of phytoplankton chlorophyll using N,N-dimethylformamide. *Journal of the Oceanographical Society of Japan*, **46**, 190–194.
- THOMAS, D.N. & PAPADIMITRIOU, S. 2003. Biogeochemistry of sea ice. In THOMAS, D.N. & DIECKMANN, G.S., eds. *Sea ice: an introduction to its physics, chemistry, biology and geology*. Oxford: Blackwell Science, 267–302.