



Tree densities in sparse graph classes

Tony Huynh and David R. Wood

Abstract. What is the maximum number of copies of a fixed forest T in an n -vertex graph in a graph class \mathcal{G} as $n \rightarrow \infty$? We answer this question for a variety of sparse graph classes \mathcal{G} . In particular, we show that the answer is $\Theta(n^{\alpha_d(T)})$ where $\alpha_d(T)$ is the size of the largest stable set in the subforest of T induced by the vertices of degree at most d , for some integer d that depends on \mathcal{G} . For example, when \mathcal{G} is the class of k -degenerate graphs then $d = k$; when \mathcal{G} is the class of graphs containing no $K_{s,t}$ -minor ($t \geq s$) then $d = s - 1$; and when \mathcal{G} is the class of k -planar graphs then $d = 2$. All these results are in fact consequences of a single lemma in terms of a finite set of excluded subgraphs.

1 Introduction

Many classical theorems in extremal graph theory concern the maximum number of copies of a fixed graph H in an n -vertex graph¹ in some class \mathcal{G} . Here, a *copy* means a subgraph isomorphic to H . For example, Turán's Theorem determines the maximum number of copies of K_2 (that is, edges) in an n -vertex K_t -free graph [94]. More generally, Zykov's Theorem determines the maximum number of copies of a given complete graph K_s in an n -vertex K_t -free graph [99]. The excluded graph need not be complete. The Erdős–Stone Theorem [35] determines, for every nonbipartite graph X , the asymptotic maximum number of copies of K_2 in an n -vertex graph with no X -subgraph. Analogues of the Erdős–Stone Theorem for the number of (induced) copies of a given graph within a graph class defined by an excluded (induced) subgraph have recently been widely studied [4–6, 36, 48–50, 55, 67, 72, 76].

For graphs H and G , let $C(H, G)$ be the number of copies of H in G . For a graph class \mathcal{G} , let

$$C(H, \mathcal{G}, n) := \max_{G \in \mathcal{G}, |V(G)|=n} C(H, G).$$

This paper determines the asymptotic behavior of $C(T, \mathcal{G}, n)$ as $n \rightarrow \infty$ for various sparse graph classes \mathcal{G} and for an arbitrary fixed forest T . In particular, we show that $C(T, \mathcal{G}, n) \in \Theta(n^k)$ for some k depending on T and \mathcal{G} .

Received by the editors December 13, 2020; revised May 11, 2021; accepted June 9, 2021.

Published online on Cambridge Core June 29, 2021.

Research supported by the Australian Research Council.

AMS subject classification: 90C35, 05C05, 05C83.

Keywords: Extremal combinatorics, trees, graph minors, sparsity.

¹All graphs in this paper are undirected, finite, and simple, unless stated otherwise. Let $\mathbb{N} := \{1, 2, \dots\}$ and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For $a, b \in \mathbb{N}_0$, let $[a, b] := \{a, a + 1, \dots, b\}$ and $[b] := [1, b]$.



It turns out that k depends on the size of particular stable sets in T . A set S of vertices in a graph G is *stable* if no two vertices in S are adjacent. Let $\alpha(G)$ be the size of a largest stable set in G . For a graph G and $s \in \mathbb{N}_0$, let

$$\alpha_s(G) := \alpha(G[\{v \in V(G) : \deg_G(v) \leq s\}]).$$

Note that for a forest T (indeed any bipartite graph), $\alpha_s(T)$ can be computed in polynomial time. See [10, 12, 13] for bounds on the size of bounded degree stable sets in forests, planar graphs, and other classes.

The first sparse class we consider are the graphs of given degeneracy.²

Theorem 1.1 *Fix $k \in \mathbb{N}$ and let \mathcal{D}_k be the class of k -degenerate graphs. Then for every fixed forest T ,*

$$C(T, \mathcal{D}_k, n) \in \Theta(n^{\alpha_k(T)}).$$

Our second main theorem determines $C(T, \mathcal{G}, n)$ for many minor-closed classes.^{3,4} Several examples of this result are given in Section 4.

Theorem 1.2 *Fix $s, t \in \mathbb{N}$ and let \mathcal{G} be a minor-closed class such that every graph with treewidth at most s is in \mathcal{G} and $K_{s+1,t} \notin \mathcal{G}$. Then for every fixed forest T ,*

$$C(T, \mathcal{G}, n) \in \Theta(n^{\alpha_s(T)}).$$

The lower bounds in Theorems 1.1 and 1.2 are proved via the same construction given in Section 2. The upper bounds in Theorems 1.1 and 1.2 are proved in Section 3. We in fact prove a stronger result (Lemma 3.3) that shows that for any fixed forest T and $s \in \mathbb{N}$ there is a particular finite set \mathcal{F} such that $C(T, G) \in O(n^{\alpha_s(T)})$ for every n -vertex graph G with $O(n)$ edges and containing no subgraph in \mathcal{F} . This result is applied in Section 5 to determine $C(T, \mathcal{G}, n)$ for various non-minor-closed classes \mathcal{G} . For example, we show a $\Theta(n^{\alpha_s(T)})$ bound for graphs that can be drawn in a fixed surface with a bounded average number of crossings per edge, which matches the known bound with no crossings.

1.1 Related results

Before continuing we mention related results from the literature. For a fixed complete graph K_s , $C(K_s, \mathcal{G}, n)$ has been extensively studied for various graph classes \mathcal{G} including: graphs of given maximum degree [2, 14, 20, 21, 31, 46, 47, 60, 97]; graphs with a given number of edges, or more generally, a given number of smaller complete graphs

²A graph G is *k-degenerate* if every subgraph of G has minimum degree at most k .

³A graph H is a *minor* of a graph G if a graph isomorphic to H can be obtained from a subgraph of G by contracting edges. A graph class \mathcal{G} is *minor-closed* if some graph is not in \mathcal{G} , and for every graph $G \in \mathcal{G}$, every minor of G is also in \mathcal{G} .

⁴A *tree decomposition* of a graph G is given by a tree T whose nodes index a collection $(B_x \subseteq V(G) : x \in V(T))$ of sets of vertices in G called *bags*, such that: (T1) for every edge vw of G , some bag B_x contains both v and w , and (T2) for every vertex v of G , the set $\{x \in V(T) : v \in B_x\}$ induces a nonempty (connected) subtree of T . The *width* of such a tree decomposition is $\max\{|B_x| - 1 : x \in V(T)\}$. The *treewidth* of a graph G , denoted by $\text{tw}(G)$, is the minimum width of a tree decompositions of G . See [57, 85] for surveys on treewidth. For each $s \in \mathbb{N}$ the class of graphs with treewidth at most s is minor-closed.

[19, 29, 30, 38, 39, 45, 58, 61, 62, 82]; graphs without long cycles [71]; planar graphs [69, 81, 97]; graphs with given Euler genus [27, 59]; and graphs excluding a fixed minor or subdivision [42–44, 66, 79, 84].

When \mathcal{J} is the class of planar graphs, $C(H, \mathcal{J}, n)$ has been determined for various graphs H including: complete bipartite graphs [3], planar triangulations without nonfacial triangles [3], triangles [56, 69, 70, 97], four-cycles [1, 69], five-cycles [53], four-vertex paths [54], and four-vertex complete graphs [3, 97]. $C(H, \mathcal{J}, n)$ has also been studied for more general planar graphs H . Perles (see [3]) conjectured that if H is a fixed three-connected planar graph, then $C(H, \mathbb{S}_0, n) \in \Theta(n)$. Perles noted the converse: If H is planar, not three-connected and $|V(H)| \geq 4$, then $C(H, \mathbb{S}_0, n) \in \Omega(n^2)$. Perles’ conjecture was proved by Wormald [98] and independently by Eppstein [32]. Recently, Huynh et al. [59] extended these results to all surfaces and all graphs H (see Section 4).

Finally, we mention a result of Nešetřil and Ossona de Mendez [76], who proved that for every infinite nowhere dense hereditary graph class \mathcal{G} and for every fixed graph F , the maximum, taken over all n -vertex graphs $G \in \mathcal{G}$, of the number of induced subgraphs of G isomorphic to F is $\Omega(n^\beta)$ and $O(n^{\beta+o(1)})$ for some integer $\beta \leq \alpha(F)$. Our results (when F is a forest and \mathcal{G} is one of the classes that we consider) imply this upper bound (since the number of induced copies of T in G is at most $C(T, G)$). Moreover, our bounds are often more precise since $\alpha_s(T)$ can be significantly less than $\alpha(T)$.

2 Lower bound

Lemma 2.1 *Fix $s \in \mathbb{N}$ and let T be a fixed forest with $\alpha_s(T) = k$. Then there exists a constant $c_{2.1}(k) := (2k)^{-k}$ such that for all sufficiently large $n \in \mathbb{N}$, there exists a graph G with $|V(G)| \leq n$ and $\text{tw}(G) \leq s$ and $C(T, G) \geq c_{2.1}(k)n^k$.*

Proof Let S be a maximum stable set in $T[\{v \in V(T) : \deg_T(v) \leq s\}]$ with $|S| = k$. Let $m := \lfloor \frac{n-|V(T)|}{k} \rfloor$. Let G be the graph obtained from T as follows: for each vertex v in S add to G a set C_v of m vertices, such that $N_G(x) := N_T(v)$ for each vertex $x \in C_v$. Observe that G has at most n vertices. Each choice of one vertex $x \in C_v$ (for each $v \in S$), along with the vertices in $V(T) \setminus S$, induces a copy of T . Thus $C(T, G) \geq m^k$, which is at least $c_{2.1}(k)n^k$ for $n \geq 2|V(T)| + 2k$.

We now show $\text{tw}(G) \leq s$. Let T_1 be a connected component of T and G_1 be the corresponding connected component of G . Since the treewidth of a graph equals the maximum treewidth of its components, it suffices to show $\text{tw}(G_1) \leq s$. We may assume $|V(T_1)| \geq 2$, as otherwise $\text{tw}(G_1) = 0$. Let T'_1 be the tree obtained from T_1 as follows: for each vertex $v \in S \cap V(T_1)$ and each vertex $x \in C_v$, add one new vertex x and one new edge xv to T'_1 . Choose $r \in V(T_1) \setminus S$ and consider T'_1 to be rooted at r . We use T'_1 to define a tree-decomposition of G_1 , where the bags are defined as follows. Let $B_r := \{r\}$. For each vertex $w \in V(T_1) \setminus (S \cup \{r\})$, if p is the parent of w in T'_1 , let $B_w := \{w, p\}$. For each vertex $v \in S \cap V(T_1)$ and each vertex x in C_v , let $B_v := N_{T_1}(v) \cup \{v\}$ and $B_x := N_{T_1}(v) \cup \{x\}$.

We now show that $(B_x : x \in V(T'_1))$ is a tree-decomposition of G_1 . The bags containing r are indexed by $N_{T_1}(r) \cup \{r\}$, which induces a (connected) subtree of T'_1 .

For each vertex $w \in V(T_1) \setminus (S \cup \{r\})$ with parent p , the bags containing w are those indexed by $\cup\{C_v \cup \{v\} : v \in N_{T_1}(w) \cap S\} \cup \{w\} \cup (N_{T_1}(w) \setminus \{p\})$, which induces a subtree of T'_1 (since $vx \in E(T'_1)$ for each $x \in C_v$ where $v \in N_{T_1}(w) \cap S$). For each vertex $v \in S$ with parent p , the bags containing v are those indexed by $N_{T_1}(v) \cup \{v\} \setminus \{p\}$, which induces a subtree of T'_1 . For each vertex $v \in S$ and $x \in C_v$, B_x is the only bag that contains x . Hence property (T1) in the definition of tree-decomposition holds. For each edge pv of T_1 where p is the parent of v , the bag B_v contains both p and v . Every other edge of G_1 joins x and w for some $v \in S$ and $x \in C_v$ and $w \in N_{T_1}(v)$, in which case B_x contains both x and w . Hence (T2) holds. Therefore $(B_x : x \in V(T'_1))$ is a tree-decomposition of G_1 . Since each bag has size at most $s + 1$, we have $\text{tw}(G_1) \leq s$. ■

3 Upper bound

To prove upper bounds on $C(T, \mathcal{G}, n)$, it is convenient to work in the following setting. For graphs G and H , an *image* of H in G is an injection $\phi : V(H) \rightarrow V(G)$ such that $\phi(u)\phi(v) \in E(G)$ for all $uv \in E(H)$. Let $I(H, G)$ be the number of images of H in G . For a graph class \mathcal{G} , let $I(H, \mathcal{G}, n)$ be the maximum of $I(H, G)$ taken over all n -vertex graphs $G \in \mathcal{G}$. If H is fixed then $C(H, G)$ and $I(H, G)$ differ by a constant factor. In particular, if $|V(H)| = h$ then

$$(3.1) \quad \begin{aligned} C(H, G) &\leq I(H, G) \leq h! C(H, G), \\ C(H, \mathcal{G}, n) &\leq I(H, \mathcal{G}, n) \leq h! C(H, \mathcal{G}, n). \end{aligned}$$

So to bound $C(T, \mathcal{G}, n)$ it suffices to work with images rather than copies.

Our proof needs two tools from the literature. The first is due to Eppstein [32]. A collection \mathcal{H} of images of a graph H in a graph G is *coherent* if for all distinct images $\phi_1, \phi_2 \in \mathcal{H}$ and for all distinct vertices $x, y \in V(H)$, we have $\phi_1(x) \neq \phi_2(y)$.

Lemma 3.1 [32] *Let H be a graph with h vertices and let G be a graph. Every collection of at least $c_{3.1}(h, t) := h!^2 t^h$ images of H in G contains a coherent subcollection of size at least t .*

We also use the following result of Erdős and Rado [34]; see [7, 9] for recent quantitative improvements. A *t-sunflower* is a collection \mathcal{S} of t sets for which there exists a set R such that $X \cap Y = R$ for all distinct $X, Y \in \mathcal{S}$. The set R is called the *kernel* of \mathcal{S} .

Lemma 3.2 (Sunflower Lemma [34]) *Every collection of at least $c_{3.2}(h, t) := h!(t - 1)^h + 1$ many h -subsets of a set contains a t -sunflower.*

Consider graphs H and G . An *H-model* in a graph G is a collection $(X_v : v \in V(H))$ of pairwise disjoint connected subgraphs of G indexed by the vertices of H , such that for each edge $vw \in E(H)$ there is an edge of G joining X_v and X_w . Each subgraph X_v is called a *branch set*. A graph G contains an *H-model* if and only if H is a minor of G . An *H-model* $(X_v : v \in V(H))$ in G is *c-shallow* if X_v has radius at most c for each $v \in V(H)$. An *H-model* $(X_v : v \in V(H))$ in G is *c-small* if $|V(X_v)| \leq c$ for each $v \in V(H)$. Shallow models are key components in the sparsity theory of Nešetřil and de Mendez [77]. Small models have also been studied [37, 75, 91, 95].




Figure 1: $H^{(3,4)}$ where $V(H) = \{a, b, c, d, e\}$.

The next lemma is the heart of the paper. To describe the result we need the following construction, illustrated in Figure 1. For a graph H , and $s, t \in \mathbb{N}$, and $v \in V(H)$ let

$$\overline{\text{deg}}_{H,s}(v) := \max\{s + 1 - \text{deg}_H(v), 0\}.$$

Then define $H^{(s,t)}$ to be the graph with vertex set

$$V(H^{(s,t)}) := \{(v, i) : v \in V(H), i \in [t]\} \cup \{(v, j)^* : v \in V(H), j \in [\overline{\text{deg}}_{H,s}(v)]\}$$

and edge set

$$E(H^{(s,t)}) := \{(v, i)(w, i) : vw \in E(H), i \in [t]\} \cup \{(v, i)(v, j)^* : v \in V(H), i \in [t], j \in [\overline{\text{deg}}_{H,s}(v)]\}.$$

Several notes about $H^{(s,t)}$ are in order:

- (A) For each $i \in [t]$, let X_i be the subgraph of $H^{(s,t)}$ induced by $\{(v, i) : v \in V(H)\}$. Then $X_i \cong H$. Contracting each X_i to a single vertex produces $K_{s',t}$ where

$$s' := \sum_{v \in V(H)} \overline{\text{deg}}_{H,s}(v) \geq \sum_{v \in V(H)} (s + 1 - \text{deg}_H(v)) = (s + 1)|V(H)| - 2|E(H)|.$$

If H is a nonempty tree then $s' \geq |V(H)|(s - 1) + 2 \geq s + 1$, implying $K_{s+1,t}$ is a minor of $H^{(s,t)}$.

- (B) Each vertex $(v, j)^*$ has degree t and each vertex (v, i) has degree $\deg_H(v) + \overline{\deg}_{H,s}(v) \geq s + 1$. In particular, if $t \geq s + 1$ then $H^{(s,t)}$ has minimum degree at least $s + 1$.
- (C) If H is connected then $\text{diameter}(H^{(s,t)}) \leq \text{diameter}(H) + 2$.

Define the *density* of a graph G to be $\rho(G) := \frac{|E(G)|}{|V(G)|}$. For a graph class \mathcal{G} , let $\rho(\mathcal{G}) := \sup\{\rho(G) : G \in \mathcal{G}\}$

Lemma 3.3 *For all $s, t, h \in \mathbb{N}$ and $\rho \in \mathbb{R}_{\geq 0}$, there exists a constant $c := c_{3.3}(s, t, h, \rho) := c_{3.1}(h, c_{3.2}(h, t))(\rho + 1)^h$ such that for every forest T with h vertices, if G is a graph with $\rho(G) \leq \rho$ and $I(T, G) \geq c|V(G)|^{\alpha_s(T)}$, then G contains $U^{(s,t)}$ as a subgraph for some (non-empty) subtree U of T .*

Proof Let $S := \{v \in V(T) : \deg_T(v) \leq s\}$. Let X be a stable set in $F := T[S]$ of size $k := \alpha_s(T)$. Since F is bipartite, by Konig’s Edge Cover Theorem [64], there is a set $Y \subseteq V(F) \cup E(F)$ with $|Y| = |X|$ such that each vertex of F is either in Y or is incident to an edge in Y . In fact, $Y \cap V(F)$ is the set of isolated vertices of F , although we will not need this property.

Let G be an n -vertex graph with $\rho(G) \leq \rho$ and $I(T, G) \geq cn^k$. Let \mathcal{J} be the set of images of T in G . So $|\mathcal{J}| \geq cn^k$. Let $\mathcal{X} := \binom{V(G) \cup E(G)}{k}$. Note that $|\mathcal{X}| \leq \binom{(\rho+1)n}{k} \leq (\rho + 1)^k n^k$. For each $\phi \in \mathcal{J}$, let

$$Y_\phi := \{\phi(x) : x \in Y \cap V(F)\} \cup \{\phi(x)\phi(y) : xy \in Y \cap E(F)\},$$

which is an element of \mathcal{X} since $|Y| = k$. For each $Z \in \mathcal{X}$, let $\mathcal{J}_Z := \{\phi \in \mathcal{J} : Y_\phi = Z\}$. By the pigeonhole principle, there exists $Z \in \mathcal{X}$ such that

$$|\mathcal{J}_Z| \geq |\mathcal{J}|/|\mathcal{X}| \geq c/(\rho + 1)^k \geq c/(\rho + 1)^h = c_{3.1}(h, c_{3.2}(h, t)).$$

By Lemma 3.1 applied to \mathcal{J}_Z , there is a coherent family $\mathcal{J}_1 \subseteq \mathcal{J}_Z$ with $|\mathcal{J}_1| = c_{3.2}(h, t)$.

We claim that the vertex sets in G corresponding to the images of T in \mathcal{J}_1 are all distinct. Suppose that $V(\phi_1(V(T))) = V(\phi_2(V(T)))$ for $\phi_1, \phi_2 \in \mathcal{J}_1$. Let x be any vertex in T . If $\phi_1(x) \neq \phi_2(x)$, then $\phi_2(y) = \phi_1(x)$ for some vertex y of T with $y \neq x$ (since $V(\phi_1(V(T))) = V(\phi_2(V(T)))$), which contradicts the definition of coherence. Thus $\phi_1(x) = \phi_2(x)$ for each vertex x of T . Thus $\phi_1 = \phi_2$. This proves our claim.

Therefore, by Lemma 3.2 applied to $\{\phi(V(T)) : \phi \in \mathcal{J}_1\}$, there is a set R of vertices in G and a subfamily $\mathcal{J}_2 \subseteq \mathcal{J}_1$ such that $\phi_1(V(T)) \cap \phi_2(V(T)) = R$ for all distinct $\phi_1, \phi_2 \in \mathcal{J}_2$, and $|\mathcal{J}_2| = t$.

Fix $\phi_0 \in \mathcal{J}_2$ and let $K := \phi_0^{-1}(R)$. Note that K does not depend on the choice of ϕ_0 . Moreover, $S \subseteq K$ because $Y_\phi = Z$ for every $\phi \in \mathcal{J}_2$, and each vertex in S is either in Y or is incident to an edge in Y . Let U be some connected component of $T - K$. Note that $V(U) \cap S = \emptyset$, since $S \subseteq K$. Thus, each vertex $v \in V(U)$ has $\deg_T(v) \geq s + 1$ and thus there is a set N_v of at least $\overline{\deg}_{U,s}(v)$ neighbors of v in K . Again by coherence, $\phi_1(N_v) = \phi_2(N_v)$ for all $\phi_1, \phi_2 \in \mathcal{J}_2$ and $v \in V(U)$. Observe that $N_{v_1} \cap N_{v_2} = \emptyset$ for distinct $v_1, v_2 \in U$, as otherwise T would contain a cycle. Thus $(\phi(U) : \phi \in \mathcal{J}_2)$ and $(\phi_0(N_v) : v \in V(U))$ define a subgraph of G isomorphic to $U^{(s,t)}$. ■

We now prove our first main result.

Proof of Theorem 1.1 Since every graph with treewidth k is in \mathcal{D}_k , Lemma 2.1 implies $C(T, \mathcal{D}_k, n) \in \Omega(n^{\alpha_k(T)})$. For the upper bound, let G be a k -degenerate graph. So $\rho(G) \leq k$. By Lemma 3.3 with $s = k$ and $t = k + 1$, if $I(T, G) \geq c|V(G)|^k$ then G contains $U^{(k, k+1)}$ as a subgraph for some subtree U of T . However, $U^{(k, k+1)}$ has minimum degree $k + 1$, contradicting the k -degeneracy of G . Hence $I(T, G) \leq c|V(G)|^k$ and $C(T, \mathcal{D}_k, n) \in O(n^{\alpha_k(T)})$ by Equation (3.1). ■

The following special case of Lemma 3.3 will be useful. Say $\{X_1, \dots, X_s; Y_1, \dots, Y_t\}$ is a (p, q) -model of $K_{s,t}$ in a graph G if:

- $X_1, \dots, X_s, Y_1, \dots, Y_t$ are pairwise disjoint connected subgraphs of G ,
- for each $i \in [s]$ and $j \in [t]$ there is an edge in G between X_i and Y_j ,
- $|V(X_i)| \leq p$ for each $i \in [s]$ and $|V(Y_j)| \leq q$ for each $j \in [t]$.

Corollary 3.4 For all $s, t, h \in \mathbb{N}$ and $\rho \in \mathbb{R}_{\geq 0}$, for every forest T with h vertices, if G is a graph with $\rho(G) \leq \rho$ and $I(T, G) \geq c_{3.3}(s, t, h, \rho)|V(G)|^{\alpha_s(T)}$, then for some $h' \in [h]$, G contains a subgraph of diameter at most $h' + 1$ that contains a $(1, h')$ -model of $K_{h'(s-1)+2, t}$. In particular, G contains a $(1, h)$ -model of $K_{s+1, t}$.

Proof By Lemma 3.3, G contains $U^{(s, t)}$ as a subgraph for some subtree U of T . The main claim follows from (A) and (C) where $h' := |V(U)|$. The final claim follows since $h' \in [h]$, implying $h'(s - 1) + 2 \geq s + 1$. ■

4 Minor-closed classes

Theorem 1.2 is implied by Lemma 2.1 and Corollary 3.4 and since every minor-closed class has bounded density [65, 93]. We now give several examples of Theorem 1.2.

Treewidth:

Let \mathcal{T}_k be the class of graphs with treewidth at most k . Then \mathcal{T}_k is a minor-closed class, and every graph in \mathcal{T}_k has minimum degree at most k , implying $\rho(\mathcal{T}_k) \leq k$ and $K_{k+1, k+1} \notin \mathcal{T}_k$. Thus Theorem 1.2 with $s = k$ implies that for every fixed forest T ,

$$C(T, \mathcal{T}_k, n) \in \Theta(n^{\alpha_k(T)}).$$

Surfaces:

Let \mathcal{S}_Σ be the class of graphs that embed⁵ in a surface Σ . Then \mathcal{S}_Σ is a minor-closed class. Huynh et al. [59] proved that for every $H \in \mathcal{S}_\Sigma$,

$$C(H, \mathcal{S}_\Sigma, n) \in \Theta(n^{f(H)}),$$

where $f(H)$ is a graph invariant called the *flap-number* of H , which is independent of Σ . Huynh et al. [59] noted that $f(T) = \alpha_2(T)$ for a forest T . So, in particular,

$$C(T, \mathcal{S}_\Sigma, n) \in \Theta(n^{\alpha_2(T)}).$$

⁵For $h \geq 0$, let \mathbb{S}_h be the sphere with h handles. For $c \geq 0$, let \mathbb{N}_c be the sphere with c cross-caps. Every surface is homeomorphic to \mathbb{S}_h or \mathbb{N}_c . The Euler genus of \mathbb{S}_h is $2h$. The Euler genus of \mathbb{N}_c is c . The Euler genus of a graph G is the minimum Euler genus of a surface in which G embeds with no crossings. See [74] for background about graphs embedded in surfaces.

This result is also implied by Theorem 1.2 since for every surface Σ of Euler genus g , Euler’s formula implies that $K_{3,2g+3}$ is not in \mathcal{S}_Σ (first observed by Ringel [86]), and

$$\rho(\mathcal{S}_\Sigma) \leq \rho_g := \max\left\{3, \frac{1}{4}(5 + \sqrt{24g + 1})\right\};$$

see [22] for a proof.

Excluding a complete bipartite minor:

Let $\mathcal{B}_{s,t}$ be the class of graphs containing no complete bipartite graph $K_{s,t}$ minor, where $t \geq s$. Since $K_{s,t}$ has treewidth s , every graph with treewidth at most $s - 1$ is in $\mathcal{B}_{s,t}$. By Theorem 1.2, for every fixed forest T ,

$$(4.1) \quad C(T, \mathcal{B}_{s,t}, n) \in \Theta(n^{\alpha_{s-1}(T)}).$$

This answers affirmatively a question raised by Huynh et al. [59].

Excluding a complete minor:

Let \mathcal{C}_k be the class of graphs containing no complete graph K_k minor. Then $K_{k-1,k-1} \notin \mathcal{C}_k$ (since contracting a $(k - 2)$ -edge matching in $K_{k-1,k-1}$ gives K_k). Every graph with treewidth at most $k - 2$ is in \mathcal{C}_k . Thus Theorem 1.2 with $s = k - 2$ implies that for every fixed forest T ,

$$C(T, \mathcal{C}_k, n) \in \Theta(n^{\alpha_{k-2}(T)}).$$

Colin de Verdière number:

The Colin de Verdière parameter $\mu(G)$ is an important graph invariant introduced by de Verdière [23, 24]; see [90, 96] for surveys. It is known that $\mu(G) \leq 1$ if and only if G is a disjoint union of paths, $\mu(G) \leq 2$ if and only if G is outerplanar, $\mu(G) \leq 3$ if and only if G is planar, and $\mu(G) \leq 4$ if and only if G is linklessly embeddable. Let $\mathcal{V}_k := \{G : \mu(G) \leq k\}$. Then \mathcal{V}_k is a minor-closed class [23, 24]. Goldberg and Berman [51] proved that $\mu(G) \leq \text{tw}(G) + 1$. So every graph with treewidth at most $k - 1$ is in \mathcal{V}_k . van der Holst et al. [96] proved that $\mu(K_{s,t}) = s + 1$ for $t \geq \max\{s, 3\}$, so $K_{k,\max\{k,3\}} \notin \mathcal{V}_k$. Thus Theorem 1.2 with $s = k - 1$ and $t = \max\{k, 3\}$ implies that for every fixed forest T ,

$$(4.2) \quad C(T, \mathcal{V}_k, n) \in \Theta(n^{\alpha_{k-1}(T)}).$$

Linkless graphs:

A graph is *linklessly embeddable* if it has an embedding in \mathbb{R}^3 with no two linked cycles [87, 89]. Let \mathcal{L} be the class of linklessly embeddable graphs. Then \mathcal{L} is a minor-closed class whose minimal excluded minors are the so-called Petersen family [88], which includes $K_6, K_{4,4}$ minus an edge, and the Petersen graph. As mentioned above, $\mathcal{L} = \mathcal{V}_4$. Thus Equation (4.2) with $k = 4$ implies for every fixed forest T ,

$$C(T, \mathcal{L}, n) \in \Theta(n^{\alpha_3(T)}).$$

Knotless graphs:

A graph is *knotlessly embeddable* if it has an embedding in \mathbb{R}^3 in which every cycle forms a trivial knot; see [83] for a survey. Let \mathcal{K} be the class of knotlessly embeddable graphs. Then \mathcal{K} is a minor-closed class whose minimal excluded minors include K_7 and $K_{3,3,1,1}$ (see [18, 40]). More than 260 minimal excluded minors are known [52], but the full list of minimal excluded minors is unknown. Since $K_7 \notin \mathcal{K}$, we have $\rho(\mathcal{K}) \leq \rho(\mathcal{C}_7) < 5$ by a theorem of Mader [73]. Shimabara [92] proved that $K_{5,5} \notin \mathcal{K}$. By Theorem 1.2,

$$C(T, \mathcal{K}, n) \in O(n^{\alpha_4(T)}).$$

This bound would be tight if every treewidth 4 graph is knotlessly embeddable, which is an open problem of independent interest.

The above results all depend on excluded complete bipartite minors. We now show that excluded complete bipartite minors determine $C(T, \mathcal{G}, n)$ for a broad family of minor-closed classes.

Theorem 4.1 *Let \mathcal{G} be a minor-closed class such that every minimal forbidden minor of \mathcal{G} is two-connected. Let s be the maximum integer such that $K_{s,t} \in \mathcal{G}$ for every $t \in \mathbb{N}$. Then for every forest T ,*

$$C(T, \mathcal{G}, n) = \Theta(n^{\alpha_s(T)}).$$

Proof Note that the condition that every minimal forbidden minor of \mathcal{G} is two-connected is equivalent to saying that \mathcal{G} is closed under the one-sum operation (that is, if $G_1, G_2 \in \mathcal{G}$, and $|V(G_1 \cap G_2)| \leq 1$, then $G_1 \cup G_2 \in \mathcal{G}$).

The proof of Lemma 2.1 shows that for all sufficiently large $n \in \mathbb{N}$ there exists an n -vertex graph G with $C(T, G) \geq cn^{\alpha_s(T)}$, where G is obtained from one-sums of complete bipartite graphs $K_{s',t}$ with $s' \leq s$. By the definition of s and since \mathcal{G} is closed under one-sums, $G \in \mathcal{G}$. Thus $C(T, \mathcal{G}, n) \in \Omega(n^{\alpha_s(T)})$.

Now we prove the upper bound. Since \mathcal{G} is minor-closed, \mathcal{G} has bounded density [65, 93]. By the definition of s , there exists $t \in \mathbb{N}$ such that $K_{s+1,t} \notin \mathcal{G}$. By (A), we have $U^{(s,t)} \notin \mathcal{G}$ for every nonempty subtree U of T . Thus $I(T, \mathcal{G}, n) \in O(n^{\alpha_s(T)})$ by Lemma 3.3. ■

Note that minor-closed classes with bounded pathwidth (that is, those excluding a fixed forest as a minor [11]) are examples not covered by Theorem 4.1. Determining $C(T, \mathcal{G}_k, n)$, where \mathcal{G}_k is the class of pathwidth k graphs, is an interesting open problem.

5 Beyond minor-closed classes

This section asymptotically determines $C(T, \mathcal{G}, n)$ for several nonminor-closed graph classes \mathcal{G} .

5.1 Shortcut systems

Dujmović et al. [28] introduced the following definition which generalizes the notion of shallow immersion [78] and provides a way to describe a graph class in terms

of a simpler graph class. Then properties of the original class are (in some sense) transferred to the new class. Let \mathcal{P} be a set of nontrivial paths in a graph G . Each path $P \in \mathcal{P}$ is called a *shortcut*; if P has endpoints v and w then it is a vw -*shortcut*. Given a graph G and a shortcut system \mathcal{P} for G , let $G^{\mathcal{P}}$ be the simple supergraph of G obtained by adding the edge vw for each vw -shortcut in \mathcal{P} . Dujmović et al. [28] defined \mathcal{P} to be a (k, d) -*shortcut system* (for G) if:

- every path in \mathcal{P} has length at most k and
- for every $v \in V(G)$, the number of paths in \mathcal{P} that use v as an internal vertex is at most d .

We use the following variation. Say \mathcal{P} is a $(k, d)^*$ -*shortcut system* (for G) if:

- every path in \mathcal{P} has length at most k and
- for every $v \in V(G)$, if M_v is the set of vertices $u \in V(G)$ such that there exists a uw -shortcut in \mathcal{P} in which v is an internal vertex, then $|M_v| \leq d$.

Clearly, every $(k, d)^*$ -shortcut system is a $(k, \binom{d}{2})$ -shortcut system (since $G^{\mathcal{P}}$ is simple), and every (k, d) -shortcut system is a $(k, 2d)^*$ -shortcut system.

The next lemma shows that if $G^{\mathcal{P}}$ contains a “small” model of a “large” complete bipartite graph, then so does G .

Lemma 5.1 *For all $s, t, d, k, p, q \in \mathbb{N}$, let $s' := (d(k-1)(p-1)+1)(s-1)+1$ and $t' := (2d(k-1)(s+q-1)+1)(t-1)+1+sd(p+(k-1)(p-1))$. Let \mathcal{P} be a $(k, d)^*$ -shortcut system for a graph G . If $G^{\mathcal{P}}$ contains a (p, q) -model of $K_{s', t'}$, then G contains a $(p+(k-1)(p-1), q+(k-1)(s+q-1))$ -model of $K_{s, t}$.*

Proof Let $(X_1, \dots, X_{s'}; Y_1, \dots, Y_{t'})$ be a (p, q) -model of $K_{s', t'}$ in $G^{\mathcal{P}}$. We may assume that each edge of G is (a path of length 1) in \mathcal{P} . Let $I := [s']$ and $J := [t']$. We may assume that X_i and Y_j are subtrees of $G^{\mathcal{P}}$ for $i \in I$ and $j \in J$.

Consider each $i \in I$. Let C_i be the set of all vertices internal to some uw -shortcut with $uw \in E(X_i)$. Since $|E(X_i)| \leq p-1$, we have $|C_i| \leq (k-1)(p-1)$. For each $i \in I$, let \hat{X}_i be the subgraph of G induced by $V(X_i) \cup C_i$. By construction, \hat{X}_i is connected and $|V(\hat{X}_i)| \leq p+(k-1)(p-1)$.

Consider the graph A with $V(A) := I$ where two vertices $i, i' \in V(A)$ are adjacent if $V(\hat{X}_i) \cap V(\hat{X}_{i'}) \neq \emptyset$. For each $ii' \in E(A)$, fix a vertex $v_{i, i'}$ in $V(\hat{X}_i) \cap V(\hat{X}_{i'})$, which is in $C_i \cup C_{i'}$ since $V(X_i) \cap V(X_{i'}) = \emptyset$. For $i \in I$ and $v \in C_i$, define $E_{v, i}$ to be the set of all edges $ii' \in E(A)$ with $v_{i, i'} = v$. If ii' is in $E_{v, i}$ and $v \notin X_{i'}$, then $|M_v \cap X_{i'}| \geq 2$. Also $|M_v \cap X_i| \geq 2$. Since v is in at most one $X_{i'}$, in total, $|M_v| \geq 2|E_{v, i}|$, implying $|E_{v, i}| \leq \frac{d}{2}$. Since $|I| = |V(A)|$ and $|C_i| \leq (k-1)(p-1)$,

$$|E(A)| \leq \sum_{i \in I} \sum_{v \in C_i} |E_{v, i}| \leq \frac{d}{2}(k-1)(p-1)|V(A)|.$$

Thus, A has average degree at most $d(k-1)(p-1)$. By Turán’s Theorem, A contains a stable set I' of size $\lceil |I|/(d(k-1)(p-1)+1) \rceil = s$. For distinct $i, i' \in I'$, the subgraphs \hat{X}_i and $\hat{X}_{i'}$ are disjoint. Let $\mathcal{X} := \cup_{i \in I'} V(\hat{X}_i)$. Note that $|\mathcal{X}| \leq s(p+(k-1)(p-1))$.

Let $Z := \cup_{x \in \mathcal{X}} M_x$. Then $|Z| \leq sd(p+(k-1)(p-1))$. Thus, Y_j intersects Z for at most $sd(p+(k-1)(p-1))$ elements $j \in J$. Hence, J contains a subset K of size $(2d(k-1)(s+q-1)+1)(t-1)+1$ such that $V(Y_j) \cap Z = \emptyset$ for each $j \in K$.

Consider each $j \in K$. Initialize $D_j := \emptyset$. For each $i \in I'$, choose $x \in V(X_i)$ and $w \in V(Y_j)$ such that $xw \in E(G^{\mathcal{P}})$, and add all the internal vertices of the xw -shortcut $P \in \mathcal{P}$ to D_j . For each edge uw of Y_j , add all the internal vertices of the uw -shortcut $P \in \mathcal{P}$ to D_j . Note that

$$|D_j| \leq (k - 1)|I'| + (k - 1)|E(Y_j)| \leq (k - 1)(s + q - 1),$$

since Y_j has at most $q - 1$ edges. Moreover, $D_j \cap \mathcal{X} = \emptyset$ since $V(Y_j) \cap Z = \emptyset$.

For each $j \in K$, let \hat{Y}_j be the subgraph of G induced by $V(Y_j) \cup D_j$. By construction, \hat{Y}_j is connected with at most $q + (k - 1)(s + q - 1)$ vertices and is disjoint from \mathcal{X} .

Consider the graph B with $V(B) := K$ where two vertices $j, j' \in V(B)$ are adjacent if $V(\hat{Y}_j) \cap V(\hat{Y}_{j'}) \neq \emptyset$. For each $jj' \in E(B)$, fix a vertex $v_{j,j'}$ in $V(\hat{Y}_j) \cap V(\hat{Y}_{j'})$, which is in $D_j \cup D_{j'}$ since $V(Y_j) \cap V(Y_{j'}) = \emptyset$. For $j \in K$ and $v \in D_j$, define $E_{v,j}$ to be the set of all edges $jj' \in E(B)$ with $v_{j,j'} = v$.

We now bound $|E(B)|$. If jj' is in $E_{v,j}$ and $v \notin Y_{j'}$, then $|M_v \cap Y_{j'}| \geq 1$. Also $|M_v \cap Y_j| \geq 1$. Since v is in at most one $Y_{j'}$, in total, $|M_v| \geq |E_{v,j}|$, implying $|E_{v,j}| \leq d$. Since $|K| = |V(B)|$ and $|D_j| \leq (k - 1)(s + q - 1)$,

$$|E(B)| \leq \sum_{j \in K} \sum_{v \in D_j} |E_{v,j}| \leq d(k - 1)(s + q - 1)|V(B)|,$$

implying B has average degree at most $2d(k - 1)(s + q - 1)$. By Turán's Theorem, B contains a stable set L of size $\lceil |K| / (2d(k - 1)(s + q - 1) + 1) \rceil = t$.

For distinct $j, j' \in L$, since L is a stable set in B , \hat{Y}_j and $\hat{Y}_{j'}$ are disjoint. For each $j \in L$, Y_j and \mathcal{X} are disjoint by assumption, and D_j and \mathcal{X} are disjoint by construction. Also, for each $i \in I'$ and $j \in L$, there is an edge between \hat{X}_i and \hat{Y}_j by construction. Thus $\{\hat{X}_i : i \in I'\}$ and $\{\hat{Y}_j : j \in L\}$ form a $(p + (k - 1)(p - 1), q + k(s + q - 1))$ -model of $K_{s,t}$ in G . ■

Lemma 5.1 with $p = 1$ implies the following result. We emphasize that the value of s does not change in the two models.

Corollary 5.2 Fix $s, t, k, d, q \in \mathbb{N}$. Let $t' := (2d(k - 1)(s + q - 1) + 1)(t - 1) + 1 + sd$. Let \mathcal{P} be a $(k, d)^*$ -shortcut system for a graph G . If $G^{\mathcal{P}}$ contains a $(1, q)$ -model of $K_{s,t'}$, then G contains a $(1, q + (k - 1)(s + q - 1))$ -model of $K_{s,t}$.

5.2 Low-degree squares of graphs

The above result on shortcut systems leads to the following extension of our results for minor-closed classes. For a graph G and $d \in \mathbb{N}$, let $G^{(d)}$ be the graph obtained from G by adding a clique on $N_G(v)$ for each vertex $v \in V(G)$ with $\deg_G(v) \leq d$. (This definition incorporates and generalizes the square of a graph with maximum degree d .) Note that $G^{(d)} = G^{\mathcal{P}}$, where \mathcal{P} is the $(2, d)^*$ -shortcut system $\{uvw : v \in V(G); \deg_G(v) \leq d; u, w \in N_G(v); u \neq w\}$. For a graph class \mathcal{G} , let $\mathcal{G}^{(d)} := \{G^{(d)} : G \in \mathcal{G}\}$. Note that $\rho(G^{(d)}) \leq \rho(G) + \binom{d}{2}$. Corollary 3.4 and Corollary 5.2 with $k = 2$ and $q = h$ imply:

Corollary 5.3 Fix $s, t, d, h \in \mathbb{N}$ and $\rho \in \mathbb{R}_{\geq 0}$. Let T be fixed forest with h vertices. Let $t' := (2d(s + h - 1) + 1)(t - 1) + 1 + sd$. Let G be a graph with $\rho(G) \leq \rho$ and

containing no $(1, 2h + s - 1)$ -model of $K_{s,t}$. Then $G^{(d)}$ contains no $(1, h)$ -model of $K_{s,t'}$, and

$$C(T, G^{(d)}) \leq I(T, G^{(d)}) \leq c_{3.3}(s - 1, t', h, \rho + \binom{d}{2}) |V(G)|^{\alpha_{s-1}(T)}.$$

With Lemma 2.1 we have:

Theorem 5.4 Fix $s, t, d, h \in \mathbb{N}$ and $\rho \in \mathbb{R}_{\geq 0}$. Let T be fixed forest with h vertices. Let $t' := (2d(s + h - 1) + 1)(t - 1) + 1 + sd$. Let \mathcal{G} be a graph class such that $\rho(\mathcal{G}) \leq \rho$, every graph with treewidth at most $s - 1$ is in \mathcal{G} , and no graph in \mathcal{G} contains a $(1, 2h + s - 1)$ -model of $K_{s,t}$. Then no graph in $\mathcal{G}^{(d)}$ contains a $(1, h)$ -model of $K_{s,t'}$, and

$$C(T, \mathcal{G}^{(d)}, n) = \Theta(n^{\alpha_{s-1}(T)}).$$

Theorem 5.4 is applicable to all the minor-closed classes discussed in Section 4. For example, we have the following extension of Equation (4.1). Recall that $\mathcal{B}_{s,t}^{(d)}$ is the class of graphs $G^{(d)}$ where G contains no $K_{s,t}$ -minor. Then for every fixed forest T ,

$$C(T, \mathcal{B}_{s,t}^{(d)}, n) = \Theta(n^{\alpha_{s-1}(T)}).$$

5.3 Map graphs

Map graphs are defined as follows. Start with a graph G_0 embedded in a surface Σ , with each face labelled a “nation” or a “lake,” where each vertex of G_0 is incident with at most d nations. Let G be the graph whose vertices are the nations of G_0 , where two vertices are adjacent in G if the corresponding faces in G_0 share a vertex. Then G is called a (Σ, d) -map graph. A (\mathbb{S}_0, d) -map graph is called a (plane) d -map graph; such graphs have been extensively studied [15–17, 25, 41]. Let $\mathcal{M}_{\Sigma,d}$ be the set of all (Σ, d) -map graphs. Since $\mathcal{M}_{\Sigma,3} = \mathcal{S}_{\Sigma}$ (see [17, 26]), map graphs provide a natural generalization of graphs embeddable in a surface.

Let $G \in \mathcal{M}_{\Sigma,d}$ where Σ has Euler genus g . Let T be a fixed forest with h vertices. Dujmović et al. [28] proved that G is a subgraph of $G_0^{\mathcal{P}}$ for some graph $G_0 \in \mathcal{S}_{\Sigma}$ and some $(2, \frac{1}{2}d(d - 3))$ -shortcut system \mathcal{P} of G_0 . Inspecting the proof in [28] one observes that \mathcal{P} is a $(2, d)^*$ -shortcut system. In the plane case, Chen [15] proved that $\rho(\mathcal{M}_{\mathbb{S}_0,d}) < d$. An analogous argument shows that $\rho(\mathcal{M}_{\Sigma,d}) \in O(d\sqrt{g+1})$. The same bound can also be concluded from Equation (5.2). Since G_0 contains no $K_{3,2g+3}$ minor, by Corollary 5.2, for each $q \in \mathbb{N}$, $G_0^{\mathcal{P}}$ and thus G contains no $(1, q)$ -model of $K_{3,t'}$ where $t' := (2d(q + 2) + 1)(2g + 2) + 1 + 3d$. With $q = h$, Corollary 3.4 then implies that $C(T, G) \leq I(T, G) \leq c_{3.3}(2, t', h, \rho) |V(G)|^{\alpha_2(T)}$. Hence

$$C(T, \mathcal{M}_{\Sigma,d}, n) \in \Theta(n^{\alpha_2(T)}),$$

where the lower bound follows from Lemma 2.1 since every graph with treewidth 2 is planar and is thus a (Σ, d) -map graph. Also note the $q = 1$ case above shows that

$$K_{3,(6d+1)(2g+2)+1+3d} \notin \mathcal{M}_{\Sigma,d}.$$

5.4 Bounded number of crossings

Here, we consider drawings of graphs with a bounded number of crossings per edge. Throughout the paper, we assume that no three edges cross at a single point in a drawing of a graph. For a surface Σ and $k \in \mathbb{N}$, let $\mathcal{S}_{\Sigma,k}$ be the class of graphs G that have a drawing in Σ such that each edge is in at most k crossings. Since $\mathcal{S}_{\Sigma,0} = \mathcal{S}_{\Sigma}$, this class provides a natural generalization of graphs embeddable in surfaces and is widely studied [22, 28, 80]. Graphs in $\mathcal{S}_{\Sigma_0,k}$ are called k -planar. The case $k = 1$ is particularly important in the graph drawing literature; see [63] for a bibliography with over 100 references.

Let T be a fixed forest with h vertices. Let $G \in \mathcal{S}_{\Sigma,k}$ where Σ has Euler genus g . Dujmović et al. [28] noted that by replacing each crossing point by a dummy vertex, we obtain a graph $G_0 \in \mathcal{S}_{\Sigma}$ such that G is a subgraph of $G_0^{\mathcal{P}}$ for some $(k + 1, 2)$ -shortcut system \mathcal{P} , which is a $(k + 1, 4)^*$ -shortcut system. Results of Ossona de Mendez et al. [22] show that $\rho(\mathcal{S}_{\Sigma,k}) \leq 2\sqrt{k + 1}\rho_g$ (see Equation (5.2) below). Since G_0 contains no $K_{3,2g+3}$ minor, by Corollary 5.2, for all $q \in \mathbb{N}$, $G_0^{\mathcal{P}}$ and thus G contains no $(1, q)$ -model of $K_{3,t'}$ where $t' := (8k(q + 2) + 1)(2g + 2) + 13$. Applying this result with $q = h$, Corollary 3.4 then implies $C(T, G) \leq I(T, G) \leq c_{3,3}(2, t', h, 2\sqrt{k + 1}\rho_g) |V(G)|^{\alpha_2(T)}$. Hence,

$$(5.1) \quad C(T, \mathcal{S}_{\Sigma,k}, n) \in \Theta(n^{\alpha_2(T)}),$$

where the lower bound follows from Lemma 2.1 since every treewidth 2 graph is planar and is thus in $\mathcal{S}_{\Sigma,k}$. Also note the $q = 1$ case above shows that

$$K_{3,(24k+1)(2g+2)+13} \notin \mathcal{S}_{\Sigma,k}.$$

5.5 Bounded average number of crossings

Here, we generalize the results from the previous section for graphs that can be drawn with a bounded average number of crossings per edge. Ossona de Mendez et al. [22] defined a graph G to be k -close to Euler genus g if every subgraph G' of G has a drawing in a surface of Euler genus at most g with at most $k |E(G')|$ crossings.⁶ Let $\mathcal{E}_{g,k}$ be the

⁶The case $g = 0$ is similar to other definitions from the literature, as we now explain. Eppstein and Gupta [33] defined the *crossing graph* of a drawing of a graph G to be the graph with vertex set $E(G)$, where two vertices are adjacent if the corresponding edges in G cross. Eppstein and Gupta [33] defined a graph to be a d -degenerate crossing graph if it admits a drawing whose crossing graph is d -degenerate. Independently, Bae et al. [8] defined a graph G to be k -gap-planar if G has a drawing in the plane in which each crossing is assigned to one of the two involved edges and each edge is assigned at most k of its crossings. This is equivalent to saying that the crossing graph has an orientation with outdegree at most k at every vertex. Hakimi [68] proved that any graph H has such an orientation if and only if every subgraph of H has average degree at most $2k$. So a graph G is k -gap-planar if and only if G has a drawing such that every subgraph of the crossing graph has average degree at most $2k$ if and only if G has a drawing such that every subgraph G' of G has at most $k |E(G')|$ crossings in the induced drawing of G' . The only difference between “ k -close to planar” and “ k -gap planar” is that a k -gap planar graph has a single drawing in which every subgraph has the desired number of crossings. To complete the comparison, the definition of Eppstein and Gupta [33] is equivalent to saying that G has a drawing in which the crossing graph has an acyclic orientation with outdegree at most k at every vertex. Thus every k -degenerate crossing graph is k -gap-planar graph, and every k -gap-planar graph is a $2k$ -degenerate crossing graph.

class of graphs k -close to Euler genus g . This is a broader class than $\mathcal{S}_{\Sigma,k}$ since it allows an average of k crossings per edge, whereas $\mathcal{S}_{\Sigma,k}$ requires a maximum of k crossings per edge. In particular, if Σ has Euler genus g , then $\mathcal{S}_{\Sigma,k} \subseteq \mathcal{E}_{g,k/2}$.

The next lemma is of independent interest.

Lemma 5.5 Fix $g, r \in \mathbb{N}_0$ and $d \in \mathbb{N}$ and $k \in \mathbb{R}_{\geq 0}$. Assume that graph $G \in \mathcal{E}_{g,k}$ contains an r -shallow H -model $(X_v : v \in V(H))$ such that for every vertex $v \in V(H)$ we have $\deg_H(v) \leq d$ or $|V(X_v)| = 1$. Then H is in $\mathcal{E}_{g,2kd^2(2r+1)}$.

Proof For each $v \in V(H)$, let a_v be the central vertex of X_v . We may assume that X_v is a Breadth-first search (BFS) spanning tree of $G[V(X_v)]$ rooted at a_v and with radius at most r . Orient the edges of X_v away from a_v .

Let H' be an arbitrary subgraph of H . For each $v \in V(H')$, let X'_v be a minimal subtree of X_v rooted at a_v , such that $(X'_v : v \in V(H'))$ is an r -shallow H' -model. By minimality, X'_v has at most $\deg_{H'}(v)$ leaves. Each edge of X'_v is on a path from a leaf to a_v , implying $|E(X'_v)| \leq r \deg_{H'}(v)$.

Let G' be the subgraph of G consisting of $\cup_{v \in V(H')} X'_v$ along with one undirected edge $y_{vw}y_{wv}$ for each edge $vw \in E(H')$, where $y_{vw} \in V(X'_v)$ and $y_{wv} \in V(X'_w)$. Let P_{vw} be the directed $a_v y_{vw}$ -path in X'_v . Note that

$$|E(G')| = |E(H')| + \sum_{v \in V(H')} |E(X'_v)| \leq |E(H')| + r \sum_{v \in V(H')} \deg_{H'}(v) = (2r + 1)|E(H')|.$$

Since G is k -close to Euler genus g , G' has a drawing in a surface of Euler genus at most g with at most $k|E(G')|$ crossings. For each $e \in E(G')$, let $\ell(e)$ be the number of crossings on e in this drawing of G' . Since each crossing contributes towards ℓ for exactly two edges,

$$\sum_{e \in E(G')} \ell(e) \leq 2k|E(G')| \leq 2k(2r + 1)|E(H')|.$$

Let G'' be the multigraph obtained from G' as follows: for each vertex $v \in V(H')$ and edge e in X'_v , let the multiplicity of e in G'' equal the number of edges $vw \in E(H')$ for which the path P_{vw} uses e . Edges of G'' inherit their orientation from G' . Note that G'' has multiplicity at most d . By replicating edges in the drawing of G' we obtain a drawing of G'' such that every edge of G'' corresponding to $e \in E(G')$ is in at most $d \ell(e)$ crossings. Since each edge $e \in E(G')$ has multiplicity at most d in G'' , the number of crossings in the drawing of G'' is at most $\sum_{e \in E(G')} d^2 \ell(e) \leq 2kd^2(2r + 1)|E(H')|$.

Note that at each vertex y in G'' , in the circular ordering of edges in G'' incident to y determined by the drawing of G'' , all the incoming edges form an interval. We now use the drawing of G' to produce a drawing of a graph G''' , which is a subdivision of H' , where each vertex $v \in V(H')$ is drawn at the location of a_v . Here is the idea (see Figure 2): First “assign” each edge $y_{vw}y_{wv}$ of G' to the edge vw of H' . Next “assign” each edge of G' arising from some X'_v to exactly one edge incident to v , such that for each edge vw of H' incident to v there is a path in G' from a_v to y_{vw} consisting of edges assigned to vw . Then each edge vw in H' is drawn by following this path.

We now provide the details of this idea. Initialize $V(G''') := V(G')$ and $E(G''') := \{y_{vw}y_{wv} : vw \in E(H)\}$. Consider each vertex $v \in V(H')$. Consider the vertices $y \in$




Figure 2: Construction of the drawing of H .

$V(X'_v) \setminus \{a_v\}$ in nonincreasing order of $\text{dist}_{X'_v}(a_v, y)$ (that is, we consider the vertices of X'_v furthest from a_v first, and then move towards the root). Let x be the parent of y in X'_v . The incoming edges at y are copies of xy . Each outgoing/undirected edge yz at y is already assigned to one edge vw incident to v . Say yz_1, \dots, yz_q are the outgoing/undirected edges of G'' incident to y in clockwise order in the drawing of G'' , where yz_i is assigned to edge vw_i . If e_1, \dots, e_q are the incoming edges at y in clockwise order, then assign e_{q-i+1} to vw_i for each $i \in [q]$. Now in G''' replace vertex y by vertices y_1, \dots, y_q drawn in a sufficiently small disc around y , where y_i is incident to e_{q-i+1} and y_iz_i in G''' . Thus the edges in G''' assigned to vw form a path from a_v to y_{vw} and a path from a_w to y_{wv} . Hence G''' is a subdivision of H' (since $y_{vw}y_{wv}$ is an edge of G'''). Each edge of G''' has the same number of crossings as the corresponding edge of G'' . Thus, the total number of crossings in the drawing of G''' is at most $2kd^2(2r+1)|E(H')|$. Since G''' is a subdivision of H' , the drawing of G''' determines a drawing of H' with the same number of crossings. Therefore H is $2kd^2(2r+1)$ -close to Euler genus g . ■

We need the following results of Ossona de Mendez et al. [22]:

$$(5.2) \quad \rho(\mathcal{E}_{k,g}) \leq 2\sqrt{2k+1}\rho_g$$

$$(5.3) \quad K_{3,3k(2g+3)(2g+2)+2} \notin \mathcal{E}_{g,k}.$$

We now reach the main result of this section.

Theorem 5.6 For fixed $k, g \in \mathbb{N}_0$ and every fixed forest T ,

$$C(T, \mathcal{E}_{g,k}, n) \in \Theta(n^{\alpha_2(T)}).$$

Proof First, we prove the lower bound. By Lemma 2.1 with $s = 2$, for all sufficiently large $n \in \mathbb{N}$, there exists a graph G with $|V(G)| \leq n$ and $\text{tw}(G) \leq 2$ and

$C(T, G) \geq c_{2.1}(\alpha_2(T)) n^{\alpha_2(T)}$. Since $\text{tw}(G) \leq 2$, G is planar and is thus in $\mathcal{E}_{g,k}$. Hence, $C(T, \mathcal{E}_{g,k}, n) \in \Omega(n^{\alpha_2(T)})$.

Now, we prove the upper bound. Let $s := 2$ and $r := |V(T)|$ and $t := 54k(2r + 1)(2g + 3)(2g + 2) + 2$. Let G be an n -vertex graph in $\mathcal{E}_{g,k}$. By Equation (5.2), $\rho(G) \leq 2\sqrt{2k + 1}\rho_g$. Suppose on the contrary that $I(T, G) \geq cn^{\alpha_2(T)}$, where $c := c_{3.3}(s, t, r, 2\sqrt{2k + 1}\rho_g)$.

Let $H := K_{3,t}$. Corollary 3.4 implies that G contains a $(1, r)$ -model $(X_v : v \in V(H))$ of H . This model is r -shallow and for every vertex $v \in V(H)$ we have $\text{deg}_H(v) \leq 3$ or $|V(X_v)| = 1$. Thus, Lemma 5.5 is applicable with $d = 3$, implying that $K_{3,t} \in \mathcal{E}_{g,18k(2r+1)}$, which contradicts Equation (5.3). ■

An almost identical proof to that of Lemma 5.5 shows the following analogous result for $\mathcal{S}_{\Sigma,k}$. This can be used to prove Equation (5.1) without using shortcut systems.

Lemma 5.7 *Fix a surface Σ and $k, r \in \mathbb{N}_0$ and $d \in \mathbb{N}$. Let G be a graph in $\mathcal{S}_{\Sigma,k}$ that contains an r -shallow H -model $(X_v : v \in V(H))$ such that for every vertex $v \in V(H)$ we have $\text{deg}_H(v) \leq d$ or $|V(X_v)| = 1$. Then H is in $\mathcal{S}_{\Sigma, kd^2(2r+1)}$.*

6 Open problems

In this paper, we determined the asymptotic behavior of $C(T, \mathcal{G}, n)$ as $n \rightarrow \infty$ for various sparse graph classes \mathcal{G} and for an arbitrary fixed forest T . One obvious question is what happens when T is not a forest?

For arbitrary graphs H , the answer is no longer given by $\alpha_s(H)$. Huynh et al. [59] define a more general graph parameter, which they conjecture governs the behavior of $C(H, \mathcal{G}, n)$. An s -separation of H is a pair (A, B) of edge-disjoint subgraphs of H such that $A \cup B = H$, $V(A) \setminus V(B) \neq \emptyset$, $V(B) \setminus V(A) \neq \emptyset$, and $|V(A) \cap V(B)| = s$. A $(\leq s)$ -separation is an s' -separation for some $s' \leq s$. Separations (A, B) and (C, D) of H are independent if $E(A) \cap E(C) = \emptyset$ and $(V(A) \setminus V(B)) \cap (V(C) \setminus V(D)) = \emptyset$. If H has no $(\leq s)$ -separation, then let $f_s(H) := 1$; otherwise, let $f_s(H)$ be the maximum number of pairwise independent $(\leq s)$ -separations in H .

Conjecture 6.1 [59] *Let $\mathcal{B}_{s,t}$ be the class of graphs containing no $K_{s,t}$ minor, where $t \geq s \geq 1$. Then for every fixed graph H with no $K_{s,t}$ minor,*

$$C(H, \mathcal{B}_{s,t}, n) \in \Theta(n^{f_{s-1}(H)}).$$

As evidence for Conjecture 6.1, Eppstein [32] proved it when $f_{s-1}(H) = 1$ and Huynh et al. [59] proved it when $s \leq 3$ (and that the lower bound holds for all $s \geq 1$). It is easy to show that $f_s(T) = \alpha_s(T)$ for all $s \geq 1$ and every forest T . Thus, if true, Conjecture 6.1 would simultaneously generalize Theorem 1.2 and results from [59].

In light of Theorem 1.1, we also conjecture the following generalization.

Conjecture 6.2 *Let \mathcal{D}_k be the class of k -degenerate graphs. Then for every fixed k -degenerate graph H ,*

$$C(H, \mathcal{D}_k, n) \in \Theta(n^{f_k(H)}).$$

Acknowledgement Many thanks to both referees for several helpful comments.

References

- [1] A. F. Alameddine, *On the number of cycles of length 4 in a maximal planar graph*. J. Graph Theory 4(1980), no. 4, 417–422.
- [2] J. Alexander, J. Cutler, and T. Mink, *Independent sets in graphs with given minimum degree*. Electron. J. Combin. 19(2012), no. 3, P37.
- [3] N. Alon and Y. Caro, *On the number of subgraphs of prescribed type of planar graphs with a given number of vertices*. In: Convexity and graph theory, North-Holland Math. Stud., 86, North-Holland, 1984, pp. 25–36.
- [4] N. Alon, A. Kostochka, and C. Shikhelman, *Many cliques in H -free subgraphs of random graphs*. J. Comb. 9(2018), no. 4, 567–597.
- [5] N. Alon and C. Shikhelman, *Many T copies in H -free graphs*. J. Combin. Theory Ser. B 121(2016), 146–172.
- [6] N. Alon and C. Shikhelman, *H -free subgraphs of dense graphs maximizing the number of cliques and their blow-ups*. Discrete Math. 342(2019), no. 4, 988–996.
- [7] R. Alweiss, S. Lovett, K. Wu, and J. Zhang, *Improved bounds for the sunflower lemma*. In: Proc. 52nd Annual ACM Symp. on Theory of Computing (STOC 2020), 2020, pp. 624–630.
- [8] S. W. Bae, J.-F. Baffier, J. Chun, P. Eades, K. Eickmeyer, L. Grilli, S.-H. Hong, M. Korman, F. Montecchiani, I. Rutter, and C. D. Tóth, *Gap-planar graphs*. Theor. Comput. Sci. 745(2018), 36–52.
- [9] T. Bell, S. Chueluecha, and L. Warnke, *Note on sunflowers*. Discrete Math. 344(2021), no. 7, 112367.
- [10] T. Biedl and D. F. Wilkinson, *Bounded-degree independent sets in planar graphs*. Theory Comput. Syst. 38(2005), no. 3, 253–278.
- [11] D. Bienstock, N. Robertson, P. D. Seymour, and R. Thomas, *Quickly excluding a forest*. J. Combin. Theory Ser. B 52(1991), no. 2, 274–283.
- [12] P. Bose, V. Dujmović, and D. R. Wood, *Induced subgraphs of bounded treewidth and bounded degree*. Contrib. Discrete Math. 1(2006), no. 1, 88–105.
- [13] P. Bose, M. Smid, and D. R. Wood, *Light edges in degree-constrained graphs*. Discrete Math. 282(2004), nos. 1–3, 35–41. MR:2059504.
- [14] Z. Chase, *The maximum number of triangles in a graph of given maximum degree*. Adv. Combin. (2020), 5–10.
- [15] Z.-Z. Chen, *Approximation algorithms for independent sets in map graphs*. J. Algorithms 41(2001), no. 1, 20–40.
- [16] Z.-Z. Chen, *New bounds on the edge number of ak -map graph*. J. Graph Theory 55(2007), no. 4, 267–290.
- [17] Z.-Z. Chen, M. Grigni, and C. H. Papadimitriou, *Map graphs*. J. ACM. 49(2002), no. 2, 127–138.
- [18] J. H. Conway and C. M. Gordon, *Knots and links in spatial graphs*. J. Graph Theory 7(1983), 445–453.
- [19] J. Cutler and A. J. Radcliffe, *Extremal problems for independent set enumeration*. Electron. J. Combin. 18(2011), no. 1, P169.
- [20] J. Cutler and A. J. Radcliffe, *The maximum number of complete subgraphs in a graph with given maximum degree*. J. Combin. Theory Ser. B 104(2014), 60–71.
- [21] J. Cutler and A. J. Radcliffe, *The maximum number of complete subgraphs of fixed size in a graph with given maximum degree*. J. Graph Theory 84(2017), no. 2, 134–145.
- [22] P. O. de Mendez, S.-i. Oum, and D. R. Wood, *Defective colouring of graphs excluding a subgraph or minor*. Combinatorica 39(2019), no. 2, 377–410.
- [23] Y. C. de Verdière, *Sur un nouvel invariant des graphes et un critère de planarité*. J. Combin. Theory Ser. B 50(1990), no. 1, 11–21.
- [24] Y. C. de Verdière, *On a new graph invariant and a criterion for planarity*. In: Graph structure theory, Contemp. Math., 147, Amer. Math. Soc., 1993, pp. 137–147.
- [25] E. D. Demaine, F. V. Fomin, M. Hajiaghayi, and D. M. Thilikos, *Fixed-parameter algorithms for (k, r) -center in planar graphs and map graphs*. ACM Trans. Algorithms 1(2005), no. 1, 33–47.
- [26] V. Dujmović, D. Eppstein, and D. R. Wood, *Structure of graphs with locally restricted crossings*. SIAM J. Discrete Math. 31(2017), no. 2, 805–824.
- [27] V. Dujmović, G. Fijavž, G. Joret, T. Sulanke, and D. R. Wood, *On the maximum number of cliques in a graph embedded in a surface*. European J. Combin. 32(2011), no. 8, 1244–1252.

- [28] V. Dujmović, P. Morin, and D. R. Wood, *Graph product structure for non-minor-closed classes*. 2019. [arXiv:1907.05168](https://arxiv.org/abs/1907.05168)
- [29] J. Eckhoff, *The maximum number of triangles in a K_4 -free graph*. *Discrete Math.* 194(1999), nos. 1–3, 95–106.
- [30] J. Eckhoff, *A new Turán-type theorem for cliques in graphs*. *Discrete Math.* 282(2004), nos. 1–3, 113–122.
- [31] J. Engbers and D. Galvin, *Counting independent sets of a fixed size in graphs with a given minimum degree*. *J. Graph Theory* 76(2014), no. 2, 149–168.
- [32] D. Eppstein, *Connectivity, graph minors, and subgraph multiplicity*. *J. Graph Theory* 17(1993), no. 3, 409–416.
- [33] D. Eppstein and S. Gupta, *Crossing patterns in nonplanar road networks*. In: Proc. 25th ACM SIGSPATIAL Int'l Conf. on Advances in Geographic Information Systems, 40, 2017, pp. 1–9.
- [34] P. Erdős and R. Rado, *Intersection theorems for systems of sets*. *J. London Math. Soc. Second Ser.* 35(1960), no. 1, 85–90.
- [35] P. Erdős and A. H. Stone, *On the structure of linear graphs*. *Bull. Amer. Math. Soc.* 52(1946), 1087–1091.
- [36] B. Ergemlidze, A. Methuku, N. Salia, and E. Győri, *A note on the maximum number of triangles in a C_5 -free graph*. *J. Graph Theory* 90(2019), no. 3, 227–230.
- [37] S. Fiorini, G. Joret, D. O. Theis, and D. R. Wood, *Small minors in dense graphs*. *Eur. J. Combin.* 33(2012), no. 6, 1226–1245.
- [38] D. C. Fisher and J. Ryan, *Conjectures on the number of complete subgraphs*. In: Proc. 20th Southeastern Conf. on Combinatorics, Graph Theory, and Computing, 70, pp. 217–219. 1990.
- [39] D. C. Fisher and J. Ryan, *Bounds on the number of complete subgraphs*. *Discrete Math.* 103(1992), no. 3, 313–320.
- [40] J. Foisy, *Intrinsically knotted graphs*. *J. Graph Theory* 39(2002), no. 3, 178–187.
- [41] F. V. Fomin, D. Lokshantov, and S. Saurabh, *Bidimensionality and geometric graphs*. In: Proc. 23rd Annual ACM-SIAM Symp. on Discrete Algorithms, 2012, pp. 1563–1575.
- [42] F. V. Fomin, S.-I. Oum, and D. M. Thilikos, *Rank-width and tree-width of H -minor-free graphs*. *Eur. J. Combin.* 31(2010), no. 7, 1617–1628.
- [43] J. Fox and F. Wei, *On the number of cliques in graphs with a forbidden minor*. *J. Combin. Theory Ser. B* 126(2017), 175–197.
- [44] J. Fox and F. Wei, *On the number of cliques in graphs with a forbidden subdivision or immersion*. *SIAM J. Discrete Math.* 34(2020), no. 4, 2556–2582.
- [45] A. Frohmader, *A Kruskal-Katona type theorem for graphs*. *J. Combin. Theory Ser. A* 117(2010), no. 1, 17–37.
- [46] D. Galvin, *Two problems on independent sets in graphs*. *Discrete Math.* 311(2011), no. 20, 2105–2112.
- [47] W. Gan, P.-S. Loh, and B. Sudakov, *Maximizing the number of independent sets of a fixed size*. *Combin. Probab. Comput.* 24(2015), no. 3, 521–527.
- [48] D. Gerbner, E. Győri, A. Methuku, and M. Vizer, *Generalized Turán problems for even cycles*. *J. Combin. Theory Ser. B* 145(2020), 169–213.
- [49] D. Gerbner, A. Methuku, and M. Vizer, *Generalized Turán problems for disjoint copies of graphs*. *Discrete Math.* 342(2019), no. 11, 3130–3141.
- [50] D. Gerbner and C. Palmer, *Counting copies of a fixed subgraph in F -free graphs*. *Eur. J. Combin.* 82(2019), 103001.
- [51] F. Goldberg and A. Berman, *On the Colin de Verdière number of graphs*. *Linear Algebra Appl.* 434(2011), no. 7, 1656–1662.
- [52] N. Goldberg, T. W. Mattman, and R. Naimi, *Many, many more intrinsically knotted graphs*. *Algebr. Geom. Topol.* 14(2014), no. 3, 1801–1823.
- [53] E. Győri, A. Paulos, N. Salia, C. Tompkins, and O. Zamora, *The maximum number of paths of length three in a planar graph*. Preprint 2019. [arXiv:1909.13539](https://arxiv.org/abs/1909.13539)
- [54] E. Győri, A. Paulos, N. Salia, C. Tompkins, and O. Zamora, *The maximum number of pentagons in a planar graph*. Preprint 2019. [arXiv:1909.13532](https://arxiv.org/abs/1909.13532)
- [55] E. Győri, N. Salia, C. Tompkins, and O. Zamora, *The maximum number of P_ℓ copies in P_k -free graphs*. *Discrete Math. Theor. Comput. Sci.* 21(2019), 14.
- [56] J. Harant, M. Horňák, and Z. Skupień, *Separating 3-cycles in plane triangulations*. *Discrete Math.* 239(2001), nos. 1–3, 127–136.
- [57] D. J. Harvey and D. R. Wood, *Parameters tied to treewidth*. *J. Graph Theory* 84(2017), no. 4, 364–385.

- [58] B. Hedman, *The maximum number of cliques in dense graphs*. Discrete Math. 54(1985), no. 2, 161–166.
- [59] T. Huynh, G. Joret, and R. D. Wood, *Subgraph densities in a surface*, Preprint, 2020, [arXiv:2003.13777](https://arxiv.org/abs/2003.13777)
- [60] J. Kahn, *An entropy approach to the hard-core model on bipartite graphs*. Combin. Probab. Comput. 10(2001), no. 3, 219–237.
- [61] N. Khadzhiivanov and N. Nenov, *Sharp estimates of the highest number of cliques of a graph*. Annuaire Univ. Sofia Fac. Math. Méc. 70(1975/76), 23–26.
- [62] R. Kirsch and A. J. Radcliffe, *Many triangles with few edges*. Electron. J. Combin. 26(2019), no. 2, P2.36.
- [63] S. G. Kobourov, G. Liotta, and F. Montecchiani, *An annotated bibliography on 1-planarity*. Comput. Sci. Rev. 25(2017), 49–67.
- [64] D. König, *Theorie der endlichen und unendlichen Graphen*. Kombinatorische Topologie der Streckenkomplexe, Akademische Verlagsgesellschaft, Leipzig, 1936.
- [65] A. V. Kostochka, *Lower bound of the Hadwiger number of graphs by their average degree*. Combinatorica 4(1984), no. 4, 307–316.
- [66] C. Lee and S.-I. Oum, *Number of cliques in graphs with a forbidden subdivision*. SIAM J. Disc. Math. 29(2015), no. 4, 1999–2005.
- [67] S. Letzter, *Many H -copies in graphs with a forbidden tree*. SIAM J. Discrete Math 33(2019), no. 4, 2360–2368.
- [68] S. Louis Hakimi, *On the degrees of the vertices of a directed graph*. J. Franklin Inst. 279(1965), 290–308.
- [69] S. Louis Hakimi and E. F. Schmeichel, *On the number of cycles of length k in a maximal planar graph*. J. Graph Theory 3(1979), no. 1, 69–86.
- [70] S. Louis Hakimi and E. F. Schmeichel, *Bounds on the number of cycles of length three in a planar graph*. Israel J. Math. 41(1982), nos. 1–2, 161–180.
- [71] R. Luo, *The maximum number of cliques in graphs without long cycles*. J. Combin. Theory Ser. B 128(2018), 219–226.
- [72] J. Ma and Y. Qiu, *Some sharp results on the generalized Turán numbers*. Eur. J. Combin. 84(2020), 103026.
- [73] W. Mader, *Homomorphiesätze für Graphen*. Math. Ann. 178(1968), 154–168.
- [74] B. Mohar and C. Thomassen, *Graphs on surfaces*. Johns Hopkins University Press, 2001.
- [75] R. Montgomery, *Logarithmically small minors and topological minors*. J. London Math. Soc. 91(2015), no. 1, 71–88.
- [76] J. Nešetřil and P. O. de Mendez, *How many F 's are there in G ?* Eur. J. Combin. 32(2011), no. 7, 1126–1141.
- [77] J. Nešetřil and P. O. de Mendez, *Sparsity*. Algorithms and Combinatorics, 28, Springer, 2012.
- [78] J. Nešetřil and P. O. de Mendez, *On low tree-depth decompositions*. Graphs Combin 31(2015), no. 6, 1941–1963.
- [79] S. Norine, P. Seymour, R. Thomas, and P. Wollan, *Proper minor-closed families are small*. J. Combin. Theory Ser. B 96(2006), no. 5, 754–757.
- [80] J. Pach and G. Tóth, *Recognizing string graphs is decidable*. Discrete Comput. Geom. 28(2002), no. 4, 593–606.
- [81] C. H. Papadimitriou and M. Yannakakis, *The clique problem for planar graphs*. Inform. Process. Lett. 13(1981), nos. 4–5, 131–133.
- [82] L. Petingi and J. Rodriguez, *A new proof of the Fisher-Ryan bounds for the number of cliques of a graph*. In: Proc. 31st Southeastern Int'l Conf. on Combinatorics, Graph Theory and Computing, Congr. Numer., 146, 2000, pp. 143–146.
- [83] J. L. Ramírez Alfonsín, *Knots and links in spatial graphs: a survey*. Discrete Math 302(2005), nos. 1–3, 225–242.
- [84] B. Reed and D. R. Wood, *A linear time algorithm to find a separator in a graph excluding a minor*. ACM Trans. Algorithms 5(2009), no. 4, 39.
- [85] B. A. Reed, *Tree width and tangles: a new connectivity measure and some applications*. In: R. A. Bailey (ed.), Surveys in combinatorics, London Math. Soc. Lecture Note Ser., 241, Cambridge Univ. Press, 1997, pp. 87–162.
- [86] G. Ringel, *Das Geschlecht des vollständigen paaren Graphen*. Abh. Math. Sem. Univ. Hamburg 28(1965), 139–150.
- [87] N. Robertson, P. Seymour, and R. Thomas, *A survey of linkless embeddings*. In: N. Robertson and P. Seymour (eds.), Graph structure theory, Proc. AMS-IMS-SIAM Joint Summer Research Conf. on Graph Minors, Contemporary Math., 147, Amer. Math. Soc., 1993, pp. 125–136.

- [88] N. Robertson, P. Seymour, and R. Thomas, *Sachs' linkless embedding conjecture*. J. Combin. Theory Ser. B 64(1995), 185–227.
- [89] H. Sachs, On a spatial analogue of Kuratowski's theorem on planar graphs — an open problem. In: M. Borowiecki, J. W. Kennedy, and M. M. Syslo (eds.), Proc. Conf. on Graph Theory, Lecture Notes in Math., 1018, Springer, 1983, pp. 230–241.
- [90] A. Schrijver, *Minor-monotone graph invariants*. In: Surveys in combinatorics, London Math. Soc. Lecture Note Ser., 241, Cambridge Univ. Press, 1997, pp. 163–196.
- [91] A. Shapira and B. Sudakov, *Small complete minors above the extremal edge density*. Combinatorica 35(2015), no. 1, 75–94.
- [92] M. Shimabara, *Knots in certain spatial graphs*. Tokyo J. Math 11(1988), no. 2, 405–413.
- [93] A. Thomason, *An extremal function for contractions of graphs*. Math. Proc. Cambridge Philos. Soc 95(1984), no. 2, 261–265.
- [94] P. Turán, *On an extremal problem in graph theory*. Mat. Fiz. Lapok 48(1941), 436–452.
- [95] W. C. van Batenburg, T. Huynh, G. Joret, and J.-F. Raymond, *A tight Erdős-Pósa function for planar minors*. Adv. Combin. 2(2019).
- [96] H. van der Holst, L. Lovász, and A. Schrijver, *The Colin de Verdière graph parameter*. In: Graph theory and combinatorial biology, Bolyai Soc. Math. Stud., 7, János Bolyai Math. Soc., 1999, pp. 29–85.
- [97] D. R. Wood, *On the maximum number of cliques in a graph*. Graphs Combin. 23(2007), no. 3, 337–352.
- [98] N. C. Wormald, *On the frequency of 3-connected subgraphs of planar graphs*. Bull. Austral. Math. Soc. 34(1986), no. 2, 309–317.
- [99] A. A. Zykov, *On some properties of linear complexes*. Mat. Sbornik N.S. 24(1949), no. 66, 163–188.

School of Mathematics, Monash University, Melbourne, Australia

e-mail: tony.bourbaki@gmail.com david.wood@monash.edu