


## Standard Paper

# *Catillaria flexuosa* (Catillariaceae), a new lichen species described from the Netherlands

Pieter P. G. van den Boom<sup>1</sup>  and Pablo Alvarado<sup>2</sup>

<sup>1</sup>Arafura 16, 5691 JA, Son, the Netherlands and <sup>2</sup>Alvalab, Dr Fernando Bongera St., Severo Ochoa Bldg S1.04, 33006 Oviedo, Spain

### Abstract

A new lichen species is described from specimens growing on *Fraxinus* trees north of Eindhoven (the Netherlands). Morphological and genetic studies suggest that the new species belongs in the genus *Catillaria*, and the name *Catillaria flexuosa* is proposed because of its flexuose apothecia. The new species is characterized by the relatively large apothecia (up to 0.9 mm diam.) and relatively thick, knobby to  $\pm$ subsquamulose, greenish, thallus. Due to their similar morphological features, *C. flexuosa* can be easily confused with *Catillaria chalybeia*, *C. fungoides* or *C. nigroclavata*, so it is therefore compared with these species. In addition, *Arthonia epiphyscia* is reported being a very rare species in the Netherlands.

**Key words:** Ascomycota, Benelux, *Leprocaulales*, mtSSU, taxonomy, 28S nrDNA

(Accepted 9 November 2020)

### Introduction

A long and intensive study of lichens in the south-east part of the Noord-Brabant Province (the Netherlands) resulted in multiple interesting taxonomic novelties (e.g. van den Boom 2004, 2015; van den Boom & van den Boom 2009). Recently, a collection was found growing on a medium-sized trunk of *Fraxinus excelsior*. It was thought to be a species of *Catillaria* A. Massal. because it presented paraphyses with black tips, a feature typically found in this genus, as well as relatively small and hyaline 1-septate ascospores.

The first genetic data from the type species of *Catillaria*, *C. chalybeia* (Borr.) A. Massal., was provided by Bendiksby & Timdal (2013), who showed that this species is closely related to the genera *Solenopsora* A. Massal. (type species *S. requienii* A. Massal.) and *Lopadium* Körb. (type species *L. pezizoideum* (Ach.) Körb.), forming a monophyletic clade outside the main group of *Lecanorales* Nannf. Miadlikowska *et al.* (2014) identified this clade as *Leprocaulales* Lendemer & B. P. Hodk., an invalid name due to the lack of a description or diagnosis in Lendemer & Hodkinson (2013), where this clade was shown to include also samples identified as *Halecania alpivaga* (Th. Fr.) M. Mayrhofer (the type species of *Halecania* M. Mayrhofer) and *Speerschneidera euploca* (Tuck.) Trevis. (the type species of *Speerschneidera* Trevis.).

According to the genetic results obtained by several authors (Andersen & Ekman 2005; Reese Næsborg *et al.* 2007; Ekman *et al.* 2008; Lee *et al.* 2008; Kistenich *et al.* 2018), the traditional concept of the genus *Catillaria* is not monophyletic. Several species belong in unrelated families according to Miadlikowska *et al.* (2014); namely *Catillaria alba* Coppins & Vězda, *C. aphana* (Nyl.) Coppins,

*C. corymbosa* (Hue) M. Lamb, *C. croatica* Zahlbr. and *C. scotinoides* (Nyl.) Coppins in *Ramalinaceae*, *C. contristans* (Nyl.) Zahlbr. in *Pilocarpaceae*, *C. erysiboides* (Nyl.) Th. Fr. in *Psilolechiaceae*, and *C. modesta* (Müll. Arg.) Coppins in *Lecideaceae*. Kelly *et al.* (2011) produced molecular data from specimens identified as *C. nigroclavata* and later Guttová *et al.* (2014) obtained data from other samples identified as *C. lenticularis*, but they could not provide any hypothesis about their placement at the family level.

Three species of *Catillaria* have been described from Australia and Tasmania recently and from the type locality only: *Catillaria austrolittoralis* Kantvilas & van den Boom (Kantvilas & van den Boom 2013) and *C. laevigata* P. M. McCarthy & Elix (Elix & McCarthy 2018), two saxicolous species with a rimose or areolate thallus, containing argopsin or pannarin; *C. gerroana* P. M. McCarthy & Elix (McCarthy & Elix 2017), which lacks these chemical compounds and has ascospores of  $(9-11.5(-15) \times (3.5-4.5(-5.5) \mu\text{m})$ . Another saxicolous species, *C. ulleungdoensis* S. Y. Kondr. *et al.*, was recently described from South Korea; it has a weakly developed thallus and rather small apothecia (0.15–0.3(–0.55) mm diam.) (Konratyuk *et al.* 2016). *Catillaria patteeana* D. P. Waters & Lendemer has brown apothecia with an opaque white to light brown margin and is sorediate (Waters & Lendemer 2019). Two corticolous species, *C. praedicta* Tretiach & Hafellner and *C. subpraedicta* M. Brand & van den Boom, known from Mediterranean areas and Lanzarote (Canary Islands) respectively, have multispored asci.

In the present work, the taxonomic identity of the new samples found with *Fraxinus* in Noord-Brabant is resolved.

### Materials and Methods

#### Morphological study and literature

All specimens collected were studied by conventional macro- and microscopical techniques after mounting hand-cut sections of the

**Author for correspondence:** Pieter P. G. van den Boom. E-mail: pvdboom@kpnmail.nl

**Cite this article:** van den Boom PPG and Alvarado P (2021) *Catillaria flexuosa* (Catillariaceae), a new lichen species described from the Netherlands. *Lichenologist* 53, 193–202. <https://doi.org/10.1017/S0024282921000050>

**Table 1.** Species names, voucher specimens and GenBank Accession numbers of sequences of *Catillaria* and related taxa used in the phylogenetic analyses. Samples in bold were sequenced in the present study. Sequences were obtained mainly from Gaya *et al.* (2012), Lendemer & Hodkinson (2013), Guttová *et al.* (2014) and Prieto & Wedin (2016).

Species name	Voucher	Accession numbers				
		ITS	LSU	mtSSU	Mcm7	Tub2
<i>Acolium inquinans</i>	Wedin 6352 (UPS)	AY450583	AY453639	AY143404	JX000161	-
<i>Acroscyphus sphaerophoroides</i>	Obermayer 6077 (UPS)	KX512898	-	KX512984	KX529029	-
<i>Allocalcium adaequatum</i>	Spribille 14143 (UPS)	KX512906	KX512859	KX512986	-	KX528996
<i>Amandinea punctata</i>	Wedin 2/3/96 (UPS) / AFTOL-ID 1306	KX512899	DQ986756	DQ986874	KX529025	-
<i>Anaptychia ciliaris</i>	Wedin 6429 (UPS)	AY143391	KX512894	-	KX529054	-
<i>Athallia scopularis</i>	Søchting 7521 (C), USE197	KC179340	KC179150	KC179480	-	-
<i>Baculifera remensa</i>	Prieto (S)	-	KX512881	KX512962	-	-
<i>Bryobilimbia diapensiae</i>	Arup L04400	HQ650660	HQ660539	HQ660564	-	-
<i>Buellia disciformis</i>	Wedin 6155 (BM)	AY143392	JX000082	AY143401	JX000152	-
<i>Calicium viride</i>	Wedin 24/4 2000 / Prieto 3015 (S)	HQ650703	AY340538	AY143402	JX000153	KX529013
<i>Calogaya decipiens</i>	USE15, 1995 Søchting (C) / Arup L06187 / JV7964	KC179344	KC179167	KC179498	-	KY748807
<i>Caloplaca cerina</i>	Elvebakk 03-084 (TROM)	KC179425	KC179168	KC179499	-	-
<b><i>Catillaria chalybeia</i></b>	<b>P.v.d. Boom 59147 (hb. P.v.d. Boom)</b>	<b>MT248987</b>	<b>MT248985</b>	-	-	-
<b><i>C. chalybeia</i></b>	<b>P.v.d. Boom 59646 (hb. P.v.d. Boom)</b>	<b>MW045826</b>	<b>MW045829</b>	<b>MW077133</b>	-	-
<i>C. chalybeia</i>	R. Haugan 7947 / 155291 (O-L)	KF360370	KF360447	-	-	-
<b><i>C. flexuosa</i></b>	<b>P.v.d. Boom 58587 (holotype, B)</b>	<b>MT248986</b>	<b>MT248984</b>	-	-	-
<b><i>C. flexuosa</i></b>	<b>P.v.d. Boom 59636 (hb. P.v.d. Boom)</b>	<b>MW045827</b>	<b>MW045830</b>	-	<b>MW049361</b>	-
<i>C. lenticularis</i>	Fackovcova 21-Oct-2011 (SAV)	KF689898	-	-	-	-
<i>C. nigroclavata</i>	EDNA09-02107 (E)	FR799313	-	-	-	-
<i>C. nigroclavata</i>	EDNA09-02108 (E)	FR799312	-	-	-	-
<i>C. nigroclavata</i>	G.M. 2015-07-09.5 (LUX)	MT912697	-	-	-	-
<b><i>Catillaria</i> sp.</b>	<b>P.v.d. Boom 55102 (hb. P.v.d. Boom)</b>	<b>MW045825</b>	<b>MW045828</b>	-	-	-
<i>Culbersonia nubila</i>	ALV14224	-	MH121319	-	-	-
<i>Cyphelium tigillare</i>	Prieto 3038 (S)	JX000104	JX000088	JX000123	JX000162	KX529002
<i>Dimelaena oreina</i>	Lendemer 4193 (S)	KX512922	KX512867	KX512976	KX529036	KX528999
<i>Diplotomma alboatrum</i>	Prieto 3034 (S)	KX512924	KX512877	KX512966	KX529043	KX529007
<i>Dirinaria applanata</i>	Seaward 109735 (S)	-	KX512856	DQ972983	-	-
<i>Fulgensia fulgens</i>	Gaya 39 (BCN)	JQ301671	JQ301567	JQ301503	-	-
<i>Halecania alpivaga</i>	195847 (O-L)	KY266832	-	AY756368	-	-
<i>Heterodermia speciosa</i>	Wetmore 88030 (S)	KX512927	JX000089	KX512975	JX000166	KX529000
<i>Leprocaulon adhaerens</i>	NY351, Lendemer 13546 (NY)	KC184100	-	-	-	-
<i>L. adhaerens</i>	NY598, Morse 16380 (NY)	KC184101	-	-	-	-
<i>L. americanum</i>	NY184, Knudsen 9605 (NY)	KC184112	-	KC183999	-	-
<i>L. americanum</i>	NY185, Lendemer 11445 (NY)	KC184111	-	-	-	-
<i>L. calcicola</i>	E2080	KX674677	-	-	-	-
<i>L. calcicola</i>	E2117, C.J.B. Hitch (type, NMW)	NR_158498	-	-	-	-
<i>L. knudsenii</i>	NY62, Lendemer 11476 (NY)	KC184115	-	-	-	-
<i>L. knudsenii</i>	NY702, Knudsen 9256 (type, NY)	KC184076	-	-	-	-
<i>L. microscopicum</i>	Ekman 3685 (BG)	AF517921	-	-	-	-

(Continued)

Table 1. (Continued)

Species name	Voucher	Accession numbers				
		ITS	LSU	mtSSU	<i>Mcm7</i>	<i>Tub2</i>
<i>L. quisquiliare</i>	98GB04	MN959918	-	-	-	-
<i>L. quisquiliare</i>	L-200884 (O)	MK812636	-	-	-	-
<i>L. quisquiliare</i>	NY1261, Buck 55960 (NY)	-	KJ766593	KC183993	-	-
<i>L. santamonicae</i>	NY1264, Knudsen 12058 (NY)	KC184105	-	KC183995	-	-
<i>L. santamonicae</i>	NY565, Lendemmer 19660 (NY)	KC184108	-	-	-	-
<i>L. terricola</i>	NY355, Knudsen 2659.2 (NY)	KC184102	-	KC184017	-	-
<i>L. terricola</i>	NY655, Knudsen 9608 (NY)	KC184103	-	-	-	-
<i>L. textum</i>	NY57, Lendemmer 11500 (NY)	KC184113	-	-	-	-
<i>Letrouitia domingensis</i>	AFTOL-ID 102	HQ650700	AY584648	AY584619	-	-
<i>Megalospora sulphurata</i>	Kantvilas 376/09 (F)	JQ693555	JQ693538	JQ693518	-	-
<i>Phaeophyscia ciliata</i>	Prieto (S)	KX512929	KX512886	KX512958	KX529051	KX529012
<i>Physcia tenella</i>	Odelvik & Hellström 0827 (S)	KX512932	KX512869	KX512974	KX529038	-
<i>Physconia muscigena</i>	17040 (BCN-Lich)	GU247161	-	GU247195	-	-
<i>Polycauliona polycarpa</i>	USE599, Söchting 10507 (C)	KC179389	KC179222	KC179558	-	-
<i>Pseudothelomma ocellatum</i>	Tehler 8063 (S)	KX512934	KX512862	KX512957	KX529028	KX529019
<i>Pyxine soreliata</i>	Wetmore 91254 (S)	KX512937	JX000093	KX512973	JX000179	KX529001
<i>Rusavskia elegans</i>	MP78, Odelvik 04532 (S)	KX512947	KX512896	KX512948	KX529056	KX529024
<i>Solenopsis candicans</i>	D. Ertz 7604 (DUKE) / Guttova (SAV)	KF689797	KJ766660	KJ766493	KF689899	-
<i>S. cesatii</i>	cesatii_19_FR / cesatii_10_FR (SAV)	KF689850	-	-	KF689933	-
<i>S. cesatii</i>	cesatii_9_SK (SAV)	KF689849	-	-	-	-
<i>S. chihuahuana</i>	BC51, L-15843b (O)	MN148640	-	-	-	-
<i>S. crenata</i>	BC67, Nash-38664b (ASU)	MN148641	-	-	-	-
<i>S. grisea</i>	grisea_7_IT (SAV)	KF689874	-	-	-	-
<i>S. grisea</i>	grisea_9_AL (BP)	KF689875	-	-	-	-
<i>S. grisea</i> subsp. <i>grisea</i>	grisea_4_MK / grisea_11_ME (SAV)	KF689871	-	-	KF689970	-
<i>S. holophaea</i>	1865	AM292708	-	-	-	-
<i>S. holophaea</i>	holophaea_1_FR (SAV)	KF689876	-	-	-	-
<i>S. liparina</i>	liparina_1_FR (SAV)	KF689877	-	-	-	-
<i>S. liparina</i>	liparina_2_FR (SAV)	KF689878	-	-	-	-
<i>S. liparina</i>	liparina_3_CZ/ 2014_6l (SAV)	KF689879	-	-	MN166068	-
<i>S. marina</i>	marina_5_TR (SAV)	KF689884	-	-	-	-
<i>S. marina</i>	Slovak & Kucera 27-May-2012 (SAV)	KF689885	-	-	KF689981	-
<i>S. olivacea</i>	olbiensis_1_IT (SAV)	KF689886	-	-	-	-
<i>S. olivacea</i>	olivacea_2_HR (SAV)	KF689888	-	-	-	-
<i>S. olivacea</i>	Timdal 21-Jul-2010 (O)	KF689889	-	-	KF689990	-
<i>S. vulturiensis</i>	0290344 (M)	MH269330	-	-	-	-
<i>S. vulturiensis</i>	98GB03	MN959915	-	-	-	-
<i>S. vulturiensis</i>	vulturiensis_3_FR (SAV)	KF689896	-	-	-	-
<i>S. vulturiensis</i>	vulturiensis_4_FR (SAV)	KF689897	-	-	KF689998	-
<i>Speerschneidera euploca</i>	Egan 14906 (F)	-	AY300862	AY300912	-	-
<i>Teloschistes flavicans</i>	E. Gaya 02.26.10-8 & F. Lutzoni (DUKE) / Frödén 1624 (LD)	KT291472	KC179255	KC179594	-	-

(Continued)

**Table 1.** (Continued)

Species name	Voucher	Accession numbers				
		ITS	LSU	mtSSU	<i>Mcm7</i>	<i>Tub2</i>
<i>Tetramelas pulverulentus</i>	Nordin 6368 (UPS)	KX512940	KX512860	KX512983	-	KX528990
<i>Texosporium sancti-jacobi</i>	Rosentreter & De Bolt 6514 (UPS)	KX512941	KX512863	JX000132	JX000187	KX528994
<i>Thelomma mammosum</i>	Tibell 23775 (UPS)	KX512942	KX512888	KX512954	KX529065	KX529016
<i>Tholurna dissimilis</i>	Wedin 6330 (UPS)	AY143397	KX512893	AY143407	KX529053	KX528992
<i>Tornabea scutellifera</i>	Tibell 23833 (UPS)	KX512946	KX512873	KX512970	KX529042	-
<i>Xanthocarpia ochracea</i>	Gaya 21, P. Navarro-Rosines (BCN)	-	JQ301558	JQ301494	-	-
<i>Xanthodactylon flammeum</i>	36/1940 (HSBG)	EU681315	-	EU680897	-	-
<i>Xanthomendoza mendozae</i>	Søchting 10209 (C)	KC179138	KC179281	KC179620	-	-
<i>Xanthoria parietina</i>	Gaya 8 (BCN)	JQ301691	JQ301589	-	-	-

material in tap water. Amyloid reactions were tested using Lugol's iodine solution (K/I). Accompanying specimens were identified. Photographs of habitus and microscopic characters are provided for the new species. Voucher specimens are kept in the herbarium of the first author and some specimens, including the holotype specimen, have been deposited in the herbarium of the Botanischer Garten und Botanisches Museum Berlin-Dahlem (B). Differences between corticolous species of *Catillaria* s. str. are summarized in Table 2 (data obtained from Kiliyas (1981), Tretiach & Hafellner (1998), van den Boom & Etayo (2001), Smith *et al.* (2009), van den Boom (2010); and from corticolous *Catillaria chalybeia* specimens in this study).

### Phylogenetic studies

Total DNA was extracted from dry specimens employing a modified protocol based on Murray & Thompson (1980). PCR reactions (Mullis & Faloona 1987) included 35 cycles with an annealing temperature of 54 °C. The primers ITS1F and ITS4 (White *et al.* 1990; Gardes & Bruns 1993) were employed to amplify the ITS rDNA region (ITS), LR0R and LR5 (Vilgalys & Hester 1990; Cubeta *et al.* 1991) were used for the 28S rDNA region (LSU), mrSSU1 and mrSSU3R for the mitochondrial small ribosomal subunit (mtSSU), and finally MCM7-709for and MCM7-1348rev for the DNA replication licensing factor (*Mcm7*). PCR products were checked in 1% agarose gels and positive reactions were sequenced with one or both PCR primers. Chromatograms were checked, searching for putative reading errors, and these were corrected. BLAST (Altschul *et al.* 1990) was used to select the most closely related sequences from the International Nucleotide Sequence Database Collaboration public database (INSDC; Cochrane *et al.* 2011). Two independent datasets were built. The first alignment comprised sequence data of ITS2, LSU, mtSSU and *Mcm7* (exons only) from the samples analyzed. Also included were homologous sequences from other species in the *Leprocaulaceae* and *Catillariaceae* families retrieved from public databases, and representatives from the orders *Caliciales* and *Teloschistales*, using *Bryobilimbia diapensiae* (Th. Fr.) Fryday *et al.* as outgroup because of its external position relative to these clades. The purpose of this analysis was to ascertain whether or not the newly generated sequences belong in *Leprocaulales*, as several species of *Catillaria* have been combined into different

families and orders in the past. In addition, *tub2* sequences available from the species in the alignment were incorporated, making a total of five partitions. Sequences (Table 1) came mainly from Gaya *et al.* (2012), Lendemer & Hodkinson (2013), Guttová *et al.* (2014) and Prieto & Wedin (2016). Each partition was aligned in MEGA 5.0 (Tamura *et al.* 2011) with its Clustal W application and then corrected manually. The final alignment included 126/266 (ITS2,  $n = 59$ ), 251/652 (LSU,  $n = 44$ ), 339/602 (mtSSU,  $n = 43$ ), 217/443 (*Mcm7*,  $n = 27$ ) and 217/624 (*tub2*,  $n = 15$ ) variable/total sites. The aligned partitions were loaded in MrBayes v.3.2.6 (Ronquist *et al.* 2012), where a Bayesian analysis was performed (data partitioned, GTR + G model for all partitions, two simultaneous runs, six chains, temperature set to 0.2, sampling every 100th generation) until convergence parameters were met after 1.27 M generations, standard deviation having fallen below 0.01. Finally, a full search for the best-scoring maximum likelihood (ML) tree was performed in RAXML v.8.2.12 (Stamatakis 2014) using the standard search algorithm (same partitions, GTRCAT model, 2000 bootstrap replications). The significance threshold was set above 0.95 for posterior probability (PP) and 70% bootstrap proportions (BP).

The second alignment included only ITS sequences of *Leprocaulaceae* and *Catillariaceae* (using *Catillaria lenticularis* as outgroup because of its external position outside *Catillaria* s. str.). The purpose of this analysis was to resolve the different species in *Catillaria* s. str. and compare their intraspecific variability with that observed in other species of these families. Sequences (Table 1) came mainly from the same sources. The sequences were first aligned in MEGA 5.0 and then subjected to GBLOCKS (Castresana 2000) to remove 345 ambiguously aligned sites, resulting in a new dataset with 204/458 variable sites among 44 sequences. This alignment was loaded in MrBayes v.3.2.6, where a Bayesian analysis was performed (data not partitioned, GTR + G + I model, two simultaneous runs, four chains, temperature set to 0.2, sampling every 100th generation) until convergence parameters were met after 0.2 M generations, standard deviation having fallen below 0.01. A full search for the best-scoring ML tree was performed in RAXML v.8.2.12 using the standard search algorithm (data not partitioned, GTRGAMMA model, 2000 bootstrap replications). The significance threshold was set again at 0.95 for posterior probability (PP) and 70% bootstrap proportions (BP).

**Table 2.** Comparison of the main characters of the six known corticolous *Catillaria* s. str. species. Information obtained from Killias (1981), Tretiach & Hafellner (1998), van den Boom & Etayo (2001), Smith et al. (2009), van den Boom (2010) and the present study.

Species	Thallus	Apothecia (mm)	Margin	Disc	Ascospores ( $\mu\text{m}$ )	Pycnidia ( $\mu\text{m}$ ); conidia ( $\mu\text{m}$ )	Ecology
<i>C. chalybeia</i>	thin, smooth to somewhat scurfy, grey-brown to very pale brown, $\pm$ slightly shiny	0.2–0.5	thin, concolorous with the disc, rarely paler	black or bluish black	9.5–12.2 $\times$ 3.1–4.1; 8 per ascus	50–80 (–100); 1.8–3.5 $\times$ 0.5–0.8 (–1.2)	saxicolous or corticolous only on <i>Fraxinus</i> trees
<i>C. flexuosa</i>	thick, uneven, knobby, $\pm$ subsquamulose, dark green, with $\pm$ pale brown tinge, matt	0.3–0.9	thick, often flexuose, concolorous with the disc	black	(6–)7–10(–11) $\times$ 2.5–3.5(–4); 8 per ascus	100–150; 2–3 $\times$ 0.9–1.2	corticolous only on trunks of <i>Fraxinus</i> trees
<i>C. fungoides</i>	very thin, smooth, whitish to pale grey, soredate, shiny	0.2–0.4	thin, concolorous or slightly paler than the disc, finally excluded	black	10–12 $\times$ 3–3.5; 8 per ascus	unknown	corticolous on a wide range of phorophytes
<i>C. nigroclavata</i>	very thin, smooth, greyish, shiny	0.1–0.3	thin, concolorous or slightly paler than the disc	dark brown to black	8–10 $\times$ (2–)2.5(–4); 8 per ascus	unknown	corticolous on a wide range of phorophytes
<i>C. praedicta</i>	smooth to rimose-areolate, pale brown to dark brown or olivaceous, matt	0.25–0.5(–0.8)	thick, concolorous with the disc	brown to almost black	(4.5–)5.5–7.2(–9) $\times$ (2–)2.2–3(–3.5); 24–32 per ascus	100–150; 2.5–3 $\times$ 1.5	corticolous or lignicolous, on a wide range of phorophytes
<i>C. subpraedicta</i>	scurfy to farinose spongy granular; pale to dark brown, matt	0.2–0.4(–0.5)	thin, somewhat paler than the disc	dark grey to blackish	6.8–7.9 $\times$ 2.7–2.8; 16 per ascus	c. 75; 2.6–2.8 $\times$ 1.1	corticolous on a wide range of phorophytes

## Results and Discussion

According to the present phylogenetic studies (Figs 1 & 2), the specimens studied are closely related to *Catillaria chalybeia*, the type species of *Catillaria*, represented by sequences obtained from saxicolous (*R. Haugan* 7947/ O-L-155291) and corticolous (*P. v. d. Boom* 59147, 59646) samples. A saxicolous collection found on the Azores Islands (*P. v. d. Boom* 55102) seems to be unrelated to the other lineages, being c. 60% similar in ITS, (475/782 bp identical), or 90% if a large 250 bp insertion in the ITS1 region is excluded, and is provisionally identified here as *Catillaria* sp. A third clade is composed of sequences of two collections with flexuose apothecia and differences of only 1.5% between them (11/784 bp), but both being c. 91% similar to sequences of *C. chalybeia* (712/781 bp). Regarding LSU, the three lineages differ in c. 2–3% (10–20/645 bp different), with a very low intraspecific variability (2/645 bp variable sites in both *C. chalybeia* and the clade with flexuose apothecia). In turn, all these samples are related to specimens identified as *C. nigroclavata* (G.M. 2015-07-09.5 (LUX), EDNA09-02107 (E) and EDNA09-02108 (E)), although the ITS sequence of EDNA09-02108 (E) seems to differ in up to 4.5% (21/477 bp) from the other two, maybe indicating that they come from different taxa. While the exact identity of *C. chalybeia* needs to be further confirmed by checking additional collections and appointing an epitype, the specimens with flexuose apothecia clearly represent a distinct taxon, a new name for which is proposed here.

All these species form a significantly supported clade that is considered here to be the core of *Catillaria*. Unfortunately, the genetic limits between *Catillaria* and the closely related genera *Halecania*, *Leprocaulon*, *Solenopsora* and *Speerschneidera* are not clear enough, probably due to the incomplete genetic data from most collections analyzed (Table 1). Synonymies between some of these genera cannot be rejected and, in this case, *Catillaria* (A. Massal. 1852) should be the priority name. In the present analysis, *C. lenticularis* seems to be unrelated to the core of the genus *Catillaria*. The limits between the families *Catillariaceae* and *Leprocaulaceae* are not clear either, maybe also being synonyms, in which case *Catillariaceae* (Hafellner 1984) should be given priority.

## Taxonomy


### *Catillaria flexuosa* van den Boom & P. Alvarado sp. nov.

Mycobank No.: MB 835063

Distinguished from *Catillaria chalybeia* by the dark green, relatively thick thallus, up to 0.4 mm, continuous to weakly rimose, areolate, with knobby granules, sometimes becoming subsquamulose. Apothecia 0.3–0.9 mm diam., often flexuose; smaller ascospores, (6–)7–10(–11)  $\times$  2.5–3.5(–4)  $\mu\text{m}$ , larger pycnidia 100–150  $\mu\text{m}$  diam., and differing conidia, ellipsoid, 2–3  $\times$  0.9–1.2  $\mu\text{m}$ .

Type: The Netherlands, Noord-Brabant Prov., S of Veghel, S of Zijtaart, *Fraxinus* trees (medium size) along road, grid ref. 45-56-51, 51°35.3'N, 5°32.5'E, 10 m, 12 August 2019, *P. van den Boom* 58587 (B—holotype; hb. van den Boom—isotype). GenBank Accession nos: MT248986 (ITS rDNA), MT248984 (28S rDNA).

(Fig. 3A–C)




**Fig. 1.** A 50% majority rule ITS rDNA-28S rDNA-mtSSU-Mcm7-tub2 consensus phylogram of the family Catillariaceae (with a selection of species from the orders Teloschistales and Caliciales, as well as *Bryobilimbia diapensiae* as outgroup) obtained in MrBayes from 23 925 sampled trees. Nodes were annotated if supported by  $\geq 0.95$  Bayesian posterior probability (PP) (left) or  $\geq 70\%$  maximum likelihood bootstrap proportions (BP) (right). Sequences newly generated in this study are in bold. In colour online.

*Thallus* relatively thick, up to 0.4 mm, continuous to weakly rimose, areolate, with knobby granules, sometimes becoming subsquamulose, growing in patches up to 1 cm wide, dark green, often with a brownish tinge, matt; *prothallus* not present; *photobiont* chlorophycean, cells 5–15  $\mu\text{m}$  diam.

*Apothecia* abundantly present, 0.3–0.9 mm diam., flat, rarely slightly convex; *proper margin* conspicuous, presenting as a

small rim especially in young apothecia, often flexuose, especially in mature apothecia (30–60  $\mu\text{m}$  wide), persistent, somewhat shiny; *disc* black, matt; *true exciple* dark brown to blackish, sometimes with greenish pigments, mainly in young apothecia; *hymenium* 40–50  $\mu\text{m}$  high, not presenting oil droplets; *epithecium* dark brown to greenish black, mainly in young apothecia, without crystals, K–; *hypothecium* dark




**Fig. 2.** A 50% majority rule ITS rDNA consensus phylogram of the family Catillariaceae (with a selection of species from Leprocaulaceae, as well as *Catillaria lenticularis* as outgroup) obtained in MrBayes from 1500 sampled trees. Nodes were annotated if supported by  $\geq 0.95$  Bayesian posterior probability (PP) (left) or  $\geq 70\%$  maximum likelihood bootstrap proportions (BP) (right). Sequences newly generated in this study are in bold. In colour online.

brown, K–; *paraphyses* simple to rarely sparingly branched, 1–1.5(–2)  $\mu\text{m}$  wide, septate, apices dark brown to blackish, widened, 3–6  $\mu\text{m}$  diam.; *asci* cylindrical-clavate to clavate, 8-spored, *Catillaria*-type, with a uniformly blue apical dome in K/I, 35–40  $\times$  8–12  $\mu\text{m}$ ; *ascospores* ellipsoid, (6–)7–10(–11)  $\times$

2.5–3.5(–4)  $\mu\text{m}$ , 1-septate, thin walled, often with oil droplets, not or only rarely slightly constricted at septum.

*Pycnidia* often present, immersed to somewhat erumpent, 100–150  $\mu\text{m}$  diam., dark brown to blackish; *conidia* ellipsoid, hyaline, 2–3  $\times$  0.9–1.2  $\mu\text{m}$ .



**Fig. 3.** *Catillaria flexuosa* (holotype, P. van den Boom 58587). A, habitus. B, section of apothecium. C, paraphyses and some ascospores. Scales: A = 1 mm; B = 100 µm; C = 20 µm. In colour online.

**Chemistry.** No chemical compounds detected by TLC. No crystals in thallus or apothecia.


**Etymology.** The epithet refers to the habitus of the apothecia.

**Ecology and distribution.** This new species is frequent in the type locality, growing on bark of wayside *Fraxinus excelsior* trees. It has a rather small distribution, being known from only five localities in the area, in the east of the province of Noord-Brabant. Other lichen species occurring at the type locality include *Candelaria concolor* (Dicks.) Stein, *Candelariella reflexa* (Nyl.) Lettau, *C. vitellina* (Hoffm.) Müll. Arg., *Catillaria nigroclavata* (Nyl.) Schuler, *Phaeophyscia nigricans* (Flörke) Moberg, *P. orbicularis* (Neck.) Moberg, *Physcia adscendens* H. Olivier, *P. caesia*

(Hoffm.) Fűrnr., *Physconia grisea* (Lam.) Poelt and *Xanthoria parietina* (L) Th. Fr. In addition, several lichenicolous fungi were found, including *Arthonia epiphyscia* Nyl. (very rare in the Netherlands), *A. parietinaria* Hafellner & A. Fleischhacker and *Lichenochora weillii* (Werner) Hafellner & R. Sant.

**Notes.** The morphological features of *Catillaria chalybeia* are described in detail by Kiliás (1981), who chose the lectotype from a collection found by Borrer, on tiles and flints, housed in BM. The thallus is usually 80–200 µm thick, rimose-areolate to verrucose-areolate, areoles 0.1–0.4 mm wide, sometimes very poorly developed, beige to olive-brown, dark brown to black, often slightly shiny, and the prothallus is often not present or dark brown to black. Apothecia are usually 0.2–0.5 mm diam.,





**Fig. 4.** *Catillaria chalybeia* (corticolous, *P. van den Boom* 59192), habitus. Scale = 0.5 mm. In colour online.

constricted at the base, the disc dark brown to black, excipulum black at outer rim, paler at inner side, asci 30–50 × 10–15 µm, ascospores 9.5–12.2 × 3.1–4.1 µm, pycnidia 50–80(–100) µm diam., and conidia bacilliform, 1.8–3.5 × 0.5–0.8 (–1.2) µm. *Kiliás (1981)* listed many records from all over Europe.

*Catillaria chalybeia* is commonly found growing saxicolous in the Netherlands, Belgium and Luxemburg (*Diederich et al. 2020*), but in the present work it was also found corticolous (*Fig. 4*). Many saxicolous collections from the study area have been checked; they usually grew on brick and on schist, always appeared with a thin, smooth, rimose areolate, greyish to greyish brown slightly shiny thallus, and apothecia that are up to 0.5 mm diam., roundish, without a flexuose margin, often clustered, some slightly deformed. Corticolous specimens differ somewhat in thallus colour, with a more brownish tinge.

*Catillaria flexuosa* could be easily confused with *C. chalybeia* but differs by its matt, relatively thick greenish thallus, larger apothecia (up to 0.9 mm wide) often with a flexuose margin, smaller ascospores, and somewhat thicker conidia. In the field, *C. flexuosa* as well as the corticolous *C. chalybeia* form small and scattered inconspicuous patches, not exceeding 1 cm in width, and with no more than a few patches on isolated tree trunks.

*Catillaria nigroclavata* is also very common in the distribution area of the new species and both grow close to each other, and with *C. chalybeia*. *Catillaria nigroclavata* has much smaller apothecia, measuring c. 0.1–0.3 mm diam., and a very thin to interspersed thallus with a greyish tinge. It is often found widely scattered on tree bark, but also on branches. *Catillaria fungoides* Etayo & van den Boom is also common in the studied area. Its apothecia resemble those of *C. flexuosa* but they have a thin, somewhat shiny thallus, and characteristic abundant black soralia; further differences to the aforementioned and to *Catillaria prae-dicta* Tretiach & Hafellner and *C. subpraedicta* M. Brand & van den Boom are provided in *Table 2*. In the field, *C. flexuosa* can be easily confused with *Amandinea punctata* (Hoffm.) Coppins & Scheid., a species with quite similar apothecia but with a more conspicuous, greyish thallus and brown ascospores.

**Additional specimens examined. The Netherlands:** Noord-Brabant: SSW of Erp, S of Keldonk, Rad to Diepers, *Fraxinus* trees (medium size) along road, grid ref. 51-16-14, 51°35'N, 5°

35.1'E, 10 m, 2019, *P. & B. van den Boom* 59120 (hb. v. d. Boom); W of St Oedenrode, S of road to Olland, road to Bobbenagelse Brug, corticolous on *Fraxinus* tree, grid ref. 51-14-34, 51°34.1'N, 5°25.8'E, 10 m, 2019, *P. van den Boom* 59211 (hb. v. d. Boom); N of Uden, De Maashorst, Menzel, Menzelsch Veld, Koudenoord, roadside *Fraxinus* trees along field, grid ref. 45-36-34, 51°41.9'N, 5°35'E, 15 m, 2020, *N. Ettema & P. van den Boom* 59636 (hb. v. d. Boom); Veghel, NE of city, W of road to Mariaheide, *Fraxinus* trees along small road, just outside housing estate, grid ref. 45-56-13, 51°37.55'N, 5°34.3'E, 10 m, 2020, *P. & B. van den Boom* 59644 (hb. v. d. Boom).

***Catillaria chalybeia* (corticolous), specimens examined. The Netherlands:** Noord-Brabant: W of St Oedenrode, S of road to Olland, road to Bobbenagelse Brug, on *Fraxinus* tree, grid ref. 51-14-34, 51°34.1'N, 5°25.8'E, 10 m, 2019, *P. van den Boom* 59147 (hb. v. d. Boom); Breugel, Eind, near crossing to centre of village and road to Lieshout, roadside trees (*Tilia* and *Fraxinus*), on *Fraxinus*, grid ref. 51-25-35, 51°31.2'N, 5°31.4'E, 2019, *P. & B. van den Boom* 59192 (hb. v. d. Boom); Veghel, NE of city, W of road to Mariaheide, on *Fraxinus* trees along small road, just outside housing estate, grid ref. 45-56-13, 51°37.55'N, 5°34.3'E, 10 m, 2020, *P. van den Boom* 59646 (hb. v. d. Boom).

***Catillaria chalybeia* (saxicolous), selected specimens examined. The Netherlands:** Noord-Brabant: Eindhoven, NNW of Geldrop, road to Urkhoven, on brick of culvert, grid. ref. 51-46-31, 51°26.0'N, 5°32.2'E, 20 m, 1993, *P. & B. van den Boom* 13805 (hb. v. d. Boom); Valkenswaard centre, churchyard since 1887, tombstones of concrete, brick, schist etc., on schist, grid. ref. 57-14-25, 51°21.1'N, 5°27.5'E, 20 m, 2009, *P. & B. van den Boom* 43474 (hb. v. d. Boom); Eindhoven centre, churchyard, gravestones of concrete and schist, on schist, grid. ref. 51-44-15, 51°27.1'N, 5°27.5'E, 20 m, 2014, *P. & B. van den Boom* 51920 (hb. v. d. Boom).

***Catillaria* sp. (saxicolous), specimen examined. Portugal:** Azores: Faial, W side of island, NNE of Praia do Norte, Baia da Ribeira das Cabras, coastal area, some scattered trees, outcrops and walls with volcanic stones along field, on volcanic stone, 38°36.72'N, 28°45.74'W, 35 m, 2016, *P. & B. van den Boom* 55102 (hb. v. d. Boom).

***Arthonia epiphyscia*, specimens examined. The Netherlands:** Noord-Brabant: S of Veghel, S of Zijtaart, *Fraxinus* trees (medium size) along road, grid ref. 45-56-51, 51°35.3'N, 5°32.5'E, 10 m, 2019, *P. van den Boom* 58591 (hb. v. d. Boom); N of Uden, De Maashorst, Menzel, Menzelsch Veld, Koudenoord, roadside *Fraxinus* trees along field, grid ref. 45-36-34, 51°41.9'N, 5°35'E, 15 m, 2020, *N. Ettema & P. van den Boom* 59640 (hb. v. d. Boom).

**Acknowledgement.** We would like to thank Harrie Sipman (Berlin), who provided TLC analysis.

**Author ORCID.**  Pieter P. G. van den Boom, 0000-0002-1929-2088.

## References

- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Andersen HL and Ekman S (2005) Disintegration of the *Micareaeaceae* (lichenized Ascomycota): a molecular phylogeny based on mitochondrial rDNA sequences. *Mycological Research* **109**, 21–30.

- Bendiksby M and Timdal E** (2013) Molecular phylogenetics and taxonomy of *Hypocnemomyce* sensu lato (Ascomycota: Lecanoromycetes): extreme polyphyly and morphological/ecological convergence. *Taxon* **62**, 940–956.
- Castresana J** (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* **17**, 540–552.
- Cochrane G, Karsch-Mizrachi I and Nakamura Y on behalf of the International Nucleotide Sequence Database Collaboration** (2011) The International Nucleotide Sequence Database Collaboration. *Nucleic Acids Research* **39**, D15–D18.
- Cubeta MA, Echandi E, Abernethy T and Vilgalys R** (1991) Characterization of anastomosis groups of binucleate *Rhizoctonia* species using restriction analysis of an amplified ribosomal RNA gene. *Phytopathology* **81**, 1395–1400.
- Diederich P, Ertz D, Stapper N, Sérusiaux E, Van den Broeck D, van den Boom P and Ries C** (2020) *The lichens and lichenicolous fungi of Belgium, Luxembourg and northern France*. [WWW resource] URL <http://www.lichenology.info> [Accessed 3 January 2020].
- Ekman S, Andersen HL and Wedin M** (2008) The limitations of ancestral state reconstruction and the evolution of the ascus in the *Lecanorales* (lichenized Ascomycota). *Systematic Biology* **57**, 141–156.
- Elix JA and McCarthy PM** (2018) Ten new lichen species (Ascomycota) from Australia. *Australasian Lichenology* **82**, 20–59.
- Gardes M and Bruns TD** (1993) ITS primers with enhanced specificity for Basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113–118.
- Gaya E, Högnabba F, Holguin Á, Molnar K, Fernández-Brime S, Stenroos S, Arup U, Söchting U, van den Boom P and Lücking R** (2012) Implementing a cumulative supermatrix approach for a comprehensive phylogenetic study of the *Teloschistales* (Pezizomycotina, Ascomycota). *Molecular Phylogenetics and Evolution* **63**, 374–387.
- Guttová A, Zozomová-Lihová J, Timdal E, Kučera J, Slovák M, Píknová K and Paoli L** (2014) First insights into genetic diversity and relationships of European taxa of *Solenopsisora* (*Catillariaceae*, Ascomycota) with implications for their delimitation. *Botanical Journal of the Linnean Society* **176**, 203–223.
- Hafellner J** (1984) Studien in Richtung einer natürlicheren Gliederung der Sammelfamilien *Lecanoraceae* und *Lecideaceae*. *Beiheft zur Nova Hedwigia* **79**, 241–371.
- Kantvilas G and van den Boom PPG** (2013) A new species of *Catillaria* (lichenised Ascomycetes: *Catillariaceae*) from southern Australia. *Journal of the Adelaide Botanic Gardens* **26**, 5–8.
- Kelly LJ, Hollingsworth PM, Coppins BJ, Ellis CJ, Harrold P, Tosh J and Yahr R** (2011) DNA barcoding of lichenized fungi demonstrates high identification success in a floristic context. *New Phytologist* **191**, 288–300.
- Kiliás R** (1981) Revision gestinsbewohnender Sippen der Flechtengattung *Catillaria* Massal. in Europa (*Lecanorales*, *Lecideaceae*). *Herzogia* **5**, 209–448.
- Kistenich S, Timdal E, Bendiksby M and Ekman S** (2018) Molecular systematics and character evolution in the lichen family *Ramalinaceae* (Ascomycota: *Lecanorales*). *Taxon* **67**, 871–904.
- Kondratyuk S, Lőkös L, Tschabanenko S, Moniri MH, Farkas E, Wang X, Oh S-O and Hur J-S** (2016) New and noteworthy lichen-forming and lichenicolous fungi: 5. *Acta Botanica Hungarica* **58**, 319–396.
- Lee JS, Lee HK, Hur J-S, Andreev M and Hong SG** (2008) Diversity of the lichenized fungi in King George Island, Antarctica, revealed by phylogenetic analysis of partial large subunit rDNA sequences. *Journal of Microbiology and Biotechnology* **18**, 1016–1023.
- Lendemer JC and Hodkinson BP** (2013) A radical shift in the taxonomy of *Lepraria* s.l.: Molecular and morphological studies shed new light on the evolution of asexuality and lichen growth form diversification. *Mycologia* **105**, 994–1018.
- McCarthy PM and Elix JA** (2017) Five new lichen species (Ascomycota) and a new record from southern New South Wales, Australia. *Telopea* **20**, 335–353.
- Miadlikowska J, Kauff F, Högnabba F, Oliver JC, Molnár K, Fraker E, Gaya E, Hafellner J, Hofstetter V, Gueidan C, et al.** (2014) A multigene phylogenetic synthesis for the class Lecanoromycetes (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. *Molecular Phylogenetics and Evolution* **79**, 132–168.
- Mullis K and Faloona FA** (1987) Specific synthesis of DNA *in vitro* via a polymerase-catalyzed chain reaction. *Methods in Enzymology* **155**, 335–350.
- Murray MG and Thompson WF** (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research* **8**, 4321–4325.
- Prieto M and Wedin M** (2016) Phylogeny, taxonomy and diversification events in the *Caliciaceae*. *Fungal Diversity* **82**, 221–238.
- Reese Næsberg R, Ekman S and Tibell L** (2007) Molecular phylogeny of the genus *Lecania* (*Ramalinaceae*, lichenized Ascomycota). *Mycological Research* **111**, 581–591.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, and Huelsenbeck JP** (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**, 539–542.
- Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW and Wolseley PA** (eds) (2009) *The Lichens of Great Britain and Ireland*. London: British Lichen Society.
- Stamatakis A** (2014) RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M and Kumar S** (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**, 2731–2739.
- Tretiaich M and Hafellner J** (1998) A new species of *Catillaria* from coastal Mediterranean regions. *Lichenologist* **30**, 221–229.
- van den Boom PPG** (2004) A long-term inventory of lichens and lichenicolous fungi of the Strabrechtse Heide and Lieropse Heide in Noord-Brabant, The Netherlands. *Österreichische Zeitschrift für Pilzkunde* **13**, 131–151.
- van den Boom PPG** (2010) Lichens and lichenicolous fungi from Lanzarote (Canary Islands), with the descriptions of two new species. *Cryptogamie, Mycologie* **31**, 183–199.
- van den Boom PPG** (2015) Lichens and lichenicolous fungi from graveyards of the area of Eindhoven (the Netherlands), with the description of two new species. *Annalen des Naturhistorischen Museums in Wien. Serie B, Botanik und Zoologie* **117**, 245–276.
- van den Boom PPG and Etayo J** (2001) Two new sorediate species of lichens in the *Catillariaceae* from the Iberian Peninsula. *Lichenologist* **33**, 103–110.
- van den Boom P and van den Boom B** (2009) Diversity of lichens and lichenicolous fungi in a primeval heathland and adjacent managed forest in southern Netherlands (Groote Heide and 't Leenderbos). *Österreichische Zeitschrift für Pilzkunde* **18**, 25–45.
- Vilgalys R and Hester M** (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**, 4238–4246.
- Waters DP and Lendemer JC** (2019) The lichens and allied fungi of Mercer County, New Jersey. *Opuscula Philolichenum* **18**, 17–51.
- White TJ, Bruns TD, Lee S and Taylor JW** (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In Innis MA, Gelfand DH, Sninsky J and White TJ (eds), *PCR Protocols: a Guide to Methods and Applications*. San Diego: Academic Press, pp. 315–322.