

Original Article

Cite this article: Jacobina UP, Torres RA, de Mello Affonso PRA, dos Santos EV, Calado LL, de Araújo Bitencourt J (2020). DNA barcoding reveals cryptic diversity and peculiar phylogeographic patterns in mojarras (Perciformes: Gerreidae) from the Caribbean and South-western Atlantic. *Journal of the Marine Biological Association of the United Kingdom* **100**, 277–283. <https://doi.org/10.1017/S0025315419001206>

Received: 25 July 2019

Revised: 25 November 2019

Accepted: 2 December 2019

First published online: 30 January 2020


Key words:

COI; cryptic species; estuaries; *Eucinostomus*; MOTU; Western Atlantic

Author for correspondence:

Uedson Pereira Jacobina,
E-mail: uedson.jacobina@penedo.ufal.br

DNA barcoding reveals cryptic diversity and peculiar phylogeographic patterns in mojarras (Perciformes: Gerreidae) from the Caribbean and South-western Atlantic

Uedson Pereira Jacobina¹ , Rodrigo Augusto Torres², Paulo Roberto Antunes de Mello Affonso³, Ewerton Vieira dos Santos¹, Leonardo Luiz Calado⁴ and Jamille de Araújo Bitencourt³

¹Laboratório de Ictiologia e Conservação, Campus-Penedo/Universidade Federal de Alagoas, Avenida Beira Rio s/n, Penedo CEP 57200-000, Alagoas, Brazil; ²Laboratório de Genômica Evolutiva e Ambiental, Departamento de Zoologia, Universidade Federal de Pernambuco, Av Prof. Nelson Chaves s/n, Cidade Universitária, CEP 50670-420, Recife, Pernambuco, Brasil; ³Department of Biological Sciences, Universidade Estadual do Sudoeste da Bahia, Av. José Moreira Sobrinho, s/n, Jequiezinho, 45206190, Jequié, Bahia, Brazil and ⁴Laboratório de Genética de Recursos Marinhos, Universidade Federal do Rio Grande do Norte, Av. Senador Salgado Filho, S/N, Campus Universitário, 59078970, Natal, Rio Grande do Norte, Brazil

Abstract

The mojarras (*Eucinostomus*) are a widespread group of coastal fishes of controversial taxonomy because of similarities in their external morphology. In the present study, we assessed the genetic diversity of species and populations of *Eucinostomus* using DNA barcodes using a systematic and phylogeographic context. In total, 416 COI sequences of all valid *Eucinostomus* representatives were analysed based on public databases and collected specimens from the north-eastern coast of Brazil (Western South Atlantic). Several cases of misidentification were detected in the barcode dataset (*E. argenteus*, *E. harengulus*, *E. gula*, *E. dowii* and *E. jonesii*) that could account for the taxonomic issues in this genus. In contrast, we identified four molecular operational taxonomic units (MOTUs), with divergence above 2% in the Western Atlantic, that correspond to cryptic forms within *E. argenteus*, *E. harengulus*, *E. gula* and *E. melanopterus*. These data suggest that Plio-Pleistocene events (rise of the Panama isthmus, Amazonas outflow and sea-level fluctuations) played a major role in the diversification of mojarras. While subtle morphological differences have been used as proxies to discriminate *Eucinostomus* species, the genetic data proved to be efficient in differentiating them and revealing potentially undescribed taxa. Therefore, we recommend that further taxonomic studies in mojarras should incorporate DNA-based evidence.

Introduction

The apparent weakness of geographic barriers over large distances in marine environments hinders the identification of cladogenesis and several cases of cryptic lineages might remain overlooked (Rocha, 2003; Luiz *et al.*, 2012; Da Silva *et al.*, 2016). Moreover, considering that speciation processes might take place without major changes in morphology, the actual number of species might be underestimated when external body features are analysed in isolation (Winker, 2005). Failure of species diagnosis has a particularly negative impact on economically important species in fisheries where distinct morphotypes putatively related to a single species might encompass multiple evolutionary units (Mayden, 1997; Winker, 2005; Da Silva *et al.*, 2018).

The mojarras (genus *Eucinostomus*, family Gerreidae) comprise 11 nominal taxa (Froese & Pauly, 2019) widespread in the Western Atlantic (WA; seven species) and Eastern Pacific (EP; four species) that have commercial relevance to local fisheries. These fish are found in coastal areas, including estuaries and hypersaline lakes because of their high osmoregulation adaptability (Nelson *et al.*, 2016). Morphologically, they are characterized by dorsal compression, protractile mouth, smooth gill bones, cycloid scales on the head and ctenoid scales over the body (Nelson *et al.*, 2016). From a systematic viewpoint, the genus *Eucinostomus* is considered one of the most problematic genera of coastal fishes from the New World since their putative interspecific morphological traits are subtle and usually overlapped, while their evolutionary interrelationships remain poorly known (Matheson & McEachran, 1984; De La Cruz-Agüero & Galvan-Magana, 1993; De La Cruz-Agüero, 2013). Because of their controversial taxonomy, some species in this group, such as *Ulaema lefroyi* (*E. lefroyi*), are regarded as *inquirendae* (Nelson *et al.*, 2016). In addition, other taxonomic methods such as cytogenetic analyses have been inefficient in discriminating mojarra species (Calado *et al.*, 2014).

The difficulties in recognizing *Eucinostomus* species based on traditional taxonomy were recently reported in a wide study of DNA barcoding in fish species from north-eastern Brazil, WA (Brandão *et al.*, 2016). These authors verified that 82 specimens of mojarras



morphologically identified as *Eucinostomus melanopterus* or only to the genus level (*Eucinostomus* sp.) encompassed five molecular units with deep genetic divergence, representing *E. melanopterus* (N = 5), *E. lefroyi* (N = 7), *E. gula* (N = 2), *E. harengulus* (N = 64) and *E. jonesii* (N = 4). Moreover, this work represented the first report about the occurrence of *E. harengulus* and *E. jonesii* in north-eastern Brazil.

The accurate diagnosis of taxa is a key step to biodiversity conservation (Bickford *et al.*, 2007). The utilization of mitochondrial markers such as the Cytochrome c oxidase subunit I (COI) as DNA barcodes has been particularly useful in the identification of cryptic species and in the resolution of taxonomic uncertainties, with several examples in fishes (Ferreira *et al.*, 2014; Hyde *et al.*, 2014; Winterbottom *et al.*, 2014; Barreira *et al.*, 2016; Nirchio *et al.*, 2018). This approach has also been improved over the last decade by the incorporation of distinct algorithms to test the hypothesis of evolutionary independent lineages and to avoid synonyms in the dataset (Brown *et al.*, 2012; Puillandre *et al.*, 2012; Zhang *et al.*, 2013; Luo *et al.*, 2018).

Considering the problematic systematics, the wide range of mojaras and the efficiency of DNA-based identification, we assessed the genetic diversity of all *Eucinostomus* representatives with available COI sequences throughout most of their geographic distribution. Based on distinct analytical methods, including species delimitation algorithms, we reviewed their systematic relationships within a phylogeographic context, providing evidence of overlooked phylogenetic diversity and variation in the range of evolutionary units, with emphasis in the WA.

Materials and methods

Sampling

A total of 400 COI sequences of *Eucinostomus*, comprising 11 recognized taxa were analysed: *E. argenteus* (N = 90), *E. cf. gula* (N = 6), *E. currany* (N = 7), *E. dowii* (N = 6), *E. entomelas* (N = 10), *E. gracilis* (N = 2), *E. gula* (N = 49), *E. havana* (N = 2), *E. harengulus* (N = 88), *E. jonesii* (N = 86), *E. lefroyi* (N = 11) and *E. melanopterus* (N = 20), as well as other sequences named as *Eucinostomus* sp. (N = 23). These sequences were downloaded from the public datasets, for example Bold Systems and GenBank (NCBI) (supplementary material 1). We also added 16 specimens of mojaras, *E. argenteus* (N = 1), *E. gula* (N = 8), *E. jonesii* (N = 2), *E. lefroyi* (N = 5), collected in the state of Rio Grande do Norte (05°05'26''S 36°16'031W''), north-eastern Brazilian coast, that were identified according to Woodland (2006). Fish were collected by gillnets and euthanasia was accomplished by immersion in cold water for 10–15 min. Afterwards, muscle tissues were removed from each individual and stored in ethanol at –20 °C (Blessing *et al.*, 2010).

DNA isolation, amplification and sequencing

Total DNA was isolated from stored muscle tissues, using the DNeasy kit (QIAGEN). A fragment of 645 base pairs (bp) of the COI gene was amplified via PCR using the primers VF2_t1 (5'TGTAACACGACGGCCAGTCAACCAACCACAAAGACAT-TGGCAC3') and FishR2_t1 (5'CAGGAAACAGCTATGACACTCAGGGTGACCGAAGAATCAGAA3') as described by Ward *et al.* (2005). The reactions encompassed 12 µl of 2× Taq master mix (Vivantis), 2 µl of template DNA solution at 40 ng µl⁻¹, 0.5 µl of each primer (10 mM) and ultrapure water to a final volume of 25 µl. The PCR steps (adapted from Ward *et al.*, 2005) included a first denaturation at 95°C for 2 min, 35 cycles at 94°C (30 s), 57°C (30 s) and 72°C (2 min), plus a final extension at 72°C for 7 min. The PCR products were purified with

ExoSap IT enzymatic system (Affimetrix). The sequencing of COI fragments was carried out using the BigDyeTM Terminator v 3.1 Cycle Sequencing Ready Reaction kit (Applied Biosystems) using the M-13 initiator followed by reading in ABI 3130 automatic sequencer (Applied Biosystems).

The electropherograms were checked and edited in the software Geneious (Kearse *et al.*, 2012), followed by visual inspection of consensus sequences for final edition adjustments. In addition, the COI sequences were uploaded and deposited in the BOLD platform under the project 'Assessing the genetic diversity of species *Eucinostomus* – EUCI', being automatically assigned to a Barcode Index Number – BIN (group of sequences that should correspond to a single taxon), following the analytical procedures of Ratnasingham & Hebert (2013). Afterwards, these sequences were aligned with those available in Bold Systems and GenBank by using the ClustalW method.

Phylogenetic and distance analyses

Phylogenetic reconstructions were carried out based on Maximum likelihood (ML) and Bayesian inference (BI). The best evolutionary model for both ML and BI trees was HKY + I as indicated in the software PartitionFinder (Lanfear *et al.*, 2012). Branch support in ML analysis was based on 1000 bootstrap replicates using RAxML (Stamatakis *et al.*, 2012). In the case of BI, ultrametric trees following the Yule Speciation prior model were generated in the software BEAST 1.8.4 (Drummond & Rambaut, 2007), based on 20 million generations with sampling at every 2000 generations. The convergence of Markov chains was inspected in Tracer 1.6 (Drummond *et al.*, 2012). All values of Effective Sample Size (ESS) were above 200. Based on 10% of burn-in, the remaining trees were used to obtain a consensus tree and the branch support was based on the posterior probability values.

Based on BI and ML results, we calculated the genetic distance of each cluster with support values higher than 1 of probability or 95% of bootstrap to build a Neighbour-joining (NJ) tree in the software MEGA6 (Tamura *et al.*, 2013) using the Kimura 2-parameter (K2P) evolutionary model as identified by the Barcode of Life initiative (www.boldsystems.org). All trees were visualized using FigTree v. 1.4.1 (Rambaut & Drummond, 2009).

Species delimitation methods

To establish a potential threshold among *Eucinostomus* species, we built a distance matrix based on the K2P model using the function sppDistMatrix available in the R package SPIDER v1.3–0 (Brown *et al.*, 2012). Based on this matrix, we created a density object with the minimum local function which disregards any previous knowledge about the species identity to indicate potential thresholds to infer intra- and interspecific variation levels (Brown *et al.*, 2012). We also used three widely used algorithms for species delimitation from molecular data: the Bayesian Poisson Tree Process (bPTP) (Zhang *et al.*, 2013), the generalized mixed Yule-coalescent model (GMYC) (Pons *et al.*, 2006) and the Automatic Barcode Gap Discovery (ABGD) (Puillandre *et al.*, 2012).

The bPTP method was carried out in the web server <<http://species.h-its.org/ptp/>> using as input a non-ultrametric tree based on ML inference. This analysis was performed with 500,000 generations with sampling at every 500 generations and 10% of burn-in. The GMYC was implemented in the web server <<http://species.h-its.org/gmyc/>>, based on an ultrametric tree obtained in BEAST (Drummond *et al.*, 2012).

The ABGD was carried out based on the pairwise genetic distances (based on p-distance, K2P and Jukes–Cantor models). The

analysis was done using a gap width value of 1.0 for all distances using the software available in <<http://wwwabi.snv.jussieu.fr/public/abgd/>>. The congruence among the delimitation of MOTUs was evaluated by comparing the clusters inferred from each algorithm.

Population analysis

Haplotype networks were built in the software Popart (Leigh & Bryant, 2015) using the Median-joining network algorithm to elucidate the genealogical relationships of *E. argenteus* and *E. gula*, because both species encompassed a high number of sequences from distinct localities throughout most of their range.

Results

The final alignment of COI sequences comprised 593 bp. Insertions, deletions or stop codons were absent, indicating that all sequences correspond to functional COI genes and not to putative pseudogenes or nuclear mitochondrial DNA segments (numts).

Phylogenetic analyses combined with distance methods and tools of molecular identification were helpful in identifying 34 potentially misidentified sequences in the analysed dataset of *Eucinostomus*, related to *E. argenteus*, *E. harengulus*, *E. jonesii*, *E. dowii* and *E. gula* (see supplementary material 2). The genetic variation in these samples ranged from 0.2–1.7%. Therefore, these sequences were reallocated to their respective groups and analysed according to their actual taxonomic classification to avoid biased evolutionary inferences. Sequences with divergence above 2%, which did not fit into any of the described taxonomic species of *Eucinostomus* were included with the acronym ‘cf.’ They are: *E. cf. argenteus*, *E. cf. melanopterus*, *E. cf. harengulus* and *E. cf. gula*.

The BI and ML analyses recovered 15 MOTUs distributed into highly supported clades 1/>>95, as follows: *E. jonesii*, *E. dowii*, *E. cf. argenteus*, *E. harengulus*, *E. cf. harengulus*, *E. gracilis*, *E. gula*, *E. cf. gula*, *E. entomelas*, *E. argenteus*, *E. lefroyi*, *E. melanopterus*, *E. cf. melanopterus*, *E. currani* and *E. havana* (Figure 1). The species delimitation algorithms (GMYP, bPTP, BINs and ABGD) also separated these clusters, besides revealing four additional MOTUs within *E. gula*, *E. argenteus*, *E. melanopterus* and *E. harengulus* (Figure 1). However, only the barcode index (BINs) was not found for *E. cf. harengulus*, since these sequences are only available in NCBI.

According to the distance method based on the K2P evolutionary model, the highest genetic distance was observed between *E. havana* and *E. argenteus* (21.8%), while the lowest value was detected between *E. cf. melanopterus* and *E. melanopterus* (2.9%). Two sequences erroneously assigned to *E. argenteus* were closely related to *E. harengulus* with 3.3% genetic divergence, being thus referred to as *E. cf. harengulus*. The highest values of intraspecific genetic variation were observed in *E. gula* (1.6%) and *E. cf. gula* (1.2%), while values close to 0% were observed in most of the *Eucinostomus* representatives (Table 1). The threshold potential from intra- to interspecific variation in mojarras as inferred by the minimal local function using Spider was established as 2.7% (Figure 1). This value was in agreement with the species delimitation algorithms where sequences with genetic distances above 2.7% were recovered as distinct MOTUs.

Haplotype networks based on the most representative taxa with information about their geographic origin (*E. argenteus* and *E. gula*) revealed two haplogroups within each. The haplogroups composed of *E. argenteus* + *E. cf. argenteus* and *E. gula* + *E. cf. gula* were separated by 103 mutation steps (genetic distance = 20.7%) and 23 mutation steps (genetic distance = 4.4%), respectively (Figure 2). Particularly within *E. argenteus*/*E. cf. argenteus*, the range of each haplogroup corresponds to the Caribbean and Brazilian provinces.

Discussion

Besides identifying cryptic diversity, DNA barcoding analyses have been particularly effective in detecting synonyms and misidentifications among fish (Ferreira *et al.*, 2014; Hyde *et al.*, 2014). Accordingly, the pairwise divergence in COI sequences of *Eucinostomus* associated with comparative tools (BLASTn and Species Level Barcode Records) and phylogenetic inferences detected several cases of misidentification in the sequences available from Bold Systems and GenBank, as observed in *E. argenteus*, which has four BINs representing different species with the same morphological identification. It is possible to observe that two of these BINs include the species *E. jonesii*, *E. dowii*, *E. harengulus* and *Eucinostomus* sp. as well as a possible cryptic lineage for the coast of Brazil. Two clusters were also found for *E. gula*, *E. melanopterus* and *E. harengulus* species. For the latter sequences are only available on GenBank (see supplementary material and Figure 1). Furthermore, some sequences were




Fig. 1. Species delimitation using GMYP, bPTP, ABGD and BINs within *Eucinostomus*. The density related to the genetic distances among representatives of the genus *Eucinostomus* is shown below on the left side.

Table 1. Matrix of genetic distance based on K2P model among *Eucinostomus* species

Distance between species																
Species	Distance within species	HAR	HAR1	JON	GUL	GUL1	ARG	ARG1	ENT	DOW	HAV	CUR	LEF	MEL	MEL1	GRA
HAR	0.007	*														
HAR1	0.007	0.033	*													
JON	0.003	0.116	0.120	*												
GUL	0.016	0.144	0.134	0.138	*											
GUL1	0.012	0.151	0.141	0.143	0.044	*										
ARG	0.004	0.197	0.182	0.181	0.167	0.161	*									
ARG1	0.001	0.139	0.121	0.127	0.171	0.173	0.207	*								
ENT	0.001	0.165	0.164	0.150	0.175	0.165	0.103	0.181	*							
DOW	0.001	0.128	0.117	0.110	0.159	0.158	0.185	0.117	0.157	*						
HAV	n/c	0.203	0.183	0.185	0.211	0.201	0.218	0.207	0.190	0.190	*					
CUR	0.001	0.177	0.168	0.203	0.182	0.180	0.163	0.178	0.206	0.206	0.215	*				
LEF	0.004	0.152	0.145	0.146	0.162	0.175	0.165	0.155	0.155	0.166	0.207	0.176	*			
MEL	0.005	0.178	0.167	0.194	0.209	0.203	0.193	0.200	0.193	0.202	0.202	0.135	0.175	*		
MEL1	0.001	0.132	0.169	0.191	0.185	0.187	0.204	0.198	0.198	0.198	0.202	0.178	0.178	0.029	*	
GRA	0.000	0.170	0.127	0.135	0.129	0.142	0.182	0.168	0.132	0.132	0.192	0.151	0.151	0.182	0.174	*

HAR, *E. harengulus*; HAR1, *E. cf. harengulus*; JON, *E. jonesii*; ARG, *E. argenteus*; ARG1, *E. cf. argenteus*; GUL, *E. gula*; GUL1, *E. cf. gula*; CUR, *E. currani*; MEL, *E. melanopterus*; MEL1, *E. cf. melanopterus*; DOW, *E. dowii*; ENT, *E. entomelas*; GRA, *E. gracilis*.




Fig. 2. Map and haplotype network based on COI sequences in populations of *E. gula*, *E. cf. gula*, *E. argenteus* and *E. cf. argenteus* throughout their range in the Western Atlantic. The bars over lines indicate the number of mutations between the haplotypes.

identified only at genus level. In general, these sequences are derived from reports about larvae and juveniles of Caribbean fish without any previous information about COI sequences for comparative analysis, thus hindering their precise identification (Valdez-Moreno *et al.*, 2010; Weigt *et al.*, 2012). Indeed, even the morphological identification of adult stages in *Eucinostomus* species is still under debate (Matheson & McEachran, 1984; De La Cruz-Agüero & Galvan-Magana, 1993).

Another issue commonly observed in studies using DNA barcodes is the restricted range of sampling from putative widespread taxa (Ward *et al.*, 2005; de Ribeiro *et al.*, 2012; Brandão *et al.*, 2016). This issue restrains further biogeographic inferences, including the potential detection of phylopatric evolutionary lineages (Neves *et al.*, 2016). In this sense, the phylogenetic and species delimitation analyses carried out in the present study were particularly informative in assessing the actual diversity in *Eucinostomus*, since 15 distinct lineages (MOTUs) were clearly identified. Four MOTUs were related to cryptic forms in the formal taxa *E. argenteus*, *E. gula*, *E. melanopterus* and *E. harengulus* from the Western Atlantic (Caribbean and Brazilian provinces). Moreover, the tree topologies suggest distinct evolutionary histories for these lineages (Figure 1), as also supported by their high genetic diversity (20.7% for *E. argenteus*, 4.4% for *E. gula*, 3.3% for *E. harengulus* and 2.9% for *E. melanopterus*).

Similarly, molecular studies in distinct animal groups have shown the remarkable isolation of Caribbean lineages when compared with other regions (De Biasse *et al.*, 2016; Fields *et al.*, 2016; Hurtado *et al.*, 2017). Furthermore, recent reports have revealed that the biogeographic patterns in fishes from the Brazilian Province are quite heterogeneous, with growing evidence for high levels of endemism according to body size, dispersal routes and environmental features (Argolo *et al.*, 2018; Pinheiro *et al.*, 2018). Therefore, some of these vicariant effects could be responsible for the distinction among the lineages within the complex *E. argenteus*. In the Caribbean province, the great biodiversity reflects ancient and recent patterns of circulation and topography from the Miocene, when the Americas were not yet connected. Climatic changes during this period conditioned the opening and closing of currents at different depths that modified the

circulation pattern, and influenced global climate changes during the Pliocene (Williams & Duda, 2008; Williams *et al.*, 2013; Thacker, 2017). In addition, change in the flow of the Amazon on the north coast of South America between 10–6 mya and closure of the Isthmus of Panama between 5.5–3.6 mya might have, although temporarily, affected other taxa, such as *E. gula*, *E. melanopterus* and *E. harengulus*. Therefore, major diversification events during the Miocene–Pliocene could have become more and more accentuated during Pleistocene sea level fluctuations, leading to population isolation and divergence of cryptic MOTUs in *E. cf. argenteus*, *E. cf. melanopterus*, *E. cf. gula* and *E. cf. harengulus* (Lambeck & Chappell, 2001; Müller *et al.*, 2008).

It should also be pointed out that the divergence between the cryptic lineages in *E. argenteus* was seven times higher than that established as the threshold from intra- to interspecific variation (2.7%). Similar cases of deep variation in COI sequences have also been reported in marine fish such as *Scorpaena nonata* (18%; Landi *et al.*, 2014), lantern fishes (17–25%; Pappalardo *et al.*, 2015), Gonostomatidae, Sternoptychidae and Myctophidae (16–23%; Kenchington *et al.*, 2017). In fact, *E. argenteus* is a controversial taxon within *Eucinostomus*, being regarded as a putative species complex (Matheson & McEachran, 1984). Moreover, the present data diagnosed that most of misidentified sequences in public datasets are related to *E. argenteus*, bringing more noise to their taxonomic status that remains to be resolved. Besides these aspects, the close relationship between the cryptic lineages of *E. argenteus* from the Atlantic in relation to *E. entomelas* and *E. downi* from the Pacific (~10–12% of divergence) is also intriguing. This pattern reinforces the role of the Panama isthmus in cladogenetic events of the ichthyofauna from the Eastern Pacific and Western Atlantic (Bacon *et al.*, 2015; O’Dea *et al.*, 2016; Thacker, 2017).

In *Eucinostomus gula* a different scenario was observed. Even though the sampled region of *E. gula* overlaps with the cryptic lineages of *E. argenteus*, the genetic difference between the lineages in the former is 50% higher than the threshold of 2.7% for interspecific distinction, suggesting a more recent evolutionary history for the cryptic lineages of *E. gula*. The close relationship between both lineages in this formal taxon indicates a recent expansion of their range in the Brazilian Province.

The lack of shared haplotypes in the complex *E. gula* provides strong evidence for phylogeographic disjunction caused by allopatry. Therefore, *E. gula* is likely to encompass two Evolutionary Significant Units (ESUs), one of them representing an undescribed taxon since a divergence of 4.4% was observed in COI sequences in relation to the other lineages. Putatively, a Brazilian endemic lineage evolved in allopatry as a result of major past vicariant events previously mentioned for *E. argenteus*. A similar pattern was also reported in *Chromis multilineata* (Rocha et al., 2008), a reef fish species of similar distribution to *E. gula*.

The hypothesis of allopatric evolution of distinct lineages in *E. gula* could be weakened by the lack of geographic isolation of both lineages. However, the evidence combined with reciprocal monophyly, geographic coexistence and lack of a shared haplotype reinforces that both ESUs of *E. gula* in the tropical Atlantic evolved in allopatry followed by dispersal that determined their secondary contact. This evidence combined with reciprocal monophyly, geographic coexistence and lack of a shared haplotype reinforces that both ESUs of *E. gula* in the tropical Atlantic evolved in allopatry followed by dispersal that determined their secondary contact.

Unfortunately, the lack of georeferenced data in the other cryptic lineages of *E. melanopterus* and *E. harengulus* restrains further inferences about their diversification processes. Therefore, further phylogeographic studies focusing on both formal taxa are highly recommended.

Final remarks

Our phylogenetic analyses combining species molecular delimitation in a biogeographic context revealed new operational taxonomic molecular units (MOTUs) in *Eucinostomus* species from the Atlantic. Based on extensive barcode datasets we were able to recover the intraspecific diversity for this genus, estimating a potential threshold of 2.7%. Our data also corroborated the efficiency of COI markers in detecting cryptic lineages of mojarra that could be useful in resolving the taxonomic uncertainties in this fish group by providing a database to further support integrative approaches involving morphology, ecological features and other molecular markers. Subtle and overlapped morphological differences have served as proxies for species discrimination within *Eucinostomus*, however, in future systematic studies are strongly encouraged to consider genetic evidence as well.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0025315419001206>.

Financial support. UPJ thanks Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE) for research funding (BCT-0125-2.04/15), Conselho Nacional de Desenvolvimento Científico e Tecnológico CNPq (425080/2016-1) and Fundação de Amparo do Estado de Alagoas, FAPEAL (APQ – 1026/2016).

References

Argolo LA, Ramos RTC, Barreto SB, Bitencourt JA, Sampaio I, Schneider H and Affonso PRAM (2018) The flounder next door: closer evolutionary relationship between allopatric than sympatric *Bothus* (Rafinesque, 1810) species (Pleuronectiformes, Bothidae). *Zoologischer Anzeiger* 277, 131–142.

Bacon CD, Silvestro D, Jaramillo C, Smith BT, Chakrabarty P and Antonelli A (2015) Biological evidence supports an early and complex emergence of the Isthmus of Panama. *Proceedings of the National Academy of Sciences USA* 112, 6110–6115.

Barreira AS, Lijtmaer DA and Tubaro PL (2016) The multiple applications of DNA barcodes in avian evolutionary studies. *Genome* 59, 899–911.

Bickford D, Lohman DJ, Sodhi NS, Ng PKL, Meier R, Winker K, Ingram KK and Das I (2007) Cryptic species as a window on diversity and conservation. *Trends in Ecology & Evolution* 22, 148–155.

Blessing JJ, Marshall JC and Balcombe SR (2010) Humane killing of fishes for scientific research: a comparison of two methods. *Journal of Fish Biology* 76, 2571–2577.

Brandão JHSG, de Bitencourt JA, Santos FB, Watanabe LA, Schneider H, Sampaio I and de Affonso PRAM (2016) DNA barcoding of coastal ichthyofauna from Bahia, northeastern Brazil, South Atlantic: high efficiency for systematics and identification of cryptic diversity. *Biochemical Systematics and Ecology* 65, 214–224.

Brown SDJ, Collins RA, Boyer S, Lefort M, Malumbres-Olarte J, Vink CJ and Cruickshank RH (2012) Spider: an R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. *Molecular Ecology Resources* 12, 562–565.

Calado LL, Bertollo LAC, Cioffi MB, Costa GWWF, Jacobina UP and Molina WF (2014) Evolutionary dynamics of rDNA genes on chromosomes of the *Eucinostomus* fishes: cytotoxic and karyoevolutive implications. *Genetics and Molecular Research* 13, 9951–9959.

Da Silva WC, Marceniuk AP, Sales JBL and Araripe J (2016) Early Pleistocene lineages of *Bagre bagre* (Linnaeus, 1766) (Siluriformes: Ariidae), from the Atlantic coast of South America, with insights into the demography and biogeography of the species. *Neotropical Ichthyology* 14, e150184.

Da Silva TF, Schneider H, Sampaio I, Angulo A, Brito MFG, de Santos ACA, Santos JA, de Carvalho-Filho A and Santos S (2018) Phylogeny of the subfamily Stelliferinae suggests speciation in Ophioscion Gill, 1863 (Sciaenidae: Perciformes) in the western South Atlantic. *Molecular Phylogenetics and Evolution* 125, 51–61.

De Biasse MB, Richards VP, Shivji MS and Hellberg ME (2016) Shared phylogeographical breaks in a Caribbean coral reef sponge and its invertebrate commensals. *Journal of Biogeography* 43, 2136–2146.

De La Cruz-Agüero J (2013) The caudal skeleton of mojarra from genus *Eucinostomus* (Perciformes: Gerreidae) and related species. *Journal of Ichthyology* 53, 994–1006.

de la Cruz-Agüero J and Galvan-Magana F (1993) Morphological discrimination of *Eucinostomus* spp. from the Pacific coast of Mexico. *Bulletin of Marine Science* 52, 819–824.

de Ribeiro AO, Caires RA, Mariguela TC, Pereira LHG, Hanner R and Oliveira C (2012) DNA barcodes identify marine fishes of São Paulo State, Brazil. *Molecular Ecology Resources* 12, 1012–1020.

Drummond AJ and Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7, 214.

Drummond AJ, Suchard MA, Xie D and Rambaut A (2012) Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29, 1969–1973.

Ferreira M, Kavalco KF, de Almeida-Toledo LF and Garcia C (2014) Cryptic diversity between two *Imparfinis* species (Siluriformes, Heptapteridae) by cytogenetic analysis and DNA barcoding. *Zebrafish* 11, 306–3017.

Fields AT, Feldheim KA, Gelsleichter J, Pfoertner C and Chapman DD (2016) Population structure and cryptic speciation in bonnethead sharks *Sphyrna tiburo* in the south-eastern USA and Caribbean. *Journal of Fish Biology* 89, 2219–2233.

Froese R and Pauly D (2019) Fish Base. www.fishbase.org.

Hurtado LA, Mateos M and Liu S (2017) Phylogeographic patterns of a lower intertidal isopod in the Gulf of California and the Caribbean and comparison with other intertidal isopods. *Ecology and Evolution* 7, 346–357.

Hyde JR, Underkoffler KE and Sundberg MA (2014) DNA barcoding provides support for a cryptic species complex within the globally distributed and fishery important opah (*Lampris guttatus*). *Molecular Ecology Resources* 14, 1239–1247.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P and Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)* 28, 1647–1649.

Kennington EL, Baillie SM, Kennington TJ and Bentzen P (2017) Barcoding Atlantic Canada's mesopelagic and upper bathypelagic marine fishes. *PLoS ONE* 12, e0185173.

Lambeck K and Chappell J (2001) Sea level change through the last glacial cycle. *Science* 292, 679–686.

Landi M, Dimech M, Arculeo M, Biondo G, Martins R, Carneiro M, Carvalho GR, Lo Brutto S and Costa FO (2014) DNA barcoding for

- species assignment: the case of Mediterranean marine fishes. *PLoS ONE* **9**, e106135.
- Lanfear R, Calcott B, Ho SYW and Guindon S** (2012) Partitionfinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* **29**, 1695–1701.
- Leigh JW and Bryant D** (2015) Popart: full-feature software for haplotype network construction. *Methods in Ecology and Evolution* **6**, 1110–1116.
- Luiz OJ, Madin JS, Robertson DR, Rocha LA, Wirtz P and Floeter SR** (2012) Ecological traits influencing range expansion across large oceanic dispersal barriers: insights from tropical Atlantic reef fishes. *Proceedings of the Royal Society of London B: Biological Sciences* **279**, 1033–1040.
- Luo A, Ling C, Ho SYW and Zhu C-D** (2018) Comparison of methods for molecular species delimitation across a range of speciation scenarios. *Systematic Biology* **67**, 830–846. doi: 10.1093/sysbio/syy011.
- Matheson Jr RE and McEachran JD** (1984) Taxonomic studies of the *Eucinostomus argenteus* complex (Pisces: Gerreidae): preliminary studies of external morphology. *Copeia* **4**, 893–902.
- Mayden RL** (1997) A hierarchy of species concepts: the denouement in the saga of the species problem. In Claridge MF, Dawah HA and Wilson MR (eds), *Species: The Units of Diversity*. London: Chapman & Hall, pp. 381–423.
- Müller RM, Sdrolia M, Gaina C, Steinberger B and Heine C** (2008) Long-term sea-level fluctuations driven by ocean basin dynamics. *Science* **319**, 1357–1362.
- Nelson JS, Grande TC and Wilson MVH** (2016) *Fishes of the World*. New York, NY: John Wiley & Sons.
- Neves JMM, Lima SMQ, Mendes LF, Torres RA, Pereira RJ and Mott T** (2016) Population structure of the rockpool blenny *Entomacrodus vomerinus* shows source-sink dynamics among ecoregions in the tropical Southwestern Atlantic. *PLoS ONE* **11**, e0157472.
- Nirchio M, Paim FG, Milana V, Rossi AR and Oliveira C** (2018) Identification of a new mullet species complex based on an integrative molecular and cytogenetic investigation of *Mugil hospes* (Mugilidae: Mugiliformes). *Frontiers in Genetics* **9**, 17.
- O'Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-Moreno SA, Cione AL, Collins LS, de Queiroz A, Farris DW, Norris RD, Stallard RF, Woodburne MO, Aguilera O, Aubry M-P, Berggren WA, Budd AF, Cozzuol MA, Coppard SE, Duque-Caro H, Finnegan S, Gasparini GM, Grossman EL, Johnson KG, Keigwin LD, Knowlton N, Leigh EG, Leonard-Pingel JS, Marko PB, Pyenson ND, Rachello-Dolmen PG, Soibelzon E, Soibelzon L, Todd JA, Vermeij GJ and Jackson JBC** (2016) Formation of the Isthmus of Panama. *Science Advances* **2**, e1600883.
- Pappalardo AM, Cuttitta A, Sardella A, Musco M, Maggio T, Patti B, Mazzola S and Ferrito V** (2015) DNA barcoding and COI sequence variation in Mediterranean lanternfishes larvae. *Hydrobiologia* **749**, 155–167.
- Pinheiro HT, Rocha LA, Macieira RM, Carvalho-Filho A, Anderson AB, Bender MG, Carlos FDD, Ferreira EL, Figueiredo-Filho J, Francini-Filho R, Gasparini JL, Joyeux J, Luiz OJ, Mincarone MM, Moura RL, de Nunes JACC, Quimbayo JP, Rosa RS, Sampaio CLS, Sazima I, Simon T, Vila-Nova DA, Floeter SR and Treml E** (2018) South-western Atlantic reef fishes: zoogeographical patterns and ecological drivers reveal a secondary biodiversity centre in the Atlantic Ocean. *Diversity and Distributions* **24**, 951–965.
- Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD and Vogler AP** (2006) Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology* **55**, 595–609.
- Puillandre N, Lambert A, Brouillet S and Achaz G** (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology* **21**, 1864–1877.
- Rambaut A and Drummond AJ** (2009) FigTree version 1.3. 1. <http://Tree.Bio.Ed.Ac.Uk/Software/Figtree/>.
- Ratnasingham S and Hebert PDN** (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PLoS ONE* **8**, e66–213.
- Rocha LA** (2003) Patterns of distribution and processes of speciation in Brazilian reef fishes. *Journal of Biogeography* **30**, 1161–1171.
- Rocha L, Claudia R, Ross RD and Brian B** (2008) Comparative phylogeography of Atlantic reef fishes indicates both origin and accumulation of diversity in the Caribbean. *BMC Evolutionary Biology* **8**, 157.
- Stamatakis A, Aberer AJ, Goll C, Smith SA, Berger SA and Izquierdo-Carrasco F** (2012) RAxML-Light: a tool for computing terabyte phylogenies. *Bioinformatics (Oxford, England)* **28**, 2064–2066.
- Tamura K, Stecher G, Peterson D, Filipiński A and Kumar S** (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* **30**, 2725–2729.
- Thacker CE** (2017) Patterns of divergence in fish species separated by the Isthmus of Panama. *BMC Evolutionary Biology* **17**, 111.
- Valdez-Moreno M, Vásquez-Yeomans L, Elías-Gutiérrez M, Ivanova NV and Hebert PDN** (2010) Using DNA barcodes to connect adults and early life stages of marine fishes from the Yucatan Peninsula, Mexico: potential in fisheries management. *Marine and Freshwater Research* **61**, 655–671.
- Ward RD, Zemlak TS, Innes BH, Last PR and Hebert PD** (2005) DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **360**, 1847–1857.
- Weigt LA, Baldwin CC, Driskell A, Smith DG, Ormos A and Reyier EA** (2012) Using DNA barcoding to assess Caribbean reef fish biodiversity: expanding taxonomic and geographic coverage. *PLoS ONE* **7**, e41059.
- Williams ST and Duda Jr TF** (2008) Did tectonic activity stimulate Oligo-Miocene speciation in the Indo-West Pacific? *Evolution: International Journal of Organic Evolution* **62**, 1618–1634.
- Williams ST, Smith LM, Herbert DG, Marshall BA, Warén A, Kiel S, Dyal P, Linse K, Vilvens C and Kano Y** (2013) Cenozoic climate change and diversification on the continental shelf and slope: evolution of gastropod diversity in the family Solariellidae (Trochoidea). *Ecology and Evolution* **3**, 887–917.
- Winker K** (2005) Sibling species were first recognized by William Derham (1718). *The Auk* **122**, 706–707.
- Winterbottom R, Hanner R, Burridge M and Zur M** (2014) A cornucopia of cryptic species – a DNA barcode analysis of the gobiid fish genus *Trimma* (Percomorpha, Gobiiformes). *ZooKeys* **381**, 79–111. <https://doi.org/10.3897/zookeys.381.6445>.
- Woodland DJ** (2006) Gerreidae. Silver-biddies. In Carpenter KE and Niem V (eds), *FAO species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific*, 6th Edn. Rome: FAO, pp. 2946–2960.
- Zhang J, Kapli P, Pavlidis P and Stamatakis A** (2013) A general species delimitation method with applications to phylogenetic placements. *Bioinformatics (Oxford, England)* **29**, 2869–2876.