

Giardia intestinalis can interact, change its shape and internalize large particles and microorganisms

Research Article

Cite this article: Benchimol M (2021). *Giardia intestinalis* can interact, change its shape and internalize large particles and microorganisms. *Parasitology* **148**, 500–510. <https://doi.org/10.1017/S0031182020002292>

Received: 2 August 2020
Revised: 5 October 2020
Accepted: 25 November 2020
First published online: 7 December 2020

Key words:

Endocytosis; *Giardia duodenalis*; *Giardia lamblia*; microorganism's interaction

Author for correspondence:

Marlene Benchimol,
E-mail: marlenebenchimol@gmail.com

Marlene Benchimol^{1,2} 

¹UNIGRANRIO-Universidade do Grande Rio-Duque de Caxias-Rio de Janeiro, Rio de Janeiro, Brazil and ²UFRJ-Universidade Federal do Rio de Janeiro-Instituto de Biofísica Carlos Chagas Filho-Laboratório de Ultraestrutura Celular Hertha Meyer, and Instituto Nacional de Ciência e Tecnologia-INBEB, Centro Nacional de Biologia Estrutural e Bioimagens-CENABIO, Rio de Janeiro, Brazil

Abstract

Giardia intestinalis is a parasitic protozoan that inhabits its vertebrate hosts' upper small intestine and is the most common cause of waterborne diarrhoea worldwide. *Giardia* trophozoites present few organelles, and among them, they possess peripheral vesicles (PVs), which are considered an endosomal–lysosomal system. All experimental procedures carried out until now indicate that *Giardia* ingests macromolecules by fluid-phase and receptor-mediated endocytic pathways. Still, there is no description concerning the interaction and ingestion of large materials. Here, we tested *Giardia*'s capacity to interact with large particles; once, *in vivo*, it inhabits an environment with a microbiota. We tested protozoan interaction with yeasts, bacteria, latex beads, ferritin and albumin, in different times of interaction and used several microscopy techniques (light microscopy, scanning electron microscopy and transmission electron microscopy) to follow their fate. *Giardia* interacted with all of the materials we tested. Projections of the plasma membrane similar to pseudopods were seen. As albumin, small markers were found in the PVs while the larger materials were not seen there. Large vacuoles containing large latex beads were detected intracellularly. Thus, we observed that: (1) *Giardia* interacts with large materials; (2) *Giardia* can display an amoeboid shape and exhibit membrane projections when in contact with microorganisms and large inorganic materials; (3) the region of the exit of the ventral flagella is very active when in contact with large materials, although all cell surface also present activity in the interactions; (4) intracellular vacuoles, which are not the PVs, present ingested large beads.

Introduction

Giardia intestinalis is a parasite protozoan that inhabits its vertebrate hosts' upper small intestine and is the most common cause of waterborne diarrhoea worldwide. Giardiasis' clinical manifestations vary from an asymptomatic or mild infection to an acute or chronic disease associated with diarrhoea and malabsorption. *Giardia* infection initiates with the cyst form's ingestion, which excysts in the upper small intestine where *Giardia* stabilizes, and the parasite affects the host. Parasite–host relationships are of great importance, and how *Giardia* nourishes itself, causing adverse effects on the host, has been the object of interest in scientific research.

Giardia has two nuclei and a glycogen-rich cytoplasm. The membrane system of *G. intestinalis* is considered very simple, with a unique set of vesicles, named peripheral vesicles (PVs), the endoplasmic reticulum network and mitosomes. Recently, Acosta-Virgen *et al.* (2018) suggested, using several approaches, the presence of peroxisomes. The absence of a typical Golgi complex and a canonical lysosome is a matter of discussion (Kattenbach *et al.*, 1991; Tai *et al.*, 1993; Lanfredi-Rangel *et al.*, 1998, 1999; Wampfler *et al.*, 2014).

The PVs are oval organelles 100–200 nm in size, located in the cell periphery, below the plasma membrane, and considered the endosomal/lysosomal system in *Giardia* (Feely and Dyer, 1987; Kattenbach *et al.*, 1991; Rivero *et al.*, 2010). The PVs participate in the endocytosis process, digestion and retrograde transport of the parasite (Lanfredi-Rangel *et al.*, 1998). Previous studies demonstrated the presence of resident enzymes such as acid phosphatases, sulphur-binding proteins (SH) and glucose-6-phosphatase in the PVs as well as the accumulation macromolecules such as albumin, peroxidase, transferrin, low-density lipoprotein, and accumulating fluid-phase and membrane-bound molecules and in PVs (Kattenbach *et al.*, 1991; Tai *et al.*, 1993; Lanfredi-Rangel *et al.*, 1998). The uptake of soluble material from the environment into PV is not selective, in contrast to further retrograde transport that allows some substances to rapidly cross over into the proximal endoplasmic reticulum (ER) (Abodeely *et al.*, 2009). The presence of protein complexes associated with the endocytic machinery in *Giardia* has been reported, as clathrin heavy chain (*GI*CHC), subunits of the AP2 heterotetramer (*GI*AP2) and *Giardia* dynamin-related protein (*GI*DRP) with a discrete localization by fluorescence microscopy (Gaechter *et al.*, 2008; Rivero *et al.*, 2010). Besides, a protein interactome of *GI*CHC revealed all of the conserved factors in addition to a novel protein, a putative clathrin light chain (Zumthor *et al.*, 2016).

The endoplasmic reticulum in *Giardia* is an extensive reticular system distributed from the nuclear envelopes throughout the cell body. The presence of a tubule-vesicular contiguous compartment in *Giardia* that functions as endoplasmic reticulum and endosomes/lysosome with connections with the PVs was proposed (Abodeely *et al.*, 2009). The presence of some specific marker proteins in this organelle, similar to those of higher eukaryotic cells, such as glucose-6-phosphate, a resident ER enzyme, detected by electron microscopy cytochemistry was reported (Lanfredi-Rangel *et al.*, 1998). Previous studies demonstrated that although *G. intestinalis* has a conventional secretory trafficking, some elements, such as the post-translational modification machinery, are missing entirely. An extensive genomic and biochemical studies indicated that the parasite lacks several nucleotide sugar transporters (Banerjee *et al.*, 2008). Thus, in *G. intestinalis*, the ER, in addition to protein synthesis, has different functions such as endocytic activities and extracellular material degradation, being a pluripotent compartment (Abodeely *et al.*, 2009).

Giardia does not present mitochondria; however, double-membrane small organelles, 200 nm in size, presenting some mitochondria characteristics named mitosomes, were described (Tovar *et al.*, 1999; Tovar *et al.*, 2003). The mitosomes are distributed over the cytoplasm and also placed between the flagellar axonemes. Mitochondrial characteristics such as biosynthesis of Fe-S clusters, a TOM and TIM protein family transport machinery were found in mitosomes (Tachezy and Smid, 2008). However, mitosomes do not present several mitochondrial characteristics, such as ATP synthesis, the citric acid cycle, oxidative phosphorylation, heme biosynthesis, presence of DNA, lipid metabolism and the amino acid and urea cycles (Makiuchi and Nozaki, 2014). One important finding was the presence of an iron-sulphur complex (IscS and IscU proteins) (Tachezy *et al.*, 2001). Chaperones, such as Cnp60 and HSP70, are also present (Midlej *et al.*, 2016).

During the differentiation of trophozoite to cysts, two types of membrane-bounded vesicles appear, the encystation-specific vesicles (ESVs) (Faubert *et al.*, 1991) and the encystation carbohydrate-positive vesicles (Midlej *et al.*, 2013), and both act to build the cyst wall (CW), a constitutive and essential structure of cyst. The ESVs are large vesicles (1 µm of size), participate in CW formation and appear at the beginning of the encystation process (Faubert *et al.*, 1991). The protein content of the ESVs is the cyst-wall proteins 1–3, which originate in the endoplasmic reticulum (Hehl and Marti, 2004).

The Golgi complex is an organelle characterized by stacked cisternae, often located around the nucleus and close to the endoplasmic reticulum. However, an organelle that fits with these criteria has not been identified in *G. intestinalis*. Some groups proposed a similarity between the *Giardia*'s ESVs and the Golgi complex (Luján *et al.*, 1995; Marti *et al.*, 2003; Stefanic *et al.*, 2009; Ebnetter *et al.*, 2016), but this is still very controversial. Besides, classical Golgi markers such as GM130, galactosyl transferases or the trans-Golgi network marker Rab6 were not detected in the parasite. Thus, a typical Golgi in *G. intestinalis* is not present (Dacks and Field, 2007).

Because the endocytosis is an essential function in all cells and especially in parasites, investigations concerning the ingestion of molecules and other elements in the *Giardia* habitat and its traffic are necessary and still incomplete.

Giardia nutrition has been the subject of several works, mainly focusing on endocytosis by the parasite. In the literature, experimental designs start with *Giardia* growth in axenic culture media, which contains essential nutritional elements.

In vivo, *Giardia* inhabits the human gut, which contains 500–1000 microorganism's species, and its diversity is known as 'microbiome'. Thus, in natural conditions in the intestine of a human, *Giardia* has the company of thousands of other

organisms. The possibility of nourishing itself from this flora raises an important question: Do our laboratory experiments, where we grow *Giardia* in an axenic environment, answer the right questions to its mode of nutrition? Using the culture medium in the laboratory experiments, we can observe only fluid-phase endocytosis, since the medium only contains macromolecules. Thus, several questions regarding this endocytic pathway remain unanswered: would *Giardia* be able to phagocytize large particles or microorganisms present in its natural habitat? In the present study, we decided to evaluate *Giardia*'s behaviour in an environment where microorganisms and large materials are present. Our observations show clearly that *Giardia*, once appropriately stimulated, can ingest large particles and even whole microorganisms.

Materials and methods

Cell culture

Trophozoites of *G. intestinalis* WB strain (ATTC, USA, No. 30957) were cultured for 48 h at 37°C in TYI-S-33 medium (Keister, 1983) enriched with 10% fetal bovine serum and supplemented with 0.1% bovine bile.

Endocytic assays

Incubation with ferritin and albumin

Cells were washed and transferred to phosphate-buffered saline (PBS). They were incubated for periods varying from 1 to 60 min in a solution containing 5% native ferritin in PBS, pH 7.2 or 1 mg mL⁻¹ bovine serum albumin (BSA; Sigma, USA), at 37°C. After incubation, the parasites were washed twice with PBS and fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2, and processed for scanning electron microscopy (SEM) and transmission electron microscopy (TEM) as described below.

Incubation with polystyrene microspheres

For TEM polystyrene microspheres, sizes of 10 nm, 110 nm or 0.9 µm in diameter were used (Polysciences, USA), according to a previously published protocol (Benchimol *et al.*, 1990). They were suspended in PBS and allowed to interact with the parasites for 1, 5, 20, 30 and 60 min. Afterward, the cells were washed twice with PBS and fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2, plus 5 mM CaCl₂, and postfixed in 1% OsO₄, in 0.1 M cacodylate buffer plus 0.8% potassium ferrocyanide. After that, the cells were washed in PBS, dehydrated in acetone, and embedded in Epon. Besides, in some experiments, latex particles were coated with cationized ferritin (1 mg mL⁻¹, final concentration of about 10⁵ particles mL⁻¹) or albumin (1 mg mL⁻¹ BSA, Sigma) and used in interactions with the parasites.

Incubation with FluoSpheres

Living *Giardia* was incubated with FluoSpheres™ carboxylate-modified 100 nm, red (580/605), from Invitrogen (ThermoFisher Scientific), diluted 5000 times, and observed at 1, 5, 15, 30, 60, 120 min in medium without serum. The FluoSpheres were coated with BSA 1% before the incubation, sonicated and centrifuged to remove particle clusters. The samples were observed both alive and fixed for 10 min with 4% formaldehyde diluted in 0.1 M cacodylate buffer, pH 7.2. The fluorescence microscope used was Zeiss Elyra (Germany) in Confocal mode (LSM 710 system), Laser 543 nm, Objective Lens EC Plan-apochromat 63×/1.4.

Incubation with *Saccharomyces cerevisiae*

Saccharomyces cerevisiae was obtained from yeast powder (Fleischmann & Royal, Jundiaí, SP, Brazil), and diluted in warm water (37°C). Yeast cells were washed three times with warm 0.1 M sterile PBS (pH 7.2) for 5 min at 3000 g and suspended in fresh TYM medium without serum, the concentration of 5×10^7 cells mL⁻¹. The yeast cells were used immediately for interaction assays following Pereira-Neves and Benchimol (2007) protocol. Co-incubation of *G. intestinalis* (WB strain) with yeast cells was performed in medium without serum at parasite/yeast cell ratios of 1:10 and 1:40 for different times (1, 5, 20, 30, 60 min) at 37°C. Triplicate preparations were analysed by differential interference contrast (DIC) microscopy. After interaction, samples were processed for TEM and SEM.

Incubation with bacteria

Escherichia coli, strain 346, was grown at 37°C for 24 h in BT broth (Bacto-tryptone, Difco) medium. *Giardia* was collected by centrifugation at 6000 g for 5 min. Next, they were allowed to interact with the bacteria for periods varying from 1 to 60 min using bacteria: protozoa ratio of 10:1. After the interaction, the cells were fixed for 2 h at room temperature in a solution containing 2.5% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.2) and processed for SEM and TEM as described below. In some experiments, a mixture of bacteria, yeasts and albumin was used.

Light microscopy

To evaluate the parasites' behaviour in contact with different materials after interaction, living cells were observed under light microscopy, in DIC and phase contrast mode (Zeiss Aphiort light microscopy, Germany).

Scanning electron microscopy

Cells were allowed to adhere to Poly-L-lysine-coated glass coverslips and then fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2 for 60 min. After that, the cells were postfixed for 5 min in 1% OsO₄, dehydrated in ethanol, critical point dried with CO₂, sputter-coated with gold-palladium and examined in a Jeol 5800X SEM operating at 15 KV.

Transmission electron microscopy

Samples were fixed in 2.5% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.2). They were then postfixed in 1% osmium tetroxide (OsO₄) containing 0.8% potassium ferrocyanide in 0.1 M cacodylate buffer for 40 min, dehydrated in acetone and embedded in epoxy resin. Ultrathin sections (60–70 nm thick) were collected and stained with uranyl acetate and lead citrate. At the end of processing, the samples were observed using a Jeol 1210 (Jeol, Japan).

Cytochemistry for carbohydrate detection (Thiéry, 1967): Thiéry's technique (periodic acid-thiosemicarbazide-silver proteinate)

After a routine fixation and Epon embedding procedure, the ultrathin sections were collected on gold grids, treated for 20 min with 1% periodic acid, incubated for 48–72 h in 1% thiosemicarbazide, and exposed to 1% silver proteinate for 30 min. The sections were analysed using a JEOL 1210 TEM. The controls were performed by omitting the periodic acid step.




Fig. 1. Incubation with 100 nm red FluoSpheres BSA coated for 30–60 min and observation by phase contrast (a) or confocal laser-scanning microscope and DIC (b–e). (a) General field view at low magnification to observe the living trophozoites during FluoSpheres interaction for 30 min using phase-contrast microscopy. Vacuoles containing the fluorescent beads are seen in the *Giardia* cytosol as bright spots. (b–e) Overlay of DIC and confocal fluorescence images after *Giardia* interaction with 100 nm red FluoSpheres. In (b, d, e), *Giardia* was allowed to interact for 60 min, fixed and observed, whereas (c) is a living cell after 30 min interaction, and the image is presented in RGB overlay. The 100 nm beads sizes are below the light microscopy resolution; thus, the red spots are vacuoles with beads concentration. Virtual sections were made to check if the beads are on or in the cell. (d, e) Fluorescence (d) and DIC overlay (e) of the same section.

Results

Light microscopy

To observe whether there would be interaction and possible ingestion of large materials at the light microscopy level, living *Giardia* were incubated with BSA-coated FluoSpheresTM carboxylate-modified 100 nm, red (580/605) and observed at different periods. Video microscopy, phase contrast microscopy, DIC mode and




Fig. 2. *Giardia* SEM after incubation with live yeasts (*S. cerevisiae*) (a–f) and also together with latex beads (a) or *E. coli* (b) for 5 min. Notice that flagella adhere to all these materials. Figures (e, f) show that the parasite's plasma membrane exhibits projections (arrows) in the areas of contact with the yeasts. Y, yeast; B, bacteria; L, latex beads. Artificially coloured.

confocal laser scanning microscopy were used (Fig. 1). The samples were observed either alive (Fig. 1a and c) or fixed for 10 min with 4% formaldehyde (Fig. 1b, d and e). Different focal planes were analysed (Z-level) to check if the beads were outside or inside the cell. After a few min, we observed red spots outside the cells and on the cell's surface. Living cells showed an intense beating of ventral flagella (not shown), and beads were also found in the ventral flagella exit region. After 30 min, several cells' compartments exhibited bright fluorescence, some of them around the nucleus (Fig. 1b).

Scanning electron microscopy

Since by light microscopy we observed an intense flagella movement and preferential interaction of the large particles at the region where the ventral flagella exit, we decided to check using SEM (Figs 2, 3 and 6a). The other flagella also contact the large materials, as seen by SEM (Fig. 2a and c). *Giardia*'s whole-cell surface also interacted with all material used here (Fig. 2f), even the bare zone (Fig. 3f) of the ventral disc. Large materials as yeasts, which measures 4–7 μm, were also seen in the process of interaction (Figs 2 and 3e), and the anterior flagella seemed to participate in this process (Fig. 2a and c). In the first 5 min of incubation, we noted plasma membrane ripples (Fig. 1, asterisks; Fig. 3b and c) and large cell projections (Fig. 4a–c). All materials tested, large or small, such as bacteria, yeasts, coated latex beads, albumin and ferritin, were taken by the cells. We did not observe the ingestion of uncoated latex beads. When we used short periods of incubation, smaller markers were preferentially seen in the region where the ventral flagella exit (Figs 3 and 6a). Large extensions and invaginations on *Giardia*'s cell surface were observed on the parasite's dorsal region after incubation with cells such as bacteria, yeasts or polystyrene microspheres (Figs 4, 6–8). An

interesting observation was the change in the *Giardia* cell surface: cells grown in axenic culture medium presented a smooth and uniform texture (Fig. 5a) when observed by routine SEM whereas, after endocytosis, *Giardia*'s surface showed clear signs of irregularities and wrinkles, probably due to the intake of large materials as observed after 20 min of incubation with *E. coli* (Fig. 5b).

Transmission electron microscope

After different incubation times with *Giardia* trophozoites, ultra-thin sections were analysed using a TEM. Small markers, such as free albumin and ferritin, were found in the PVs. We observed markers' entry through all cell membrane places, especially where the ventral flagella emerge (Fig. 6), confirming observations made by light microscopy and SEM. After 5 min of co-incubation, membrane's expansions, similar to pseudopods, were observed when coated latex beads or bacteria were used (Figs 7a–c, 8b and c, 9a and b). A process of adhesion of ferritin or albumin-coated latex beads with these projections were noted (Fig. 7b and inset). After incubation with yeasts, a sharp interaction between *Giardia* and yeast was observed (Fig. 7d). After 30 min of incubation, incorporated elements such as coated latex beads were found in a vacuole (Fig. 7e). When we carried out simultaneous incubations with bacteria and yeasts, or bacteria and latex beads (Figs 8 and 9), we noticed that the parasites present drastic changes in shape, and membrane projections appeared. In Fig. 8c, we show a latex bead already incorporated while the parasite presents a dramatic change in shape, apparently trying to reach a bacterium, forming a pseudopod-like extension (Figs 8b and 9). One of these projections clearly shows filamentous structures (Fig. 9b). In some situations, *Giardia* has changed its shape profoundly, assuming an amoeboid form (Figs 8c and 9).




Fig. 3. *Giardia* trophozoites observed by SEM after incubations for 10 min with different materials: (a, b) with ferritin-coated latex beads presenting two different sizes (LF); (c) with *E. coli* (B) and uncoated latex beads (L); (d) with albumin-coated latex beads; (e) with a mixture of ferritin-coated latex beads (L), bacteria (B) and yeasts (Y); (f) with ferritin-coated latex beads (L). Notice in (a) that there is adhesion of the bead with one flagellum. The small particles over the cell are free ferritin. In (b), the arrowhead points to an opening below the ventral disc, whereas the arrows indicate a large compartment below the plasma membrane in images (b, c). In (d, e), the uncoated latex beads (L) are in the process of internalization precisely in the region of the emergence of the ventral flagella (arrows). In (f), two uncoated latex beads have adhered to the ventral disc bare zone. Images were artificially coloured.

We decided to compare fluid-phase endocytosis with large particle phagocytosis and then incubated the cells with albumin (Fig. 8d). The difference was striking since albumin incorporation occurred only in PVs. Besides, we noticed an intense concentration of albumin in the ventral flagella exit region, where many PVs containing albumin were also seen (Fig. 8d). After 10 min of incubation, yeasts were found inside a vacuole (Fig. 8e).

When we searched for intracellular localization of materials tested here, we were able to find BSA-coated latex beads in vacuoles, which are not the PVs by its size and localization (Fig. 10a–f and h). Single large beads were seen in a large vacuole (Fig. 10a–e), whereas several small beads were clustered in other vacuoles (Fig. 10f–h).

Cytochemistry

Using the periodic acid-thiosemicarbazide-silver proteinate (Thiéry, 1967), a cytochemical technique associated with electron

microscopy, carbohydrates were detected in a large vacuole. After incubation with bacteria, there was material in a vacuole under an apparent process of degradation (Fig. 10g).

Discussion

Giardia intestinalis is a parasite whose endocytic pathway has been previously studied. There is consensus that the PVs found below the plasma membrane of this parasite forms an interconnected network, which includes part of the endoplasmic reticulum, correspond to *Giardia's* early and late endosomes and the lysosomal system (Feely and Dyer, 1987; Kattenbach *et al.*, 1991; Rivero *et al.*, 2010). In all these studies, trophozoites were grown in an axenic medium, where essential macromolecules are supplied to the parasite during its *in vitro* growth. All markers used until now were restricted to follow the fluid-phase or receptor-mediated endocytosis.




Fig. 4. SEM. After 5 min of incubation with *E. coli* (B) and free ferritin (spots on the membrane), *Giardia* trophozoites' cell surface. Significant changes on the cell surface were observed in the parasite's dorsal region, similar to the formation of large pseudopods (arrows). The bacteria were artificially coloured.




Fig. 5. Scanning electron microscopy of *G. intestinalis* under two different conditions: in (a) a cell grown in an axenic culture medium and (b) a cell after 20 min of incubation with *E. coli* (B) and with free ferritin. There is a significant change in the cell surface, which is much wrinkled after the materials' internalization.

Fig. 6. (a) Scanning electron microscopy of small latex beads coated with BSA when in interaction with *G. intestinalis* for 5 min. Notice that the beads are located in the area of the ventral flagella (VF) exit (arrow). D, ventral disc. (b) Transmission electron microscope (TEM). *Giardia* trophozoite after incubation for 5 min with yeasts (Y). Note the contact of the ventral flagella (VF) with the microorganism and the yeast location in the region of exit of the ventral flagella, just below the area of the ventral disc. N, nucleus; F, flange; Ax, axonemes.





Fig. 7. Images obtained with the transmission electron microscope. In (a, b), albumin-coated latex beads (L) are observed adhered to on *Giardia*'s cell surface. Expansions of the membrane (arrows), similar to pseudopods, and even a contact with one of the latex beads are noted (b). An adhesion point between the *Giardia* membrane and a latex bead's surface is observed in the inset. In figure (c), an endocytosis experiment was performed where tiny latex beads were coated with ferritin. There are evident expansions of the membrane (arrows) forming pseudopods and even a closure of one of these expansions keeping the material inside. Notice a compartment (C) in the cytosol. In figure (d), a large yeast (Y) is in phagocytosis by *Giardia*. In figure (e), an incubation with latex beads coated with ferritin was made, and one of the particles (L) is observed in a large internal vacuole of the trophozoite. N, nucleus.



It is essential to consider that *in vivo*, *Giardia* is found in the small intestine where there is an enormous population of other microorganisms that are part of the intestinal

microbiota. In this way, it was missing an analysis of this parasite's behaviour in experimental conditions closer to those found *in vivo*.




Fig. 8. TEM of several different incubations of *G. intestinalis* with bacteria (a, b, c) coated BSA latex beads (c), and yeasts (a, e). In (d), incubation was performed only with albumin, and it is observed that the peripheral vesicles (PV) present this protein inside. A high concentration of albumin in the process of endocytosis is seen in the region of the disc (D) where the ventral flagella exit (arrows) and the peripheral vesicles containing albumin are present. In (a, e), the experiments were carried out with yeasts (Y), and the presence of three yeasts is already observed inside a vacuole in (e). B, bacteria; D, ventral disc; N, nucleus.

In 1979, Nemanic *et al.* reported bacterial-like structures, considered as endosymbionts, in the cytoplasm of *Giardia muris* and *G. lamblia* collected from mice intestines. The endosymbionts found by the authors were both trophozoites and cysts. Subsequently, Sogayar and Gregorio (1989) showed internalized

bacteria in *Giardia duodenalis* found in rat intestine (*Rattus rattus*). The authors did not characterize either the species of *Giardia* or the bacteria found in the gut. They published that the bacteria were in phagocytic vacuoles without signs of bacteria digestion. The authors also claimed that the bacteria were




Fig. 9. TEM of *G. intestinalis* after interaction with bacteria and small latex beads for 5 min. The parasite has drastically changed its shape, presenting an amoeboid form with short and long projections (arrowheads). One long membrane projection, similar to a pseudopod, is seen in contact with a bacterium (B). In this membrane projection, filamentous structures (asterisks) can be noted. The arrows point to 100 nm latex beads. Figure (b) is a higher magnification of (a). D, ventral disc; F, flagellum; N, nucleus.

observed attached to the trophozoite membrane until their total incorporation by phagocytosis. Here, three essential points caught my attention: (1) there is no mention of the participation of PVs in the endocytic mechanism described; (2) *Giardia* were collected from a naturally infected animal, which means that the parasites were found in a natural community with several microorganisms; (3) the work indicated the capacity of ingestion of a large material by *Giardia*. These observations lead us to explore further, using a more controlled experimental system, *G. intestinalis* trophozoites' ability to ingest large particles.

Preferential place for uptake

Our present observations showed that *Giardia* uptake of ferritin and other materials occurred throughout the cell surface, but preferentially through the ventral flagella's exit, just below the ventral disc. Abodeely *et al.* (2009) made a similar observation when they used 40 nm biotin-conjugated FluoSpheres. The authors also found that clathrin, an essential protein involved in receptor-mediated endocytosis, was abundant near the ventral flagella (Abodeely *et al.*, 2009). We also noted the same preferential place, the ventral flagella exit, although the whole *Giardia* cell surface showed contact with all materials tested.

Despite extensive EM studies of the *Giardia* endocytic system, a detailed morphological characterization of the incorporated

molecules' destination and its digestive process was not carried out. This parasite has been described as presenting an endomembrane system restricted to a few structures such as the endoplasmic reticulum and PVs, a contiguous collaborative compartment working in harmony (Abodeely *et al.*, 2009). Although numerous articles have been published with excellent results on endocytosis mechanisms in *Giardia*, they focused on using small markers and their incorporation by fluid-phase endocytosis or/and clathrin-mediated receptor (Hernandez *et al.*, 2007; Zumthor *et al.*, 2016). The previous studies are unanimous in describing the participation and importance of PVs as sites for incorporating and possibly digestion of the ingested material. Based on the localization of lysosomal enzymes such as acid phosphatase and cysteine proteases, PVs have been described as having lysosomal properties (Feely and Dyer, 1987; Lindmark, 1988; McCaffery and Gillin, 1994), which means they acidify and behave as digestive organelles. Cytochemical, biochemical or molecular-tag analyses using the search for acid phosphatase has been carried out by several authors, who invariably found this enzyme in some PVs and also in the endoplasmic reticulum (Lindmark, 1988; Kattenbach *et al.*, 1991; Lanfredi-Rangel *et al.*, 1998; Touz *et al.*, 2004; Rivero *et al.*, 2012). A previous work raised the possibility of an alternative endosomal compartment that lies adjacent to the ER or in direct contact with it (Abodeely *et al.*, 2009).

In the present work, occasional cells with membrane projections resembling pseudopods-like structures were noticed, when trophozoites were incubated in the presence of large materials such as coated latex beads, bacteria and yeasts. One interesting observation was the drastic parasite change in shape and an extensive projection towards one bacterium. This dramatic change in shape has not been described previously and suggests that the protozoan can detect signals indicating the necessity to internalize a microorganism. The presence of filaments in these projections corroborates the idea that they correspond to a true pseudopod involved in a phagocytic process, a fact not yet described for *Giardia*.

Also, we were able to find several images of the large beads in cellular structures that do not correspond to the PVs, either by size or location. These vacuoles have not yet been described in the *Giardia* literature. They may represent either a new organelle with a lysosomal function or a path that has not yet been fully elucidated in *Giardia*. These vacuoles were found in different places in the cell, but preferably close to the nucleus. The presence of a perinuclear structure as an alternative place of incorporating external materials from PV had already been suggested in Abodeely *et al.*'s work (Abodeely *et al.*, 2009). Further studies are needed to follow a probable endocytic alternative pathway in *Giardia*, and new experiments are underway to add new information in this regard.

Several studies have shown the relationship between bacteria and *E. histolytica* growth and virulence. They have noted that the bacterial microbiota may influence the behaviour and virulence of pathogens, their immune response, and lead to variability in the outcome of parasitic infections (Marie and Petri, 2014; Bär *et al.*, 2015). Thus, it would be important to study further the relationships between *Giardia* and other microorganisms in the same habitat to see if it affects the parasite's virulence. A greater understanding of *Giardia*'s eating habits and its relationship with intestinal microbiota may provide knowledge for new treatments and expand knowledge in the field of intestinal parasitism.

In conclusion, our observations indicate that in addition to a classic fluid-phase and receptor-mediated endocytosis, *Giardia* is also able, when properly stimulated, to ingest large particles and microorganisms through a phagocytic-like process, as previously shown for other pathogenic protozoa such as *Trichomonas vaginalis* (Pereira-Neves and Benchimol, 2007) and *Entamoeba histolytica* (Orozco *et al.*, 1983). Ingestion




Fig. 10. Intracellular localization of BSA-coated latex beads. *Giardia intestinalis* were incubated with polystyrene microspheres (L), sizes of 10 nm, 110 nm or 0.9 μm and fixed after 30 min (a–f, h). Some cells presented beads internalization (L). The large beads were seen alone in a large vacuole (a–e), whereas several small beads are together in a vacuolar structure (f, arrow). Figure (h) is a higher magnification of (f). Figure (g) is an image of the parasite after incubation with bacteria for 30 min using cytochemistry [periodic acid-thiosemicarbazide-silver proteinate (Thiéry, 1967) for carbohydrates detection]. A positive reaction is observed in one large vacuole where a possible biological material degradation is observed (arrow). Ax, flagella axonemes; D, ventral disc; N, nucleus; F, flagellum.

takes place, preferably at the ventral flagella exit. The ingested material most probably follows another pathway and is accumulated into large vacuoles.

Financial support. This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), Grant number E-26/202.824/2017.

Conflict of interest. None.

Ethical standards. Not applicable.

References

Abodeely M, DuBois KN, Hehl A, Stefanic S, Sajid M, De Souza W, Attias M, Engel JC, Hsieh I, Fetter RD and McKerrrow JH (2009) A contiguous

compartment functions as endoplasmic reticulum and endosome/lysosome in *Giardia lamblia*. *Eukaryotic Cell* **8**, 1665–1676.

Acosta-Virgen K, Chávez-Munguía B, Talamás LD, Guillén DL, Martínez-Higuera A, Lazcano A, Martínez-Palomo A and Espinosa-Cantellano M (2018) *Giardia lamblia*: identification of peroxisomal-like proteins. *Experimental Parasitology* **191**, 36–43.

Banerjee S, Cui J, Robbins PW and Samuelson J (2008) Use of *Giardia*, which appears to have a single nucleotide-sugar transporter for UDP-GlcNAc, to identify the UDP-Glc transporter of *Entamoeba*. *Molecular Biochemical Parasitology* **159**, 44–53.

Bär AK, Phukan N, Pinheiro J and Simoes-Barbosa A (2015) The interplay of host, and parasitic protozoans at mucosal interfaces: implications for the outcomes of infections and diseases. *PLoS Neglected Tropical Diseases* **9**, e0004176.

Benchimol M, Batista C and De Souza W (1990) Fibronectin – and – laminin mediated endocytic activity in the parasitic protozoa *Trichomonas vaginalis* and *Tritrichomonas foetus*. *Journal of Submicroscopy and Cytology Pathology* **22**, 39–45.

- Dacks JB and Field MC** (2007) Evolution of the eukaryotic membrane trafficking system: origin, tempo and mode. *Journal of Cell Science* **120**, 2977–2985.
- Ebneter JA, Heusser SD, Schraner EM, Hehl AB and Faso C** (2016) Cyst wall-protein-1 is fundamental for Golgi-like organelle neogenesis and cyst wall biosynthesis in *Giardia lamblia*. *Nature Communication* **15**, 13859.
- Faubert G, Reiner DS and Gillin FD** (1991) *Giardia lamblia*: regulation of secretory vesicle formation and loss of ability to reattach during encystation *in vitro*. *Experimental Parasitology* **72**, 345–354.
- Feely DE and Dyer JK** (1987) Localization of acid phosphatase activity in *Giardia lamblia* and *Giardia muris* trophozoites. *Journal of Protozoology* **34**, 80–83.
- Gaechter V, Schraner E, Wild P and Hehl AB** (2008) The single dynamin family protein in the primitive protozoan *Giardia lamblia* is essential for stage conversion and endocytic transport. *Traffic (Copenhagen, Denmark)* **9**, 57–71.
- Hehl AB and Marti M** (2004) Secretory protein trafficking in *Giardia intestinalis*. *Molecular Microbiology* **53**, 19–28.
- Hernandez Y, Castillo C, Roychowdhury S, Hehl A, Aley SB and Das S** (2007) Clathrin-dependent pathways and the cytoskeleton network are involved in ceramide endocytosis by a parasitic protozoan, *Giardia lamblia*. *International Journal for Parasitology* **37**, 21–32.
- Kattenbach WM, Pimenta PFP, de Souza W and Pinto da Silva P** (1991) *Giardia duodenalis*: a freeze-fracture, fracture-flip and cytochemistry study. *Parasitology Research* **77**, 651–658.
- Keister DB** (1983) Axenic culture of *Giardia lamblia* in TYI-S-33 medium supplemented with bile. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **77**, 487–488.
- Lanfredi-Rangel A, Attias M, Carvalho TMU, Kattenbach WM and De Souza W** (1998) The peripheral vesicles of trophozoites of the primitive protozoan *Giardia lamblia* may correspond to early and late endosomes and lysosomes. *Journal Structural Biology* **123**, 225–235.
- Lanfredi-Rangel A, Kattenbach WM, Diniz JAJ and de Souza W** (1999) Trophozoites of *Giardia lamblia* may have a Golgi-like structure. *FEMS Microbiology Letters* **181**, 245–251.
- Lindmark DG** (1988) *Giardia lamblia*: localization of hydrolase activities in lysosome-like organelles of trophozoites. *Experimental Parasitology* **65**, 141–147.
- Luján HD, Marotta A, Mowatt MR, Sciaky N, Lippincottschwartz J and Nash TE** (1995) Developmental induction of Golgi structure and function in the primitive eukaryote *Giardia lamblia*. *Journal of Biological Chemistry* **270**, 4612–4618.
- Makiuchi T and Nozaki T** (2014) Highly divergent mitochondrion-related organelles in anaerobic parasitic protozoa. *Biochimie* **100**, 3–17.
- Marie C and Petri WA Jr** (2014) Regulation of virulence of *Entamoeba histolytica*. *Annual Review of Microbiology* **68**, 493–520.
- Marti M, Li Y, Schraner EM, Wild P, Kohler P and Hehl AB** (2003) The secretory apparatus of an ancient eukaryote: protein sorting to separate export pathways occurs before formation of transient Golgi-like compartments. *Molecular Biology of the Cell* **14**, 1433–1447.
- McCaffery JM and Gillin FD** (1994) *Giardia lamblia*: ultrastructural basis of protein transport during growth and encystation. *Experimental Parasitology* **79**, 220–235.
- Midlej V, Meinig I, de Souza W and Benchimol M** (2013) A new set of carbohydrate-positive vesicles in encysting *Giardia lamblia*. *Protist* **164**, 261–271.
- Midlej V, Penha L, Silva R, de Souza W and Benchimol M** (2016) Mitosomal chaperone modulation during the life cycle of the pathogenic protist *Giardia intestinalis*. *European Journal of Cell Biology* **95**, 531–542.
- Nemanic PC, Owen RL, Stevens DP and Mueller JC** (1979) Ultrastructural observations on *Giardiasis* in a mouse model. II. Endosymbiosis and organelle distribution in *Giardia muris* and *Giardia lamblia*. *Journal of Infectious Diseases* **140**, 222–228.
- Orozco E, Guarneros G, Martínez-Palomo A and Sánchez T** (1983) *Entamoeba histolytica*. Phagocytosis as a virulence factor. *The Journal of Experimental Medicine* **158**, 1511–1521. <https://doi.org/10.1084/jem.158.5.1511>.
- Pereira-Neves A and Benchimol M** (2007) Phagocytosis by *Trichomonas vaginalis*: new insights. *Biology of the Cell* **99**, 87–101.
- Rivero MR, Vranich CV, Bisbal M, Maletto BA, Ropolo AS and Touz MC** (2010) Adaptor protein 2 regulates receptor-mediated endocytosis and cyst formation in *Giardia lamblia*. *Biochemical Journal* **428**, 33–45.
- Rivero MR, Miras SL, Feliziani C, Zamponi N, Quiroga R, Hayes SF, Rópolo AS and Touz MC** (2012) Vacuolar protein sorting receptor in *Giardia lamblia*. *PLoS ONE* **7**, e43712.
- Sogayar MIL and Gregorio EA** (1989) Uptake of bacteria by trophozoites of *Giardia duodenalis* (Say). *Annals of Tropical Medicine and Parasitology* **83**, 63–66.
- Stefanic S, Morf L, Kulangara C, Regös A, Sonda S, Schraner E, Spycher C, Wild P and Hehl AB** (2009) Neogenesis and maturation of transient Golgi-like cisternae in a simple eukaryote. *Journal of Cell Science* **122**, 2846–2856.
- Tachezy J and Smid O** (2008) Mitosomes in parasitic protists. In Tachezy J (ed.), *Hydrogenosomes and Mitosomes: Mitochondria of Anaerobic Eukaryotes*. Berlin: Springer-Verlag, pp. 201–230.
- Tachezy J, Sanchez LB and Muller M** (2001) Mitochondrial type iron-sulfur cluster 15 assembly in the amitochondriate eukaryotes *Trichomonas vaginalis* and *Giardia intestinalis*, as indicated by the phylogeny of IscS. *Molecular Biology Evolution* **18**, 1919–1928.
- Tai JH, Ong SJ, Chang SC and Su HM** (1993) *Giardia* virus enters *Giardia lamblia* WB trophozoite *via* endocytosis. *Experimental Parasitology* **76**, 165–174.
- Thiéry JP** (1967) Mise en évidence des polysaccharides sur coupes fines en microscopie électronique. *Journal of Microscopy* **6**, 987–1018.
- Touz MC, Kulakova L and Nash TE** (2004) Adaptor protein complex 1 mediates the transport of lysosomal proteins from a Golgi-like organelle to peripheral vacuoles in the primitive eukaryote *Giardia lamblia*. *Molecular Biology of the Cell* **15**, 3053–3060.
- Tovar J, Fischer A and Clark CG** (1999) The mitosome, a novel organelle related to mitochondria in the amitochondrial parasite *Entamoeba histolytica*. *Molecular Microbiology* **32**, 1013–1021.
- Tovar J, León-Avila G, Sánchez LB, Sutak R, Tachezy J, van der Giezen M, Hernández M, Müller M and Lucocq JM** (2003) Mitochondrial remnant organelles of *Giardia* Function in iron-sulphur protein maturation. *Nature* **426**, 172–176.
- Wampfler PB, Tosevski V, Nanni P, Spycher C and Hehl AB** (2014) Proteomics of secretory and endocytic organelles in *Giardia lamblia*. *PLoS ONE* **9**, e94089.
- Zumthor JP, Cernikova L, Rout S, Kaech A, Faso C and Hehl AB** (2016) Static clathrin assemblies at the peripheral vacuole-plasma membrane interface of the parasitic protozoan *Giardia lamblia*. *PLoS Pathogens* **12**, e1005756.