

## ON AN ESTIMATE FOR SOLUTIONS OF NONLINEAR ELLIPTIC VARIATIONAL INEQUALITIES<sup>1)</sup>

HARUO NAGASE

### Introduction

Let  $\Omega$  be a bounded domain in  $R^n$  with the boundary  $\partial\Omega$  of class  $C^{0,1}$  and  $E$  be a compact subset (resp. a compact subset on an  $(n-1)$ -dimensional hypersurface of class  $C^{0,1}$ ) in  $\Omega$ . We assume that the usual function spaces  $C^k(\bar{\Omega})$ ,  $C_0^k(\Omega)$ ,  $L^p(\Omega)$ ,  $W^{1,p}(\Omega)$  and  $W_0^{1,p}(\Omega)$  are known.

The first constraint condition is given by the following set:

$$(1) \quad K_1 = \{v \in W_0^{1,p}(\Omega); v(x) \in K(x) \text{ a.e. (resp. p.p.) } x \in E\},$$

where  $K(x)$  is a closed convex set in  $R^1$  depending on  $x$ .

Next let  $\partial_1\Omega$  and  $\partial_2\Omega$  be two disjoint open subsets of  $\partial\Omega$  such that  $\partial\Omega = \overline{\partial_1\Omega} \cup \overline{\partial_2\Omega}$  and  $\partial_1\Omega \neq \emptyset$ . We set

$$C_{(0)}^1(\bar{\Omega}) = \{v \in C^1(\bar{\Omega}); v = 0 \text{ in a neighborhood of } \overline{\partial_1\Omega}\}.$$

The completion of  $C_{(0)}^1(\bar{\Omega})$  with respect to the norm  $\|u\|_{1,p} = \|u\|_p + \|\nabla u\|_p$ <sup>2)</sup> is denoted by  $W_{(0)}^{1,p}(\Omega)$ . The following set  $K_2$  defines the second constraint condition:

$$(2) \quad K_2 = \{v \in W_{(0)}^{1,p}(\Omega); v(x) \in k(x) \text{ p.p. } x \in \overline{\partial_2\Omega}\},$$

where  $k(x)$  is also a closed convex set in  $R^1$  depending on  $x$ .

The aim of this paper is to establish an estimate for the solution  $u \in K_i$  of the following variational inequality:

$$(3) \quad \sum_{j=1}^n \langle a_j(x, \nabla u), \partial x_j(u - v) \rangle + (a_0(x, u), u - v) \leq (f, u - v)$$

for any  $v \in K_i$ ,

---

Received March 24, 1986.

<sup>1)</sup> The content of this paper was lectured by the author at the Nonlinear P.D.E. Symposium, held at Tokyo University from February 24 to 26, 1986.

<sup>2)</sup> For the sake of simplicity we write  $\|\cdot\|_{L^p(\Omega)} = \|\cdot\|_p$  and  $(\partial_1 u, \partial_{x_2} u, \dots, \partial_{x_n} u) = \nabla u$ .

where we describe the pairing between  $W_0^{1,p}(\Omega)(W_{(0)}^{1,p}(\Omega))$  and its dual by  $\langle \cdot, \cdot \rangle$  and the inner product of  $L^2(\Omega)$  by  $(\cdot, \cdot)$ .

At first we note that the norm  $\|u\|_{1,p}$  is equivalent to the norm  $\|\nabla u\|_p$  in  $W_0^{1,p}(\Omega)$  or  $W_{(0)}^{1,p}(\Omega)$  by Poincaré’s inequality.

In the case when  $K = K_1$  (resp.  $K = K_2$ ) and  $E = \Omega$  (resp.  $\partial_1\Omega = \emptyset$ ), many authors obtained a number of results with respect to the properties of solutions of (3) for operators  $a_j$  and  $K(x)$  (resp.  $k(x)$ ) of various types. Here we do not refer explicitly to such cases.

Before stating our theorem, we will refer to results related to our problems. D. Kinderlehrer ([23]) proved the existence of a Lipschitz continuous solution in  $R^2$  under the assumption that  $a_j = a_j(\eta)^{3)}$  satisfies  $\sum_{j=1}^n (a_j(\eta) - a_j(\eta'))(\eta_j - \eta'_j) \geq \nu|\eta - \eta'|^2$  ( $\nu > 0$ ) and  $E$  is a segment. In [13] J. Frehse discussed the continuity of the first order derivatives of solutions when  $a_j$  satisfies  $\sum_{i,j=1}^n (\partial a_j / \partial \eta_i)(\eta) \xi_i \xi_j \geq \nu|\xi|^2$  ( $\nu > 0$ ) and  $E$  is an  $(n - 1)$ -dimensional manifolds. Also G. H. Williams ([38]) proved the existence of a Lipschitz continuous solution for the case that  $a_j$  satisfies the inequality  $\sum_{j=1}^n (a_j(x, z, \eta) - a_j(x, z', \eta'))(\eta_j - \eta'_j) + (a_0(x, z, \eta) - a_0(x, z', \eta'))(z - z') \geq C|\eta - \eta'|^2$  ( $C > 0$ ) and  $E$  is a subset in  $\Omega$ . In the above three cases  $K(x)$  is only of the type  $K(x) = \{u(x) \geq \phi(x)\}$  for a given function  $\phi(x)$ . Additionally when  $n = 2$  and  $\partial_2\Omega = (a, b) \times \{x_2 = 0\}$  particularly, D. Kinderlehrer ([24]) showed that solutions belong to class  $C^{1,\alpha}(\Omega \cup \partial_2\Omega)$  under the same condition on  $a_j$  as in [23] and with the assumption that  $k(x) = \{u(x) \geq \psi(x)\}$  for some function  $\psi(x)$ .

Secondly we mention results for the cases when  $a_j$  degenerate. To begin with we impose the next conditions on  $a_j$  ( $j = 0, 1, \dots, n$ ):

$$(4) \quad \begin{cases} \sum_{j=1}^n a_j(x, z, \eta)\eta_j \geq a|\eta|^p - b(x)|z|^p - l(x), \\ |a_j(x, z, \eta)| \leq a'|\eta|^{p-1} + g(x)|z|^{p-1} + h(x), \quad j = 1, \dots, n \\ |a_0(x, z, \eta)| \leq d(x)|\eta|^{p-1} + e(x)|z|^{p-1} + m(x). \end{cases}$$

Assuming that  $a_j$  ( $j = 1, \dots, n$ ) are independent of  $z$  and that  $b, g, h$ , and  $l$  equal to zero and that furthermore  $a_0 = |z|^{p-2}z$ , H. Beirão da Veiga ([35]) obtained the  $L^\infty(\Omega)$  estimate for solutions of variational inequalities under the constraint condition that  $u(x) \geq \phi_i(x)$  on  $E_i$  ( $i = 1, \dots, m$ ). In [35] the boundary condition is of either Dirichlet, Neumann or Signorini’s

<sup>3)</sup> In place of the operator  $a_j(x, u, \nabla u)$  we consider the function  $a_j(x, z, \eta)$  defined on  $\Omega \times R^1 \times R^n$ .

type, furthermore the mixed type of these conditions is also treated. Moreover in [36] he established the  $L^\infty(\Omega)$  estimate and the  $C^\alpha(\bar{\Omega})$  estimate under the assumption (4) for the constraint condition  $K = \{v \in W^{1,p}(\Omega); v(x) \geq \psi(x) \text{ on } \partial\Omega, v(x) = \psi(x) \text{ on } \Gamma\}$ . H. Beirão da Veiga-F. Conti ([37]) proved that solutions belong to  $C^\alpha(\bar{\Omega})$  when  $a_j (j = 1, \dots, n)$  are independent of  $z$  and  $a_0 = 0$  in (4), in addition  $a_j$  satisfies the condition  $\sum_{j=1}^n (a_j(x, \eta) - a_j(x, \eta'))(\eta_j - \eta'_j) > 0$ . Here the constraint condition is of type:  $u(x) \geq \phi(x)$  on  $E$ . When  $b = e = g = 0$  in (4), A. Domarkas ([9], [10]) showed that solutions belong to  $C^\alpha(\bar{\Omega})$  for the constraint condition  $K: K = \{v \in W_0^{1,p}(\Omega); v(x) \geq \phi_1(x) \text{ on } E_1 \text{ and } v(x) \leq \phi_2(x) \text{ on } E_2\}$  or  $K = \{v \in W^{1,p}(\Omega); v(x) \geq \psi_1(x) \text{ on } \Gamma_1 \text{ and } v(x) \leq \psi_2(x) \text{ on } \Gamma_2\}$ . In the above works  $E$  and  $E_i$  are subsets of  $\Omega$  and  $\Gamma, \Gamma_i$  are subsets of  $\partial\Omega$ , additionally  $\phi_i(x), \psi(x)$  and  $\psi_i(x)$  are some given functions.

Throughout this paper let  $p \geq 2$  and let us assume that  $\alpha \geq 0$  if  $p \geq n, 0 \leq \alpha < n(p - 1)/(n - p) - 1$  if  $p < n$ .

Now we impose the following assumptions on  $a_j (j = 0, 1, \dots, n)$ :

ASSUMPTION A.

(I) (i)  $a_j = a_j(x, \eta) \in C^0(\Omega \times R^n) \cap C^1(\Omega \times (R^n - \{0\}))$ ,  $a_j(x, 0) = 0$ ,  
 $j = 1, \dots, n$ .

(ii)  $\sum_{i,j=1}^n \frac{\partial a_j}{\partial \eta_i}(x, \eta) \xi_i \xi_j \geq \gamma(\kappa_0 + |\eta|^{p-2})|\xi|^2$ ,

$$\left| \frac{\partial a_j}{\partial \eta_i}(x, \eta) \right| \leq A(\kappa_0 + |\eta|^{p-2}), \quad i = 1, \dots, n,$$

$$\left| \frac{\partial a_j}{\partial x_k}(x, \eta) \right| \leq A(\kappa_0|\eta| + |\eta|^{p-1}), \quad k = 1, \dots, n.$$

(II) (i)  $a_0 = a_0(x, z) \in C^0(\Omega \times R^1) \cap C^1(\Omega \times (R^1 - \{0\}))$ ,  $a_0(x, 0) = 0$ .

(ii)  $\frac{\partial a_0}{\partial z}(x, z) \geq 0$ ,  $\left| \frac{\partial a_0}{\partial z}(x, z) \right| \leq A(|z|^\alpha + 1)$ ,

$$\left| \frac{\partial a_0}{\partial x_k}(x, z) \right| \leq A(|z|^{\alpha+1} + 1), \quad k = 1, \dots, n.$$

in the above  $x \in \Omega, z \in R^1 - \{0\}, \eta \in R^n - \{0\}$  and  $\xi \in R^n$ . And  $\kappa_0$  is a non-negative constant,  $\gamma$  and  $A$  are some positive constants.

Hereafter we write by the same  $C$  all constants independent of  $u$  and  $f$ , which appear in (3). We define the following function  $\Phi(x)$  in  $R^n$ :

$$\Phi(x) = \begin{cases} \text{dis}(x, E \cup \partial\Omega) & \text{for the case of } K = K_1 \\ \text{dis}(x, \partial\Omega) & \text{for the case of } K = K_2 \end{cases}$$

where  $\text{dis}(A, B)$  is the distance between  $A$  and  $B$ .

Our theorem is as follows.

**THEOREM.** *Under the assumption A if  $f \in W^{1,p^*}(\Omega)$  (resp.  $L^2(\Omega)$ ) in the case of  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ), the following assertions hold for the solution  $u$  of the nonlinear variational inequality (3).*

(I) *In any case  $\Phi(x)a_j(x, \nabla u) \in W^{1,p^*}(\Omega)$ ,  $\Phi(x)|\nabla u|^{p/2} \in W^{1,2}(\Omega)$ ,  $j = 1, \dots, n$ .*

(II) *If  $\kappa_0 = 0$ ,*

$$\|\Phi(\cdot)a_j(\cdot, \nabla u)\|_{1,p^*}^{p^*} + \|\Phi(\cdot)|\nabla u|^{p/2}\|_{1,2}^2 \leq C(1 + \|f\|_{1,p^*}^{p^*} + \|f\|_{p^*}^{(\alpha+2)/(p-1)}).$$

(III) *If  $\kappa_0 \neq 0$ ,  $\Phi(x)\nabla u \in W^{1,2}(\Omega)$  and*

$$\|\Phi(\cdot)\nabla u\|_{1,2}^2 + \|\Phi(\cdot)a_j(\cdot, \nabla u)\|_{1,p^*}^{p^*} + \|\Phi(\cdot)|\nabla u|^{p/2}\|_{1,2}^2 \leq C(1 + \|f\|_2^{2\beta/p}),$$

where  $\beta = \max(p, \alpha + 2)$  and  $p^*$  is the dual number of  $p$ , i.e.,  $1/p + 1/p^* = 1$ .

In connection to the estimate in our theorem the following results were obtained by G. H. Yakovlev. He gave first the estimate of  $\|\partial_{x_j}(|\partial_{x_i} u|^{p/2})\|_{L^2(\Omega_\delta)}$  for solutions of the variational problem which is derived from the functional of the form  $I(v) = \int_\Omega \left\{ \sum_{j=1}^n |\partial_{x_j} v|^p + a(x)v \right\} dx$  with the Dirichlet boundary condition ([40]). Here  $\Omega_\delta = \{x \in \Omega; \text{dis}(x, \partial\Omega) > \delta\}$ .

In [43] also he obtained estimates of

$$\|a_j(\cdot, u, \nabla u)\|_{W^{1,p^*}(\Omega_\delta)} \quad \text{and} \quad \left\| \frac{\partial a_j}{\partial x_k}(\cdot, u, \nabla u) \right\|_{W^{1,p^*}(\Omega_\delta)}$$

for weak solutions of the nonlinear elliptic equation  $\sum_{j=1}^n \partial_{x_j}(a_j(x, u, \nabla u)) = a_0(x, u, \nabla u)$  under weaker assumptions than ours. The method in [43] is to use the quotient of differentials with the usual parallel transformation, namely,

$$h^{-1}[u(x_1, x_2, \dots, x_{i-1}, x_i + h, x_{i+1}, \dots, x_n) - u(x)],$$

which implies that the estimate of the norm is obliged to be restricted in  $\Omega_\delta$ . In this paper we use a transformation with weight function  $\Phi(x)$ , so we can estimate the norm in the whole  $\Omega$ . Our estimates are very similar to that of G. H. Yakovlev ([41]), where a nonlinear elliptic equation with the Dirichlet boundary condition was treated. He prepared the estimate for smooth solutions and applied the Galerkin's method, so his technique can not be applied to our variational inequality.

Finally we refer to the regularity of weak solutions of the nonlinear elliptic equation. Let us put  $v = u \pm \phi$  in (3) for a solution  $u$  and any function  $\phi$  in  $C_0^\infty(\Omega \setminus E)$  (resp.  $C_0^\infty(\Omega)$ ) in the case of  $K = K_1$  (resp.  $K = K_2$ ),

then we see that  $u$  is a weak solution of the nonlinear elliptic equation

$$(5) \quad - \sum_{j=1}^n \partial_{x_j}(a_j(x, \nabla u)) + a(x, u) = f$$

in  $\Omega \setminus E$  (resp.  $\Omega$ ).

J. C. Evans ([11]) and J. L. Lewis ([26]) proved that weak solutions of (5) belong to  $C_{loc}^{1,\alpha}(\Omega \setminus E)$  (resp.  $C_{loc}^{1,\alpha}(\Omega)$ ) when  $a_j = |\nabla u|^{p-2} \partial_{x_j} u$  and  $a_0 = 0$ . Besides E. DiBenedetto ([8]) and P. Tolksdorf ([33]) showed the same results under weaker assumptions on  $a_j$  and  $a_0$  than those of [11] and [26].

1.

The existence of a unique solution for the variational inequality (3) is derived from Lemma 1.2 in this section.

First we will prepare the following lemma which will be frequently used in this paper:

LEMMA 1.1. *Under the assumption A the following assertions hold:*

(i)  $a_j (j = 0, 1, \dots, n)$  are estimated in such a way that

$$(1.1) \quad |a_j(x, \eta)| \leq C(\kappa_0 + |\eta|^{p-2})|\eta|, \quad j = 1, \dots, n,$$

$$(1.2) \quad |a_0(x, z)| \leq C(1 + |z|^q)|z|.$$

(ii) (P. Tolksdorf [33, p. 129] and P. Lindqvist [28, p. 310]) *There exists a positive number  $\gamma_0$  depending only on  $\gamma$  and  $p$  such that*

$$(1.3) \quad \sum_{j=1}^n (a_j(x, \eta) - a_j(x, \eta'))(\eta_j - \eta'_j) \geq \gamma_0(\kappa_0 + |\eta|^{p-2} + |\eta'|^{p-2})|\eta - \eta'|^p,$$

$$(1.4) \quad (a_0(x, z) - a_0(x, z'))(z - z') \geq 0.$$

Here  $x \in \Omega$ ,  $\eta, \eta' \in R^n$  and  $z, z' \in R^1$ .

*Proof.*

(i) The estimates (1.1) and (1.2) are derived from (I)-(ii) and (II)-(ii) in the assumption A and the equalities

$$a_j(x, \eta) = \int_0^1 \sum_{i=1}^n \frac{\partial a_j}{\partial \eta_i}(x, t\eta) \eta_i dt, \quad a_0(x, z) = \int_0^1 \frac{\partial a_0}{\partial z}(x, tz) z dt.$$

(ii) Without loss of the generality we may assume that  $|\eta| \leq |\eta'|$ . According to simple calculations and (I)-(ii) in the assumption A, we see that

$$\begin{aligned}
 (1.5) \quad & \sum_{j=1}^n (a_j(x, \eta) - a_j(x, \eta')) (\eta_j - \eta'_j) \\
 &= \sum_{j=1}^n \int_0^1 \frac{d}{dt} a_j(x, t\eta + (1-t)\eta') (\eta_j - \eta'_j) dt \\
 &\geq \int_0^{1/4} \sum_{i,j=1}^n \frac{\partial a_j}{\partial \eta_i} (x, t\eta + (1-t)\eta') (\eta_i - \eta'_i) (\eta_j - \eta'_j) dt \\
 &\geq \gamma \int_0^{1/4} (\kappa_0 + |t\eta + (1-t)\eta'|^{p-2}) |\eta - \eta'|^2 dt.
 \end{aligned}$$

On the other hand it holds that

$$\begin{aligned}
 (1.6) \quad & |t\eta + (1-t)\eta'| \geq (1-t)|\eta'| - t|\eta| \\
 &= (3/4 - t)|\eta'| - (t + 1/4)|\eta| + 1/4(|\eta'| + |\eta|) \\
 &\geq (t + 1/4)(|\eta'| - |\eta|) + 1/4(|\eta'| + |\eta|) \\
 &\geq 1/4(|\eta'| + |\eta|),
 \end{aligned}$$

for all  $t \in [0, 1/4]$ . The inequalities (1.5) and (1.6) imply (1.3). The estimate (1.4) can be proved more easily. Q.E.D.

*Remark.* In (1.6) we note that the estimate  $|t\eta + (1-t)\eta'| \geq (1/4) \cdot |\eta' - \eta|$  holds for all  $t \in [0, 1/4]$ . Thus we have the following estimate in place of (1.3):

$$(1.7) \quad \sum_{j=1}^n (a_j(x, \eta) - a_j(x, \eta')) (\eta_j - \eta'_j) \geq \gamma_0 (\kappa_0 + |\eta - \eta'|^{p-2}) |\eta - \eta'|^2.$$

This is due to K. L. Kuttler Jr. ([25]).

**LEMMA 1.2.** For  $u, v \in W_0^{1,p}(\Omega) (W_{(0)}^{1,q}(\Omega))$  we define the operator  $A$  in such a way that  $(A(u), v) = \sum_{j=1}^n \langle a_j(x, \nabla u), \partial_{x_j} v \rangle + (a_0(x, u), v)$ , then the operator  $A$  is pseudo-monotone and coercive from  $W_0^{1,p}(\Omega) (W_{(0)}^{1,p}(\Omega))$  to its dual space.

*Proof.* We write  $W_0^{1,p}(\Omega) (W_{(0)}^{1,p}(\Omega))$  simply by  $V$  and its dual space by  $V'$ .

In order to prove the pseudo-monotonicity of the operator  $A$ , it is enough to show that the operator  $A$  is (i) bounded, (ii) hemi-continuous and (iii) monotone (J.L. Lions [29], p. 179).

(i) Boundedness.

According to the definition

$$(1.8) \quad |(A(u), v)| \leq \sum_{j=1}^n |\langle a_j(x, \nabla u), \partial_{x_j} v \rangle| + |(a_0(x, u), v)|.$$

We estimate each term on the right-hand side of (1.8).

At first we obtain

$$(1.9) \quad \begin{aligned} |\langle a_j(x, \nabla u), \partial_{x_j} v \rangle| &\leq C \int_{\Omega} (\kappa_0 + |\nabla u|^{p-2}) |\nabla u| \cdot |\nabla v| \, dx \\ &\leq C(\|\nabla u\|_{p^*} \|\nabla v\|_p + \|\nabla u\|_p^{p-1} \|\nabla v\|_p), \end{aligned}$$

by (1.1) and Hölder’s inequality. Since  $p \geq 2$ ,  $p$  and  $p^*$  satisfy  $1 < p^* \leq 2 \leq p$ . Thus the inequality  $\|\nabla u\|_{p^*} \leq C\|\nabla u\|_p$  is verified. Therefore from (1.9) we deduce the estimate

$$(1.10) \quad |\langle a_j(x, \nabla u), \partial_{x_j} v \rangle| \leq C(\|u\|_V + \|u\|_V^{p-1}) \|v\|_V.$$

On account of Lemma 1.1 and Hölder’s inequality, we obtain similarly the inequality

$$(1.11) \quad |(a_0(x, u), v)| \leq \{C\|u\|_{p^*} \|v\|_p + (\|u\|_{(\alpha+1)p^*})^{\alpha+1} \|v\|_p\}.$$

By Poincaré’s inequality, we get

$$(1.12) \quad \|u\|_{p^*} \leq C\|\nabla u\|_{p^*} \leq C\|\nabla u\|_p \leq C\|u\|_V,$$

$$(1.13) \quad \|v\|_p \leq C\|\nabla v\|_p \leq C\|v\|_V.$$

And Sobolev’s imbedding theorem yields the inequality

$$(1.14) \quad \|u\|_{(\alpha+1)p^*} \leq C\|u\|_{1,p},$$

because  $0 \leq \alpha < n(p - 1)/(n - p) - 1$  for  $p < n$ . From (1.11)–(1.14) it follows that

$$(1.15) \quad |(a_0(x, u), v)| \leq C(\|u\|_V + \|u\|_V^{\alpha+1}) \|v\|_V.$$

Combining (1.8), (1.10) with (1.15), we derive the following estimate from the definition of the dual norm of  $A(u)$  in  $V'$ :

$$(1.16) \quad \|A(u)\|_{V'} \leq C(\|u\|_V^{p-1} + \|u\|_V^{\alpha+1} + \|u\|_V).$$

(ii) Hemi-continuity.

Let us suppose that  $|\lambda| \leq \lambda_0$ . For  $u, v$  and  $w \in V$ , it is valid that

$$(1.17) \quad (A(u + \lambda v), w) = \int_{\Omega} \left\{ \sum_{j=1}^n a_j(x, \nabla(u + \lambda v)) \partial_{x_j} w + a_0(x, u + \lambda v) w \right\} dx,$$

by the definition of the operator  $A$ . Applying the inequalities (1.1) and (1.2) to each term of the integrand in (1.17), we have

$$|a_j(x, \nabla(u + \lambda v))\partial_{x_j} w| \leq C\{\kappa_0(|\nabla u| + \lambda_0|\nabla v|) + |\nabla u|^{p-1} + \lambda_0^{p-1}|\nabla v|^{p-1}\}|\nabla w|,$$

$$|a_0(x, u + \lambda v)w| \leq C\{(|u| + \lambda_0|v|) + |u|^{\alpha+1} + \lambda_0^{\alpha+1}|v|^{\alpha+1}\}|w|.$$

By the same way as in the proof of (i), we can prove that each term on the right-hand sides is integrable. Hence  $(A(u + \lambda v), w)$  is continuous with respect to  $\lambda$  by Lebesgue’s convergence theorem.

(iii) Monotonicity.

From the remark after Lemma 1.1, it holds for any  $u, v \in V$  that

$$(1.18) \quad (A(u) - A(v), u - v) \geq \gamma_0(\kappa_0\|\nabla(u - v)\|_2^2 + \|\nabla(u - v)\|_p^p) \geq 0.$$

Consequently, we have shown the pseudo-monotonicity of the operator  $A$ .

Finally we prove that the operator  $A$  is coercive. Setting  $v = 0$  in (1.18), we have

$$(A(u), u) \geq \gamma_0(\kappa_0\|\nabla u\|_2^2 + \|\nabla u\|_p^p).$$

Hence

$$\frac{(A(u), u)}{\|u\|_V} \geq C\gamma_0\|u\|_V^{p-1} \longrightarrow \infty \quad \text{as } \|u\|_V \longrightarrow \infty.$$

Therefore the operator  $A$  is coercive.

Q.E.D.

As we have mentioned at the beginning of this section, a solution of the variational inequality (3) exists for any  $f \in V'$  (J. L. Lions [29, p. 247]). The uniqueness of solutions follows immediately from (1.18).

The estimate for the gradient of the solution  $u$  is carried out in the next lemma.

LEMMA 1.3. *Under the assumption A if  $f \in L^{p^*}(\Omega)$  (resp.  $L^2(\Omega)$ ) and  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ), the gradient of the solution  $u$  of the variational inequality (3) is estimated as follows:*

$$(1.19) \quad \kappa_0\|\nabla u\|_2^2 + \|\nabla u\|_p^p \leq C(1 + \|f\|_{p^*}^{p^*}) \quad (\text{resp. } C(1 + \|f\|_2^2)).$$

*Proof.* We denote by  $u_0$  the particular solution of (3) for  $f = 0$ . We easily see that

$$(1.20) \quad \sum_{j=1}^n \langle a_j(x, \nabla u) - a_j(x, \nabla u_0), \partial_{x_j}(u - u_0) \rangle + (a_0(x, u) - a_0(x, u_0), u - u_0) \leq (f, u - u_0).$$

Applying the inequalities (1.4) and (1.7) to the left-hand side, we get



$$(1.21) \quad \gamma_0(\kappa_0 \|\nabla u - \nabla u_0\|_2^2 + \|\nabla u - \nabla u_0\|_p^p) \leq |(f, u - u_0)|.$$

Using Hölder’s inequality and Poincaré’s inequality for the right-hand side of (1.21), we obtain the estimate (1.19). Q.E.D.

We give a sufficient condition to assure that a sequence of functions converges weakly in  $L^q(\Omega)$ .

**LEMMA 1.4.** *Let  $u$  be a distribution in  $\Omega$  and let  $\{u_\nu\}_{\nu=1}^\infty$  be a sequence in  $L^q(\Omega)$  ( $1 < q < \infty$ ) such that the norms  $\|u_\nu\|_q$  are uniformly bounded. If for any  $\phi \in C_0^\infty(\Omega)$ ,*

$$(u_\nu, \phi) \longrightarrow (u, \phi) \quad \text{as } \nu \longrightarrow \infty,$$

*then  $u$  belongs to  $L^q(\Omega)$  and the sequence  $u_\nu$  converges weakly to  $u$  in  $L^q(\Omega)$ .*

*Proof.* From the assumption it holds for any  $\phi \in C_0^\infty(\Omega)$ ,

$$(1.22) \quad |(u, \phi)| \leq (\overline{\lim}_{\nu \rightarrow \infty} \|u_\nu\|_q) \|\phi\|_{q^*}.$$

For arbitrary  $v \in L^{q^*}(\Omega)$  we take a sequence  $\{\phi_k\}_{k=1}^\infty$  in  $C_0^\infty(\Omega)$  such that  $\|\phi_k - v\|_{q^*} \rightarrow 0$  as  $k \rightarrow \infty$ , then from (1.22) the sequence  $\{(u, \phi_k)\}_{k=1}^\infty$  is a Cauchy sequence. Accordingly,  $\lim_{k \rightarrow \infty} (u, \phi_k)$  exists and from (1.22) it is trivial that the limit depends only on  $v$  and does not depend on any choice of the sequence. Hence we can express  $\lim_{k \rightarrow \infty} (u, \phi_k) = l_u(v)$ . It is easy to verify that  $l_u$  is a linear functional on  $L^{q^*}(\Omega)$ , so there exists an element  $L_u$  in  $L^q(\Omega)$  ( $= (L^{q^*}(\Omega))'$ ) such that  $l_u(v) = (L_u, v)$  for all  $v \in L^{q^*}(\Omega)$ . The definition of  $l_u(v)$  ensures the equality  $(u, \phi) = (L_u, \phi)$  for any function  $\phi$  in  $C_0^\infty(\Omega)$ , which is dense in  $L^{q^*}(\Omega)$ . Thus we can conclude that  $u = L_u$  and therefore  $u \in L^q(\Omega)$ . The remainder of the proof is due to Theorem 3 in [44, p. 121]. Q.E.D.

**2.**

We introduce a coordinate transformation with the weight function  $\Phi(x)$  and prepare some results with respect to it. We refer to some lemmas in [20].

Let  $h$  be a non-zero vector in  $R^n$  with the length  $h = |h|$ . Hereafter  $h$  is assumed to be sufficiently small. As mentioned in the introduction we put  $\Phi(x) = \text{dis}(x, \partial\Omega \cup E)$  (resp.  $\text{dis}(x, \partial\Omega)$ ) for  $K = K_1$  (resp.  $K = K_2$ ) and we consider the transformation of the coordinates:

---

<sup>4)</sup>  $q^*$  is the dual number of  $q$ , i.e.,  $q^* = q/(q-1)$ .

$$(2.1) \quad \Phi_h: y = x + \Phi(x)h.$$

We write  $h = (h_1, h_2, \dots, h_n)$  and  $J = \partial(y_1, y_2, \dots, y_n)/\partial(x_1, x_2, \dots, x_n)$ , then we have

$$(2.2) \quad J = \begin{bmatrix} 1 + h_1\partial_{x_1}\Phi & h_2\partial_{x_1}\Phi & \cdots & h_n\partial_{x_1}\Phi \\ h_1\partial_{x_2}\Phi & 1 + h_2\partial_{x_2}\Phi & \cdots & h_n\partial_{x_2}\Phi \\ \vdots & \vdots & \ddots & \vdots \\ h_1\partial_{x_n}\Phi & h_2\partial_{x_n}\Phi & \cdots & 1 + h_n\partial_{x_n}\Phi \end{bmatrix}.$$

Let us put  $e = h^{-1}h$  and let  $e$  be arbitrarily fixed. Noting that  $|\partial_{x_j}\Phi| \leq 1$  ([20, p. 57]), we see that the determinant  $J$  of  $J$  is not zero for sufficiently small  $h$ , therefore the mapping  $\Phi_h$  and its inverse  $\Phi_h^{-1}$  are both one-to-one from  $R^n$  onto itself. If we set  $\Psi(y) = -\Phi(x) (= -\Phi(\Phi_h^{-1}(y)))$ , it is written

$$(2.3) \quad \Phi_h^{-1}: x = y + \Psi(y)h.$$

Here we remark that from (2.2) we can put  $J = 1 + hJ_1$  and the determinant  $J^{-1}$  of the Jacobian  $J^{-1}$  connected with the inverse transformation  $\Phi_h^{-1}$  can be described in the form  $J^{-1} = 1 + hJ_2$ , where  $J_1$  and  $J_2$  are uniformly bounded in  $x \in \Omega$  and  $h \in R^n$ . Furthermore the transformation  $\Phi_h$  maps  $x \in \Omega$  to  $y \in \Omega$  and  $x \in \Omega^c$  to  $y \in \Omega^c$  respectively, so it is a one-to-one mapping from  $\Omega$  onto itself.

Now we define

$$(2.4) \quad \begin{cases} (S_h u)(x) = u(x + \Phi(x)h), & (T_h u)(y) = u(y + \Psi(y)h), \\ (P_h u)(x) = h^{-1}[(S_h u)(x) - u(x)], \\ (Q_h u)(y) = h^{-1}[(T_h u)(y) - u(y)]. \end{cases}$$

Hereafter we write simply by  $S_h u, T_h u, \dots$  the functions  $(S_h u)(x), (T_h u)(y), \dots$ , respectively.

LEMMA 2.1 ([20, p. 58, 59]).

(i) We have

$$(2.5) \quad \begin{cases} \nabla_x(S_h u) = S_h \nabla_x u + h(e \cdot S_h \nabla_x u) \nabla_x \Phi, \\ \nabla_y(T_h u) = T_h \nabla_y u + h(e \cdot T_h \nabla_y u) \nabla_y \Psi, \\ \nabla_x(P_h u) = P_h \nabla_x u + (e \cdot S_h \nabla_x u) \nabla_x \Phi, \\ \nabla_y(Q_h u) = Q_h \nabla_y u + (e \cdot T_h \nabla_y u) \nabla_y \Psi. \end{cases}$$

(ii) If  $u \in W^{1,q}(\Omega)$  ( $1 < q < \infty$ ), there exists a constant  $C$  independent of  $h$  and  $u$  such that

$$(2.6) \quad \|P_h u\|_q, \quad \|Q_h u\|_q \leq C \|\nabla u\|_q.$$

We do not repeat the proof of the above lemma, since it is parallel to that of [20].

The following lemma is as important as Lemma 1.4 for the proof of our theorem.

LEMMA 2.2. *If  $u \in L^q(\Omega)$  ( $1 < q < \infty$ ), then for any function  $\phi$  in  $C_0^\infty(\Omega)$  it holds that*

$$(2.7) \quad (P_h u - J_2 u, \phi) \longrightarrow ((e \cdot \nabla)(\Phi u), \phi) \quad \text{as } h \longrightarrow 0,$$

where the derivative  $(e \cdot \nabla)(\Phi u)$  of  $\Phi u$  is in the sense of the distribution.

*Proof.* For any  $u \in L^q(\Omega)$  and  $\phi \in L^{q^*}(\Omega)$  the formula

$$(2.8) \quad (P_h u, \phi) - (J_2 u, \phi) = - (S_h u, P_h \phi)$$

holds. The function  $\phi$  belongs to  $C_0^\infty(\Omega)$  in this case, so we can prove the following convergence by the similar technique as in [20] (Lemma 5 in p. 59):

$$(2.9) \quad P_h \phi \longrightarrow \Phi(e \cdot \nabla)\phi \quad \text{in } L^{q^*}(\Omega) \quad \text{as } h \longrightarrow 0.$$

On the other hand it is an immediate consequence that

$$(2.10) \quad S_h u \longrightarrow u \quad \text{in } L^q(\Omega) \quad \text{as } h \longrightarrow 0,$$

from the fact that  $u \in L^q(\Omega)$ . Therefore the right-hand side on (2.8) tends to  $-(u, \Phi(e \cdot \nabla)\phi)$  by virtue of (2.9) and (2.10). In this way we arrive at the assertion of this lemma. Q.E.D.

### 3.

This section is devoted to the statement and the proof of the main proposition in this paper. It is very important for the proof of our theorem.

PROPOSITION. *Let us assume the assumption A. Then the following estimates hold for the solution  $u$  of the variational inequality (3):*

(i) *If  $\kappa_0 = 0$  and  $f \in W^{1,p^*}(\Omega)$ ,*

$$\| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \leq C(1 + \|f\|_{1,p^*}^{p^*} + \|f\|_{p^*}^{(\alpha+2)/(p-1)}).$$

(ii) *If  $\kappa_0 \neq 0$  and  $f \in L^2(\Omega)$ ,*

$$\| |S_h \nabla u \|_2^2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \leq C(1 + \|f\|_2^{2\beta/p}).$$

where  $\beta = \max(p, \alpha + 2)$ .

*Proof.* From now on we write simply  $\partial_{x_j} = \partial_j$  or  $\partial_{y_j} = \partial_j$  and we abbreviate the notation of sums.

As  $S_h u$  and  $T_h u$  belong to  $K_i$  ( $i = 1$  or  $2$ ) (see p. 60 in [20]), so we can put  $v = S_h u$  and  $T_h u$  in (3). Hence we obtain the inequalities

$$\begin{aligned} \langle a_j(x, \nabla u), \partial_j(u - S_h u) \rangle + (a_0(x, u), u - S_h u) &\leq (f, u - S_h u), \\ \langle a_j(y, \nabla u), \partial_j(u - T_h u) \rangle + (a_0(y, u), u - T_h u) &\leq (f, u - T_h u). \end{aligned}$$

Adding these two inequalities, we have

$$\begin{aligned} (3.1) \quad &\langle a_j(x, \nabla u), \partial_j(u - S_h u) \rangle + \langle a_j(y, \nabla u), \partial_j(u - T_h u) \rangle \\ &+ (a_0(x, u), u - S_h u) + (a_0(y, u), u - T_h u) \\ &\leq (f, u - S_h u) + (f, u - T_h u). \end{aligned}$$

Denoting each term on the left-hand side by  $I_j$  ( $j = 1, 2, 3, 4$ ) in turn from the left, we write  $I_1$  and  $I_2$  as follows by Lemma 2.1:

$$\begin{aligned} I_1 &= \langle a_j(x, \nabla u), \partial_j u - S_h \partial_j u \rangle - h \langle a_j(x, \nabla u), (\mathbf{e} \cdot S_h \nabla u) \partial_j \Phi \rangle, \\ I_2 &= \langle (1 + hJ_1) S_h a_j(x, \nabla u), S_h \partial_j u - \partial_j u \rangle - h \langle a_j(y, \nabla u), (\mathbf{e} \cdot T_h \nabla u) \partial_j \Psi \rangle. \end{aligned}$$

Consequently, we get

$$\begin{aligned} (3.2) \quad I_1 + I_2 &= \langle S_h a_j - a_j, S_h \partial_j u - \partial_j u \rangle + h \langle J_1 S_h a_j, S_h \partial_j u - \partial_j u \rangle \\ &- h \langle a_j, (\mathbf{e} \cdot T_h \nabla u) \partial_j \Psi \rangle - h \langle a_j, (\mathbf{e} \cdot S_h \nabla u) \partial_j \Phi \rangle. \end{aligned}$$

After putting the right-hand side of (3.2) by  $\sum_{j=5}^8 I_j$ , we estimate each term  $I_j$ . We rewrite  $I_5$  in the form

$$\begin{aligned} (3.3) \quad I_5 &= \langle a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u), S_h \partial_j u - \partial_j u \rangle \\ &+ \langle a_j(x, S_h \nabla u) - a_j(x, \nabla u), S_h \partial_j u - \partial_j u \rangle. \end{aligned}$$

For the first term in (3.3), the following estimate holds:

$$\begin{aligned} (3.4) \quad &|\langle a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u), S_h \partial_j u - \partial_j u \rangle| \\ &\leq Ch^2 \int_{\Omega} (\kappa_0 |S_h \nabla u| \cdot |P_h \nabla u| + |S_h \nabla u|^{p-1} |P_h \nabla u|) dx \\ &\leq Ch^2 (\kappa_0 \|\nabla u\|_2 \|P_h \nabla u\|_2 + \|\nabla u\|_p^{p/2} \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2). \end{aligned}$$

Here we have used the equality

$$a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u) = \int_0^1 \sum_{k=1}^n \frac{\partial a_j}{\partial x_k} (x + \theta \Phi(x) \mathbf{h}, S_h \nabla u) h_k \Phi(x) d\theta$$

and the inequality in the assumption A.

Secondly the remaining term in (3.3) is estimated from below by (1.3) as follows:

$$(3.5) \quad \langle a_j(x, S_h \nabla u) - a_j(x, \nabla u), S_h \partial_j u - \partial_j u \rangle \geq h^2 \gamma_0 \{ \kappa_0 \|P_h \nabla u\|_2^2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \}.$$

In this way (3.3), (3.4) and (3.5) yield that

$$(3.6) \quad I_5 \geq \frac{\gamma_0 h^2}{2} \{ \kappa_0 \|P_h \nabla u\|_2^2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \} - Ch^2 (\kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^2).$$

Next we see by Lemma 1.1 that

$$(3.7) \quad |I_6| \leq Ch^2 \int_{\Omega} (\kappa_0 + |S_h \nabla u|^{p-2}) |S_h \nabla u| \cdot |P_h \nabla u| dx \leq Ch^2 \{ \kappa_0 \|\nabla u\|_2 \|P_h \nabla u\|_2 + \|\nabla u\|_p^{p/2} \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2 \}.$$

Before proceeding to successive terms  $I_7$  and  $I_8$ , we note that we can write  $\nabla_y \Psi = \nabla_x \Psi(\mathbf{J}^{-1}) = -(\nabla_x \Phi)(I + hH)$ , where  $I$  is the unit matrix and each component of the matrix  $H$  is essentially bounded (see p. 64 in [20]). Consequently, it is seen that

$$(3.8) \quad \begin{aligned} I_7 + I_8 &= -h \{ \langle a_j, (\mathbf{e} \cdot T_h \nabla u) [(-\nabla_x \Phi(x))(I + hH)]_j \rangle \\ &\quad + \langle a_j, (\mathbf{e} \cdot S_h \nabla u) \partial_j \Phi \rangle \}^{5)} \\ &= -h \{ -\langle (1 + hJ_1) S_h a_j, (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle + \langle a_j, (\mathbf{e} \cdot S_h \nabla u) \partial_j \Phi \rangle \} \\ &\quad + h^2 \langle a_j, (\mathbf{e} \cdot T_h \nabla u) (\nabla_x \Phi H)_j \rangle \\ &= h \{ \langle S_h a_j, (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle - \langle a_j, (\mathbf{e} \cdot S_h \nabla u) \partial_j \Phi \rangle \} \\ &\quad + h^2 \langle J_1 S_h a_j, (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle + h^2 \langle a_j, (\mathbf{e} \cdot T_h \nabla u) (\nabla_x \Phi H)_j \rangle. \end{aligned}$$

Again we write  $I_7 + I_8 = \sum_{j=9}^{11} I_j$  in (3.8). We rewrite  $I_9$  in the same way as in (3.3), i.e.,

$$(3.9) \quad \begin{aligned} I_9 &= h \langle a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u), (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle \\ &\quad + h \langle a_j(x, S_h \nabla u) - a_j(x, \nabla u), (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle \\ &\quad + h \langle a_j(x, \nabla u), \mathbf{e} \cdot (\nabla u - S_h \nabla u) \partial_j \Phi \rangle. \end{aligned}$$

By applying the similar technique as in (3.4) to the first term of (3.9), we get the inequality

$$(3.10) \quad h \langle a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u), (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle \leq Ch^2 (\kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^2).$$

<sup>5)</sup> We write the  $j$ -th component of a vector  $U$  by  $U_j$ .

The second term of (3.9) is estimated as follows:

$$\begin{aligned}
 (3.11) \quad & h|\langle a_j(x, S_h \nabla u) - a_j(x, \nabla u), (\mathbf{e} \cdot \nabla u) \partial_j \Phi \rangle| \\
 & \leq Ch^2 \int_{\Omega} (\kappa_0 + |\nabla u|^{p-2} + |S_h \nabla u|^{p-2}) |P_h \nabla u| \cdot |\nabla u| dx \\
 & \leq Ch^2 \{ \kappa_0 \|\nabla u\|_2 \|P_h \nabla u\|_2 + \|\nabla u\|_p^{p/2} (\|\nabla u\|^{(p-2)/2} \|P_h \nabla u\|_2 \\
 & \quad + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2) \}.
 \end{aligned}$$

In the above estimates we have used the assumption A and the equality

$$a_j(x, S_h \nabla u) - a_j(x, \nabla u) = \int_0^1 \sum_{i=1}^n \frac{\partial a_j}{\partial \eta_i} (x, \theta S_h \nabla u + (1 - \theta) \nabla u) (S_h \partial_i u - \partial_i u) d\theta.$$

According to Lemma 1.1 the last term of (3.9) is estimated as follows:

$$\begin{aligned}
 (3.12) \quad & h|\langle a_j, \mathbf{e} \cdot (\nabla u - S_h \nabla u) \partial_j \Phi \rangle| \leq Ch^2 \int_{\Omega} (\kappa_0 + |\nabla u|^{p-2}) |\nabla u| \cdot |P_h \nabla u| dx \\
 & \leq Ch^2 (\kappa_0 \|\nabla u\|_2 \|P_h \nabla u\|_2 + \|\nabla u\|_p^{p/2} \| | \nabla u |^{(p-2)/2} P_h \nabla u \|_2).
 \end{aligned}$$

By virtue of (3.9)–(3.12) we get

$$\begin{aligned}
 (3.13) \quad & |I_9| \leq Ch^2 \{ \kappa_0 \|\nabla u\|_2 \|P_h \nabla u\|_2 + \kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^p \\
 & \quad + \|\nabla u\|_p^{p/2} (\|\nabla u\|^{(p-2)/2} \|P_h \nabla u\|_2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2) \}.
 \end{aligned}$$

On the other hand by Lemma 1.1,  $I_{10}$  and  $I_{11}$  are immediately estimated, that is,

$$(3.14) \quad |I_{10}|, |I_{11}| \leq Ch^2 (\kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^2).$$

Consequently (3.8), (3.13) and (3.14) yield that

$$\begin{aligned}
 (3.15) \quad & |I_7 + I_8| \leq Ch^2 \{ \kappa_0 (\|\nabla u\|_2 \|P_h \nabla u\|_2 + \|\nabla u\|_2^2) + \|\nabla u\|_p^p \\
 & \quad + \|\nabla u\|_p^{p/2} (\|\nabla u\|^{(p-2)/2} \|P_h \nabla u\|_2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2) \}.
 \end{aligned}$$

With the aid of (3.2), (3.6), (3.7) and (3.15) we obtain the inequality

$$\begin{aligned}
 (3.16) \quad & I_1 + I_2 \geq \frac{\gamma_0 h^2}{4} \{ \kappa_0 \|P_h \nabla u\|_2^2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \\
 & \quad + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \} - Ch^2 (\kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^2).
 \end{aligned}$$

Now we estimate the sum of  $I_3$  and  $I_4$ . It is written in the form

$$\begin{aligned}
 (3.17) \quad & I_3 + I_4 = (a_0, u - S_h u) + ((1 + hJ_1) S_h a_0, S_h u - u) \\
 & = (a_0(S_h x, S_h u) - a_0(x, S_h u), S_h u - u) \\
 & \quad + (a_0(x, S_h u) - a_0(x, u), S_h u - u) + h(J_1 S_h a_0, S_h u - u).
 \end{aligned}$$

Similarly as in (3.4) we can estimate the first term of (3.17) as follows:

$$\begin{aligned}
 (3.18) \quad & |(a_0(S_h x, S_h u) - a_0(x, S_h u), S_h u - u)| \\
 & \leq Ch^2 \int_{\rho} (1 + |S_h u|^{\alpha+1}) |P_h u| dx \\
 & \leq Ch^2 \{ \|P_h u\|_p + \|P_h u\|_p (\|u\|_{(\alpha+1)p^*})^{\alpha+1} \} \\
 & \leq Ch^2 (\|\nabla u\|_p + \|\nabla u\|_p^{\alpha+2}).
 \end{aligned}$$

Here we have used Lemma 2.1 and (1.14).

Because of Lemma 1.1 the second term of (3.17) is non-negative, i.e.,

$$(3.19) \quad (a_0(x, S_h u) - a_0(x, u), S_h u - u) \geq 0.$$

For the last term in (3.17) we attain the following estimate from Lemma 1.1 and Hölder’s inequality:

$$\begin{aligned}
 (3.20) \quad & h |(J_1 S_h a_0, S_h u - u)| \leq Ch^2 \int_{\rho} (1 + |S_h u|^{\alpha}) |S_h u| \cdot |P_h u| dx \\
 & \leq Ch^2 \{ \|P_h u\|_p \|S_h u\|_{p^*} + \|P_h u\|_p (\|S_h u\|_{(\alpha+1)p^*})^{\alpha+1} \}.
 \end{aligned}$$

Similarly as in (3.18) we obtain the following inequality from (3.20):

$$(3.21) \quad h |(J_1 S_h a_0, S_h u - u)| \leq Ch^2 (\|\nabla u\|_p^2 + \|\nabla u\|_p^{\alpha+2}).$$

In this way we arrive at the inequality

$$(3.22) \quad I_3 + I_4 \geq -Ch^2 (\|\nabla u\|_p^{\alpha+2} + \|\nabla u\|_p + \|\nabla u\|_p^2)$$

from (3.17), (3.18), (3.19) and (3.21).

Finally we estimate the right-hand side of (3.1). By simple calculations it holds that

$$(3.23) \quad (f, u - T_h u) + (f, u - S_h u) = h^2 \{ (P_h f, P_h u) + (J_1 S_h f, P_h u) \}.$$

Let us suppose that  $\kappa_0 = 0$ . From (3.23) and (ii) in Lemma 2.1 we have

$$(3.24) \quad |\text{the right-hand side of (3.1)}| \leq Ch^2 \|\nabla u\|_p \|f\|_{1,p^*}.$$

Combining (3.1), (3.16), (3.22) with (3.24), we deduce that if  $\kappa_0 = 0$ ,

$$\begin{aligned}
 & \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \\
 & \leq C (\|\nabla u\|_p^2 + \|\nabla u\|_p^{\alpha+2} + \|\nabla u\|_p + \|\nabla u\|_p^2 + \|f\|_{1,p^*}).
 \end{aligned}$$

Therefore if  $\kappa_0 = 0$ , our proposition is correct in virtue of Lemma 1.3.

Secondly we suppose that  $\kappa_0 \neq 0$ . In this case we transform the first term in the brackets on the right-hand side of (3.23) into the following

form by the expression (2.8) in the proof of Lemma 2.2:

$$(P_h f, P_h u) = - (S_h f, P_h P_h u) + (J_2 f, P_h u).$$

Taking account of (3.23) and (ii) in Lemma 2.1, we obtain the inequality

$$(3.25) \quad |\text{the right-hand side of (3.1)}| \leq Ch^2 \|f\|_2 (\|P_h \nabla u\|_2 + \|\nabla u\|_2).$$

Hence from (3.1), (3.16), (3.22) and (3.25) it follows that

$$\begin{aligned} & \gamma_0 \{ \kappa_0 \|P_h \nabla u\|_2^2 + \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 \} \\ & \leq C(\kappa_0 \|\nabla u\|_2^2 + \|\nabla u\|_p^p + \|\nabla u\|_p^{\alpha+2} + \|\nabla u\|_p + \|\nabla u\|_p^2 + \|f\|_2^2). \end{aligned}$$

Thus the proof of our proposition is accomplished by the straight-forward application of Lemma 1.3. Q.E.D.

4.

In this section we prove our theorem.

At first we show that  $\Phi(x)a_j(x, \nabla u)$  belongs to  $W^{1,p^*}(\Omega)$  and we estimate its  $W^{1,p^*}(\Omega)$  norm. Considering Lemma 1.4 and Lemma 2.2, we first give the uniform  $L^{p^*}(\Omega)$  estimate for the sequence  $\{P_h(a_j(x, \nabla u)) - J_2 a_j(x, \nabla u)\}_{h>0}$ , where  $h$  is sufficiently small. We write  $P_h(a_j(x, \nabla u))$  in the form

$$(4.1) \quad \begin{aligned} P_h(a_j(x, \nabla u)) &= [a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u)]h^{-1} \\ &\quad + [a_j(x, S_h \nabla u) - a_j(x, \nabla u)]h^{-1}. \end{aligned}$$

Similarly as in (3.4) we see that

$$(4.2) \quad \|[a_j(S_h x, S_h \nabla u) - a_j(x, S_h \nabla u)]h^{-1}\|_{p^*} \leq C(\kappa_0^{p^*} \|\nabla u\|_{p^*}^{p^*} + \|\nabla u\|_2^2).$$

For the second term of (4.1), we get the inequality

$$(4.3) \quad \begin{aligned} & \|[a_j(x, S_h \nabla u) - a_j(x, \nabla u)]h^{-1}\|_{p^*} \\ & \leq C \left\{ \kappa_0^{p^*} \|P_h \nabla u\|_{p^*}^{p^*} + \int_{\Omega} (|P_h \nabla u|^{p^*} |S_h \nabla u|^{p^*(p-2)} + |P_h \nabla u|^{p^*} |\nabla u|^{p^*(p-2)}) dx \right\} \end{aligned}$$

by the same way as in (3.11).

Because of  $1 < p^* \leq 2 \leq p$ , we have  $\|P_h \nabla u\|_{p^*} \leq C\|P_h \nabla u\|_2$ . And the integral terms are estimated in the following way:

$$\begin{aligned} & \int_{\Omega} |P_h \nabla u|^{p^*} |S_h \nabla u|^{p^*(p-2)} dx \\ & \leq \left[ \int_{\Omega} \left\{ |P_h \nabla u|^{p^*} |S_h \nabla u|^{p^*(p-2)/2} \right\}^{2/p^*} dx \right]^{p^*/2} \left[ \int_{\Omega} |S_h \nabla u|^{(p^*(p-2)/2)(1-p^*/2)-1} dx \right]^{(1-p^*/2)} \end{aligned}$$



$$\begin{aligned}
 &= \left( \int_{\Omega} |P_h \nabla u|^2 |S_h \nabla u|^{p-2} dx \right)^{p^*/2} \left( \int_{\Omega} |S_h \nabla u|^p dx \right)^{(1-p^*/2)} \text{ }^6 \\
 &\leq C \{ \| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \|\nabla u\|_p^2 \},
 \end{aligned}$$

where we use Hölder’s inequality and Minkowski’s inequality. Another integral term is similarly estimated.

Hence we conclude from (4.3) that

$$\begin{aligned}
 (4.4) \quad &\| [a_j(x, S_h \nabla u) - a_j(x, \nabla u)] h^{-1} \|_{p^*} \\
 &\leq C \{ \kappa_0^{p^*} \| P_h \nabla u \|_2^{p^*} + \|\nabla u\|_p^p + (\| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2) \}.
 \end{aligned}$$

Because of (4.1), (4.2) and (4.4) it holds that

$$\begin{aligned}
 (4.5) \quad &\| P_h(a_j(x, \nabla u)) \|_{p^*} \leq C \{ \kappa_0^{p^*} (\|\nabla u\|_p^{p^*} + \| P_h \nabla u \|_2^{p^*}) + \|\nabla u\|_p^p \\
 &\quad + (\| |S_h \nabla u|^{(p-2)/2} P_h \nabla u \|_2^2 + \| |\nabla u|^{(p-2)/2} P_h \nabla u \|_2^2) \}.
 \end{aligned}$$

Therefore if  $\kappa_0 = 0$ ,

$$\| P_h(a_j(x, \nabla u)) \|_{p^*} \leq C(1 + \|f\|_{1,p^*}^{p^*} + \|f\|_{p^*}^{(\alpha+2)/(p-1)}),$$

from (4.5), Proposition and Lemma 1.3. We write the right-hand side by  $C\gamma_f^0$ . Moreover if  $\kappa_0 \neq 0$ , we have similarly

$$\| P_h(a_j(x, \nabla u)) \|_{p^*} \leq C(1 + \|f\|_2^{2\beta/p}),$$

where  $\beta = \max(p, \alpha + 2)$ . The right-hand side is written by  $C\gamma_f$ .

On the other hand the functions  $J_2$  are uniformly bounded in  $x$  and  $h$ . Thus by Lemma 1.1

$$(4.6) \quad \| J_2 a_j(x, \nabla u) \|_{p^*} \leq C(\kappa_0^{p^*} \|\nabla u\|_{p^*}^{p^*} + \|\nabla u\|_p^p).$$

Applying the estimate for  $\nabla u$  in Lemma 1.3 to each term on the right-hand side of (4.6), the  $L^{p^*}(\Omega)$  norms of the functions  $J_2 a_j(x, \nabla u)$  are estimated as follows: if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$\| J_2 a_j(x, \nabla u) \|_{p^*} \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

From the above the  $L^{p^*}(\Omega)$  norms of the sequence  $\{P_h(a_j(x, \nabla u)) - J_2 a_j(x, \nabla u)\}_{h>0}$  are estimated. That is, if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$(4.7) \quad \| P_h(a_j(x, \nabla u)) - J_2 a_j(x, \nabla u) \|_{p^*} \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

Because  $a_j(x, \nabla u) \in L^{p^*}(\Omega)$ , we conclude on account of Lemma 2.2 that for any  $\phi \in C_0^\infty(\Omega)$

<sup>6)</sup> Indeed,  $(p^*(p-2)/2)(1-p^*/2)^{-1} = p$ .

$$(P_h(a_j(x, \nabla u)) - J_2 a_j(x, \nabla u), \phi) \longrightarrow ((e \cdot \nabla)(\Phi a_j(x, \nabla u)), \phi),$$

as  $h \longrightarrow 0$ .

Hence Lemma 1.4 yields that the distribution  $(e \cdot \nabla)(\Phi a_j(x, \nabla u))$  belongs to  $L^{p^*}(\Omega)$  and that

$$(4.8) \quad P_h(a_j(x, \nabla u)) - J_2 a_j(x, \nabla u) \longrightarrow (e \cdot \nabla)(\Phi a_j(x, \nabla u)) \quad \text{in } L^{p^*}(\Omega),$$

as  $h \longrightarrow 0$ ,

where “ $\longrightarrow$ ” means the weak convergence.

Therefore from (4.7) and (4.8) we derive that if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$(4.9) \quad \|(e \cdot \nabla)(\Phi a_j(x, \nabla u))\|_{p^*}^0 \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

The assertion with respect to  $\Phi(x)|\nabla u|^{p/2}$  can be treated more easily. By simple calculations we deduce the inequality

$$|P_h(|\nabla u|^{p/2})| \leq C|P_h \nabla u|(|S_h \nabla u|^{(p-2)/2} + |\nabla u|^{(p-2)/2}).$$

From this inequality and the estimate in Proposition we have for the case of  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ )

$$\|P_h(|\nabla u|^{p/2})\|_2^2 \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

Since the functions  $J_2$  are uniformly bounded, we get from Lemma 1.3

$$\|J_2|\nabla u|^{p/2}\|_2^2 \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f),$$

if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ).

Thus the  $L^2(\Omega)$  norms of the sequence  $\{P_h(|\nabla u|^{p/2}) - J_2|\nabla u|^{p/2}\}_{h>0}$  are estimated as follows: if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$(4.10) \quad \|P_h(|\nabla u|^{p/2}) - J_2|\nabla u|^{p/2}\|_2^2 \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

With the aid of a priori estimate (4.10) we can select a sequence  $\{h_\nu\}_{\nu=1}^\infty$  with  $h_\nu \rightarrow 0$  ( $\nu \rightarrow \infty$ ) and choose a function  $v \in L^2(\Omega)$  such that

$$(4.11) \quad P_{h_\nu}(|\nabla u|^{p/2}) - J_2|\nabla u|^{p/2} \longrightarrow v \quad \text{in } L^2(\Omega) \quad \text{as } \nu \longrightarrow \infty,$$

and besides if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$(4.12) \quad \|v\|_2^2 \leq C\gamma_f^0 \quad (\text{resp. } C\gamma_f).$$

By Lemma 2.2 we see that for any function  $\phi$  in  $C_0^\infty(\Omega)$

$$(4.13) \quad (P_h(|\nabla u|^{p/2}) - J_2|\nabla u|^{p/2}, \phi) \longrightarrow ((e \cdot \nabla)(\Phi|\nabla u|^{p/2}), \phi)$$

as  $h \longrightarrow 0$ .

From (4.11), (4.12) and (4.13) it holds that  $v = (\mathbf{e} \cdot \mathcal{V})(\Phi|\nabla u|^{p/2})$  and if  $\kappa_0 = 0$  (resp.  $\kappa_0 \neq 0$ ),

$$(4.14) \quad \|(\mathbf{e} \cdot \mathcal{V})(\Phi|\nabla u|^{p/2})\|_2^2 \leq C\gamma_0^0 \quad (\text{resp. } C\gamma_j).$$

By virtue of (4.9) and (4.14) we have proved the required for  $\Phi(x)a_j(x, \nabla u)$  and  $\Phi(x)|\nabla u|^{p/2}$ . And the proof of the part for  $\Phi(x)\nabla u$  is left. We briefly explain it. On account of the estimate (ii) in Proposition,  $\|P_h u\|_{1,2}$  are uniformly bounded in  $h$ , more precisely,

$$\|P_h u\|_{1,2} \leq C\gamma_j^{1/2},$$

if  $\kappa_0 \neq 0$ .

Accordingly there are a sequence  $\{h_\mu\}_{\mu=1}^\infty$  with  $h_\mu \rightarrow 0 (\mu \rightarrow \infty)$  and an element  $w$  in  $W^{1,2}(\Omega)$ , to which the sequence  $P_{h_\mu} u$  converges weakly in  $W^{1,2}(\Omega)$  as  $\mu \rightarrow \infty$ . On the other hand the sequence  $P_{h_\mu} u$  converges strongly to  $\Phi(\mathbf{e} \cdot \mathcal{V})u$  in  $L^2(\Omega)$  as  $\mu \rightarrow \infty$ , which is shown by the same way as in Lemma 5 in [20]. Consequently we have that  $w = \Phi(\mathbf{e} \cdot \mathcal{V})u$  and

$$\|\Phi(\mathbf{e} \cdot \mathcal{V})u\|_{1,2} \leq C\gamma_j^{1/2}.$$

Thus the proof of our theorem is finished.

Q.E.D.

*Remark.* The same conclusion is obtained if we replace  $a_j(x, \nabla u)$  by  $|\nabla u|^{p-2} \partial_{x_j} u$  or  $|\partial_{x_j} u|^{p-2} \partial_{x_j} u$  in our theorem.

#### BIBLIOGRAPHY

- [ 1 ] A. A. Arkhipova, A problem with discontinuous obstruction for uniformly elliptic equation, *Vestn. Lenigr. Univ.*, **19** (1974), 154–155.
- [ 2 ] ———, On limiting smoothness of the solution of a problem with a two-side barrier, *Vestn. Lenigr. Univ. Mat.*, **17** (1984), 5–9.
- [ 3 ] H. Brézis-D. Kinderlehrer, The smoothness of solutions to nonlinear variational inequalities, *Indiana Univ. Math. J.*, **17** (1974), 831–844.
- [ 4 ] ——— -G. Stampacchia, Sur la régularité de la solution d'inéquations elliptiques, *Bull. Soc. Math. France*, **96** (1968), 153–180.
- [ 5 ] K. C. Chang, The obstacle problem and partial differential equations with discontinuous nonlinearities, *Comm. Pure Appl. Math.*, **38** (1980), 117–146.
- [ 6 ] F. Cooper, Existence and regularity of solutions with bounded gradient of variational inequalities of a certain class, *Quart. J. Math. Oxford*, **26** (1975), 203–214.
- [ 7 ] J. P. Dias, Une classe de problèmes variationnels non linéaires de type elliptique ou parabolique, *Ann. Mat. Pura Appl.*, **92** (1972), 263–322.
- [ 8 ] E. DiBenedetto,  $C^{1+\alpha}$  local regularity of weak solutions of degenerate elliptic equations, *Nonlinear Anal. T.M.A.*, **7** (1983), 827–850.
- [ 9 ] A. Domarkas, Regularity of solutions of quasilinear elliptic equations with unilateral boundary conditions, *Lithu. Math. J.*, **20** (1980), 8–13.

- [10] —, Unilateral problems for quasilinear elliptic equation, *ibid.*, **21** (1981), 317–327.
- [11] L. C. Evans, A new proof of local  $C^{1,\alpha}$  regularity for solutions of certain degenerate elliptic P.D.E., *J. Differential Equations*, **45** (1982), 356–373.
- [12] J. Frehse, Two dimensional variational problems with thin obstacles, *Math. Z.*, **143** (1975), 279–288.
- [13] —, On Signorini's problem and variational problems with thin obstacles, *Ann. Scuola Norm. Sup. Pisa*, **4** (1977), 343–362.
- [14] —, On the smoothness of solutions of variational inequalities with obstacles, *Banach Center Publ.*, **10** (1983), 87–128.
- [15] — -U. Mosco, Irregular obstacles and quasi-variational inequalities of stochastic impulse control, *Ann. Scuola Norm. Sup. Pisa*, **9** (1982), 105–157.
- [16] M. G. Garroni, Regularity of a nonlinear variational inequality with obstacle on the boundary, *B.U.M.I. Anal. Fun. Appl. Suppl.*, **1** (1980), 267–286.
- [17] C. Gerhardt, Regularity of solutions of nonlinear variational inequalities, *Arch. Rational Mech. Anal.*, **52** (1973), 389–393.
- [18] —, Global  $C^{1,1}$ -regularity for solutions of quasilinear variational inequalities, *ibid.*, **89** (1985), 83–92.
- [19] M. Giaquinta-G. Modica, Regolarità Lipschitziana per la soluzione di alcuni problemi di minimo con vincola, *Ann. Mat. Pura Appl.*, **106** (1975), 95–117.
- [20] K. Hayasida-H. Nagase, On solutions of variational inequalities constrained on a subset of positive capacity, *Nagoya Math. J.*, **97** (1985), 51–69.
- [21] G. Huisken,  $C^{1,1}$ -regularity of solutions to variational inequalities, *Proc. Centre Math. Anal. Austral. Nat. Univ.*, **5** (1984), 85–90.
- [22] D. Kinderlehrer, The regularity of the solution to a certain variational inequality, *Proc. Symp. Pure Appl. Math.*, **23**, A.M.S. Providence.
- [23] —, Variational inequalities with lower dimensional obstacles, *Israel J. Math.*, **16** (1971), 339–348.
- [24] —, The smoothness of the solution of the boundary obstacle problem, *J. Math. Pures Appl.*, **60** (1980), 193–212.
- [25] K. L. Kuttler Jr., Degenerate variational inequalities of evolution, *Nonlinear Anal. T.M.A.*, **8** (1984), 837–850.
- [26] J. L. Lewis, Regularity of the derivatives of solutions to certain degenerate elliptic equations, *Indiana Univ. Math. J.*, **32** (1983), 849–858.
- [27] H. Lewy-G. Stampacchia, On existence and smoothness of solutions of some non-coercive variational inequalities, *Arch. Rational Mech. Anal.*, **41** (1971), 241–252.
- [28] P. Lindqvist, On the growth of the solutions of the differential equation  $\operatorname{div}(|\nabla u|^{p-2}\nabla u)=0$  in  $n$ -dim. space, *J. Differential Equations*, **58** (1985), 307–317.
- [29] J. L. Lions, *Quelques méthodes de résolution des problèmes aux limites nonlinéaires*, Dunod Gautyier-Villars, 1969.
- [30] T. H. Michel, Regularity for solutions to obstacle problems, *Proc. Centre, Math. Anal. Austral. Nat. Univ.*, **8** (1984), 29–37.
- [31] R. Schaaf, Regularity of solutions to linear and quasilinear variational inequalities with two obstacles, *Analysis*, **2** (1982), 337–345.
- [32] G. Stampacchia, Regularity of solutions of some variational inequalities *Proc. Symp. Pure Math.*, **18**, A.M.S. Providence.
- [33] P. Tolksdorf, Regularity for a more general class of quasilinear elliptic equation, *J. Differential Equations*, **51** (1984), 126–150.
- [34] N. N. Ural'tseva, Regularity of the solutions of variational inequalities, *J. Soviet Math.*, **3** (1975), 565–573.
- [35] H. Beirão da Veiga, Proprietà di sommabilità e di limitatezza per soluzioni di

- disequazioni variazionali ellittiche, *Rend. Sem. Nat. Padova*, **46** (1971), 141–171.
- [36] —, Sur la régularité des solutions de l'équation  $\operatorname{div} A(x, u, \nabla u) = B(x, u, \nabla u)$  avec des conditions aux limites unilatérales et mêlées, *Ann. Mat. Pura Appl.*, **93** (1972), 173–230.
- [37] — -F. Conti, Equazioni ellittiche non lineari con ostacoli sottili applicazioni allo studio dei punti regolari, *Ann. Scuola Norm. Sup. Pisa*, **26** (1972), 533–562.
- [38] G. H. Williams, Lipschitz continuous solutions for nonlinear obstacle problems, *Math. Z.*, **154** (1977), 51–65.
- [39] —, Smooth solutions for variational inequalities with obstacles, *Indiana Univ. Math. J.*, **31** (1982), 317–325.
- [40] G. H. Yakovlev, A variational problem, *Differential Equations*, **5** (1969), 960–966.
- [41] —, The first boundary value problem for quasilinear elliptic equations of second order, *Proc. Steklov Inst. Math.*, **117** (1972), 381–403.
- [42] —, Properties of solutions of a class of second-order quasilinear elliptic equations in divergence form, *ibid.*, **131** (1974), 242–252.
- [43] —, Some properties of solutions of quasilinear elliptic equations, *ibid.*, **134** (1975), 441–458.
- [44] K. Yosida, *Functional analysis*, Springer-Verlag, 1968.

*Suzuka College of Technology*  
*Suzuka 510-02*  
*Japan*