

# Molecular characterization of *Babesia* and *Cytauxzoon* species in wild South-African meerkats

SARAH LECLAIRE<sup>1,2,3,4,\*</sup>, SANDIE MENARD<sup>6</sup> and ANTOINE BERRY<sup>5,6</sup>

<sup>1</sup> Department of zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

<sup>2</sup> Laboratoire Evolution & Diversité Biologique, UMR 5174 (CNRS, Université Paul Sabatier, ENFA), 118 route de Narbonne, 31062 Toulouse, France

<sup>3</sup> Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS, 1919 route de Mende, 34293 Montpellier, France

<sup>4</sup> Kalahari Meerkat Project, Kuruman River Reserve, 8467 Van Zylsrus, Northern Cape, South Africa

<sup>5</sup> Service de Parasitologie-Mycologie, Centre Hospitalier Universitaire Purpan, 31059 Toulouse, France

<sup>6</sup> Centre de Physiopathologie de Toulouse-Purpan, INSERM UMR1043–CNRS UMR5282 – Université de Toulouse III, BP 3028, 31024 Toulouse cedex 3, France

(Received 20 June 2014; revised 25 August 2014; accepted 28 August 2014; first published online 6 November 2014)

## SUMMARY

Piroplasms, including *Babesia*, *Cytauxzoon* and *Theileria* species, frequently infect domestic and wild mammals. At present, there is no information on the occurrence and molecular identity of these tick-borne blood parasites in the meerkat, one of South Africa's most endearing wildlife celebrities. Meerkats live in territorial groups, which may occur on ranchland in close proximity to humans, pets and livestock. Blood collected from 46 healthy meerkats living in the South-African Kalahari desert was screened by microscopy and molecular methods, using PCR and DNA sequencing of 18S rRNA and ITS1 genes. We found that meerkats were infected by 2 species: one species related to *Babesia* sp. and one species related to *Cytauxzoon* sp. Ninety one percent of the meerkats were infected by the *Cytauxzoon* and/or the *Babesia* species. Co-infection occurred in 46% of meerkats. The pathogenicity and vectors of these two piroplasm species remains to be determined.

Key words: piroplasms, *Babesia*, *Cytauxzoon*, meerkats, molecular phylogeny.

## INTRODUCTION

Piroplasms, including *Babesia*, *Theileria* and *Cytauxzoon* species, are one of the most common blood parasites of mammals. These tick-transmitted protozoans have been reported in many domestic animals (Ristic, 1988) where they can cause asymptomatic infection or severe disease, including fever, anaemia, depression, diarrhoea, lethargy, anorexia and abortion (Homer *et al.* 2000; Zintl *et al.* 2003). Piroplasms are thus of considerable economic, medical and veterinary importance. Because most zoonotic *Babesia* are maintained in wildlife reservoir, surveillance of wild mammals living in area where undescribed *Babesia* are emerging has been suggested (Yabsley and Shock, 2013).

In a wide range of wildlife, *Babesia* have been molecularly described (Schnittger *et al.* 2012; Yabsley and Shock, 2013). In mongooses (Herpestidae), although several piroplasms have been detected, they all have been described based entirely on morphology (Grewal, 1957; Bandyopadhyay and Ray, 1985; Penzhorn and Chaparro, 1994). The meerkat (*Suricata suricatta*), one of South Africa's most endearing wildlife celebrities, is a small burrow-living mongoose (van Staaden, 1994) that

can be infested by several ixodid tick species, including *Haemaphysalis zumpti* and *Rhipicephalus theileri* (Hoogstraal and El Kammah, 1974; Horak *et al.* 1999, 2000). Meerkats can live on ranchlands, in close proximity to humans, pets or livestock, which increases the potential for the movement of pathogens, such as *Babesia*, between species. The presence of piroplasms in meerkats has never been studied. Here we investigated the occurrence and molecular phylogeny of piroplasms infecting wild meerkats using 18S rRNA and internal transcribed spacer 1 (ITS1) rRNA assay.

## METHODS

### Animal sources

This study was conducted on a wild population of meerkats at the Kalahari Meerkat Project (Kuruman River Reserve, Northern Cape; 26°58'S, 21°49'E), on ranchland in the South African Kalahari. Thirty two meerkats from 10 social groups were sampled in March 2011, while 18 meerkats from 8 social groups were sampled in November 2011. All meerkats in the study population were individually recognizable (Jordan *et al.* 2007) and habituated to close observation. Meerkats were blood sampled in the morning (between 6:00 AM and 9:00 AM) while they were sunning themselves at a burrow entrance.

\* Corresponding author. CEFE-CNRS, 1919 Rte de Mende, 34293 Montpellier cedex 5, France. E-mail: sarah.leclaire@free.fr




Fig. 1. Photomicrograph of a Giemsa-stained thin blood film from a meerkat showing an intra-erythrocytic piroplasm (*Babesia* or *Cytauxzoon* sp.). Original magnification:  $\times 1000$ .

Meerkats were caught by gently but firmly lifting them by the tail base, and they were immediately placed in a dark-coloured cotton bag. The meerkat was supported by placing an arm under the bag while it was carried away from the rest of the group to a sampling area approximately 100 m away. Anaesthesia induction and maintenance was by isoflurane inhalation in air administered by face mask. Blood (2 mL) was withdrawn from the jugular vein using a 25-gauge needle and a 2-mL syringe and placed in a heparinized tube. Just after collection, a drop of blood was either transferred on FTA® Elute Cards (Whatman) with a glass capillary or smeared on glass slide (Fig. 1).

European captive-born meerkats have presumably never been in contact with the parasites, given their several generations in captivity and no documentation of contact with African ticks during this time. They were, therefore, used as negative controls. We used blood stored in EDTA-tubes from 3 captive-born meerkats housed at Mulhouse Zoological Garden (France), and 2 captive-born meerkats housed at Thoiry Zoological Garden (France).

#### DNA extraction and PCR

DNA from filter papers of wild meerkats was extracted using Whatman DNA extraction procedure. DNA from blood smears and from blood of captive meerkats was extracted using DNA extraction kits (DNeasy Blood and Tissue kit, QIAGEN, Southern Cross Biotechnologies, SA), following the manufacturer's protocol.

Primary outside amplification for 18S rRNA gene was conducted with 5·1 (Yabsley *et al.* 2005) and B primers (Fig. 2; Medlin *et al.* 1988). One  $\mu\text{L}$  of DNA was added to 9  $\mu\text{L}$  of a master mix containing 1X Green GoTaq reaction Buffer (Promega), 200  $\mu\text{M}$  of each dNTP (Promega), 0·4  $\mu\text{M}$  of each primer and 0·05 unit of GoTaq DNA Polymerase (Promega). Cycling conditions were 95 °C for 3 min followed by 40 cycles of 95 °C for 30 s, 48 °C for 30 s and 72 °C for 2 min, and a final extension at 72 °C for 5 min.

To obtain near full-length 18S rRNA gene sequence, overlapping sequences were amplified using 3 secondary reactions. We used 5·1 (Yabsley *et al.* 2005) and RLB-R (Gubbels *et al.* 1999) primers, or RLB-F (Gubbels *et al.* 1999) and BabRev (Blaschitz *et al.* 2008) primers, or BabFor (Blaschitz *et al.* 2008) and B (Medlin *et al.* 1988) primers (Fig. 2). To obtain sequence of the hypervariable region of the 18S rRNA gene, secondary reaction was realized using RLB-F and BabRev primers. All secondary reactions were realized using 1  $\mu\text{L}$  of primary product as a template in a 20  $\mu\text{L}$  reaction. PCR master mix composition and cycling conditions were the same as primary reaction except the annealing temperature was 44 °C, and cyclic extensions were for 1 min.

Primary outside amplification for ITS1 gene was conducted using ITS-15C and ITS-13B primers (Bostrom *et al.* 2008). PCR master mix composition was the same as for primary outside amplification for 18S rRNA gene. Cycling conditions were 95 °C for 3 min followed by 40 cycles of 95 °C for 30 s, 55 °C for 30 s and 72 °C for 1 min and a final extension at 72 °C for 10 min. Secondary amplification was performed using ITS-15D and ITS-13C primers (Bostrom *et al.* 2008) with the same condition as primary reaction except the annealing temperature was 49 °C.

All amplifications included a negative control consisting of sterile molecular-grade water in the primary reaction and the negative-control primary PCR product for the secondary reaction. The PCR products were purified and sequenced by Eurofins MWG Operon (Ebersberg, Germany) or MilleGen (Labège, France) sequencing services.

#### Phylogenetic analyses

Phylogenetic analyses were conducted on the near full-length 18S rRNA sequences obtained during the present study and additional sequences retrieved from GenBank. Sequences were aligned using Bioedit Sequence Alignment Editor. Before generating the phylogenetic trees, a series of likelihood ratio tests were completed using MEGA 6 (Tamura *et al.* 2013) to determine the best nucleotide substitution model to use for phylogeny analyses. The best model predicted using the Bayesian Information Criterion (BIC) was a Tamura-Nei model (TN93) with a




Fig. 2. Schematic illustration of the directions and combinations of the different primary and secondary primers used to obtain the near full-length 18S rRNA gene sequences.

proportion of invariant sites (I) and among site heterogeneity (G). A maximum-likelihood phylogenetic tree based on the Tamura-Nei model with evolutionary rate differences among sites modelled using a discrete Gamma distribution (number of categories = 5), a neighbour joining tree and a maximum parsimony tree model were then inferred using MEGA 6 (Tamura *et al.* 2013).

## RESULTS

### Molecular characterization

For 1 meerkat, DNA was extracted both from filter paper and from blood smear. ITS1 sequences were identical suggesting that sequences collected from the 2 DNA extraction procedures can be reliably compared. Blood from 4 meerkats was collected both in March 2011 and November 2011. For 2 meerkats, species and strain detection based on ITS1 sequences gave similar results across the 2 seasons. For the 2 remaining meerkats, infection was different between the 2 seasons, with meerkats being infected by the 2 piroplasm species in 1 season, but infected by a single species in the other season. However, for each of these 2 meerkats, species-specific sequences were identical between the 2 seasons.

All the 5 captive-born meerkats were PCR negative, while only 9% ( $n = 4$ ) of wild meerkats were. The 4 PCR-negative wild meerkats belonged to 4 different social groups.

We amplified a hypervariable region of the 18S rRNA gene ( $\sim 780$  pb) in 13 wild meerkats, the near full-length 18S rRNA gene ( $\sim 1700$  pb) in 6 of these 13 wild meerkats, and the ITS1 region ( $\sim 540$  pb) in all wild meerkats ( $n = 51$  samples). The 18S rRNA and ITS1 sequences of wild meerkats separated into 2 groups, suggesting infection by 2 piroplasm species (Fig. 3). Based on the ITS1 sequences, co-infection with the 2 species was detected in 46% of individuals ( $n = 21$ ). Prevalence of co-infection may, however, be underestimated because primers designed to amplify all piroplasms may preferentially amplify the more common template in the exponential phase of amplification.

Four wild meerkats had a 100% similar amplicon of the 18S rRNA hypervariable region. The near full-length 18S rRNA gene was sequenced for 2 of these individuals and was shown to be 100% identical to one another (GenBank accession number: KM025200). By BLAST analysis (Altschul *et al.* 1990), they shared the highest homology with *Cytauxzoon felis* (identity score: 95%; query cover: 100%; Fig. 3) and *Cytauxzoon manul* (identity score: 95%, query cover: 96%; Fig. 3). The ITS1 region of this novel *Cytauxzoon* sp. was detected in 57% of individuals ( $n = 26$ ; Table 1). However, because many meerkats were co-infected, ITS1 sequences of *Cytauxzoon* sp. were reliably sequenced for 7 individuals only. Within the 485-pb ITS1 region, we found 1 single nucleotide polymorphism (SNP) (GenBank accession number: KM025207 and KM025208). Out of the 7 individuals, 5 individuals had a single ITS1 sequence, while the remaining 2 individuals had the 2 ITS1 sequences. All meerkat social groups were infected by the *Cytauxzoon* species, except 1 group for which only 1 meerkat was sampled (Table 1).

Nine wild meerkats had similar sequence of the 18S rRNA hypervariable region. The near full-length 18S rRNA gene was sequenced for 4 of these individuals and was shown to be 100% identical to each other (GenBank accession number: KM025199). They shared the highest homology with *Babesia lengau* (identity score: 99%; query cover: 93%; Fig. 3). Based on the ITS1 sequence, infection by this *Babesia* sp. was detected in 80% of samples ( $n = 40$ ), and all meerkat social groups were infected (Table 1). Eighty-three percent of meerkats had more than one *Babesia* ITS1 sequence. Based on the electropherograms containing *Babesia* sequences only, 13 SNPs, one 1-nucleotide insertion and one 2-nucleotide deletions were detected in the 480-pb ITS1 sequence of *Babesia* sp. The 2-nucleotide deletion sequence was detected in 51% of *Babesia*-infected meerkats, while the 1-nucleotide insertion sequence was detected in 50% of *Babesia*-infected meerkats. From the unambiguous electropherograms, we reliably characterized 6 distinct ITS1




Fig. 3. (a) Maximum-likelihood, neighbour-joining and (b) maximum parsimony bootstrap consensus trees showing the phylogenetic relationships of the new *Babesia* sp. and *Cytauxzoon* sp. with other piroplasmids based on full-length 18S rRNA sequences. The numbers represent the percentage of 1000 replicates (bootstrap support) for which the same branching patterns were obtained for (a) maximum likelihood and neighbour joining analyses, respectively and (b) maximum-parsimony analysis. *Toxoplasma gondii* was used as an out-group. Sequences obtained during the present study are shown in bold. The number after the species designation refers to the GenBank number (indicated in the first tree only for clarity).

sequences (GenBank accession number: KM025201 to KM025206).

### Microscopy

In all blood smears where piroplasmids were observed, the organisms were single intraerythrocytic inclusions, small in size (measuring ca. 1  $\mu$ m in diameter; Fig. 1) and few in number. No morphological differences were discernible between the *Babesia* and the *Cytauxzoon* species.

### DISCUSSION

Meerkats were found to be infected by 2 different piroplasm species: the first one closely related to the *Cytauxzoon* species infecting Felidae (Wagner, 1976; Peixoto *et al.* 2007; Shock *et al.* 2011), and the second one closely related to *Babesia lengau*, a species infecting the South-African cheetah (*Acinonyx jubatus*) (Bosman *et al.* 2010). Several piroplasm species have been previously described in mongooses (reviewed in Penzhorn, 2006), but all have been described only from morphology. In particular,

*Babesia cynicti* has been found in South-African yellow mongooses (*Cynictis penicillata*) (Penzhorn and Chaparro, 1994), a species that can occupy the same territory and burrow, and host the same tick species as meerkats (Lynch, 1980). Molecular characterization of *Babesia cynicti* is now required to determine whether it is the same species as the ones found in meerkats.

The prevalence of infection with *Babesia* and *Cytauxzoon* in this population of meerkats was high. Ninety-one percent of meerkats were infected, including 46% that were infected by the 2 species. Similar prevalence (88%) was detected for *Babesia cynicti* in yellow mongooses (Penzhorn and Chaparro, 1994). All sampled meerkats appeared, however, clinically normal and showed no signs of disease (personal observations). Although *Babesia* and *Cytauxzoon* can cause severe disease in domestic animals (Wagner, 1976; Homer *et al.* 2000), high prevalence and subclinical infection are common in wild animals (Schnittger *et al.* 2012) where clinical disease manifests only when the host is stressed (Penzhorn, 2006). Clinical babesiosis in the sable antelope (*Hippotragus niger*), Grevy's zebra

Table 1. Percentage of wild meerkats infected by the *Babesia* species, the *Cytauxzoon* species and being co-infected within each social group

	Group ID and sample size											
	A n = 9 (%)	B n = 7 (%)	C n = 5 (%)	D n = 5 (%)	E n = 4 (%)	F n = 3 (%)	G n = 3 (%)	H n = 3 (%)	I n = 2 (%)	J n = 2 (%)	K n = 2 (%)	L n = 1 (%)
<i>Babesia</i> sp.	78	86	80	100	50	67	100	100	100	50	50	100
<i>Cytauxzoon</i> sp.	78	57	40	40	75	67	33	33	100	50	50	0
Co-infection	67	43	20	40	50	33	33	33	100	50	50	0
Total (%)												

(*Equus grevyi*), black rhinoceros (*Diceros bicornis*) and Eastern grey kangaroo (*Macropus giganteus*) appears to be triggered by the stress of capture, restraint and translocation (Dennig, 1964; Nijhof *et al.* 2003; Nijhof *et al.* 2005; Dawood *et al.* 2013; reviewed in Penzhorn, 2006). The *Babesia* and *Cytauxzoon* species identified in our study may be benign parasites, and infection does not appear to result in clinical morbidity or mortality under normal conditions.

Meerkats are infected by several ixodid ticks (Lynch, 1980), which are potential vectors of *Babesia* and *Cytauxzoon*. Preliminary characterizations of ticks collected in our meerkat population suggest that individuals are infected by *R. theileri* (Prof Ivan Horak, personal communication), a burrow associated ixodid tick that is associated with the hot and arid climate of the Northern Cape Province (Horak *et al.* 1999). The meerkat is a highly social species living in family groups which share the same burrows (Lynch, 1980). These features could facilitate the transmission of *Babesia* and *Cytauxzoon* by ticks among members of the same group and would explain the high infection prevalence in meerkats. Examination of genetic variations within the ITS region of the rRNA genes in relation to meerkat group membership and spatial location should help elucidate the transmission route of piroplasms in meerkats. In our study, ITS1 sequence analysis was complicated due to the co-infection of meerkats with different species and/or multiple strains of a single species, or due to the presence of several rRNA gene copies in the genome. Future works conducted with species-specific ITS1 primers or including cloning prior sequencing are needed to associate specific parasite genotype with host life history characteristics.

In summary, 2 piroplasm species were found in a high percentage of Kalahari meerkats. None of meerkats had evidence of disease compatible with babesiosis; thus these piroplasms appear to have low pathogenicity for meerkats. However, infection with either of them in combination with stress or other factors such as immunosuppression could possibly lead to disease.

ACKNOWLEDGMENTS

We thank T. Clutton-Brock and M. Manser for providing access to the animals at the KMP, and N. Thavarajah and J. Samson for their help in the field. We are grateful to the Kotze family and Northern Cape Conservation for allowing us to conduct research in the Kalahari. We are grateful to Dr B. Quintard at Mulhouse Zoological Garden and Dr A. Leclerc at Thoiry Zoological Garden for providing us with blood from captive meerkats. We thank M. Yabsley and E. Huchard for helpful discussion.

FINANCIAL SUPPORT

S.L. was supported by a Fondation Fyssen post-doctoral grant and an ASAB research grant.

## REFERENCES

- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Bandyopadhyay, S. and Ray, R. (1985). *Babesia bengalensis* sp (apicomplexa, babesiidae), a new piroplasmid from a common Indian mongoose, *Herpestes edwardsi* (Geoffroy). *Acta Protozoologica* **24**, 147–152.
- Blaschitz, M., Narodslavsky-Gfoeller, M., Kanzler, M., Stanek, G. and Walochnik, J. (2008). Babesia species occurring in Austrian Ixodes ricinus ticks. *Applied and Environmental Microbiology* **74**, 4841–4846.
- Bosman, A.-M., Oosthuizen, M. C., Peirce, M. A., Venter, E. H. and Penzhorn, B. L. (2010). *Babesia lengau* sp. nov., a novel Babesia species in cheetah (*Acinonyx jubatus*, Schreber, 1775) populations in South Africa. *Journal of Clinical Microbiology* **48**, 2703–2708.
- Bostrom, B., Wolf, C., Greene, C. and Peterson, D. (2008). Sequence conservation in the rRNA first internal transcribed spacer region of *Babesia gibsoni* genotype Asia isolates. *Veterinary Parasitology* **152**, 152–157.
- Dawood, K. E., Morgan, J. A. T., Busfield, F., Srivastava, M., Fletcher, T. I., Sambono, J., Jackson, L. A., Venus, B., Philbey, A. W. and Lew-Tabor, A. E. (2013). Observation of a novel *Babesia* spp. in Eastern Grey Kangaroos (*Macropus giganteus*) in Australia. *International Journal for Parasitology: Parasites and Wildlife* **2**, 54–61.
- Dennig, H. (1964). The isolation of Babesia species from wild animals. In *Proceedings of the First International Congress of Parasitology*, pp. 24–26. Rome.
- Grewal, M. S. (1957). Two new piroplasms, *Babesia* (Babesiella) *heischi* and *Babesia* (Babesia) *hoarei* from Peter's pigmy mongoose, *Helogale undulata rufula* (demonstrated by Professor Garnham). *Transactions of the Royal Society of Tropical Medicine and Hygiene* **51**, 290–291.
- Gubbels, J. M., De Vos, A. P., Van der Weide, M., Viseras, J., Schouls, L. M., De Vries, E. and Jongejan, F. (1999). Simultaneous detection of bovine *Theileria* and *Babesia* species by reverse line blot hybridization. *Journal of Clinical Microbiology* **37**, 1782–1789.
- Homer, M. J., Aguilar-Delfin, I., Telford, S. R., Krause, P. J. and Persing, D. H. (2000). Babesiosis. *Clinical Microbiology Reviews* **13**, 451–469.
- Hoogstraal, H. and El Kammah, K. M. (1974). Notes on African *Haemaphysalis* ticks. XII. *H. (Rhipistoma) zumpti* sp. n. a parasite of small carnivores and squirrels in southern Africa (ixodoidea: ixodidae). *The Journal of Parasitology* **60**, 188–197.
- Horak, I. G., Chaparro, F., Beaucournu, J. and Louw, J. (1999). Parasites of domestic and wild animals in South Africa. XXXVI. Arthropod parasites of yellow mongooses, *Cynictis penicillata* (G. Cuvier, 1829). Onderstepoort. *Journal of Veterinary Research* **66**, 33–38.
- Horak, I. G., Braack, L. E. O., Fourie, L. J. and Walker, J. B. (2000). Parasites of domestic and wild animals in South Africa. XXXVIII. Ixodid ticks collected from 23 wild carnivore species. *Onderstepoort Journal of Veterinary Research* **67**, 239–250.
- Jordan, N. R., Cherry, M. I. and Manser, M. B. (2007). Latrine distribution and patterns of use by wild meerkats: implications for territory and mate defence. *Animal Behaviour* **73**, 613–622.
- Lynch, C. D. (1980). Ecology of the suricate, *Suricata suricatta* and yellow mongoose, *Cynictis penicillata* with special reference to their reproduction. In *Memoirs van die Nasionale Museum*, pp. 145. Bloemfontein, South Africa.
- Medlin, L., Elwood, H. J., Stickel, S. and Sogin, M. L. (1988). The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding regions. *Gene* **71**, 491–499.
- Nijhof, A., Pillay, V., Steyl, J., Prozesky, L., Stoltz, W., Lawrence, J., Penzhorn, B. and Jongejan, F. (2005). Molecular characterization of *Theileria* species associated with mortality in four species of African antelopes. *Journal of Clinical Microbiology* **43**, 5907–5911.
- Nijhof, A. M., Penzhorn, B. L., Lynen, G., Mollé, J. O., Morkel, P., Bekker, C. P. and Jongejan, F. (2003). *Babesia bicornis* sp. nov. and *Theileria bicornis* sp. nov.: tick-borne parasites associated with mortality in the black rhinoceros (*Diceros bicornis*). *Journal of Clinical Microbiology* **41**, 2249–2254.
- Peixoto, P., Soares, C., Scofield, A., Santiago, C., Franca, T. and Barros, S. (2007). Fatal cytauxzoonosis in captive-reared lions in Brazil. *Veterinary Parasitology* **145**, 383–387.
- Penzhorn, B. L. (2006). Babesiosis of wild carnivores and ungulates. *Veterinary Parasitology* **138**, 11–21.
- Penzhorn, B. L. and Chaparro, F. (1994). Prevalence of *Babesia cynictis* infection in 3 populations of yellow mongooses (*Cynictis penicillata*) in the Transvaal, South-Africa. *Journal of Wildlife Diseases* **30**, 557–559.
- Ristic, M. (1988). *Babesiosis of Domestic Animals and Man*. CRC Press, Inc, Boca Raton, FL.
- Schnittger, L., Rodriguez, A. E., Florin-Christensen, M. and Morrison, D. A. (2012). *Babesia*: a world emerging. *Infection, Genetics and Evolution* **12**, 1788–1809.
- Shock, B. C., Murphy, S. M., Patton, L. L., Shock, P. M., Olfenbuttel, C., Beringer, J., Prange, S., Grove, D. M., Peek, M. and Butfiloski, J. W. (2011). Distribution and prevalence of *Cytauxzoon felis* in bobcats (*Lynx rufus*), the natural reservoir, and other wild felids in thirteen states. *Veterinary Parasitology* **175**, 325–330.
- Tamura, K., Stecher, G., Peterson, D., Filipowski, A. and Kumar, S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30**, 2725–2729.
- van Staaden, M. (1994). *Suricata suricatta*. *Mammalian Species* **483**, 1–8.
- Wagner, J. (1976). A fatal cytauxzoonosis-like disease in cats. *Journal of the American Veterinary Medical Association* **168**, 585–588.
- Yabsley, M. J. and Shock, B. C. (2013). Natural history of zoonotic *Babesia*: role of wildlife reservoirs. *International Journal for Parasitology: Parasites and Wildlife* **2**, 18–31.
- Yabsley, M. J., Davidson, W. R., Stallknecht, D. E., Varela, A. S., Swift, P. K., Devos, J. C. and Dubay, S. A. (2005). Evidence of tick-borne organisms in mule deer (*Odocoileus hemionus*) from the western United States. *Vector-Borne and Zoonotic Diseases* **5**, 351–362.
- Zintl, A., Mulcahy, G., Skerrett, H. E., Taylor, S. M. and Gray, J. S. (2003). *Babesia divergens*, a bovine blood parasite of veterinary and zoonotic importance. *Clinical Microbiology Reviews* **16**, 622–636.