

Polychaete diversity and assemblage structure in the Oualidia Lagoon, Moroccan Atlantic coast

FATIMA EL ASRI^{1,2}, HAKIMA ZIDANE², AHMED ERRHIF¹, MOHAMED-NAOUFAL TAMSOURI²,
MOHAMED MAANAN³, MOHAMED MALOULI IDRISSE² AND DANIEL MARTIN⁴

¹Health and Environment Laboratory, Faculty of Sciences Ain Chock, University Hassan II, B.P. 5366 Maârif, 20100 Casablanca, Morocco, ²Department of Fisheries Resources, National Institute for Fisheries Research (INRH), Route Sidi Abderrahmane Club Équestre Ould Jmel – Casablanca, Morocco, ³UMR 6554 LETG-Nantes, University of Nantes, BP 81227, 44312 Nantes, France, ⁴Centre for Advanced Studies of Blanes (CEAB-CSIC), Accés a la Cala St. Francesc, 14 Blanes 17300, Girona, Spain

The polychaete assemblages inhabiting the coastal waters of the Oualidia lagoon were studied during winter 2013 and summer 2013. Taxonomic composition and diversity were determined at 43 sample sites. Among the 13 species of polychaetes recorded, 10 were newly reported for Oualidia lagoon. Hediste diversicolor and Capitella sp. were the most abundant taxa in both seasons. Temperature and salinity were higher, and chl-a and OM were lower, in summer than in winter. The structure of the polychaete assemblages was characterized by forming three main clusters, either based on sampling stations or on polychaete species. These clusters were organized according to a downstream gradient, with the stations having fine sediments and a H. diversicolor assemblage in the inner lagoon being replaced by stations with medium grain-sized sediment and a Capitella sp. assemblage in the mid-lagoon, which were in turn replaced by stations having sandy sediments and assemblages dominated by Glycera alba (winter) and P. africana (summer) in the areas closer to the lagoon inlets. The shift was, in fact, from a classical, brackish, lacunar assemblage to two different, temporal aspects of a marine assemblage (close to the inlets), with a transition assemblage in between. This corresponded with a typically paralic spatial structure whose main descriptors responded to a confinement gradient. Despite the absence of a river, the organization of the polychaete assemblages closely resembled that of an estuarine system, with the tidal regime playing a major driving role.

Keywords: Polychaete diversity, seasonal variability, assemblage structure, Oualidia lagoon, Morocco, Atlantic coast

Submitted 3 October 2016; accepted 1 March 2017; first published online 17 April 2017

INTRODUCTION

Coastal lagoons are, in general, highly productive areas acting as transitional zones between land and sea. Nevertheless, anthropogenic pressure often tends to concentrate near coastal lagoons, thus becoming a major contributor to a decrease in water quality leading to habitat degradation (Newton *et al.*, 2003).

This is certainly the case of Oualidia, a lagoon located in the Atlantic coast of Morocco. Oualidia lagoon is a Natural Park providing a valuable refuge to a rich variety of birds, fish, plants and other wildlife, being the most important wintering area for migratory birds in Morocco. Therefore, it is one of the 24 Moroccan sites considered by the RAMSAR Convention as wetlands of international interest. Through the practices of boating, bathing, camping and traditional fisheries (including fish and molluscs), the lagoon and the surrounding area supports many aspects of the local economy, culture and community relationships and, consequently, a rapid urbanization has been generated during the recent years (Maanan *et al.*, 2014).

In turn, the different land uses around the lagoon (mainly agricultural, but also industrial) and its watershed produce various kinds of pollution, all of them affecting, to some extent, the lagoon environmental conditions. Among these are heavy metals (Zourarah *et al.*, 2007; Idardare *et al.*, 2008; Maanan *et al.*, 2014) and faecal pollution (Hennani *et al.*, 2012; Hassou *et al.*, 2014).

The macrofauna is often used in coastal monitoring studies and, particularly in soft-bottoms, the polychaetes are a key group (Giangrande *et al.*, 2005). The members of this group play an important role in the functioning of the benthic ecosystems, where they may represent up to 70% of the total abundance and biomass (Gray, 1974). The polychaetes are widely used as a key taxon in bioenvironmental studies to assess natural and human-induced disturbances, as they contain pollution sensitive and tolerant species that show differential distributions along pollution gradients (e.g. Pocklington & Wells, 1992; Gray, 2002; Venturini *et al.*, 2004).

Previous studies on the lagoon functioning, including the macrofaunal component, have already been carried out in some of the few Mediterranean Moroccan lagoons. These include the deep Nador lagoon, also known as Marchica (Guelorget *et al.*, 1987; El Kamcha *et al.*, 2015) and the shallower Smir lagoon (Chaouti & Bayed, 2008). In the Atlantic coast, the macrofauna has also been studied in a few

Corresponding author:
F. El Asri
Email: fatimaelasri25@gmail.com

lagoons, such as Merja Zerga (Bazaïri *et al.*, 2003), Khnifiss (Bayed *et al.*, 1988; Dakki & de Ligny, 1988; Lefrere *et al.*, 2015) and Oualidia. In this lagoon, however, the studies have mainly focused on macrobenthic taxa, such as molluscs, shellfish or echinoderms (Chbicheb, 1996; El Asri *et al.*, 2015), while the polychaete diversity was analysed nearby, but in open coastal areas, Jerf Asfar and El Jadida, located at 55 and 70 km far from Oualidia (Sif *et al.*, 2012).

Our study is the first one focusing specifically on the biodiversity and structure of the polychaete fauna in Oualidia lagoon, both under spatial and temporal approaches. Accordingly, we analyse the composition and structure of the assemblages, as well as their relationships with the main environmental variables explaining both their spatial distribution and their temporal variability in the enclosed ecosystem of the Oualidia coastal lagoon.

MATERIALS AND METHODS

Study area

Oualidia lagoon, 34°47'N 6°13'W and 34°52'N 6°14'W, is located on the Atlantic coast of Morocco (Figure 1). The lagoon measures 7 km long and about 1 km wide. The basin occupies a north-south depression bordered by a continental cliff and a coastal consolidated dune ridge and water exchanges with the ocean occur through a major inlet about 150 m wide. During spring tides there is also a secondary, shallower inlet about 50 m wide. The lagoon morphology is characterized by lateral channels, connected to a meandering main channel, with an average depth of 2 m and a maximum depth (during flood tides) that does not exceed 5 m (Carruesco, 1989). Flood tides cover more than 75%

(2.25 km²) of the lagoon surface, bringing salt water up to the most confined, inner lagoon region, as well as into a saline marsh beyond the inner dam (Figure 1).

Sampling and data analysis

Forty-three stations (Figure 1) were sampled in winter (March 2013) and summer (July 2013). All samples (two replicates) were collected using a Van Veen grab (0.0625 m² in surface area), except at station 43, which was located in a rocky zone and was thus sampled by scraping sixteen 25 × 25 cm quadrats (until a total of 1 m² of surface area). The samples were sieved *in situ* through a 1 mm pore size mesh. The material retained on the mesh was transferred to containers and fixed in a 10% formalin solution.

Each station was characterized by its distance from the lagoon opening (in km), water salinity (‰) and water temperature (°C). An additional sediment sample was collected to analyse grain size, and organic matter and chlorophyll-a (chl-a) contents. Grain size was measured with a laser granulometer (Malvern, Mastersizer) at the LETG (UMR 6554, University of Nantes) and expressed as mean grain size, in µm (to be used in the correlation analyses) and as relative proportions of sand, silt and clay, in percentages (to be used in the Canonical Correspondence Analysis, CCA). The percentage of organic matter (OM, %) was obtained as the weight losses of dried samples (24 h, 60°C) after ignition (4 h, 450°C). The chl-a content (mg m⁻²) was determined according to the Lorenzen method (Holm-Hansen *et al.*, 1965).

The macroinvertebrates were sorted under a binocular microscope, and the polychaetes were identified to species level, whenever possible, following Gil (2011). Currently accepted species names were checked at the World Polychaete Database (Read & Fauchald, 2017). Selected




Fig. 1. Map of the study area, showing the location of sampling sites in Oualidia lagoon.

specimens of the most relevant species have been deposited in the CEAB collections (reference numbers CEAB.AP.856A and CEAB.AP.856B to CEAB.AP.868).

To analyse the structure of the assemblages, the following indices were calculated: (1) species richness (S: number of species per sample); (2) species abundance (N: individuals m^{-2}); (3) diversity as indicated by the Shannon index (H' , \log_2 basis) (Shannon, 1948); and the evenness index (J') (Pielou, 1966).

The relationships between environmental variables, species density and descriptors of the assemblage structure with the distance from the lagoon opening were analysed by Pearson correlations, which were performed with the XLSTAT software (2015.5.01.23039, copyright by Addinsoft 1995–2016).

Hierarchical Ascending Classification analyses (HAC) were done both on sampling stations and polychaete species, based on the Euclidean distance and Ward's method using $\log_{10}(x+1)$ to limit the influence of the most dominant taxa (Vakharia & Wemmerlöv 1995; Cao *et al.*, 1997). Communities were identified by the IndVal index (Dufrene & Legendre, 1997). Wilcoxon tests were used to determine the significance ($P=0.05$) of the difference between seasons. All these analyses were carried out using the STATISTICA version 8 for Windows software.

Canonical Correspondence Analyses (CCA) were performed, using the PAST 3.0 free software package, to analyse the relationships between environmental variables

and polychaete assemblages. Environmental variables and polychaete densities were $\log_{10}(x+1)$ transformed prior to analysis. The significance of these relationships, as well as those of the assemblage descriptors with the environmental variables was assessed by Pearson correlation (performed with the XLSTAT software).

RESULTS

Environmental parameters

Temperature was significantly higher ($r=0.00006$; $P<0.05$) in summer (16.1°C and 26.3°C) than in winter (16.9 and 19.9°C) (Figure 2). Salinity was also significantly higher ($r=0.000215$, $P<0.05$) in summer (10.5‰ and 39.6‰) than in winter (10.1‰ and 39.5‰) (Figure 2). Chl-a showed marked changes among the studied stations. Contrary to temperature and salinity, the chl-a temporal pattern, as well as that of OM, showed significantly higher ($r=0.00008$, $P<0.05$ and $r=0.005282$, $P<0.05$) values in winter (1.19 and 23.41 $mg\ m^{-2}$ and 1.94–31.97%) than in summer (0.85 and 8.79 $mg\ m^{-2}$ and 1.73–15.1%) (Figure 2).

All environmental parameters analysed except salinity and chl-a showed significant correlations with the distance from




Fig. 2. Winter and summer differences in water temperature, water salinity, organic matter, chlorophyll-a content and granulometry (Silt, Clay and Sand contents) in the sediment of Oualidia lagoon. Mean \pm SD.

the lagoon opening both in March and July, positive for the temperature (Pearson coeff. = 0.758, $P < 0.001$ and Pearson coeff. = 0.518, $P < 0.001$) and OM (Pearson coeff. = 0.440, $P = 0.003$ and Pearson coeff. = 0.471, $P = 0.001$) and negative for the granulometry (Pearson coeff. = -0.338 ; $P = 0.027$ and Pearson coeff. = -0.448 , $P = 0.003$). Although non-significant, the salinity tended to decrease with the increasing distance from the opening both in March and July, while the chl-a tended to increase in March and to decrease in July.

Polychaete abundance and structure

A total of 3835 polychaete specimens belonging to 13 species were collected during this study (Table 1). Among them, 1051 individuals from 7 species occurred in winter and 2784 from 13 species in summer.

The three most abundant species in winter were *Hediste diversicolor* (72.4%), *Capitella* sp. (22.9%) and *Nephtys hombergii* (3.2%). The first two were also dominant in summer (64.7% and 17%, respectively), followed by *Panousea africana* (5.2%).

In winter, only *H. diversicolor* increased significantly its density significantly together with the increasing distance from the lagoon opening (Pearson coeff. = 0.337, $P = 0.027$). Conversely, in summer, there were no significant relationships with the distance from the lagoon opening, except for *P. africana*, which showed a negative correlation (Pearson coeff. = -0.340 , $P < 0.026$).

Polychaete densities were significantly higher ($r = 0.0007$; $P < 0.05$) in summer (0–392 individuals m^{-2}) than in winter (0–316 individuals m^{-2}). The number of species ranged from 0 to 4 (winter) or 8 (summer). However, there were non-significant differences ($r = 0.190$; $P > 0.05$) (Figure 3).

The Shannon diversity was similar in winter (0–1.09 bits) than in summer (0–1.57 bits) ($r = 0.091$; $P > 0.05$) and, similarly, there were non-significant differences in evenness between both seasons ($r = 0.241$, $P > 0.05$) (Figure 3).

The spatial distribution of the assemblage descriptors in winter revealed that only the polychaete densities were significantly, and positively, correlated with the distance for the

lagoon opening (Pearson coeff. = 0.381, $P = 0.012$). The remaining descriptors (S, J, H'), although non-significant, tended to increase with the increasing distance. Conversely, in summer, all descriptors tended to decrease with the distance from the lagoon opening. However, the relationships were non-significant for N and J, while they were significant in the cases of S and H' (Pearson coeff. = -0.397 ; $P = 0.008$ and Pearson coeff. = -0.437 , $P = 0.003$, respectively).

Three station clusters were obtained both during winter and summer in the HAC analyses, both for sampling stations and for polychaete species. Stations cluster 1 in winter (Figure 4A) included 15 stations mainly from the inner region of the lagoon, characterized by having sediments mainly composed by silt and clay in different proportions, high temperatures and OM and a low average polychaete density (43.4 individuals m^{-2}). The dominant species was *H. diversicolor*, followed by *Capitella* sp. (Table 2). Stations cluster 2 consisted of six stations occupying mostly the inner-central region of the lagoon, having silty sandy and clayey sandy sediments, a moderately high OM, temperature and salinity, and a high average density (56.8 individuals m^{-2}). The dominant species was *Capitella* sp., followed by *H. diversicolor* (Table 2). Stations cluster 3 consisted of 13 stations mainly close to the lagoon opening, with predominantly sandy sediments, low temperatures and OM, and a relatively low average density (4.5 individuals m^{-2}). The dominant species were *N. hombergii* and *Phyllodoce* sp. (Table 2).

During summer (Figure 4B), stations cluster 1 was the largest one. It included 17 stations, mainly located in the inner central region of the lagoon, characterized by having relatively high temperatures, high OM, clay and silty clay sediments and a high average density (91.8 individuals m^{-2}). The dominant species was *H. diversicolor* (Table 2). Stations cluster 2 included 10 stations, located all along the lagoon, but mostly in the central part, with moderately high % OM and temperature, silty sand and clayey sand sediments and a high average density (102.4 individuals m^{-2}). As it occurred during winter, the dominant polychaetes were *Capitella* sp., followed by *H. diversicolor*. Stations cluster 3 included seven stations located mostly near the lagoon opening, characterized by having low temperatures and OM, very low chl-a, a high

Table 1. Comparative analysis of the polychaetes identified in our study with those identified by Chbicheb (1996) in Oualidia lagoon.

Family	Species	This study		1996 study
		Winter	Summer	
Nereididae	<i>Hediste diversicolor</i> (Müller, 1776)	*	*	*
Capitellidae	<i>Capitella</i> sp.	*	*	
	<i>Capitella capitata</i> (Fabricius, 1780)	*		*
Ampharetidae	<i>Alkmaria romijni</i> Horst, 1919	*		
Sabellidae	<i>Panousea africana</i> Rullier & Amoureux, 1970		*	
Nephtyidae	<i>Nephtys hombergii</i> Savigny in Lamarck, 1818	*	*	*
	<i>Nephtys kersivalensis</i> McIntosh, 1908		*	
	<i>Nephtys caeca</i> (Fabricius, 1780)			*
Phyllodocidae	<i>Phyllodoce</i> sp.	*	*	
Onuphidae	<i>Diopatra</i> cf. <i>marocensis</i> Paxton et al., 1995	*	*	
Glyceridae	<i>Glycera alba</i> (Müller, 1776)	*	*	
	<i>Glycera</i> cf. <i>tridactyla</i> Schmarda, 1861		*	
Lumbrineridae	<i>Lumbrineris coccinea</i> (Renier, 1804)		*	
Polynoidae	<i>Harmothoe</i> sp.		*	
Pectinariidae	<i>Lagis</i> cf. <i>koreni</i> Malmgren, 1866		*	
Oweniidae	<i>Owenia fusiformis</i> Delle Chiaje, 1841, 1844			*




Fig. 3. Winter and summer differences in the main descriptors of the structure of the polychaete assemblages. (A) Density (ind. m⁻²). (B) Species richness. (C) Shannon diversity. (D) Evenness.




Fig. 4. Winter (A) and summer (B) dendrograms showing the three station clusters obtained in the Hierarchical Ascending Classification analysis, and the location of the respective stations in the lagoon.

percentage of sand, and the lowest recorded average density (28.6 individuals m⁻²). The dominant species were *P. africana*, *Diopatra* cf. *morocensis* and *Nephtys kersivalensis*.

The HAC analyses based on species also revealed the presence of three clusters in winter as well as in summer (Figure 5A, B). Both seasons coincided in showing a species cluster 1 including *H. diversicolor* only, and a species cluster 2 including *Capitella* sp. only. Conversely, species cluster 3 included five species in winter and 11 in summer.

Relationships between the descriptors of the polychaetes assemblages and the environmental variables

In winter, *Capitella* sp. and *N. hombergii* were positively correlated with OM (Pearson coeff. = 0.367, $P = 0.015$) and chl-a (Pearson coeff. = 0.385, $P = 0.011$), respectively. In summer, *P. africana* showed a negative correlation with OM (Pearson coeff. = -0.313, $P < 0.041$), while *H. diversicolor* and *N. kersivalensis* were positively correlated with OM

(Pearson coeff. = 0.341, $P < 0.025$) and the granulometry (Pearson coeff. = 0.480, $P = 0.001$).

In winter the first two CCA axes accounted for 85.93% of the observed variance. The species composition was mainly related to silt, OM and chl-a contents (Figure 6A). In summer, the first two CCA axes accounted for 87.91% of the relationships, with the most influencing environmental variables being OM, clay and sand (Figure 6B).

DISCUSSION

The structure of the assemblages inhabiting Oualidia largely responds to the estuarine processes and habitat mixing determined, at any given time, by the physical morphology relative to the tidal elevation inside the lagoon (El Asri *et al.*, 2015). As with all along the Moroccan Atlantic coast, tides and wind-generated waves are the dominant (natural) processes governing the morphological developments (Kalloul *et al.*, 2012). In Oualidia, however, the combined meteorological and riverine inputs are also shaping the environmental features. The

Table 2. List of the main species of each polychaete assemblage according to the IndVal index. Assemblages are named according to the species showing the highest IndVal (in bold).

Season	Cluster	Species	IndVal
Winter	1	Hediste diversicolor	238.4
		<i>Panousea africana</i>	19.1
		<i>Diopatra cf. marocensis</i>	19.1
	2	Capitella sp.	651.2
		<i>Hediste diversicolor</i>	114.0
		<i>Alkmaria romijnii</i>	29.9
	3	Glycera alba	94.5
		<i>Phyllodoce sp.</i>	63.0
		<i>Nephtys hombergii</i>	50.0
Summer	1	Hediste diversicolor	169.8
		<i>Alkmaria romijnii</i>	5.9
		<i>Harmothoe sp.</i>	3.9
	2	Capitella sp.	340.0
		<i>Glycera alba</i>	120.9
		<i>Lumbrineris coccinea</i>	87.4
	3	Panousea africana	179.6
		<i>Diopatra cf. marocensis</i>	138.8
		<i>Nephtys kersivalensis</i>	55.5

salinity tends to be lower during winter as a response to the increasing precipitation, and rises in summer due to higher evaporation rates, as well as to the lowering of the

inland freshwater inputs. In addition, there was a marked upstream-downstream gradient, with the salinities being much lower upstream due to the arrival of inland freshwater (Figure 2), as previously reported (Hennani et al., 2012; Hassou et al., 2014).

The winter increases in organic matter are linked to the arrival of a thick layer of mud following the rainwater runoff. The organic matter levels were higher in Oualidia Lagoon than those recorded at Moroccan lagoons, either Atlantic Sidi Moussa (3.6 and 12.3%) (Maanan et al., 2004) or Mediterranean Nador (0, 1 and 6.3%) (El Alami et al., 1998), and also higher than those in well studied European lagoons, such as the Venice lagoon (0.43 and 1.09%) (Bellucci et al., 2002). In turn, the high concentrations of chl-a originated from the leaching of agricultural lands in the watershed, these being so rich in nutrients that the leaching into the lagoon results in a significant increase in chl-a.

Our study represents a significant contribution to the knowledge on the polychaete macrofauna living the Oualidia lagoon, as the number of species reported is almost three times greater than those reported by Chbicheb (1996) during four seasons from December 1992 to November 1993 (Table 1). On the other hand, 11 out of the 13 species recorded in 2013 were not listed in the previous survey, while two found in 1996 appear not to be currently present (*Nephtys caeca* and *Owenia fusiformis*). These differences in the structure and diversity of the polychaete assemblages can probably be explained by the increasing levels of organic matter in the sediment, which may also be related to the changes in the hydrodynamics of the lagoon due to the construction of a pit upstream. Despite its relatively small size, Oualidia lagoon supports rather diverse polychaete assemblages, but also of molluscs (El Asri et al., 2015), when compared with other African lagoons: Aby lagoon (9 species) (Koaudio et al. 2008); Smir lagoon (12 species) (Chaouti & Bayed, 2005); Epe lagoon (10 species) (Uwadiae, 2009); and Khnifiss lagoon (17 species) (Lefrere et al., 2015). The fact that some species could not be fully identified was due mainly to the poor preservation status of the collected materials (e.g. in the case of *L. cf. koreni* or *G. cf. tridactila*). In addition, there were only anterior fragments of *D. cf. marocensis*, the specimens of *Phyllodoce* were probably juveniles too small to distinguish the key taxonomic characters or the specimens of *Harmothoe* were completely lacking the elytra. Although we were certain of the specimens' assignment to individual species, a requisite for biodiversity analyses, further effort should be made to allow more detailed taxonomic studies that would help in confirming the identity of the doubtful species. If possible these studies would include samples for molecular studies, which would be particularly helpful in identifying species such as those belonging to the *Capitella* sibling species complex (Tomioka et al., 2016). Despite their intrinsic interest, these studies are far beyond the scope of the present paper.

In the Oualidia lagoon, the three most abundant species were *H. diversicolor*, *Capitella sp.* and *N. hombergii*. The first two were also the most dominant in summer (64.7 and 17%, respectively), followed by *P. africana* (5.2%). This pattern of dominance differs completely from that reported in the Khnifess lagoon, where the most representative species were *Diopatra marocensis*, *Terebella lapidaria* and *Nicomache lumbricalis* (Artemis et al., 2006). *Diopatra marocensis* was also present in Oualidia, but its abundance was very




Fig. 5. Winter (A) and summer (B) dendrograms showing the three species clusters obtained in the Hierarchical Ascending Classification analysis.




Fig. 6. Canonical Correspondence Analysis plots. (A) Winter. (B) Summer.

low. In the Smir lagoon, the three most dominant species were *Streblospio shrubsolii* (reported as *S. dekhuyzenii*), *H. diversicolor* and *Alkmaria romijni* (Chaouti & Bayed, 2008). Coastal lagoons are highly variable in terms of environmental conditions, not only between their different areas, but also seasonally. This may certainly contribute to explain the reported changes in composition and dominance of the assemblages inhabiting different lagoons, but also to drastic changes in the species composition through time, particularly if there are associated changes in anthropogenic pressures (Hernández-Guevara *et al.*, 2008). These changes, however, are not only related to the ability of the species to respond to the environmental changes, but may also result from the existing biological interactions such as competition or predation, or may depend on intrinsic characteristics of the species, as previously discussed for similar environments (Artemis *et al.*, 2006).

Despite the temporal environmental differences found in the lagoon, and, to some extent, the differences in composition, the structure of the assemblages was the same during the two study periods. Environmental descriptors, particularly

those related to the sediment granulometry, showed a regular gradient, which was mirrored by the polychaete assemblages. In fact, the *H. diversicolor* assemblage found in the fine sediments of the inner lagoon were replaced by the *Capitella* sp. assemblage in the medium grain-sized sediment of the middle of the lagoon, and by the *Glycera alba* (winter) and *P. africana* (summer) ones in the sandy sediments closer to the inlets. The shift was, in fact, from a classical, brackish, lacunar assemblage to two different, temporal aspects of a marine assemblage (close to the inlets), with a transition assemblage in between.

In Oualidia, continental and marine environments are characteristically interwoven, as typically occurs in littoral lagoons (Amanieu *et al.*, 1980), the so-called paralic environments *sensu* Guelorget & Perthuisot (1992). Moreover, the structure of the lagoon and the associated benthic communities defined on the basis of our data agrees with the previous findings based on a physical oceanography approach that defined the lagoon as 'an estuary without a river' (Hilmi *et al.*, 2005). This structure, as well as the functioning and the influence of the nearby agricultural fields reaching the

lagoon through fresh groundwater inflows, closely resembles that described for Alfacs Bay in the Ebre's Delta (Iberian Peninsula). In this lagoon, both meio- and macrofaunal organisms are overall arranged in three main assemblages (marine, transition and brackish) from the opening to the sea to the most confined part (Palacín *et al.*, 1991, 1992). In this bay, these authors also suggested an estuarine regime in absence of a river. The main difference, from an oceanographic point of view, is that the leading force in Oualidia are tides, while in Alfacs, the main currents are wind and density generated, as Mediterranean tides are virtually negligible (Camp & Delgado, 1987). On the other hand, the highest surface and water volume, together with the largest area not in confined conditions, allows the maintenance of diverse assemblages in Alfacs Bay. This is particularly evident when specifically analysing the polychaetes: three assemblages in Oualidia vs six in Alfacs, 13 species and maximum densities of fewer than 200 individuals m^{-2} in Oualidia vs 100 species and up to 23,000 individuals m^{-2} in Alfacs (Martin *et al.*, 2000).

Despite its relatively small size, Oualidia lagoon is revealed to be, in many different senses (from oceanography to benthic ecology), an interesting environment. This, combined with the numerous human activities that have developed in the surroundings, as well as in the lagoon itself, and the declaration as a Natural Park, clearly target this lagoon as an interesting monitoring objective. Future surveys will not only show the expected changes triggered by the increasing anthropogenic influence, but may also be important in assessing the health of the lagoon ecosystem, a key issue to promote initiatives allowing to maintain or even improve the added values of the lagoon both from a naturalistic point of view (e.g. as a wintering area for migratory birds) and as a service provider (e.g. sailing, bathing, birdwatching, another touristic activities).

ACKNOWLEDGEMENTS

This paper is part of a research programme on coastal resources of the LERL/DRH-Casablanca (National Fisheries Research Institute), and contribution of DM to the Consolidated Research Group on Marine Benthic Ecology of the Generalitat de Catalunya (2014SGR120). Dr Joao Gil helped us with some of the most problematic identifications. We are deeply grateful to Mr Brahim Moutaki (INRH-Oualidia) and Mr Mohamed Amine Kaddioui (PhD student) for their help during sampling.

REFERENCES

- Amanieu M., Ferraris J. and Guélorget O. (1980) Organisation et évolution des peuplements des lagunes littorales. Application à la macrofaune benthique des étangs palavasiens. In Barbault R., Blandin P. and Meyer J.A. (eds) *Recherches d'écologie théorique. Les stratégies adaptatives*. Paris: Maloin, pp. 251–277.
- Artemis N., Petrou K., Kormas K.A. and Reizopoulou S. (2006) Inter-annual variability of soft bottom macrofaunal communities in two Ionian Sea lagoons. *Hydrobiologia* 555, 89–98.
- Bayed A., El Agbani M.A., Fekhaoui M. and Schouten J. (1988) Benthos of soft substrates in the intertidal zone of the Khnifiss lagoon. In Dakki M. and de Ligny W. (ed.) *The Khnifiss Lagoon and its surrounding environment* (Province of La'youne, Morocco). *Travaux Institut Scientifique Rabat, hors-série*, 71–80.
- Bazairi H., Bayed A., Glémarec M. and Hily C. (2003) Spatial organisation of macrozoobenthic communities in response to environmental factors in a coastal lagoon of the NW African coast (Merja Zerga, Morocco). *Oceanologica Acta* 26, 457–471.
- Bellucci L.G., Frignani M., Paolucci D. and Ravanelli M. (2002) Distribution of heavy metals in sediments of the Venice Lagoon: the role of the industrial area. *Science of the Total Environment* 295, 35–49.
- Camp J. and Delgado M. (1987) Hidrografía de las bahías del Delta del Ebro. *Revista Cubana de Investigaciones Pesqueras* 51, 351–369.
- Cao Y., Bark A.W. and Williams W.P. (1997) A comparison of clustering methods for river benthic community analysis. *Hydrobiologia* 347, 24–40.
- Carruesco C. (1989) Genese et evolution de trios lagunes du littoral atlantique depuis l'holocène: Oualidia – Moulay Bouselham (Maroc) et Arcachon (France), Tome I, pp. 24–100. Thèse de doctorat en Sciences.
- Chaouti A. and Bayed A. (2005) Diversité taxonomique et structure de la macrofaune benthique des substrats meubles de la lagune de Smir. *Travaux de l'Institut Scientifique* 4, 33–42.
- Chbicheb A. (1996) *Organisation biogéologique d'un bassin paralique: la lagune d'Oualidia (Côte Atlantique Marocaine)*. PhD thesis. University of Nantes, Nantes, France.
- Chaouti A. and Bayed A. (2008) Spatial patterns of soft-bottom macroinvertebrates and relationships with environmental conditions in a north African coastal lagoon (Smir lagoon, Morocco). *Vie et Milieu* 58, 25–35.
- Dakki M. and de Ligny W. (1988) The Khnifiss lagoon and its surrounding environment (Province of La'youne, Morocco). *Travaux Institut Scientifique de Rabat, Hors-série*. 1–172.
- Delle Chiaje S. (1841 & 1844) [Polychaeta context] *Descrizione e Notomia degli Animali Invertebrati della Sicilia Citeriore osservati vivi negli anni 1822–1830*. [Tome 1, 98 pp.; Tome 2, 146 pp.; Tome 3, 142 pp. [Molluschi Acefali, Bracciopedi, Cirropedi, Crostacei, Anellosi]; Tome 4, 142 pp.; Tome 5, 165 pp. [1841, incl. Annelosi Afroditici]; Tome 6–7, pl. 1–173 [1841], pl. 174–181 [1844]; Tome 8 (Appendici, osservazioni critiche, indice generale), 48 pp. [1844] Napoli, Stabilimento Tipografico di C. Batelli e Co.
- Dufrène M. and Legendre P. (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Society of America* 67, 345–366.
- El Alami M., Mahjoubi R., Damnati B., Kamel S., Icole M. and Taieb M. (1998) Sédimentologie et géochimie organique des sédiments superficiels de la lagune de Nador (Maroc nord-oriental). *Journal of African Earth Sciences* 26, 249–259.
- El Asri F., Zidane H., Maanan M., Tamsouri M. and Errhif A. (2015) Taxonomic diversity and structure of the molluscan fauna in Oualidia lagoon (Moroccan Atlantic coast). *Environmental Monitoring and Assessment* 187, 1–10.
- El Kamcha R., Bououarour O., Boutahar L., El Fatouani Z., El Adnani A., Zourarah B., Sghaier Y.R., Benhoussa A. and Bazairi H. (2015) Assessment of the environmental health status of the Marchica lagoon (Morocco, Mediterranean) using *Cymodocea nodosa* meadows and its associated macrofauna: preliminary results. *PeerJ PrePrints* 3, e1325.
- Giangrande A., Licciano M. and Musco L. (2005) Polychaetes as environmental indicators revisited. *Marine Pollution Bulletin* 50, 1153–1162.

- Gil J. (2011) *The European fauna of Annelida Polychaeta*. Doctoral thesis. Universidade de Lisboa, Lisboa. <http://hdl.handle.net/10451/4600>.
- Gray J. S. (1974) Animal-sediment relationships. *Oceanography and Marine Biology. An Annual Review* 12, 223–261.
- Gray J. S. (2002) Species richness of marine soft sediments. *Marine Ecology Progress Series* 244, 285–297.
- Guelorget O. and Perthuisot J.P. (1992) Paralic ecosystems. *Vie et Milieu* 42, 215–251.
- Guelorget O., Perthuisot J., Frisoni G. and Monti D. (1987) Le rôle du confinement dans l'organisation biogéologique de la lagune de Nador (Maroc). *Oceanologica Acta* 10, 435–444.
- Hassou N., Maanan M., Hennani M., Zourarah B. and Assobhei O. (2014) Spatial and temporal variation of faecal pollution indicators (*Escherichia coli* and faecal streptococci) and physico-chemical parameters at the Oualidia lagoon and its watershed (Morocco). *International Journal of Current Microbiology and Applied Sciences* 3, 675–694.
- Hennani M., Maanan M., Robin M., Chedad K. and Assobhei O. (2012) Temporal and spatial distribution of faecal bacteria in a Moroccan lagoon. *Polish Journal of Environmental Studies* 21, 627–634.
- Hernández-Guevara N.A., Pech D. and Ardisson P. (2008) Temporal trends in benthic macrofauna composition in response to seasonal variation in a tropical coastal lagoon, Celestun, Gulf of Mexico. *Marine and Freshwater Research* 59, 772–779.
- Hilmi K., Koutitonsky V., Orbi A., Lakhdar J. and Chagdali M. (2005) Oualidia lagoon, Morocco: an estuary without a river. *African Journal of Aquatic Science* 30, 1–10.
- Holm-Hansen O., Lorenzen C.J., Holmes R.W. and Strickland J.D.H. (1965) Fluorometric determination of chlorophyll. *Journal of Marine Science* 30, 3–15.
- Horst R. (1919) Twee sedentaire Polychaeten uit he brakke water var Nederland. *Zoologische Mededeelingen (Leiden)* 5, 100–102.
- Idardare Z., Chiffolleau J.F., Moukrim A., Ait Alla A., Auger D., Lefrere L. and Rozuel E. (2008) Metal concentrations in sediment and *Nereis diversicolor* in two Moroccan lagoons: Khnifiss and Oualidia. *Chemistry and Ecology* 24, 329–340.
- Kalloul S., Hamid W., Maanan M., Robin M. and Sayouty E.H. (2012) Source contributions to heavy metal fluxes into the Loukous Estuary (Moroccan Atlantic Coast). *BioOne* 28, 174–183.
- Koaudio K.N., Diomandé D., Ouattara A., Koné Y.J.M. and Gourène G. (2008) Taxonomic diversity and structure of benthic macroinvertebrates in Aby Lagoon (Ivory Coast, West Africa). *Pakistan Journal of Biological Science* 11, 2224–2230.
- Lamarck J.B. (1818) *Histoire naturelle des Animaux sans Vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie*. Volume 5. Paris: Déterville & Verdière.
- Lefrere L., Ouassas M., Guillois B., Gillet P. and Moukrim A. (2015) Macrobenthic community structure of soft-bottom sediments in the Khnifiss lagoon, South of Morocco. *Journal of Material and Environmental Science* 6, 2226–2236.
- Maanan M., Ruiz-Fernández A.C., Maanan M., Fattal P., Zourarah B. and Sahabi M. (2014) A long-term record of land use change impacts on sediments in Oualidia lagoon, Morocco. *International Journal of Sediment Research* 29, 1–10.
- Maanan M., Zourarah B., Carruesco C., Aajjane A. and Naud J. (2004) The distribution of heavy metals in the Sidi Moussa lagoon sediments (Atlantic Moroccan Coast). *Journal of African Earth Sciences* 39, 473–483.
- McIntosh W.C. (1908) *A monograph of the British annelids*. London: Ray Society.
- Malmgren A.J. (1866) Nordiska Hafs-Annulater. *Öfversigt af Königlich Vetenskapsakademiens förhandlingar, Stockholm*, 22, 51–110.
- Martin D., Pinedo S. and Sardá R. (2000) Distribution patterns and trophic structure of soft-bottom Polychaete assemblages in a North-Western Mediterranean shallow-water bay. *Ophelia* 53, 1–17.
- Müller O.F. (1776) *Zoologicae Danicae Prodrum, seu Animalium Daniae et Norvegiae indigenarum characteres, nomina et synonyma imprimis popularium*. Havniae [Copenhagen]: Hallageriis.
- Newton T.J., Allran J.W., O'Donnell J.A., Bartsch M.R. and Richardson W.B. (2003) Effects of ammonia on juvenile unionids (*Lampsilis cardium*) in laboratory sediment toxicity tests. *Environmental Toxicology and Chemistry* 22, 2554–2560.
- Palacín C., Gili J.M. and Martin D. (1992) Evidence for coincidence of meiofauna spatial heterogeneity with eutrophication processes in a shallow-water Mediterranean bay. *Estuarine, Coastal and Shelf Science* 35, 1–16.
- Palacín C., Martin D. and Gili J.M. (1991) Features of spatial distribution of benthic infauna in a Mediterranean shallow-water bay. *Marine Biology* 110, 315–321.
- Paxton H., Fadlaoui S. and Lechapt J.-P. (1995) *Diopatra marocensis*, a new brooding species of Onuphidae (Annelida: Polychaeta). *Journal of the Marine Biological Association of the United Kingdom* 75, 949–955.
- Pielou E. C. (1966) Shannon's formula as a measure of specific diversity: its use and measure. *American Naturalist* 100, 463–465.
- Pocklington P. and Wells P.G. (1992) Polychaetes. Key taxa for marine environmental quality monitoring. *Marine Pollution Bulletin* 24, 593–598.
- Read G. and Fauchald K. (eds) (2017) World Polychaeta database. Available at <http://www.marinespecies.org/polychaeta> (accessed 16 January 2017).
- Renier S.A. (1804) *Prospetto della Classe dei Vermi, nominati el ordinati secondo il Sistema de Bosc*. Padua.
- Rullier F. and Amoureux L. (1970) Nouvelle contribution a l'étude de la faune des Annélides Polychètes du Maroc. *Bulletin de la Société des Sciences Naturelles et Physiques du Maroc* 49, 109–142.
- Schmarda L.K. (1861) *Neue wirbellose Thiere beobachtet und gesammelt auf einer Reise un die Erdr 1853 bis 1857. Erster Band (zweite halfte) Turbellarian, Rotatorien un Anneliden*. Leipzig: Wilhelm Engelmann.
- Shannon C.E. (1948) A mathematical theory of communication. *Bell System Technical Journal* 27, 379–423.
- Sif J., Rouhi A., Gillet P. and Moncef M. (2012) Diversité et écologie des Annélides Polychètes du littoral atlantique de la région d'El Jadida (Maroc). *Bulletin de l'Institut Scientifique, Rabat, section Sciences de la Vie* 34, 95–106.
- Tomioka S., Kondoh T., Sato-Okoshi W., Ito K., Kakui K. and Kajihara H. (2016) Cosmopolitan or cryptic species? A case study of *Capitella teleta* (Annelida: Capitellidae). *Zoological Science* 33, 545–554.
- Uwadiae R.E. (2009) Response of benthic macroinvertebrate community to salinity gradient in a sandwiched coastal lagoon. *Report and Opinion* 1, 45–55.
- Vakharia A.J. and Wemmerlöv U. (1995) A comparative investigation of hierarchical clustering techniques and dissimilarity measures applied

to the cell formation problem. *Journal of Operational Management* 13, 117–138.

Venturini N., Tommasi L.R., Bicego M. C. and Martins C. C. (2004) Characterization of the benthic environment of a coastal area adjacent to an oil refinery, Todos os Santos Bay (NE-Brazil). *Brazilian Journal of Oceanography* 52, 123–134.

and

Zourarah B., Maanan M., Carruesco C., Aajjane A., Mehdi K. and Conceição Freitas M. (2007) Fifty-year sedimentary record of heavy

metal pollution in the lagoon of Oualidia (Moroccan Atlantic coast). *Estuarine, Coastal and Shelf Science* 72, 359–369.

Correspondence should be addressed to:

F. El Asri
Health and Environment Laboratory,
Faculty of Sciences Ain Chock, University Hassan II,
B.P. 5366 Maârif, 20100 Casablanca, Morocco
email: fatimaelasri25@gmail.com