

Associations among child abuse, mental health, and epigenetic modifications in the proopiomelanocortin gene (*POMC*): A study with children in Tanzania

TOBIAS HECKER,^{a,d} KARL M. RADTKE,^b KATHARIN HERMENAU,^{b,d} ANDREAS PAPASSOTIROPOULOS,^c
AND THOMAS ELBERT^{b,d}

^aUniversity of Zurich; ^bUniversity of Konstanz; ^cUniversity of Basel; and ^dvivo international

Abstract

Child abuse is associated with a number of emotional and behavioral problems. Nevertheless, it has been argued that these adverse consequences may not hold for societies in which many of the specific acts of abuse are culturally normed. Epigenetic modifications in the genes of the hypothalamus–pituitary–adrenal axis may provide a potential mechanism translating abuse into altered gene expression, which subsequently results in behavioral changes. Our investigation took place in Tanzania, a society in which many forms of abuse are commonly employed as disciplinary methods. We included 35 children with high exposure and compared them to 25 children with low exposure. Extreme group comparisons revealed that children with high exposure reported more mental health problems. Child abuse was associated with differential methylation in the proopiomelanocortin gene (*POMC*), measured both in saliva and in blood. Hierarchical clustering based on the methylation of the *POMC* gene found two distinct clusters. These corresponded with children's self-reported abuse, with two-thirds of the children allocated into their respective group. Our results emphasize the consequences of child abuse based on both molecular and behavioral grounds, providing further evidence that acts of abuse affect children, even when culturally acceptable. Furthermore, on a molecular level, our findings strengthen the credibility of children's self-reports.

Child abuse is commonly defined as any act of commission by a parent or any other caregiver that results in harm, potential for harm, or threat of harm to a child (Leeb, Paulozzi, Melanson, Simon, & Arias, 2008). Child abuse may result in emotional and behavioral problems that begin in childhood and can persist throughout adolescence and adulthood (Carr, Martins, Stingel, Lemgruber, & Juruena, 2013). For example, child abuse increases the risk of developing depression, anxiety disorders, posttraumatic stress disorder (PTSD), substance abuse, reduced self-esteem, suicidal behavior, conduct disorder, and aggressive or delinquent behavior (Catani, Jacob, Schauer, Kohila, & Neuner, 2008; Dube et al., 2003; Hermenau, Hecker, Elbert, & Ruf-Leuschner, 2014; Sugaya et al., 2012), as confirmed by numerous longitudinal studies

(Kaplan et al., 1998; Widom, DuMont, & Czaja, 2007). Most abused children have been exposed to multiple forms of abuse, and the greater the number of different forms of abuse, the higher the likelihood of subsequent psychopathologies (Teicher, Samson, Polcari, & Mcgreenery, 2006). Furthermore, abused individuals with a psychiatric disorder are characterized by earlier onset of disease, increased symptom severity, increased comorbidity, increased risk of suicide, poorer treatment response, and shorter interval before recurrence than are individuals with the same diagnoses who were not abused (Harkness, Bagby, & Kennedy, 2012; Nanni, Uher, & Danese, 2012; Teicher & Samson, 2013). Finally, child abuse is a major burden not only upon the affected individual but also upon the society at large due to the high costs associated with the utilization of healthcare, educational, welfare, and law enforcement services (Fang, Brown, Florence, & Mercy, 2012).

It has been argued that the aforementioned adverse consequences may not hold for societies or communities in which many of the specific acts of child abuse are culturally normed and highly prevalent. In other words, abused individuals in communities that deem such practices to be socially acceptable and legal would find the effects to be less harmful than would those living in societies in which such practices are unacceptable or illegal. Lansford et al. (2005) empirically tested this idea in six countries. They found that more frequent corporal punishment is related to more aggression and more anx-

This research was supported by the Deutsche Forschungsgemeinschaft, by the European Research Council, and by the nongovernmental organization vivo international. We are grateful to all the children who participated in this study for their readiness to participate and willingness to discuss often intimate and painful subjects. We are very grateful to our very motivated and reliable research assistants, including Huruma Kipagile, Lulu Nziku, and Heike Riedke. We also thank Danie Meyer-Parlapanis, James Moran, and Justin Preston who critically reviewed the manuscript. The first and second authors contributed equally to this work.

Address correspondence and reprint requests to: Tobias Hecker, Division of Psychopathology & Clinical Intervention, Department of Psychology, University of Zurich, Binzmuehlestrasse 14/17, Zurich 8050, Switzerland; E-mail: t.hecker@psychologie.uzh.ch.

ity in all six countries. However, the strength of the relation did vary by the perceived normativeness across countries. Many other studies demonstrated detrimental consequences for the psychological well-being and development of abused children, regardless of whether the surrounding society deems such practices acceptable (Ani & Grantham-McGregor, 1998; Hecker, Hermenau, Isele, & Elbert, 2014; Hermenau et al., 2011).

There are many countries in which many of the acts constituting child abuse are legal and socially accepted. In Tanzania, for example, a national survey with a representative sample of more than 3,700 youths revealed that the great majority (almost 75%) of both girls and boys had experienced physical abuse and more than one-quarter faced emotional abuse prior to the age of 18 (UNICEF, 2011). In concordance, we and others reported the use of harmful physical acts and psychological tactics on behalf of caregivers toward children to be highly prevalent in Tanzanian families and schools (Feinstein & Mwahombela, 2010; Hecker et al., 2014). In April 2013, the Tanzanian government reportedly confirmed that the use of corporal punishment in public schools persists (Tanzania Daily News, 2013). Given such high prevalence of child abuse, it is vital for both individuals and societies to have a better understanding of the potential effects of abuse. In particular, we must study whether the negative consequences of physical and emotional abuse of children are diminished in societies where such acts are legal and socially accepted.

Most studies on mental health problems have been conducted in Western samples. However, findings from the Democratic Republic of the Congo, Ethiopia, and Nigeria have shown that various mental health problems such as anxiety disorders, affective disorders, and hyperactivity are also common phenomena in sub-Saharan Africa (Adelekan, Ndom, Ekpo, & Oluboka, 1999; Kashala, Elgen, Sommerfelt, & Tylleskar, 2005). Adelekan et al. (1999) indicated a prevalence rate of internalizing problems of 7.3% and of externalizing problems of 8% in a representative sample from Nigeria. Kashala et al. (2005) compared their findings in a study with a representative sample in the Congo (Goodman, Meltzer, & Bailey, 1998) with prior findings from Great Britain. They found that the mean scores on all subscales of the Strength and Difficulties Questionnaire (SDQ) were significantly higher than the British mean scores of a comparable sample. Hence, Cortina, Sodha, Fazel, and Ramchandani (2012) concluded that child and adolescent mental health problems are also common in sub-Saharan Africa.

Child Abuse and the Hypothalamus–Pituitary–Adrenal (HPA) Axis

The HPA axis, when functioning properly, helps us to deal with crises. It describes a set of interactions between the hypothalamus, the pituitary gland, and the adrenal gland, which results in the release of its effector cortisol (Chrousos & Gold, 1992; de Kloet, Joëls, & Holsboer, 2005). Upon stress per-

ception, corticotropin-releasing hormone (CRH) and arginine vasopressin (AVP) are released from the hypothalamic paraventricular nucleus to activate the synthesis of proopiomelanocortin (POMC) in the anterior pituitary. POMC is processed into several peptides including adrenocorticotropin hormone (ACTH). Finally, ACTH is released into the blood stream and triggers the secretion of cortisol from the adrenal cortex. At each organizational level, the HPA axis is tightly regulated by negative feedback loops mediated by glucocorticoid receptors (GRs). After binding their ligand, cortisol, GRs dampen HPA axis activity.

Child abuse is translated into negative long-term mental health outcomes via the HPA axis. It plays a central role, because it is tuned to experiences occurring early in life, making it highly susceptible to early childhood adversities (Heim & Nemeroff, 2001). For example, adults with a history of childhood maltreatment displayed altered ACTH and cortisol responses following exposure to an acute stressor (Carpenter et al., 2007; Heim et al., 2000). HPA axis dysregulation is a key feature of a range of psychopathological symptoms (Chrousos & Gold, 1992; de Kloet et al., 2005). Both human and animal studies show that unremitting threat or stress weakens the immune response and increases abdominal fat, mental ill health, and depression via alterations of HPA functioning (McEwen & Lasley, 2002). HPA axis function, and with it behavioral changes, may be stably altered through aberrant epigenetic modifications, established as the result of child abuse.

Epigenetic Modifications of HPA Axis Genes

Of the various and complex mechanisms leading to epigenetic modification, DNA methylation is currently being studied most extensively. In humans, the relationship between early life adversities and the methylation of the GR has been extensively studied. GR promoter methylation is associated with both child abuse and psychopathology (Dammann et al., 2011; Hompes et al., 2013; Labonte, Azoulay, Yerko, Turecki, & Brunet, 2014; McGowan et al., 2009). Suicide victims with a history of childhood abuse displayed increased GR methylation in brain tissue (Labonte et al., 2012; McGowan et al., 2009). Higher GR methylation in peripheral blood mononuclear cells has been observed in patients suffering from borderline personality disorder (i.e., in individuals who have usually been exposed to severe forms of abuse during development). Disruption or lack of adequate nurturing, as measured by child maltreatment and inadequate parental care, was also associated with increased GR promoter methylation (Perroud et al., 2011; Tyrka, Price, Marsit, Walters, & Carpenter, 2012). In addition, epigenetic changes in the *POMC* gene may promote HPA axis dysfunction. Recent studies suggest epigenetic programming of *POMC* operates through nutritional cues, such as being underweight (Ehrlich et al., 2010), while other research suggests an association with alcohol abuse and dependence via increased craving (Muschler et al., 2010). Animal models

demonstrate epigenetic programming of additional HPA axis genes such as *CRH* (Mueller & Bale, 2008) and *AVP* (Murtagroyd et al., 2009). Thus, current research has highlighted epigenetic modifications in genes associated with the HPA axis as being a possible driving force producing child abuse-induced disorders.

In the present study we investigated associations of child abuse with both the phenotype and the methylation status of genes related to the HPA axis in Tanzanian children. We limited our analyses of DNA methylation to the genes coding for the main components of the HPA axis, that is, the genes coding for *AVP*, *CRH*, and *POMC*, from which *ACTH* is cleaved. In addition, we included the gene encoding *GR*, nuclear receptor subfamily 3, group C, member 1 (*NR3C1*), because several studies demonstrated its methylation status as being predictive for childhood abuse (Labonte et al., 2012; McGowan et al., 2009; Perroud et al., 2011; Tyrka et al., 2012). We hypothesized that exposed children (a) report more emotional and behavioral problems and (b) display altered epigenetic modifications in the genes related to HPA axis functioning.

Methods

Procedure

In the context of a larger research project, a team of Tanzanian and German psychologists conducted structured interviews with a sample of Tanzanian school children ($N = 409$). Interviewers were taught in interview skills during a 2-week training session. Furthermore, the Tanzanian interviewers were trained to translate from English to Swahili and back in order to assist the German researchers. All instruments were translated in written form to Swahili. A valid and accurate translation into English was ensured through the use of a written, blind backtranslation. In the total sample, 33 interviews were rated by two independent assessors to determine high interrater reliability. Prior to the interviews, we sent a written informed consent form to all parents or caregivers of the children from Class 2 to 7 (age 6–15) explaining the purpose of the study.

Based on these structured interviews, we selected children who had been exposed to high levels of physical and emotional abuse in their homes and those who had been exposed to only low levels of physical and emotional abuse. An a priori power analysis ($\alpha = 0.05$, power = 0.80, $d = 0.80$) using G*Power software (Faul, Erdfelder, Lang, & Buchner, 2007) indicated a required sample size of $n = 26$ per group to detect significant group differences. Therefore, we aimed for two groups from the extreme ends of the abuse continuum (no abuse vs. high levels of abuse) of 30 children each. Because many children, particularly younger children, reported a strong fear of drawing blood, due to harmful experiences in the Tanzanian health system, we decided not to include children of 8 years or younger. We sent an invitation and informed consent form to 96 parents and caregivers of the selected children clarifying that donating blood and saliva samples would be entirely voluntary and no monetary com-

pensation would be offered. In total, 64% ($n = 61$ of 96) of the parents and caregivers signed the informed consent. We were unable to recruit enough children who had never been exposed to any type of abuse. This is not too surprising, given that several acts of child abuse are culturally normed and highly prevalent in Tanzania. Almost 75% reported exposure to physical abuse in a nationally representative sample (UNICEF, 2011). Nevertheless, our sampling approach resulted in two extreme groups: one group ($n = 35$) reporting high levels of child abuse (i.e., six or more different types) and one group ($n = 25$) reporting low levels of child abuse (i.e., four or fewer different types). In the total sample, 173 (42%) children reported low levels of child abuse with only 8 (2%) reporting no exposure to any form of child abuse. In contrast, 175 (43%) children of the total sample reported high levels of child abuse. Only children with an informed consent signed by their caregivers and who also assented themselves orally were included in the study (only 1 child refused to participate despite parents' informed consent). A trained nurse from the University of Konstanz with extensive work experience in East Africa collected the blood and saliva samples. The Ethical Review Board of the University of Konstanz approved the study. Other, nonepigenomic parts of the data gathered for the total sample are presented in reports by Hecker et al. (2014); Hermenau, Eggert, Landolt, and Hecker (2015); and Hermenau et al. (2014).

Participants

The children participating in this study were enrolled at a primary school in a city of approximately 150,000 inhabitants in southern Tanzania. The high-exposure group consisted of $n = 35$ children (60% girls) who were on average 11.31 years old ($SD = 1.47$, range = 9–15). The low-exposure group consisted of $n = 25$ children (56% girls) who were on average 11.76 years old ($SD = 1.20$, range = 10–14).

Measures

All instruments were applied as a structured interview in Swahili. The first part of the interview consisted of sociodemographic information, including age, grade, and gender.

Child abuse. We assessed exposure to abuse at home using the Maltreatment and Abuse Chronology of Exposure—Pediatric Version (Isele et al., 2015; Teicher & Parigger, 2015). The Maltreatment and Abuse Chronology of Exposure is a structured interview for children consisting of 45 dichotomous (yes/no) questions, measuring witnessed or self-experienced forms of child maltreatment throughout the lifetime. In this study, we only used the 14 items covering possible forms of physical and emotional abuse (see Table 1) by an adult person living in the same household (e.g., parent, relative, or caregiver) or by a minor living in the same household (e.g., housemaid or sibling). In Tanzania many children grow up not only with their parents in one household but also with

Table 1. Occurrence of physical and emotional abuse during the children's lifetime

| | Exposure % (n) | |
|---|----------------|---------|
| | High | Low |
| Physical abuse | | |
| 1. Has any adult intentionally pinched, slapped, punched, or kicked you? | 80 (28) | 48 (12) |
| 2. Has any adult spanked you with the palm of his/her hand on buttocks, arms, or legs? | 74 (26) | 24 (6) |
| 3. Has any adult spanked you with an object such as a belt, stick, tube, or wooden spoon? | 89 (31) | 60 (15) |
| 4. Has any adult hit you so hard that you were injured? | 40 (14) | 4 (1) |
| 5. Has any minor intentionally pinched, slapped, punched, or kicked you? | 74 (26) | 24 (6) |
| 6. Has any minor spanked you with the palm of his/her hand on buttocks, arms, or legs? | 51 (18) | 4 (1) |
| 7. Has any minor spanked you with an object such as a belt, stick, tube, or wooden spoon? | 31 (11) | 0 (0) |
| 8. Has any minor hit you so hard that you were injured? | 46 (16) | 0 (0) |
| Emotional abuse | | |
| 9. Has any adult called you names or said hurtful things (e.g., fat, ugly, stupid)? | 51 (18) | 32 (8) |
| 10. Has any adult yelled or screamed at you? | 86 (30) | 68 (17) |
| 11. Has any adult locked you in a dark & narrow place (e.g. basement, closet)? | 20 (7) | 0 (0) |
| 12. Has any minor called you names or said hurtful things (e.g. fat, ugly, stupid)? | 77 (27) | 4 (1) |
| 13. Has any minor yelled or screamed at you? | 60 (21) | 8 (2) |
| 14. Has any minor locked you in a dark and narrow place (e.g. basement, closet)? | 3 (1) | 0 (0) |

Note: Adult, Person living in the same household (e.g., parent, relative, or caregiver); minor, person under the age of 18 living in the same household (e.g., housemaid or sibling).

other members of their extended families. We also focused on minors in the household because in urban Tanzania many children are raised by an underaged housemaid (12–17) as primary caregiver, while both parents have to work. Using an event checklist, we assessed the presence of different types of abuse but not the frequency. We calculated an abuse score by totaling up all of the question responses. The possible score ranges from 0 to 14. The Cohen κ coefficient measuring the interrater reliability was >0.99 (0.99–1).

Mental health. The self-evaluation of internalizing and externalizing problems was assessed with the SDQ (Goodman, Ford, Simmons, Gatward, & Meltzer, 2000; Goodman et al., 1998). We used the self-report version for children in interview form, which consists of 25 statements. The total difficulties score is generated by summing the scores of all items, except the items for prosocial behavior, and ranges from 0 to 40. A score over 16 indicates an enhanced level of internalizing and externalizing problems. In the present sample, the Cronbach α coefficient was 0.71 and the Cohen κ coefficient was 0.99 (0.94–1).

The University of California at Los Angeles PTSD Reaction Index for Children DSM-IV (Steinberg, Brymer, Decker, & Pynoos, 2004) was used to screen for symptoms of PTSD, again in interview form. For each DSM-IV symptom, the frequency of occurrence within the last month is scored. The PTSD severity score ranges from 0 to 68. In the present sample, the Cronbach α was 0.92 and the Cohen κ was 0.98 (0.82–1).

The severity of depressive symptoms was assessed by means of the Children's Depression Inventory (Kovacs, 2001; Sitarenios & Kovacs, 1999), which has already been successfully implemented and validated in Tanzanian settings (Traube, Dukay, Kaaya, Reyes, & Mellins, 2010; Wallis & Dukay, 2009).

For each of its 27 items, the children were offered three statements and asked to choose the one which best describes their situation. The maximum score possible is 54. A threshold of 12 has been established as being ideal for identifying children at risk of depression (Kovacs, 2001; Kovacs, Goldstein, & Gastonis, 1993; Traube et al., 2010). In the present sample, the Cronbach α was 0.81 and the Cohen κ was 0.99 (0.92–1).

DNA methylation. Lymphocytes from blood were isolated via a Ficoll gradient and stored in a preservation solution (DNAgard[®] Tissues & Cells, Biomatrix, San Diego, CA) in order to ensure recovery of high-quality DNA. In addition, saliva samples were collected and stored using the Oragene •DISCOVER (OGR-500) saliva collection kit (DNA Genotek Inc., Ontario, Canada). The tissue samples were subjected to DNA-extraction (DNeasy[®] Blood & Tissue Kit, Qiagen, Hilden, Germany). Genome-wide analysis of DNA methylation was then conducted at the Barts and the London Genome Centre (Queen Mary University of London). One microgram (1 μ g) of genomic DNA was bisulfite converted (EZ DNA Methylation Kit, Zymo) and applied to the Human Methylation 450K array (Illumina). The raw data were pre-processed using both the R package lumi (Du, Kibbe, & Lin, 2008) and beta mixture quantile dilation as suggested elsewhere (Marabita et al., 2013). After preprocessing, DNA methylation was assessed for all of the 41, 26, 14, and 14 cytosine nucleotide–phosphate–guanine nucleotide (CpG) sites associated with the *GR* gene (*NR3C1*), the *POMC* gene, the *CRH* gene, or the *AVP* gene, respectively.

Transcription factor binding site (TFBS). To reveal potential functional properties associated with the CpG sites included in our study, the respective sequences were submitted

Table 2. Demographic characteristics of children with high and low exposure to child abuse

| | High Exposure (<i>n</i> = 35) | | Low Exposure (<i>n</i> = 25) | | <i>t</i> | Adj. <i>p</i> | <i>d</i> |
|-------------|-----------------------------------|-----------|----------------------------------|-----------|----------|---------------|----------|
| | <i>M</i> | <i>SD</i> | <i>M</i> | <i>SD</i> | | | |
| Abuse types | 7.80 | 1.26 | 2.64 | 1.29 | 15.47 | <.001 | 4.05 |
| SDQ score | 12.31 | 5.83 | 7.48 | 4.83 | 3.50 | <.001 | 0.81 |
| UCLA score | 9.77 | 11.47 | 2.04 | 5.04 | 3.55 | <.001 | 0.77 |
| CDI score | 9.14 | 5.59 | 3.64 | 3.33 | 4.76 | <.001 | 1.00 |

Note: *t*, Test statistics based on Welch *t* test; Adj. *p*, adjusted *p* value based on Welch *t* test corrected for alpha inflation due to multiple testing; *d*, Cohen effect size; SDQ, Strengths and Difficulties Questionnaire; UCLA, University of California at Los Angeles PTSD Reaction Index for Children DSM-IV; CDI, Children's Depression Inventory.

to the Jaspas database (Mathelier et al., 2014) in order to predict known TFBSs. A conservative threshold of 90% sequence identity was applied.

Data analysis

For the analyses regarding either mental health or exposure to abuse, parametric Welch *t* tests were performed. For DNA methylation, individual 2 (Abuse) × 2 (Gender) analyses of variance (ANOVAs) for each CpG site were performed using exposure to abuse and gender as between-group factors. We included gender in these analyses in order to account for potential effects arising from gender on DNA methylation. For three CpGs in blood (cg27107893, cg02079741, cg09916783) and one in saliva (cg23035419), the models did not fulfill the requirement of homogeneity of variances, as indicated by a significant Levene test (Fox & Weisberg, 2011) and are thus not reported. Nonparametric tests could not be performed because these would not control for the potential influence of gender. In addition, we computed individual 2 (tissue) × 2 (gender) ANOVAs for each CpG site using tissue and gender as between-group factors. Because of the heterogeneity of variances, 25 probes were excluded from the analyses (*NR3C1*: cg06613263, cg08818984, cg08845721, cg10847032, cg18998365, cg19457823, cg26720913, cg27107893; *POMC*: cg02079741, cg03560973, cg08030082, cg09527270, cg09672383, cg09916783, cg13025668, cg16302441, cg20387815, cg20807790; *CRH*: cg00603617, cg23027580; and *AVP*: cg03279206, cg04360210, cg14065127, cg23035419, cg24257309). Nonparametric tests could not be performed because these would not control for the potential influence of gender.

All analyses used a two-tailed $\alpha = 0.05$. Our metric for a small effect size was $d \geq 0.20$ or $\eta^2 \geq 0.01$, for a medium effect $d \geq 0.50$ or $\eta^2 \geq 0.06$, and for a large effect $d \geq 0.80$ or $\eta^2 \geq 0.13$. To adjust for multiple testing (for three mental health variables and across the CpG sites for each gene), *p* values were computed according to Benjamin–Hochberg (Benjamini & Hochberg, 1995) applying a false discovery rate of 0.05. In an exploratory approach, we also considered the unadjusted *p* values. All statistical analyses were performed using IBM SPSS Statistics version 21 for Mac or R for Mac version 3.0.3.

Results

Mental health

Table 2 displays the descriptive statistics for both groups. In concordance with the sample selection, the high-exposure group reported a substantially higher number of different abuse types than the low-exposure group. The differences between the two groups are especially notable for the items indicating that a minor in the household was the perpetrator of the abuse (see Table 1). All mental health variables (SDQ, UCLA PTSD Reaction Index, and Children's Depression Inventory) differed significantly between groups with medium to large effects (see Table 2). In total, $n = 11$ (31%) children in the high-exposure group showed an enhanced level of internalizing and externalizing problems compared to $n = 1$ (4%) in the low-exposure group. Accordingly, $n = 9$ (26%) children in the high-exposure group fulfilled the clinical diagnosis for PTSD compared to $n = 2$ (8%) in the low-exposure group. In addition, $n = 10$ (29%) children in the high-exposure group were at risk of suffering from depression compared to $n = 1$ (4%) in the low-exposure group.

DNA methylation of genes associated with the HPA axis

We found a group difference between the high-exposure and low-exposure groups in *POMC* with higher DNA methylation in children with high exposure. This effect was particularly evident in saliva. In the saliva of the high-exposure group, one CpG site was significantly hypermethylated in one-tailed tests at an adjusted significance level of .05, and three additional CpG sites would be significantly hypermethylated an adjusted significance level of .10 (Figure 1, Figure 2, and Table 3). Considering unadjusted *p* values as well, three additional CpG sites belonging to *POMC* were differentially methylated in the saliva of the high-exposure group. All of the aforementioned CpG sites displayed medium to large effect sizes. In saliva, two more CpG sites in *POMC* displayed moderate effect sizes, although unadjusted *p* values exceeded the significance level of .05. In blood, six CpG sites in *POMC* were differentially methylated if unadjusted *p* values are considered. These six CpG sites displayed medium to large effect sizes. In blood, one additional CpG




Figure 1. (Color online) Mean methylation differences in high- and low-exposure groups. The effect size and the level of significance are color coded (online only) or depicted by the shape, respectively. *AVP*, Arginine-vasopressin gene; *CRH*, corticotropin-releasing hormone gene; *NR3C1*, nuclear receptor subfamily 3, group C, member 1 glucocorticoid receptor gene; *POMC*, proopiomelanocortin gene.

site in *POMC* displayed a moderate effect size, although unadjusted p values exceeded the significance level of .05.

For the remaining HPA axis genes investigated, we did not find a clear group difference in DNA methylation. In saliva, four CpG sites were differently methylated in *GR* and one in *CRH*, displaying moderate effect sizes and unadjusted p values below .05. In the blood of the high-exposure group, one CpG was hypermethylated in *CRH* at an adjusted significance level of .05, displaying a large effect. If uncorrected p values were considered, one additional differentially methylated CpG could be found in *AVP*, displaying a medium effect. If only effect sizes were considered, two additional CpG sites associated with *GR* and one associated with *AVP* differed between the groups in blood, displaying moderate effect sizes, but no significant p values were obtained.

Because we found the most pronounced effects in *POMC*, we inspected the 7 and 9 CpGs, which differed between the groups with at least moderate effect size in blood and saliva, respectively, in more detail. A comparison with the Jaspar database (Fox & Weisberg, 2011) revealed that 5 and 6 of these CpGs are either located in or directly flanking a potential TFBS. The potential TFBSs included TFAP2A, ZEB1, THAP1, YY1, BRCA1, E2F, ZNF354C, MZF1, and SPIB. It is interesting that all of these CpGs are located in the 5' promoter, whose methylation status has been shown to modulate

transcriptional activity of the *POMC* gene (Newell-Price, King, & Clark, 2001). Our analyses covered 11 and 12 CpGs in this region in blood and saliva, respectively. In blood, 1 CpG site in this region was excluded from the analyses due to heterogeneity of variances. Thus, about one-half of the CpGs in this region differed in their methylation by means of child abuse, and are associated with TFBSs.

DNA methylation of the POMC gene strengthens children's self-reports

Post hoc we hypothesized that we could replicate, on the molecular level, the group allocation that was originally based on children's self-reports. We performed unsupervised hierarchical clustering on methylation of the 26 CpG sites representing the *POMC* gene using the Euclidean distance metric and the ward clustering method in the *hclust* package in R. To account for the dispersion differences across the methylation of the CpG sites, data were z -standardized prior to cluster analysis. Both in blood and in saliva, two distinct clusters reflecting the high-exposure and low-exposure groups could be detected (Figure 3). In blood, the analysis allocated $n = 39$ (68%) children into their respective group, and in saliva $n = 35$ (60%). A chi-square test confirmed the significant concordance between the group allocation based on children's self-report and based on methylation




Figure 2. (Color online) DNA methylation of hypothalamic–pituitary–adrenal axis genes. Mean methylation of all analyzed cytosine nucleotide–phosphate–guanine nucleotide (CpG) sites. CpG sites are ordered according to their genomic location (not drawn to scale). For visual purposes, the data were mean centered. Beneath the scatterplots, the respective CpG sites and their positions in the gene model are displayed. CpG sites, which revealed at least moderate effect sizes comparing the high- and low-exposure groups are highlighted in black and bold font. *AVP*, Arginine-vasopressin gene; *CRH*, corticotropin-releasing hormone gene; *NR3C1*, nuclear receptor subfamily 3, group C, member 1 glucocorticoid receptor gene; *POMC*, proopiomelanocortin gene. ● Adjusted $p < .1$, *adjusted $p < .05$; **adjusted $p < .01$; ***adjusted $p < .001$. Black asterisks/dots depict tissue comparisons, red asterisks/dots (online only) depict comparisons in relation to child abuse in blood, and blue asterisks/dots (online only) depict comparisons in relation to child abuse in saliva.

Table 3. Analyses of variance analyzing the effect of childhood abuse on DNA methylation in CpGs associated with the *AVP*, *POMC*, *NR3C1*, and *CRH* genes

| Gene | CpG | Blood | | | | Saliva | | | |
|--------------|------------|----------|-------------|----------------|----------------|----------|-------------|----------------|----------------|
| | | <i>F</i> | η^2 | <i>p</i> | Adj. <i>p</i> | <i>F</i> | η^2 | <i>p</i> | Adj. <i>p</i> |
| <i>AVP</i> | cg03279206 | 5.82 | 0.09 | <.05 | >.10 | 0.22 | 0.00 | >.10 | >.10 |
| <i>AVP</i> | cg14065127 | 3.56 | 0.06 | <.10 | <.10 | 0.82 | 0.01 | >.10 | >.10 |
| <i>CRH</i> | cg21240762 | 0.00 | 0.00 | >.10 | >.10 | 4.19 | 0.07 | <.05 | >.10 |
| <i>CRH</i> | cg23027580 | 9.29 | 0.14 | <.01 | <.05 | 0.25 | 0.00 | >.10 | >.10 |
| <i>NR3C1</i> | cg04111177 | 3.56 | 0.06 | <.10 | >.10 | 0.23 | 0.00 | >.10 | >.10 |
| <i>NR3C1</i> | cg06521673 | 0.37 | 0.01 | >.10 | >.10 | 4.32 | 0.07 | <.05 | >.10 |
| <i>NR3C1</i> | cg07528216 | 0.03 | 0.00 | >.10 | >.10 | 5.53 | 0.09 | <.05 | >.10 |
| <i>NR3C1</i> | cg18849621 | 0.05 | 0.00 | >.10 | >.10 | 4.25 | 0.07 | <.05 | >.10 |
| <i>NR3C1</i> | cg19457823 | 1.09 | 0.02 | >.10 | >.10 | 4.54 | 0.08 | <.05 | >.10 |
| <i>NR3C1</i> | cg26464411 | 3.68 | 0.06 | <.10 | >.10 | 0.25 | 0.00 | >.10 | >.10 |
| <i>POMC</i> | cg00674304 | 4.88 | 0.08 | <.05 | >.10 | 1.30 | 0.02 | >.10 | >.10 |
| <i>POMC</i> | cg01926269 | 8.04 | 0.13 | <.01 | >.10 | 4.49 | 0.07 | <.05 | >.10 |
| <i>POMC</i> | cg02716646 | 2.38 | 0.04 | >.10 | >.10 | 3.40 | 0.06 | <.10 | >.10 |
| <i>POMC</i> | cg09916783 | NA | NA | NA | NA | 6.51 | 0.10 | <.05 | <.10 |
| <i>POMC</i> | cg11894631 | 0.32 | 0.01 | >.10 | >.10 | 7.93 | 0.13 | <.01 | <.10 |
| <i>POMC</i> | cg13025668 | 4.55 | 0.08 | <.05 | >.10 | 10.94 | 0.16 | <.01 | <.05 |
| <i>POMC</i> | cg14170547 | 4.18 | 0.07 | <.05 | >.10 | 0.71 | 0.01 | >.10 | >.10 |
| <i>POMC</i> | cg17736230 | 2.32 | 0.04 | >.10 | >.10 | 8.22 | 0.13 | <.01 | <.10 |
| <i>POMC</i> | cg20387815 | 7.23 | 0.12 | <.01 | >.10 | 5.65 | 0.09 | <.05 | >.10 |
| <i>POMC</i> | cg22900229 | 3.26 | 0.06 | <.10 | >.10 | 0.94 | 0.01 | >.01 | >.10 |
| <i>POMC</i> | cg24425171 | 4.20 | 0.07 | <.05 | >.10 | 5.28 | 0.08 | <.05 | >.10 |
| <i>POMC</i> | cg24718866 | 0.18 | 0.00 | >.10 | >.10 | 3.68 | 0.06 | <.10 | >.10 |

Note: *F*, *F* statistic for abuse; Adj. *p*, adjusted *p* value; η^2 , eta square effect size. The *p* values below .05, Adj. *p* values below .10, and effect sizes above .06 are highlighted in bold. *AVP*, Arginine-vasopressin gene; *CRH*, corticotropin-releasing hormone gene; *NR3C1*, glucocorticoid receptor gene; *POMC*, proopiomelanocortin gene. Only CpGs, which are differentially either in blood or saliva, are displayed.

value in blood ($\chi^2 = 5.95$, *df* = 1, *p* = .015) and showed a trend in saliva ($\chi^2 = 3.49$, *df* = 1, *p* = .062).

DNA methylation of HPA axis genes

We also compared DNA methylation in the four HPA axis genes between the two tissues. Generally, blood tended to show stronger signals of DNA methylated than saliva (Figure 2). The only exception was seen in *AVP*, in which the pattern was reversed and saliva was characterized by elevated DNA methylation levels compared to blood. This tendency was also revealed in the ANOVAs, as we found 3, 8, 18, and 11 CpG sites in *AVP*, *CRH*, *POMC*, and *NR3C1*, respectively, which displayed differential methylation between the tissues (online-only supplementary Table S.1).

Discussion

Child abuse is known to impair mental health across the entire life span (Carr et al., 2013). However, it has been claimed that the effects of specific forms of child abuse are not as harmful when they take place in societies or cultural groups in which such practices are common, socially accepted, and legal. Lansford et al. (2005), for example, demonstrated that the relation between corporal punishment and mental health prob-

lems varied with the perceived normativeness of corporal punishment in the respective country. However, we and others have already demonstrated the detrimental effects of child abuse in such societies (Ani & Grantham-McGregor, 1998; Hecker et al., 2014). In concordance, in the present study children with high exposure to child abuse showed decreased psychological well-being. Furthermore, we demonstrated that this link manifests itself on a molecular level that cannot be manipulated by the subject: child abuse was strongly associated with the methylation of the *POMC* gene in both blood and saliva. To date, research incorporating child abuse and the methylation of HPA axis genes has focused mainly on the *GR* gene (de Kloet et al., 2005; Labonte et al., 2014; McGowan et al., 2009; Perroud et al., 2011). Little is known about the physiological and phenotypic consequences of *POMC* methylation. The *POMC* gene is characterized by a 5' CpG islands, located at exon 1 and the promoter region, and a 3' CpG island more downstream around the intron 2 and exon 3 boundary (Gardiner-Garden & Frommer, 1994). Research investigating various disease traits or stress exposure has mainly reported differential methylation at the 5' CpG island (Mizoguchi et al., 2007; Muschler et al., 2010; Newell-Price et al., 2001; Stevens et al., 2010), but effects on the 3' CpG island (Kuehnen et al., 2012) have also been reported. In cancer tissue that did not belong to the pituitary




Figure 3. (Color online) Hierarchical clustering dendrogram. Based on the methylation of 26 cytosine nucleotide–phosphate–guanine nucleotide sites present in the proopiomelanocortin gene a hierarchical cluster analysis has been performed. Two distinct clusters were formed in both blood (a) and saliva (b) significantly replicating the two groups that are based on children’s self-reports regarding exposure to child abuse. The parts of the dendrograms highlighted in red (online only) represent the clusters containing mainly children exposed to high levels of child abuse while the turquoise highlighted segments denote the clusters containing mainly children with low exposure. The colored boxes (online only) next to the final branches denote the exposure to childhood abuse based on the self-reports (red, high exposure; turquoise, low exposure).

gland that caused Cushing syndrome (hypercortisolism), differential *POMC* methylation at the 5′ CpG island and increased ACTH levels were reported, suggesting HPA axis dysregulation, a key feature of many mental diseases (Mizoguchi et al., 2007; Newell-Price et al., 2001). Our research supports these previous findings, because the majority of differentially methylated CpGs in our study were located in the 5′ CpG island. Moreover, the respective CpGs collocate with TFBSs, suggesting transcriptional regulation. These TFBSs include an E2F response element, methylation of which has been shown to suppress *POMC* promoter activity in vitro (Newell-Price et al., 2001).

In addition to ACTH, the functionally relevant peptides β -endorphin and α -melanocyte stimulating hormone (α MSH) are cleaved from the prohormone *POMC*. Thus, the possible impairment of other systems than the HPA axis through *POMC* methylation has to be considered. β -Endorphin has antinociceptive effects that are essential for stress, in particular, the fight–flight situations. It also was reported to have rewarding properties and is considered as a factor in stress-related psychiatric disorders (Merenlender-Wagner, Dikshtein, & Yadid, 2009) and drug abuse (Roth-Deri, Green-Sadan, & Yadid, 2008). *POMC* methylation was associated with alcohol craving in patients suffering from alcohol dependence

(Muschler et al., 2010). Thus, differential *POMC* methylation by means of child abuse, as found in our study, may heighten the risk of the development of abuse-related mental illness (Carr et al., 2013), including drug abuse (Dube et al., 2003). Abuse seems to affect the methylation of the *POMC* gene and may lead to increased emotional and behavioral problems in the children, which then increase the likelihood for further abuse. In short, settings of frequent abuse would generate a vicious cycle of further abuse and behavioral problems. Because of the nature of our study, it was not possible to test this idea statistically. Future studies using larger samples and longitudinal designs should test this hypothesis empirically. Nevertheless, our findings are congruent with prior findings that child abuse is related to worse child mental health, even in a society in which specific acts of child abuse are common practice.

DNA methylation profiles appear to be tissue specific (Ollikainen et al., 2010), and several studies indicated a clear separation of samples derived from saliva and blood (Smith et al., 2014; Thompson et al., 2013; Wu et al., 2014). Accordingly, we found significantly different methylation profiles between saliva and blood. Moreover, there was a general trend of hypermethylation in saliva, which has previously been demonstrated. However, despite tissue-specific methylation, we demonstrate

that childhood abuse is associated with DNA methylation in both saliva and blood. Thus, methylation evoked by adverse experiences seems to be preserved across tissues.

Moreover, parents and caregivers often argue that children tend to overreport the exposure to abuse and the resulting harm. Thus the children's perception of their experiences is often ignored, because children are not regarded as being mature enough to accurately gauge their situation (Qvortrup, Bardy, Sgritta, & Wintersberger, 1994). Hierarchical clustering based on the methylation of *POMC*, however, allocated two-thirds of children into their respective group and a subsequent chi-square test confirmed the significant concordance between the group allocations based on children's self-report and based on methylation value. Therefore, our results strengthen the credibility of children's self-reports on a molecular level and support the conclusion that children are capable of accurately reporting their exposure to abuse. In the school context of our data assessment, we were unable to include parents' reports for logistical reasons. Furthermore, we deliberately focused on the credibility of children's reports, because their view has been often neglected in research thus far. While it is possible that the inclusion of parents' reports could have further strengthened our findings, previous studies in resource-poor countries cast doubt on the validity of parents' knowledge about their children's suffering (Elbert et al., 2009).

The methodologies employed by our study present some limitations. Our data are correlational in nature and thus cannot prove a causal relationship between child abuse and methylation patterns or decreased psychological well-being. Nevertheless, even if certain methylation patterns might increase the likelihood of child abuse, the data still confirm the credibility of children's subjective reports and with it a wealth of data showing that abused children are more likely to suffer. However, the sample size and our study design using extreme group comparisons limit the generalizability of our findings. In the school context of our data assessment, we were unable to include parents' reports for logistical reasons. Therefore, we could not gather information regarding the socioeconomic status of our sample. It remains to be tested whether socioeconomic status can impact

DNA methylation through other pathways than abuse. Furthermore, it has been suggested that probes containing single nucleotide polymorphisms (SNPs) might result in a biased signal (Price et al., 2013). Based on the 1000 Genomes Project's database (1000 Genomes Project Consortium, 2012) eight SNPs colocalize with the target sequence of probes associated with *POMC*. However, the majority of those are very rare in African populations with minor allele frequencies below 0.2% and are thus considered not relevant to our sample. Excluding one CpG, whose respective probe contained a SNP in their target sequence at higher minor allele frequency (i.e., 1.0%), did not markedly change the results (data not shown). Furthermore, this SNP was located more than 10 base pairs away from its target CpG, which does not seem to evoke biased signals (Price et al., 2013). Therefore, we consider our findings to reflect the epigenome of the participants and not as artifacts of their genotype.

In summary, we provide further evidence that in societies or cultural groups in which many specific acts of child abuse are common, legal, and socially accepted, child abuse is nevertheless detrimental for the psychological well-being of affected children. Our evidence for such a link is strengthened by the inclusion of epigenetic information from both blood and saliva. This is the first study reporting the link between child abuse and modifications of DNA methylation of *POMC*. Epigenetic modifications provide a promising mechanism through which child abuse could act to influence psychological well-being. In addition, on a molecular level, our study strengthens the credibility of children's self-reports evaluating their exposure to abuse. All in all, our findings underscore the need to inform the population at large about the adverse consequences associated with various forms of child abuse, both those societally accepted and those not. This holds especially true in societies in which such practices are commonly employed and generally regarded as effective.

Supplementary Material

To view the supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0954579415001248>.

References

- Adekan, M., Ndom, R., Ekpo, M., & Oluboka, O. (1999). Epidemiology of childhood behavioural disorders in Ilorin, Nigeria—Findings from parental reports. *West African Journal of Medicine*, 18, 29–48.
- Ani, C., & Grantham-McGregor, S. (1998). Family and personal characteristics of aggressive Nigerian boys: Differences from and similarities with Western findings. *Journal of Adolescent Health*, 23, 311–317. doi:10.1016/S1054-139X(98)00031-7
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate—A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57, 289–300.
- Carpenter, L. L., Carvalho, J. P., Tyrka, A. R., Wier, L. M., Mello, A. F., Mello, M. F., et al. (2007). Decreased adrenocorticotropic hormone and cortisol responses to stress in healthy adults reporting significant childhood maltreatment. *Biological Psychiatry*, 62, 1080–1087. doi:10.1016/j.biopsych.2007.05.002
- Carr, C. P., Martins, C. M. S., Stingel, A. M., Lemgruber, V. B., & Juruena, M. F. (2013). The role of early life stress in adult psychiatric disorders: A systematic review according to childhood trauma subtypes. *Journal of Nervous and Mental Disease*, 201, 1007–1020. doi:10.1097/NMD.0000000000000049
- Catani, C., Jacob, N., Schauer, E., Kohila, M., & Neuner, F. (2008). Family violence, war, and natural disasters: A study of the effect of extreme stress on children's mental health in Sri Lanka. *BMC Psychiatry*, 8, 33. doi:10.1186/1471-244X-8-33
- Chrousos, G., & Gold, P. (1992). The concepts of stress and stress system disorders—Overview of physical homeostasis. *Journal of the American Medical Association*, 267, 1244–1252. doi:10.1001/jama.267.9.1244
- Cortina, M. A., Sodha, A., Fazel, M., & Ramchandani, P. G. (2012). Prevalence of child mental health problems in Sub-Saharan Africa. *Archives of Pediatrics & Adolescent Medicine*, 166, 276–281. doi:10.1001/archpediatrics.2011.592

- Dammann, G., Teschler, S., Haag, T., Altmüller, F., Tuczek, F., & Dammann, R. H. (2011). Increased DNA methylation of neuropsychiatric genes occurs in borderline personality disorder. *Epigenetics*, *6*, 1454–1462. doi:10.4161/epi.6.12.18363
- de Kloet, E. R., Joëls, M., & Holsboer, F. (2005). Stress and the brain: From adaptation to disease. *Nature Reviews Neuroscience*, *6*, 463–475. doi:10.1038/nrn1683
- Du, P., Kibbe, W. A., & Lin, S. M. (2008). Lumi: A pipeline for processing Illumina microarray. *Bioinformatics*, *24*, 1547–1548. doi:10.1093/bioinformatics/btn224
- Dube, S. R., Felitti, V. J., Dong, M., Chapman, D. P., Giles, W. H., & Anda, R. F. (2003). Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: The adverse childhood experiences study. *Pediatrics*, *111*, 564–572. doi:10.1542/peds.111.3.564
- Ehrlich, S., Weiss, D., Burghardt, R., Infante-Duarte, C., Brockhaus, S., Muschler, M. A., et al. (2010). Promoter specific DNA methylation and gene expression of POMC in acutely underweight and recovered patients with anorexia nervosa. *Journal of Psychiatric Research*, *44*, 827–833. doi:10.1016/j.jpsychires.2010.01.011
- Elbert, T., Schauer, M., Schauer, E., Huschka, B., Hirsh, M., & Neuner, F. (2009). Trauma-related impairment in children—A survey in Sri Lankan provinces affected by armed conflict. *Child Abuse & Neglect*, *33*, 238–246. doi:10.1016/j.chiabu.2008.02.008
- Fang, X., Brown, D. S., Florence, C. S., & Mercy, J. A. (2012). The economic burden of child maltreatment in the United States and implications for prevention. *Child Abuse & Neglect*, *36*, 156–165. doi:10.1016/j.chiabu.2011.10.006
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*, 175–191. doi:10.3758/BF03193146
- Feinstein, S., & Mwachombela, L. (2010). Corporal punishment in Tanzania's schools. *International Review of Education*, *56*, 399–410. doi:10.1007/s11159-010-9169-5
- Fox, J., & Weisberg, S. (2011). *An {R} companion to applied regression* (2nd ed.). Thousand Oaks, CA: Sage.
- Gardiner-Garden, M., & Frommer, M. (1994). Transcripts and CpG islands associated with the pro-opiomelanocortin gene and other neurally expressed genes. *Journal of Molecular Endocrinology*, *12*, 365–382. doi:10.1677/jme.0.0120365
- Goodman, R., Ford, T., Simmons, H., Gatward, R., & Meltzer, H. (2000). Using the Strengths and Difficulties Questionnaire (SDQ) to screen for child psychiatric disorders in a community sample. *British Journal of Psychiatry*, *177*, 534–539. doi:10.1192/bjp.177.6.534
- Goodman, R., Meltzer, H., & Bailey, V. (1998). The Strengths and Difficulties Questionnaire: A pilot study on the validity of the self-report version. *European Child & Adolescent Psychiatry*, *7*, 125–130. doi:10.1007/s007870050057
- Harkness, K. L., Bagby, R. M., & Kennedy, S. H. (2012). Childhood maltreatment and differential treatment response and recurrence in adult major depressive disorder. *Journal of Consulting and Clinical Psychology*, *80*, 342–353. doi:10.1037/a0027665
- Hecker, T., Hermenau, K., Isele, D., & Elbert, T. (2014). Corporal punishment and children's externalizing problems: A cross-sectional study of Tanzanian primary school students. *Child Abuse & Neglect*, *38*, 884–892. doi:10.1016/j.chiabu.2013.11.007
- Heim, C., & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of mood and anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*, *49*, 1023–1039. doi:10.1016/S0006-3223(01)01157-X
- Heim, C., Newport, D. J., Heit, S., Graham, Y., Wilcox, M., Bonsall, R., et al. (2000). Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Journal of the American Medical Association*, *284*, 592–597. doi:10.1001/jama.284.5.592
- Hermenau, K., Eggert, I., Landolt, M. A., & Hecker, T. (2015). Neglect and perceived stigmatization impact psychological distress of orphans in Tanzania. *European Journal of Psychotraumatology*, *6*, 28617. doi:10.3402/ejpt.v6.28617
- Hermenau, K., Hecker, T., Elbert, T., & Ruf-Leuschner, M. (2014). Maltreatment and mental health in institutional care—Comparing early- and late-institutionalized children in Tanzania. *Infant Mental Health Journal*, *35*, 102–110. doi:10.1002/imhj.21440
- Hermenau, K., Hecker, T., Ruf, M., Schauer, E., Elbert, T., & Schauer, M. (2011). Childhood adversity, mental ill-health and aggressive behavior in an African orphanage: Changes in response to trauma-focused therapy and the implementation of a new instructional system. *Child and Adolescent Psychiatry and Mental Health*, *5*, 29. doi:10.1186/1753-2000-5-29
- Hompes, T., Izzi, B., Gellens, E., Morreels, M., Fieuws, S., Pexsters, A., et al. (2013). Investigating the influence of maternal cortisol and emotional state during pregnancy on the DNA methylation status of the glucocorticoid receptor gene (NR3C1) promoter region in cord blood. *Journal of Psychiatric Research*, *47*, 880–891. doi:10.1016/j.jpsychires.2013.03.009
- Isele, D., Hecker, T., Hermenau, K., Elbert, T., Ruf-Leuschner, M., Moran, J., et al. (2015). *Assessing childhood adversities: The Pediatric Maltreatment and Abuse Chronology of Exposure Interview*. Manuscript submitted for publication.
- Kaplan, S. J., Pelcovitz, D., Salzinger, S., Weiner, M., Mandel, F. S., Lesser, M. L., et al. (1998). Adolescent physical abuse: Risk for adolescent psychiatric disorders. *American Journal of Psychiatry*, *155*, 954–959. doi:10.1176/ajp.155.7.954
- Kashala, E., Elgen, I., Sommerfelt, K., & Tylleskar, T. (2005). Teacher ratings of mental health among school children in Kinshasa, Democratic Republic of Congo. *European Child & Adolescent Psychiatry*, *14*, 208–215. doi:10.1007/s00787-005-0446-y
- Kovacs, M. (2001). *Children's Depression Inventory (CDI): Technical manual*. North Tonawanda, NY: Multi Health Systems.
- Kovacs, M., Goldstein, D., & Gastonis, C. (1993). Suicidal behaviors and childhood-onset depressive disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*, *32*, 8–20. doi:10.1097/00004583-199301000-00003
- Kuehnen, P., Mischke, M., Wiegand, S., Sers, C., Horsthemke, B., Lau, S., et al. (2012). An alu element-associated hypermethylation variant of the POMC gene is associated with childhood obesity. *PLOS Genetics*, *8*, 1–12. doi:10.1371/journal.pgen.1002543
- Labonte, B., Azoulay, N., Yerko, V., Turecki, G., & Brunet, A. (2014). Epigenetic modulation of glucocorticoid receptors in posttraumatic stress disorder. *Translational Psychiatry*, *4*, e368. doi:10.1038/tp.2014.3
- Labonte, B., Yerko, V., Gross, J., Mechawar, N., Meaney, M., Szyf, M., et al. (2012). Differential glucocorticoid receptor exon 1(B), 1(C), and 1(H) expression and methylation in suicide completers with a history of childhood abuse. *Biological Psychiatry*, *72*, 41–48. doi:10.1016/j.biopsych.2012.01.034
- Lansford, J. E., Dodge, K. A., Malone, P. S., Oburu, P., Palme, K., Bacchini, D., et al. (2005). Physical discipline and children's adjustment: Cultural normativeness as a moderator. *Child Development*, *76*, 1234–1246. doi:10.1111/j.1467-8624.2005.00847.x
- Leeb, R. T., Paulozzi, L., Melanson, C., Simon, T., & Arias, I. (2008). *Child maltreatment surveillance: Uniform definitions for public health and recommended data elements*. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Retrieved from <http://www.cdc.gov/violenceprevention/pub/cmp-surveillance.html>
- Marabita, F., Almgren, M., Lindholm, M. E., Ruhmann, S., Fagerström-Billai, F., Jagodic, M., et al. (2013). An evaluation of analysis pipelines for DNA methylation profiling using the Illumina HumanMethylation450 BeadChip platform. *Epigenetics*, *8*, 333–346. doi:10.4161/epi.24008
- Mathelier, A., Zhao, X., Zhang, A. W., Parcy, F., Worsley-Hunt, R., Arenillas, D. J., et al. (2014). JASPAR 2014: An extensively expanded and updated open-access database of transcription factor binding profiles. *Nucleic Acids Research*, *42*, 1–6. doi:10.1093/nar/gkt997
- McEwen, B. S., & Lasley, E. N. (2002). *The end of stress as we know it*. Washington, DC: Joseph Henry Press.
- McGowan, P. O., Sasaki, A., D'Alessio, A. C., Dymov, S., Labonté, B., Szyf, M., et al. (2009). Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nature Neuroscience*, *12*, 342–348. doi:10.1038/nn.2270
- Merenlender-Wagner, A., Dikshtein, Y., & Yadid, G. (2009). The beta-endorphin role in stress-related psychiatric disorders. *Current Drug Targets*, *10*, 1096–1108. doi:10.2174/13894500978973514
- Mizoguchi, Y., Kajiume, T., Miyagawa, S., Okada, S., Nishi, Y., & Kobayashi, M. (2007). Steroid-dependent ACTH-produced thymic carcinoid: Regulation of POMC gene expression by cortisol via methylation of its promoter region. *Hormone Research*, *67*, 257–262. doi:10.1159/000098548
- Mueller, B. R., & Bale, T. L. (2008). Sex-specific programming of offspring emotionality after stress early in pregnancy. *Journal of Neuroscience*, *28*, 9055–9065. doi:10.1523/jneurosci.1424-08.2008
- Murgatroyd, C., Patchev, A. V., Wu, Y., Micalé, V., Bockmühl, Y., Fischer, D., et al. (2009). Dynamic DNA methylation programs persistent adverse

- effects of early-life stress. *Nature Neuroscience*, *12*, 1559–1566. doi:10.1038/nn.2436
- Muschler, M. A. N., Hillemecher, T., Kraus, C., Kornhuber, J., Bleich, S., & Frieeling, H. (2010). DNA methylation of the POMC gene promoter is associated with craving in alcohol dependence. *Journal of Neural Transmission*, *117*, 513–519. doi:10.1007/s00702-010-0378-7
- Nanni, V., Uher, R., & Danese, A. (2012). Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: A meta-analysis. *American Journal of Psychiatry*, *169*, 141–151. doi:10.1176/appi.ajp.2011.11020335
- Newell-Price, J., King, P., & Clark, A. J. (2001). The CpG island promoter of the human proopiomelanocortin gene is methylated in nonexpressing normal tissue and tumors and represses expression. *Molecular Endocrinology*, *15*, 338–348. doi:10.1210/me.15.2.338
- Ollikainen, M., Smith, K. R., Joo, E. J., Ng, H. K., Andronikos, R., Novakovic, B., et al. (2010). DNA methylation analysis of multiple tissues from newborn twins reveals both genetic and intrauterine components to variation in the human neonatal epigenome. *Human Molecular Genetics*, *19*, 4176–4188. doi:10.1093/hmg/ddq336
- 1000 Genomes Project Consortium. (2012). An integrated map of genetic variation from 1,092 human genomes. *Nature*, *491*, 56–65. doi:10.1038/nature11632
- Perroud, N., Paoloni-Giacobino, A., Prada, P., Olié, E., Salzmann, A., Nicastro, R., et al. (2011). Increased methylation of glucocorticoid receptor gene (NR3C1) in adults with a history of childhood maltreatment: A link with the severity and type of trauma. *Translational Psychiatry*, *1*, e59. doi:10.1038/tp.2011.60
- Price, E. M., Cotton, A. M., Lam, L. L., Farré, P., Emberly, E., Brown, C. J., et al. (2013). Additional annotation enhances potential for biologically-relevant analysis of the Illumina Infinium HumanMethylation450 BeadChip array. *Epigenetics & Chromatin*, *6*, 4. doi:10.1186/1756-8935-6-4
- Qvortrup, J., Bardy, M., Sgritta, G., & Wintersberger, H. (1994). *Childhood matters: Social theory, practice and politics*. Aldershot: Avebury.
- Roth-Deri, I., Green-Sadan, T., & Yadid, G. (2008). β -Endorphin and drug-induced reward and reinforcement. *Progress in Neurobiology*, *86*, 1–21. doi:10.1016/j.pneurobio.2008.06.003
- Sitarenios, G., & Kovacs, M. (1999). Use of the Children's Depression Inventory. In M. E. Maruish (Ed.), *The use of psychological testing for treatment planning and outcomes assessment* (2nd ed., pp. 267–298). Mahwah, NJ: Erlbaum.
- Smith, A. K., Kilaru, V., Klengel, T., Mercer, K. B., Bradley, B., Conneely, K. N., et al. (2014). DNA extracted from saliva for methylation studies of psychiatric traits: Evidence tissue specificity and relatedness to brain. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, *168*, 36–44. doi:10.1002/ajmg.b.32278
- Steinberg, A. M., Brymer, M. J., Decker, K. B., & Pynoos, R. S. (2004). The University of California at Los Angeles Post-Traumatic Stress Disorder Reaction Index. *Current Psychiatry Reports*, *6*, 96–100. doi:10.1007/s11920-004-0048-2
- Stevens, A., Begum, G., Cook, A., Connor, K., Rumball, C., Oliver, M., et al. (2010). Epigenetic changes in the hypothalamic proopiomelanocortin and glucocorticoid receptor genes in the ovine fetus after periconceptional undernutrition. *Endocrinology*, *151*, 3652–3664. doi:10.1210/en.2010-0094
- Sugaya, L., Hasin, D. S., Olfson, M., Lin, K., Grant, B. F., & Blanco, C. (2012). Child physical abuse and adult mental health: A national study. *Journal of Traumatic Stress*, *25*, 384–392. doi:10.1002/jts.21719
- Tanzania Daily News. (2013). *Tanzania: Public schools to continue using corporal punishment*. Retrieved from <http://allafrica.com/stories/201304090024.html>
- Teicher, M. H., & Parigger, A. (2015). The Maltreatment and Abuse Chronology of Exposure (MACE) Scale for the retrospective assessment of abuse and neglect during development. *PLOS ONE*, *10*, e0117423. doi:10.1371/journal.pone.0117423
- Teicher, M. H., & Samson, J. A. (2013). Childhood maltreatment and psychopathology: A case for ecophenotypic variants as clinically and neurobiologically distinct subtypes. *American Journal of Psychiatry*, *170*, 1114–1133. doi:10.1176/appi.ajp.2013.12070957
- Teicher, M. H., Samson, J. A., Polcari, A. M., & McGreenery, C. E. (2006). Sticks, stones, and hurtful words: Relative effects of various forms of childhood maltreatment. *American Journal of Psychiatry*, *163*, 993–1000. doi:10.1176/appi.ajp.163.6.993
- Thompson, T. M., Sharfi, D., Lee, M., Yrigollen, C. M., Naumova, O. Y., & Grigorenko, E. L. (2013). Comparison of whole-genome DNA methylation patterns in whole blood, saliva, and lymphoblastoid cell lines. *Behavior Genetics*, *43*, 168–176. doi:10.1007/s10519-012-9579-1
- Traube, D., Dukay, V., Kaaya, S., Reyes, H., & Mellins, C. (2010). Cross-cultural adaptation of the Child Depression Inventory for use in Tanzania with children affected by HIV. *Vulnerable Children and Youth Studies*, *5*, 174–187. doi:10.1080/17450121003668343
- Tyrka, A. R., Price, L. H., Marsit, C., Walters, O. C., & Carpenter, L. L. (2012). Childhood adversity and epigenetic modulation of the leukocyte glucocorticoid receptor: Preliminary findings in healthy adults. *PLOS ONE*, *7*, e30148. doi:10.1371/journal.pone.0030148
- UNICEF. (2011). *Violence against children in Tanzania: Results from a National Survey 2009*. Dar es Salaam, Tanzania: Author. Retrieved from http://www.unicef.org/media/files/violence_against_children_in_tanzania_report.pdf
- Wallis, A., & Dukay, V. (2009). Learning how to measure the well-being of OVC in a maturing HIV/AIDS crisis. *Journal of Health Care for the Poor and Underserved*, *20*, 170–184. doi:10.1353/hpu.0.0230
- Widom, C. S., DuMont, K., & Czaja, S. J. (2007). A prospective investigation of major depressive disorder and comorbidity in abused and neglected children grown up. *Archives of General Psychiatry*, *64*, 49–56. doi:10.1001/archpsyc.64.1.49
- Wu, H. C., Wang, Q., Chung, W. K., Andrusis, I. L., Daly, M. B., John, E. M., et al. (2014). Correlation of DNA methylation levels in blood and saliva DNA in young girls of the LEGACY Girls study. *Epigenetics*, *9*, 929–933. doi:10.4161/epi.28902