

# Expression and characterization of a phospholipid hydroperoxide glutathione peroxidase gene in *Schistosoma japonicum*

YING ZHANG<sup>1</sup>, YUAN HE<sup>2</sup>, LI HE<sup>2</sup>, HONGYING ZONG<sup>2</sup> and GUOBIN CAI<sup>2\*</sup>

<sup>1</sup>Department of Medical Genetics, Wuhan University School of Basic Medical Sciences, Wuhan 430071, China

<sup>2</sup>Department of Medical Parasitology, Wuhan University School of Basic Medical Sciences, Wuhan 430071, China

(Received 5 May 2015; revised 19 July 2015; accepted 22 July 2015; first published online 18 August 2015)

## SUMMARY

Phospholipid hydroperoxide glutathione peroxidase (PHGPx, GPx4) is a major antioxidant enzyme, which plays unique roles in the protection of cells against oxidative stress by catalysing reduction of lipid hydroperoxides. We isolated and characterized a full-length cDNA sequence encoding GPx gene from a blood fluke, *Schistosoma japonicum* (designated *SjGPx*), which contained an in-frame TGA codon for selenocysteine (Sec) and a concurrent Sec insertion sequence in its 3'-untranslated region. Protein encoded by *SjGPx* demonstrated a primary structure characteristic to the PHGPx family, including preservation of catalytic domains and absence of the subunit interaction domains. Semi-quantitative reverse transcription PCR and Western blotting showed that the *SjGPx* was mainly expressed in the female adults and eggs. RNA interference approach was employed to investigate the effects of knockdown of *SjGPx*. *SjGPx* expression level was significantly reduced on the 5th day post-RNAi. Significant reduction in GPx enzyme activities, as well as obvious changes in morphology of intrauterine eggs followed the reduction in *SjGPx* transcript level. We observed a 63.04% reduction in GPx activity and the eggs severely deformed. Our results revealed that *SjGPx* protein might be involved in the provision of enzyme activity during egg production.

**Key words:** *Schistosoma japonicum*, phospholipid hydroperoxide GPx, selenium-dependent GPx, RNA interference, egg production.

## INTRODUCTION

Reactive oxygen species (ROS), including superoxide anion radical (O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydroxyl radical (OH<sup>•</sup>), singlet oxygen (<sup>1</sup>O<sub>2</sub>) and lipid hydroperoxides are produced by incomplete reduction of oxygen during aerobic respiration and cellular metabolism. ROS play pivotal roles not only in control of cellular redox homeostasis but also in induction of cellular proliferation or apoptosis by activating signalling molecules (Hancock *et al.* 2001). However, the unbalanced generation of ROS may cause chemical damage to the cells, which might induce breakage of DNA strands, protein oxidation, polysaccharide depolymerization, membrane-lipid peroxidation and impairment of signal transduction from membrane receptors in various physiological processes (Dröge, 2002). The aerobic organisms have evolved a series of multi-layered enzymatic and non-enzymatic defence systems, which can remove the aggressive ROS and/or repair ROS-mediated cell damage (Sies, 1993). Distinct antioxidant enzymatic activities such as catalase, glutathione peroxidase (GPx) and peroxiredoxin (PRx) have been well characterized from numerous taxa. Trematode parasites

are known to lack catalase, suggesting that GPx proteins play important antioxidant roles alone or together with other proteins (Mkoji *et al.* 1988).

GPx encompasses six distinct families of multiple isoenzymes, including classical/cytosolic/cellular GPx (GPx1/c-GPx), gastrointestinal GPx (GPx2/GPx-GI), plasma GPx (GPx3/p-GPx), phospholipid hydroperoxide GPx (GPx4/PHGPx; E.C. 1.11.1.12), epididymis-specific GPx (GPx5/e-GPx) and odourant metabolizing GPx (GPx6/o-GPx), which catalyse the reduction of H<sub>2</sub>O<sub>2</sub>, organic hydroperoxides and lipid hydroperoxides using glutathione as a reducing agent. These GPx families share similar structural and enzymatic properties with one another (Arthur, 2000). PHGPx proteins have many distinct features compared to the other GPx families, which perform functions in a monomeric form because they lack the subunit interaction domains (Epp *et al.* 1983; Brigelius-Flohé *et al.* 1994; Arthur, 2000). These PHGPx molecules directly reduce hydroperoxidized phospholipids integrated into membranes (Ursini and Bindoli, 1987). Members of the other GPx families, however, interfere with lipid peroxidation only via a concerted operation with phospholipase, which implies that PHGPxs are deeply associated with the repair of disrupted biomembranes (Imai and Nakagawa, 2003). GPxs might also constitute the front line of enzymatic defence to ensure their survival against host

\* Corresponding author. Department of Medical Parasitology, Wuhan University School of Basic Medical Sciences, Wuhan 430071, China. E-mail: [gbcail2011@whu.edu.cn](mailto:gbcail2011@whu.edu.cn)

immune cell-derived ROS in parasitic helminths, which cause chronic infections (Zelck and von Janowsky, 2004). The parasitic helminth GPx families show a certain degree of biased distribution across taxa. GPx proteins homologous to mammalian GPx3 have been characterized in the filarial nematodes, including *Brugia pahangi*, *Dirofilaria immitis* and *Wuchereria bancrofti* (Henkle-Dührsen and Kampkötter, 2001); whereas information on the PHGPx (GPx4) members is obtainable from trematodes such as *Schistosoma mansoni* (Mei and LoVerde, 1995; Mei *et al.* 1996), *Clonorchis sinensis* (Cai *et al.* 2008) and *Paragonimus westermani* (Kim *et al.* 2009). Observations regarding the tissue distribution of the *S. mansoni*, *C. sinensis* and *P. westermani* PHGPxs were shown to be largely localized in vitellocytes within vitelline follicles and eggs (Roche *et al.* 1996; Cai *et al.* 2008; Kim *et al.* 2009).

RNA interference (RNAi), which results in post-transcriptional gene silencing in a wide variety of organisms, was originally identified in *Caenorhabditis elegans* (Tabara *et al.* 1998), and has since been used in human parasitic worms, including schistosome (Mourão *et al.* 2009; Cheng *et al.* 2009; Tchoubrieva *et al.* 2010). In eukaryotic cells, double-stranded RNA (dsRNA) can be recognized by the RNA-induced silencing complex which effects degradation of the target mRNA, and abolishes expression of the corresponding protein (Zamore *et al.* 2000). dsRNAs, which can be synthesized in large quantities and transfected into target cells in the presence of lipofectamine reagents, are the most commonly used reagents for RNAi in cultured parasites (Morales *et al.* 2008).

*Schistosoma japonicum* is a major schistosome species in Asia, infecting not only humans but also wild or domestic animals. It is estimated that nearly one million people in China are presently afflicted with schistosomiasis, and 30 million are at risk of infection with *S. japonicum* (McManus *et al.* 2004). Adult worms of *S. japonicum* settle in the mesenteric veins via the hepatic portal system. Most eggs lodge in the fine venules of intestine wall; others are carried retrograde by the portal venous system to the liver, in which they become trapped in the portal triad. The most prominent pathologic consequences of schistosomiasis are caused by inflammatory responses around the eggs which become trapped in the liver and other tissues of the host (Wyler, 1992). The parasite might be continuously exposed to oxidative stresses, from both ROS generated by endogenous metabolism and ROS generated by host inflammatory and immune cells (Selkirk *et al.* 1998; Zelck and von Janowsky, 2004). In order to overcome these stressful conditions and to maintain its lifespan, the fluke is equipped with the antioxidant enzymes, such as peroxiredoxin (Kumagai *et al.* 2009).

In this study, we isolated and characterized a novel PHGPx gene (*SjGPx*) from the blood fluke *S. japonicum*. Molecular characters of the gene were described, including selenocysteine (Sec) insertion sequence (SECIS) element, amino acid (aa) primary structure and phylogenetic position among various members of GPx families. The native *S. japonicum* glutathione peroxidase (*SjGPx*) was analysed by semi-quantitative reverse transcription PCR, Western blotting and enzyme relative activity assay. Finally, we employed a lipofectamine transfection procedure to introduce *SjGPx* dsRNA into cultured schistosome, and investigated the inhibitory effect of dsRNA-mediated gene knock-down of *SjGPx* on intrauterine egg formation in *S. japonicum*.

## MATERIALS AND METHODS

### Parasite

Positive *Oncomelania hupensis* snails were collected from the epidemic area of schistosomiasis in Hunan Province, People's Republic of China. *Schistosoma japonicum* (Chinese mainland strain) was maintained in *O. hupensis*. The 10-week-old specific pathogen free (SPF) female BALB/c mice, which were obtained from the Department of Animal, Hubei Academy of Medical Science, China, were infected percutaneously through shaven abdomen with 20 cercariae of *S. japonicum* by an adaptation of the ring method. The mice were sacrificed on the 42th day after infection, male and female adult worms were recovered by perfusion of the hepatic portal veins, and meantime, the eggs of *S. japonicum* were collected from the livers of infected mice (He *et al.* 2012). Then, all samples were washed repeatedly to remove host cells and debris with physiological saline. Eventually, some samples were either frozen and kept in liquid nitrogen or immediately used for nucleic acid extraction. Some adult worms collected for RNAi *in vitro* experiments were washed with sterile 0.9% NaCl solution containing 1000 U mL<sup>-1</sup> penicillin and 1 mg mL<sup>-1</sup> streptomycin and then cultured in the RPMI 1640 medium. The use of animals was approved by the Animal Ethics Committee of Wuhan University.

### Isolation of *S. japonicum* gene encoding GPx homologues

The non-redundant GenBank database at National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov>) was screened with the nucleotide sequence of *S. mansoni* GPx (accession no. AY729668) used as query in the BLAST searches. A clone from *S. japonicum* database (accession no. CV697435) encoding GPx protein was

selected based on BLAST algorithms. A cDNA library of *S. japonicum* was constructed using SMART<sup>®</sup> cDNA Library Construction Kit (Clontech, CA, USA) under the manufacturer's instruction. The *S. japonicum* cDNA library was screened by standard plaque lift hybridization using the DNA fragment as a probe. The insert cDNA from positive clone was amplified by PCR using universal T3 and T7 promoter primers, cloned into pMD18-T vector (Takara, Shiga, Japan) and subjected to nucleotide sequencing from both strands to obtain a full-length cDNA. The clone CV697435 was designated *SjGPx*. The coding profile and deduced aa sequence of the *SjGPx* gene was determined using the Open Reading Frame (ORF) Finder program at NCBI (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). The SECIS was scanned using the SECISearch program (ver2.19; <http://genome.unl.edu/SECISearch.html>). The Compute pI/Mw program ([http://www.expasy.ch/tools/pi\\_tool.html](http://www.expasy.ch/tools/pi_tool.html)) was used for the calculation of the theoretical molecular weight (Mr) and isoelectric point (pI) value. The putative N-terminal hydrophobic signal peptide was predicted using the SignalP program (<http://www.cbs.dtu.dk/services/SignalP>).

#### *The aa sequence alignment and phylogenetic analysis*

The translated aa sequence of *SjGPx* was used as query in a series of BLAST searches, to retrieve the closely matched sequences from a variety of GenBank genomic databases. A total of 42 sequences were retrieved, according to their homology values and the taxonomical distributions of their donors. The aa sequences were aligned using ClustalX and optimized with GeneDoc (<http://www.psc.edu./biomed/genedoc>) programs, respectively. A phylogenetic analysis was conducted by the neighbour-joining (NJ) or maximum parsimony algorithm integrated into the PHYLIP package (ver3.6b). Alignment gaps were removed as missing data and the trees were displayed with TreeView. The statistical significance of each branching point was evaluated with 100 random samplings of the initial input using SEQBOOT software.

#### *Semi-quantitative reverse transcription PCR (RT-PCR)*

Forty-second-day live adult worms were chilled on ice for several minutes to facilitate separation of mated pairs. Total RNAs were extracted from eggs, fresh cercariae, male and female adults using Trizol reagents (Gibco, Carlsbad, CA). The RNA samples were treated with RNase-free DNase (GIBCO BRL, Rockville, MD, USA). Poly(A)<sup>+</sup> RNAs were prepared from the total RNAs by oligo (dT)-affinity chromatography (Qiagen, Valencia,

CA). The first strand cDNAs were synthesized from 1 µg of poly(A)<sup>+</sup> RNA and oligo-dT primers using a RNA PCR Kit (AMV) Ver. 3.0 (TaKaRa, Shiga, Japan) under the manufacturer's instruction. The semi-quantitative RT-PCR primers for *SjGPx* were designed based on the partial sequence. The primer sequences were *SjGPx*-F, 5'-GAT CACTGGAAGTTCCGCAATG-3' and *SjGPx*-R, 5'-CTTCTGTTTCAATAGCTCCATG-3'. The gene transcript was amplified by PCR with the specific primers using the cDNA as a template. The PCR was conducted with a thermal cycling profile of 94 °C for 4 min, 25 cycles of 50 s at 94 °C, 40 s at 59 °C, 1 min at 72 °C and a final extension of 10 min at 72 °C. The reaction product was resolved by 2% agarose gel electrophoresis and visualized by ethidium bromide staining. Reaction with a primer pair for the *S. japonicum alpha-tubulin* gene (GenBank accession no. AY815746, *SjTubulin*-F, 5'-CTACTGTAGTGGATGAAGTGCGAAC-3' and *SjTubulin*-R, 5'-CAGCTGAAATTACTGGTGCA TAAG-3') was utilized as an internal quantity control, and the absence of any contaminating chromosomal DNA was verified via the preparation of reactions without reverse transcriptase during the first round of cDNA synthesis.

#### *Cloning and expression of recombinant SjGPx protein*

The gene segments spanning the ORF region of *SjGPx* was amplified by PCR using the first strand cDNA as templates, as described above, and then cloned into pGEM T-Easy vector (Promega). The unusual Sec codon (TGA) of *SjGPx* was converted into a sense Cys codon (TGC), using a complementary primer pair 5'-CTTGCAAGTGCGGC GCAACGGACAAAATTATC-3' and 5'-GAT AATTTTTGTCCGTTGCGCCGCACTTGCA AG-3', and the QuikChange Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA, USA). The mutated plasmid was transformed into competent *Escherichia coli* DH5α cells and the nucleotide sequence was verified by sequencing. Nucleotides corresponding to the mature domain of *SjGPx* were amplified from the plasmid construct using primers containing restriction enzyme sites for *EcoRI* and *Xho I* (5'-CCGAATTCGCCAAC TTCAGAAGATGCAC-3' and 5'-GGCTCGA GTCACCTCTGTTTCAATAGCTCC-3'). After purification with a QIAquick PCR Purification Kit (Qiagen, Valencia, CA, USA) and digestion with the corresponding enzymes, the DNA was cloned into the pET-28a-c (+) vector (Novagen, Madison, WI, USA). The plasmid was then transformed into *E. coli* DH5α cells and the accuracy of the nucleotide sequence was verified by sequencing. The plasmid DNA exhibiting the correct codons was introduced into *E. coli* BL21 (DE3) cells. Expression of the recombinant protein was

induced by treating the transformants with 0.5 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG). The bacteria cells were sonicated and the recombinant proteins were purified by Ni-NTA agarose chromatography (Amersham Biosciences). The purified recombinant SjGPx (rSjGPx) protein was analysed by SDS-12% PAGE and used to generate specific antibody.

#### *Generation of polyclonal antibody and Western blotting*

Six-week-old, SPF female BALB/c mice were immunized subcutaneously 3 times (2-week interval) with the purified rSjGPx protein (30  $\mu$ g each) mixed with Freund's adjuvant (Sigma). The mice finally received intravenously boosters with the proteins (10  $\mu$ g). The mice were sacrificed 7 days after the final booster and blood was collected by heart puncture. The blood was centrifuged for 10 min at 3000 g at 4 °C and the antisera were stored at -70 °C until use.

The egg, cercariae and male and female adult worms of the parasite were ground with a Teflon pestle-homogenizer in phosphate-buffered saline (PBS, 140 mM NaCl, 2.7 mM KCl, 10 mM Na<sub>2</sub>HPO<sub>4</sub> and 1.8 mM KH<sub>2</sub>PO<sub>4</sub>; pH 7.2) containing 2% SDS and a protease inhibitor cocktail (Roche, Mannheim, Germany). After centrifugation at 15 000 rpm for 30 min at 4 °C, the supernatants were recovered. SDS-PAGE was conducted using 12% gels with the native proteins and recombinant proteins under reducing condition. The gels were stained with Coomassie Brilliant Blue or transferred onto nitrocellulose membrane (Schleicher & Schuell BioScience, Dassel, Germany). The membrane was blocked for 1 h in Tris-buffered saline containing 0.05% (v/v) Tween-20 (TBS/T) and 5% (w/v) skim milk, followed by overnight incubation with the specific antibody diluted to 1:1000 in TBS/T containing 5% skim milk at 4 °C. The membrane was subsequently incubated for additional 1 h at room temperature with horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG antibody (1:2000 dilutions; Cappel, West Chester, PA, USA). Signal was detected with an ECL system (Amersham Biosciences) under the manufacturer's instructions. An anti-SjTubulin antibody, which was produced similarly to the generation of anti-SjGPx antibody using its specific sequence, was used as an internal standard.

#### *Enzyme activities of the native GPx of S. japonicum adult worm*

Live adult worms were washed over than 10 times with cold physiological saline at 4 °C, after which homogenized with a Teflon-pestle homogenizer in PBS containing 5% Triton X-100. The homogenates

were centrifuged at 15 000 rpm for 30 min at 4 °C and the supernatants containing GPx activities were stored. The total SjGPx activity was estimated using a Glutathione Peroxidase Assay Kit (Northwest Life Science Specialities, LLC, Canada) according to the manufacturer's instructions. The measurements were conducted with triplicate reactions and expressed as Mean  $\pm$  S.D. of enzymatic unit mL<sup>-1</sup> mg protein<sup>-1</sup>.

#### *Synthesis of dsRNA of SjGPx*

dsRNA was synthesized from cDNA encoding the full length of *SjGPx* using gene-targeted primers containing T7 promoter sequences 5'-TAA TACGACTCACTATAGGGATGATCACTGG AAGTTCCGCAA-3' and 5'-TAATACGACT CACTATAGGGTCACTTCTGTTTCTTTAG CTC-3' spanning coding DNA positions of 33–563. One negative control (NC) dsRNA against Firefly Luciferase was also synthesized from plasmid pGL3-Basic (Promega, Madison, WI). RNAs were synthesized and purified using the Megascript RNAi kit (Ambion, Austin, TX) according to the manufacturer's instructions. dsRNAs were precipitated with 5 M ammonium acetate and 95% ethanol, after which the RNA pellets were dissolved in water. Integrity of the dsRNAs was verified by non-denaturing 1% agarose gel electrophoresis, and concentration and purity determined with a spectrophotometer (ND-1000, NanoDrop Technologies, Wilmington, DE).

#### *Treatment of Schistosomes with dsRNA in vitro culture*

The Lipofectamine<sup>TM</sup> 2000 transfection reagent (Invitrogen, USA) was used in the dsRNA selection experiment to improve transfection efficiency. 800  $\mu$ L of OPTI-MEM (Invitrogen, USA) containing 8  $\mu$ L of Lipofectamine<sup>TM</sup> 2000 and dsRNA was added into 3.2 mL of antibiotic-free RPMI 1640 medium, in which 50 mated worms were cultured. Then the worms were cultured for 6 h at 37 °C before media were replaced with fresh RPMI 1640 medium containing 10% (v/v) fetal bovine serum and antibiotics (He *et al.* 2012).

To examine the optimal concentration of dsRNA in mediating *SjGPx* knockdown, 10 and 50 nM of the dsRNAs were added into the culture. The RNA against Firefly Luciferase was as NC group. After 5 days of cultivation, the parasites and eggs from culture medium were collected. Parasites were used for determination of *SjGPx* transcript level using semi-quantitative RT-PCR. The protein levels of SjGPx were determined by Western blotting and GPx activity assays. The morphological characteristics of intrauterine eggs and collected eggs from the culture medium were

observed using a light microscope (Olympus, CKX41SF, Japan) from the 1st to 5th day.

## RESULTS

### *Isolation of a selenium-dependent GPx gene from S. japonicum*

A clone (accession no. CV697435) from *S. japonicum* database was selected, which exhibited a significant degree of sequence identity to PHGPx isolated from various taxa including *S. mansoni* (AAU34080 and AAA29885), *C. sinensis* (ABK58679-ABK58682), *P. westermani* (ABE68811 and ABE68812), *Homo sapiens* (CAA50793), *Mus musculus* (BAA22780) and *Drosophila melanogaster* (AAR96123) ( $E$ -values  $< 7 \times 10^{-32}$ ) during the BLASTX analysis. The complete cDNA sequence was determined by consecutive cDNA library screening using the clone as probe. The cDNA, designated *SjGPx*, was comprised of 724 bp, and contained 531 bp ORFs, from an ATG at nucleotide position 33–35 to a TGA at positions 561–563. In addition to the conventional stop codon, the *SjGPx* gene contained a second in-frame stop codon within its ORF at nucleotide position 180–182, which might decode the 21st aa, Sec (Stadtman, 1996). The SECIS predicted within the 3'-untranslated regions (UTR) also suggested that the opal TGA codon encodes Sec (Fig. 1A). The SECIS conserved secondary structures characteristic to those of selenium-dependent homologues: two helices separated by an internal loop, a quartet of non-Watson–Crick base pairs located at the base of helix II and an apical loop (Kryukov *et al.* 2003). Nucleotide signatures in the quarter were found to be ATGA\_AA\_GA in other trematode genes. *SjGPx* harboured SECIS with a different sequence conservation GTGA\_AA\_GA, in that the adenosine preceding the quarter is replaced with guanosine (Fig. 1B). The full-length cDNA sequence of *SjGPx* was registered in GenBank under accession no. KJ941008.

### *Primary structure of SjGPx and its phylogenetic position within the GPx family*

The decoded polypeptide of *SjGPx* was found to be composed of 176 amino acids with predicted Mw/pI of 19.7 kDa/9.08, and shared a significant degree of sequence identity with other trematode GPx (39–75%, Fig. 2). *SjGPx* contained three conserved functional domains found in the GPx family and the aa residues constituting the catalytic triad (Sec, Gln and Trp) were positioned separately within each of the domains (Epp *et al.* 1983). The *SjGPx* protein did not harbour the subunit interaction domains, which are required for the formation of an active tetramer in GPx isozymes other than PHGPxs (Fig. 2, Brigelius-Flohé *et al.* 1994). Amino acid

sequence of *SjGPx* was registered in GenBank under accession no. AJD79086.

The aa sequence of *SjGPx* was used in homology searches with BLAST algorithms from the GenBank database, and several hundred PHGPx-like proteins were retrieved. These sequences revealed various degrees of sequence identity to *SjGPx* ranging 35–58% ( $E$ -values  $< 1 \times 10^{-20}$ ). Information regarding the other GPx families was obtained by a series of subsequent BLAST searches using the aa sequence of human GPx3 (AAP50261). A total of 42 entries were finally selected, by considering both redundancy and taxonomical distributions and used in a phylogenetic analysis. As shown in Fig. 3, six classes of the GPx family were well separated in the phylogenetic tree constructed by the NJ method. A similar clustering pattern was also observed in a tree based on the maximum parsimony algorithm, of which bootstrap values were incorporated into the NJ tree. The members of PHGPx had a highly diverse donor source and the GPxs identified from free-living nematodes, insects and plants, as well as trematodes and vertebrates, were categorized into this group, including *SjGPx* protein. The Sec codon and the associated SECIS motif were recognized exclusively within the mRNA sequences coding for the trematode and vertebrate GPx genes (indicated by † and ‡, respectively, in Fig. 3). Among the arthropod proteins examined, only an arachnid tick PHGPx (*Boophilus microplus*, ABA25916) was expected to be Sec-dependent.

### *Expression patterns of SjGPx and the native GPx activity in S. japonicum*

Expression level of the *SjGPx* mRNA was examined by semi-quantitative RT–PCR using RNAs isolated from *S. japonicum* egg, cercariae, male and female adult worms. As shown in Fig. 4A, the amplification of the target gene clearly revealed that the *SjGPx* gene was strictly transcribed in the egg and mature female worm. The gene of cercariae and adult male worm showed a very low transcription levels. The *Sj-Tubulin*, used as an internal control, exhibited a constant expression level during the stages (Fig. 4A).

To observe their expression proteins of *SjGPx*, we conducted Western blotting, employing the egg, cercariae, male and female adult worm extracts, using anti-*SjGPx* specific mouse sera (Fig. 4B). The cercariae proteins were not found to react with the *SjGPx* antibody. However, the blot containing the female and egg whole proteins, which had been extracted in the presence of 2% SDS, exhibited strong positive reactions. The results were well matched to the mRNA expression levels. As shown in Fig. 4B, a strong band (migrated approximately 20 kDa), which might be well matched with the size of the *SjGPx* protein (the predicted theoretical




Fig. 1. Analysis of SECIS elements in *SjGPx* of *S. japonicum*. (A) SECIS elements in *SjGPx*. The 3'-UTR of *SjGPx* mRNA was predicted to have a secondary structure characteristic to SECIS, including an apical loop, an internal loop, a quartet (SECIS core) and two helices (SECISearch program, ver2.19). The common structure of SECIS, as predicted using the bovine *GPx* gene (XP\_612375), is shown for comparison (leftmost panel). The nucleotide positions are numbered at both start and end points of the elements; (B) comparison of the primary SECIS structures. Nucleotide sequences constituting the SECIS motif were retrieved from the *PHGPx* genes of *C. sinensis*, *P. westermani*, *S. mansoni*, *H. sapiens* and *M. musculus*, and aligned in accordance with the predicted secondary elements. Nucleotides conserved within the invariant quartet and apical loop are highlighted in boldface.

Mr 19.7 kDa by expasy program), was determined against *SjGPx* antisera. The *S. japonicum* alpha-tubulin protein (*SjTubulin*), used as an internal control, exhibited a constant expression level during the stages. As controls, the antibodies were strongly reactive with the recombinant *SjGPx/SjTubulin* proteins.

The relative enzymatic activity of *GPx* proteins in *S. japonicum* whole adult worm extracts was  $2.76 \pm 0.12 \text{ U mL}^{-1} \text{ mg}^{-1}$  (3 replicates) using a specific Glutathione Peroxidase Assay Kit (Fig. 4C). The specific enzymatic activity was calculated using the equation,

$$[\text{GPx}] = \{2(\text{mRate}_s - \text{mRate}_b)V_{\text{Rxm}}/2 \cdot 74V_s\} \text{ df},$$

in which  $\text{mRate}_s = 1000 \Delta\text{OD}_{340} \text{ min}^{-1}$  of sample,  $\text{mRate}_b = 1000 \Delta\text{OD}_{340} \text{ min}^{-1}$  of blank,  $V_{\text{Rxm}}$  = volume of reaction mixture,  $V_s$  = volume of sample, and  $\text{df}$  = sample dilution factor.

#### RNAi-mediated knockdown of *SjGPx* and morphological observation of intrauterine eggs

Effects of RNAi-mediated knockdown on *SjGPx* were examined using semi-quantitative RT-PCR, Western blotting and *GPx* activity assays. As shown in Fig. 5, the effect of dsRNA was dose-dependent. *SjGPx* dsRNA at concentration of 50 nM was significantly reduced the transcript level of the target gene (Fig. 5A). The reduced level of

*SjGPx* protein caused by dsRNA in schistosomes paralleled reduction in mRNA level (Fig. 5B). The whole *GPx* enzyme activities were reduced approximately 29.35 and 63.04% after adding 10 and 50 nM dsRNA, respectively, as shown in Fig. 5C.

From the 1st to 5th day after *SjGPx* gene knockdown, the morphology of intrauterine eggs was continuously observed under a light microscope. On the 1st to 2nd day, the intrauterine eggs maintained a normal shape. On the 3rd day post-treatment, intrauterine eggs resembling spindles began to appear in the *SjGPx* gene knockdown group (data not shown), and on the 5th day, the eggs from intrauterine of females and eggs from culture medium were more severely deformed. Many irregular eggs, including spindle-like eggs and eggs with invaginated eggshells adhering to each other were observed in the female uteri (Fig. 5D). The majority of eggs from the untreated group and NC group were morphologically normal.

#### DISCUSSION

Among six distinct *GPx* families, *PHGPx* (*GPx4*) is a major antioxidant enzyme, which plays unique roles in the repairing of disrupted biomembranes by directly reducing hydroperoxidized phospholipids integrated into membranes (Ursini and Bindoli, 1987; Imai and Nakagawa, 2003). *PHGPx* proteins have been isolated from most eukaryotes,




Fig. 2. Primary structure of the PHGPx protein of *S. japonicum*. The deduced aa sequence of SjGPx was aligned with those of other trematode PHGPxs and vertebrate 6 GPx members retrieved from the GenBank database, using the ClustalX program, then optimized with GeneDoc. Identical aa in the alignment are highlighted in black, while similar residues are shown in grey. Boxes A, B and C represent three highly conserved domains in GPx families. Amino acid residues forming the catalytic triad are indicated by solid arrowheads. The Sec residues in box A are denoted by ‘U’ or ‘X’. Dotted boxes I and II mark subunit-binding domains found in the multiple GPx members, rather than in PHGPxs. The abbreviated names of six GPx families are provided at the head of the alignment. The identity value of each enzyme against SjGPx, which was calculated from the alignment after removing positions with a gap as missing data, is provided at the end of the alignment. The proteins used in this analysis were selected from various trematode organisms as follows: *S. mansoni*, SmGPx1 (AAA29885) and SmGPx2 (AAU34080); *C. sinensis*, CsGPx1–4 (ABK58679–ABK58682); *P. westermani*, PwGPx1 (ABE68811) and PwGPx2 (ABE68812). The vertebrate 6 GPx families proteins were selected from *H. sapiens* (Hs) and *M. musculus* (Mm): HmGPx4 and MmGPx4 (PHGPx, CAA50793 and BAA22780); HmGPx1 and MmGPx1 (c-GPx, CAA68491 and AAH86649); HmGPx2 and MmGPx2 (GPx-GI, CAA48394 and AAH54848); HmGPx3 and MmGPx3 (p-GPx, AAP50261 and AAH37027); HmGPx5 and MmGPx5 (e-GPx, NP\_001500 and NP\_034473); HmGPx6 and MmGPx6 (o-GPx, NP\_874360 and AAH13526).

including vertebrates, insects, nematodes, trematodes and plants, whereas the other GPx isoenzymes show a more restricted taxonomical distribution. In addition to vertebrates, a number of filarial nematodes are known to express GPx3-like enzymes (Henkle-Dührsen and Kampkötter, 2001). Among the reported parasitic trematodes, two GPx proteins in *S. mansoni* (Williams *et al.* 1992; Mei and LoVerde, 1995), four GPx proteins in *C. sinensis* (Cai *et al.* 2008) and two GPx proteins in *P. westermani* (Kim *et al.* 2009), showing strong phylogenetic affinities towards the vertebrate PHGPx, have been described. In the present study, we isolated and characterized a novel PHGPx gene from a blood fluke *S. japonicum*. The deduced aa sequence of the gene also revealed the primary structure characteristic to the PHGPx (GPx4) family, including aa conservation, absence of the subunit interaction domains and well-preserved catalytic and glutathione-binding domains (Fig. 2).

In a phylogenetic analysis, SjGPx protein also showed tight relationships with the other PHGPx-related members (Fig. 3).

Many GPx proteins have been known to contain an opal UGA codon, a stop signal in the standard genetic code, encodes Sec at their catalytic site. The alternative decoding of UGA generally depends on a *cis*-factor, called SECIS, which is located in the 3'-UTR of these selenoprotein genes (Stadtman, 1996). A biased distribution of Sec-dependent PHGPxs was detected along with the taxonomical positions of their donor species (Fig. 3). The Sec codon and concurrent SECIS motif were exclusively identified within the mRNA sequences of trematode and vertebrate GPx genes, except for one sequence identified from an arachnid (ABA25916). Among the trematode GPx enzymes obtainable, only *P. westermani* GPx1 (ABE68811) was found to lose Sec-related sequence factors and was determined to be a selenium-independent




Fig. 3. Phylogenetic relationships of SjGPx among various members of GPx families. A phylogenetic analysis was conducted based on a multiple alignment of aa sequences retrieved from the GenBank database. The tree was constructed by a maximum parsimony algorithm using PHYLIP after excluding positions with gaps and was unrooted. Numbers at branching nodes indicate their percentages of appearance in 100 bootstrap replicates and values of >50% are marked. The branches separating each of the GPx families are presented with thick bars. The presences of Sec codon (+) and SECIS (‡) within each of the corresponding mRNA sequences are indicated at the end of each sequence names. The identity of each analysed sequence is distinguished by a GenBank accession number followed by the species name, from which it has been isolated.




Fig. 4. Identification of the native SjGPx in *S. japonicum*. (A) Transcription profile of *SjGPx* gene in *S. japonicum* parasite. The transcription level of *SjGPx* was determined with the gene-specific primers via a semi-quantitative RT-PCR. Total RNAs were extracted from egg, cercariae, male and female adult stages of the parasite, and used as templates to examine mRNA transcripts of *SjGPx* gene. The amplified products were separated on 2% agarose gels and stained with ethidium bromide. The *Sj-Tubulin* (GenBank accession no. AY815746) gene was also amplified using the specific primers as an internal control. The absence of any contaminated genomic DNA was confirmed by preparing reactions without reverse transcriptase during the first reverse transcription step (RT-). The position of 100-bp DNA size standards (M) was shown on the left; (B) Western blotting of the native SjGPx protein. A total of 30 µg of proteins extracted from male adult, female adult, cercariae and egg were separated on a SDS-12% PAGE gels under reducing conditions and transferred onto nitrocellulose membranes. The membranes were immuno-reacted against the recombinant SjGPx-specific mouse antisera. The rSjGPx (50 ng) was employed as a positive control. Molecular weight marker (*Mr*) in kDa was shown on the left. An antibody specific to the SjTubulin was used as an internal standard. Rec, recombinant SjGPx/SjTubulin protein; (C) detection of GPx activities in *S. japonicum*. The specific activity was assayed using a specific Glutathione Peroxidase Assay Kit as described in the section ‘Materials and methods’, and the value of GPx activity was calculated using an equation as described in the section ‘Results’. The measurements were conducted with triplicate reactions employing the worm extracts (5, 10 and 20 µg of each), and the result from one reaction is presented as a representative.






Fig. 5. (colour online). Inhibitory effects of dsRNA-mediated gene knockdown of *SjGPx*. 0, 10 and 50 nM of the dsRNAs were added into the schistosomes culture. After 5 days of RNAi, mated worms were collected for determination of (A) *SjGPx* transcript level using semi-quantitative RT-PCR with the gene-specific primers; (B) *SjGPx* protein level using Western blotting with the *SjGPx*-specific mouse antisera; (C) the GPx activity using a Glutathione Peroxidase Assay Kit, the assays were conducted in triplicate employing the worm extracts (10  $\mu$ g of each) and expressed as mean  $\pm$  s.d. \* $P$  < 0.01 by Student's *t* test; and (D) RNAi effects of 50 nM *SjGPx* dsRNA on intrauterine eggs formation in *S. japonicum* female adult worms cultured *in vitro* for 5 days. Intrauterine eggs within female adult worms and laid eggs from the culture medium in the untreated group, NC group (the RNA against Firefly Luciferase from pGL3-Basic plasmid) and *SjGPx* group were observed using a light microscope under the common light (scale bars were showed on the left side of each panel).

PHGPx (Fig. 2). Our results suggested that *SjGPx* gene is selenium-dependent PHGPx, which contain Sec codon and SECIS motif within its mRNA sequence (Fig. 1).

Helminth parasites may be continuously exposed to dual oxidative stresses, from both ROS generated by endogenous intracellular metabolism and ROS generated by host inflammatory and immune cells (Selkirk *et al.* 1998; Zelck and von Janowsky, 2004). During their lifespan in host tissues, removal of ROS and protection of macromolecules from the oxidative attacks would be inevitable. The antioxidant enzymes might also constitute a major defensive system in these causative parasites (Callahan *et al.* 1988). The adults of *S. japonicum* settle in the mesenteric veins via the hepatic portal system, and their eggs lodge in the fine venules of intestine wall and are carried retrograde by the portal venous system to the liver, where they become trapped in the portal triad. The parasite might be continuously exposed to oxidative stresses. In order to overcome these stressful conditions and to maintain its life-span, the parasite is equipped with the antioxidant enzymes. Furthermore, phenol oxidase, which is secreted from the vitellocytes, acts on eggshell precursors, thus converting tyrosine into quinone in the presence of oxygen. During this tanning process, incomplete reduction of the oxygen may result in a local increase of harmful ROS (Cordingley, 1987; Cai *et al.* 2009; Bae *et al.* 2015).

Therefore, effective antioxidant enzymes, such as GPx, might be required for the protection of macromolecules and/or the fertilized ovum. Among the reported trematode GPx enzymes, we noticed *S. mansoni* GPx proteins were shown to be localized on vitelline follicles and eggs (Roche *et al.* 1996). The PHGPx-like proteins from *C. sinensis* and *P. westermani* parasites were also found localized in vitellocytes within vitelline follicles and premature eggs (Cai *et al.* 2008; Kim *et al.* 2009). In this study, our RT-PCR and Western blots results showed that the *S. japonicum* GPx was found to be confined express in the adult female worms and eggs (Fig. 4). These results indicated that trematodes GPx proteins might play key roles during the parasite egg production/eggshell formation.

Increasing evidence indicates that loss-of-function genetic manipulation through RNAi may be generally applicable for investigation of schistosome genes (Krautz-Peterson *et al.* 2007; Morales *et al.* 2008). After processing of dsRNA by dicer, the resulting short interfering RNAs join with the effector nuclease complex which recognizes and attacks the homologous target mRNAs, leading to gene silencing (Tavernarakis *et al.* 2000). In schistosomes this can result in long-term knockdown and silencing of the target gene (Morales *et al.* 2008). To investigate the involvement of the *GPx* gene of schistosome egg formation, we performed a RNAi experiment, delivering *SjGPx* dsRNA into

schistosomes by lipofectamine transfection assay. The results of RNAi-mediated knockdown of *SjGPx* in transcript showed that the GPx enzymatic activities were specifically suppressed, and the undeveloped intrauterine eggs of females were more severely deformed. Many irregular eggs were observed in the female uteri and culture medium (Fig. 5D). Therefore, effective antioxidant enzymes, such as GPx, may be required for the protection of macromolecules and/or the fertilized ovum during the trematode egg production/maturation. Gene knockdown of *SjGPx* mRNA might be represented a novel and potentially effective strategy for combatting schistosomiasis.

#### FINANCIAL SUPPORT

This study was supported by a National Science Foundation of China (grant no. 30972568 to L.H. and G.B.C.) and also by a research grant from Health Department of Hubei Province, P. R. China (grant no. XF2010-17 to G.B.C.).

#### REFERENCES

- Arthur, J. R. (2000). The glutathione peroxidases. *Cellular and Molecular Life Sciences* **57**, 1825–1835.
- Bae, Y. A., Kim, S. H., Ahn, C. S., Kim, J. G. and Kong, Y. (2015). Molecular and biochemical characterization of *Paragonimus westermani* tyrosinase. *Parasitology* **142**, 807–815.
- Brigelius-Flohé, R., Aumann, K. D., Blöcker, H., Gross, G., Kiess, M., Klöppel, K. D., Maiorino, M., Roveri, A., Schuckelt, R., Usani, F., Wingender, E. and Flohé, L. (1994). Phospholipid-hydroperoxide glutathione peroxidase. Genomic DNA, cDNA, and deduced amino acid sequence. *Journal of Biological Chemistry* **269**, 7342–7348.
- Cai, G., Bae, Y., Zhang, Y., He, Y., Jiang, M. and He, L. (2009). Expression and characterization of two tyrosinases from the trematode *Schistosoma japonicum*. *Parasitology Research* **104**, 601–609.
- Cai, G. B., Bae, Y. A., Kim, S. H., Sohn, W. M., Lee, Y. S., Jiang, M. S., Kim, T. S. and Kong, Y. (2008). Vitellocyte-specific expression of hydroperoxide glutathione peroxidases in *Clonorchis sinensis*. *International Journal for Parasitology* **38**, 1613–1623.
- Callahan, H. L., Crouch, R. K. and James, E. R. (1988). Helminth antioxidant enzymes: a protective mechanism against host oxidants? *Parasitology Today* **4**, 218–225.
- Cheng, G., Fu, Z., Lin, J., Shi, Y., Zhou, Y., Jin, Y. and Cai, Y. (2009). *In vitro* and *in vivo* evaluation of small interference RNA-mediated gynaecophoral canal protein silencing in *Schistosoma japonicum*. *The Journal of Gene Medicine* **11**, 412–421.
- Cordingley, J. S. (1987). Trematode eggshells: novel protein biopolymers. *Parasitology Today* **22**, 341–344.
- Dröge, W. (2002). Free radicals in the physiological control of cell function. *Physiological Reviews* **82**, 47–95.
- Epp, O., Ladenstein, R. and Wendel, A. (1983). The refined structure of the selenoenzyme glutathione peroxidase at 0.2-nm resolution. *European Journal of Biochemistry* **133**, 51–69.
- Hancock, J. T., Desikan, R. and Neill, S. J. (2001). Role of reactive oxygen species in cell signalling pathways. *Biochemistry Society Transactions* **29**, 345–350.
- Henkle-Dührsen, K. and Kampkötter, A. (2001). Antioxidant enzyme families in parasitic nematodes. *Molecular and Biochemical Parasitology* **114**, 129–142.
- He, Y., Cai, G., Ni, Y., Li, Y., Zong, H. and He, L. (2012). siRNA-mediated knockdown of two tyrosinase genes from *Schistosoma japonicum* cultured *in vitro*. *Experimental Parasitology* **132**, 394–402.
- Imai, H. and Nakagawa, Y. (2003). Biological significance of phospholipid hydroperoxide glutathione peroxidase (PHGPx, GPx4) in mammalian cells. *Free Radical Biology and Medicine* **34**, 145–169.
- Kim, S. H., Cai, G. B., Bae, Y. A., Lee, E. G., Lee, Y. S. and Kong, Y. (2009). Two novel phospholipid hydroperoxide glutathione peroxidase genes of *Paragonimus westermani* induced by oxidative stress. *Parasitology* **136**, 553–565.
- Krautz-Peterson, G., Radwanska, M., Ndegwa, D., Shoemaker, C. B. and Skelly, P. J. (2007). Optimizing gene suppression in schistosomes using RNA interference. *Molecular and Biochemical Parasitology* **2**, 194–202.
- Kryukov, G. V., Castellano, S., Novoselov, S. V., Lobanov, A. V., Zehtab, O., Guigó, R. and Gladyshev, V. N. (2003). Characterization of mammalian selenoproteomes. *Science* **300**, 1439–1443.
- Kumagai, T., Osada, Y., Ohta, N. and Kanazawa, T. (2009). Peroxiredoxin-1 from *Schistosoma japonicum* functions as a scavenger against hydrogen peroxide but not nitric oxide. *Molecular and Biochemical Parasitology* **164**, 26–31.
- McManus, D. P., Hu, W., Brindley, P. J., Feng, Z. and Han, Z. G. (2004). Schistosome transcriptome analysis at the cutting edge. *Trends in Parasitology* **20**, 301–304.
- Mei, H. and LoVerde, P. T. (1995). *Schistosoma mansoni*: cloning the gene encoding glutathione peroxidase. *Experimental Parasitology* **80**, 319–322.
- Mei, H., Thakur, A., Schwartz, J. and LoVerde, P. T. (1996). Expression and characterization of glutathione peroxidase activity in the human blood fluke *Schistosoma mansoni*. *Infection Immunity* **64**, 4299–4306.
- Mkoi, G. M., Smith, J. M. and Prichard, R. K. (1988). Antioxidant systems in *Schistosoma mansoni*: correlation between susceptibility to oxidant killing and the levels of scavengers of hydrogen peroxide and oxygen free radicals. *International Journal for Parasitology* **18**, 661–666.
- Morales, M. E., Rinaldi, G., Gobert, G. N., Kines, K. J., Tort, J. F. and Brindley, P. J. (2008). RNA interference of *Schistosoma mansoni* cathepsin D, the apical enzyme of the hemoglobin proteolysis cascade. *Molecular and Biochemical Parasitology* **157**, 160–168.
- Mourão, M. M., Dinguirard, N., Franco, G. R. and Yoshino, T. P. (2009). Phenotypic screen of early developing larvae of the blood fluke, *Schistosoma mansoni*, using RNA interference. *PLoS Neglected Tropical Diseases* **3**, e502.
- Roche, C., Liu, J. L., LePresle, T., Capron, A. and Pierce, R. J. (1996). Tissue localization and stage-specific expression of the phospholipid hydroperoxide glutathione peroxidase of *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* **75**, 187–195.
- Selkirk, M. E., Smith, V. P., Thomas, G. R. and Gounaris, K. (1998). Resistance of filarial nematode parasites to oxidative stress. *International Journal for Parasitology* **28**, 1315–1332.
- Sies, H. (1993). Strategies of antioxidant defense. *European Journal of Biochemistry* **215**, 213–219.
- Stadtman, T. C. (1996). Selenocysteine. *Annual Review of Biochemistry* **65**, 83–100.
- Tabara, H., Grishok, A. and Mello, C. C. (1998). RNAi in *C. elegans*: soaking in the genome sequence. *Science* **282**, 430–431.
- Tavernarakis, N., Wang, S. L., Dorovkov, M., Ryazanov, A. and Driscoll, M. (2000). Heritable and inducible genetic interference by double-stranded RNA encoded by transgenes. *Nature Genetics* **2**, 180–183.
- Tchoubrieva, E. B., Ong, P. C., Pike, R. N., Brindley, P. J. and Kalinna, B. H. (2010). Vector-based RNA interference of cathepsin B1 in *Schistosoma mansoni*. *Cellular and Molecular Life Sciences* **67**, 3739–3748.
- Ursini, F. and Bindoli, A. (1987). The role of selenium peroxidases in the protection against oxidative damage of membranes. *Chemistry and Physics of Lipids* **44**, 255–276.
- Williams, D. L., Pierce, R. J., Cookson, E. and Capron, A. (1992). Molecular cloning and sequencing of glutathione peroxidase from *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* **52**, 127–130.
- Wyler, D. J. (1992). Why does liver fibrosis occur in schistosomiasis? *Parasitology Today* **8**, 277–279.
- Zamore, P. D., Tuschl, T., Sharp, P. A. and Bartel, D. P. (2000). RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* **101**, 25–33.
- Zelck, U. E. and von Janowsky, B. (2004). Antioxidant enzymes in invertebrates *Schistosoma mansoni* and ROS-induced changes in expression. *Parasitology* **128**, 493–501.