


# First report of pre-Hispanic *Fasciola hepatica* from South America revealed by ancient DNA

María Ornela Beltrame<sup>1</sup> , Cesar Pruzzo<sup>2</sup>, Rodrigo Sanabria<sup>2,3</sup>, Alberto Pérez<sup>4</sup> and Matías Sebastián Mora<sup>5</sup>

## Research Article

**Cite this article:** Beltrame MO, Pruzzo C, Sanabria R, Pérez A, Mora MS (2020). First report of pre-Hispanic *Fasciola hepatica* from South America revealed by ancient DNA. *Parasitology* **147**, 371–375. <https://doi.org/10.1017/S0031182019001719>

Received: 9 September 2019  
Revised: 20 November 2019  
Accepted: 21 November 2019  
First published online: 20 December 2019

### Key words:

aDNA; DEER; *Fasciola hepatica*; palaeoparasitology

### Author for correspondence:

María Ornela Beltrame,  
E-mail: [ornelabeltrame@conicet.gov.ar](mailto:ornelabeltrame@conicet.gov.ar)

<sup>1</sup>Grupo de investigación: Paleoparasitología. Instituto de Investigaciones en Producción, Sanidad y Ambiente (IIPROSAM), Facultad de Ciencias Exactas y Naturales, UNMDP-CONICET, Mar del Plata, Buenos Aires, Argentina; <sup>2</sup>Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata (UNLP), La Plata, Argentina; <sup>3</sup>Instituto Tecnológico Chascomús (INTECH) – CONICET-Universidad Nacional de San Martín (UNSAM), Chascomús, Argentina; <sup>4</sup>Departamento de Antropología, Universidad Católica de Temuco, Campus San Francisco, Temuco, Región de La Araucanía, Chile and <sup>5</sup>Instituto de Investigaciones Marinas y Costeras (IIMyC), CONICET, Universidad Nacional de Mar del Plata (UNMDP), Mar del Plata, Argentina

### Abstract

It is generally assumed that the digenean human liver fluke, *Fasciola hepatica*, gained entry to South America during the 15th century upon arrival of Europeans and their livestock. Nonetheless in Patagonia, Argentina, digenean eggs similar to *F. hepatica* have been observed in deer coprolites dating back to 2300 years B.P. The main objective of our present study was to identify and characterize these eggs using an ancient DNA (aDNA) study. Eggs were isolated and used for aDNA extraction, amplification and sequencing of partial regions from the cytochrome *c* oxidase subunit 1 and the nicotinamide adenine dinucleotide dehydrogenase subunit 1 mitochondrial genes. Also, phylogenetic trees were constructed using Bayesian and maximum likelihood. Our results confirm the presence of *F. hepatica* in South America from at least 2300 years B.P. This is the first report and the first aDNA study of this trematode in South America prior to the arrival of the European cattle in the 15th century. The present work contributes to the study of phylogenetic and palaeobiogeographical aspects of *F. hepatica* and its settlement across America.

## Introduction

Fasciolosis is a zoonotic parasitic disease caused by the liver flukes *Fasciola hepatica* and *Fasciola gigantica* (Trematoda: Digenea). This helminthic disease is of worldwide distribution (Mas-Coma *et al.*, 2009) and is considered as an important veterinary health problem due to the substantive economic losses it gives rise to livestock husbandry. Moreover, it is of great public health importance in some countries, due to its high pathogenicity (Mas-Coma *et al.*, 2014) and is therefore included within the group of Foodborne Trematodiasis among Neglected Tropical Diseases (NTDs) by the World Health Organization (WHO, 2013).

*Fasciola hepatica* has a worldwide distribution and *F. gigantica* is found in tropical climates and restricted to Africa, the Middle East and South and East Asia. In America, fasciolosis is caused by *F. hepatica* and transmitted by many different intermediate snail hosts belonging to the family Lymnaeidae, mainly species included within the *Galba/Fossaria* group. In South America, human endemic areas have been described in Andean regions, mainly in highlands of Bolivia, Peru and Chile, and sporadic cases are reported in other countries (Mas-Coma, 2005; World Health Organization, 2013; Carmona and Tort, 2016).

It is generally assumed that entry of *F. hepatica* to America coincided with the first arrival of the Europeans and their associated livestock in the late 15th century. Throughout the 500 years since its introduction, the parasite gained new definitive hosts among native species. The South American camelids – llamas, alpacas, vicuñas and guanacos – the natural livestock of the Andean region, might have represented the first to be parasitized, since these species would graze along with the introduced livestock (Mas-Coma *et al.*, 2009; Barges *et al.*, 2017). The parasite is now widespread in livestock and can be mapped across the whole South America and certain regions of North America.

Argentina has a large livestock production, where sheep and cattle constitute important economic sources. Animal fasciolosis is currently found in spots across the country, according to official slaughterhouses, and the most important hosts are cattle and sheep. Goats, horses, pigs and some wild native and non-native mammals (deer, vicuña, guanacos, llamas, rabbits, hares and capybaras) are also found infected by *F. hepatica*. Despite the fragmented and anecdotal nature of several reports of liver flukes in South American wildlife, it is evident that diverse species can host the parasite and eventually act as reservoirs (Issia *et al.*, 2009; Carmona and Tort, 2016).

Given the medical and veterinary importance of fasciolosis, there are several multidisciplinary studies investigating parasite origins and dispersals across the world; shared objectives include elucidation of palaeobiogeographical origins and colonization within and between continents (Mas-Coma *et al.*, 2009).

Palaeoparasitology is the study of parasite remains from archaeological and palaeontological sites (Ferreira, 2014), focused on the knowledge of parasite-induced illness of humans in the past and on the palaeoecological knowledge of the environment, ecology, settlement, diet, hygiene and health in the antiquity (Reinhard, 1992). In a previous palaeoparasitological study, digenean eggs similar to *F. hepatica* were found in coprolites from one of the two endemic species of deer inhabiting the narrow Andean-Patagonian temperate forest strip in the west of southern America, the southern pudú (*Pudu puda*) and the huemul (*Hippocamelus bisulcus*). The samples were obtained from an archaeological site of Patagonia named 'Cueva Parque Diana' (CPD) and were dated around 2300 years B.P. (Late Holocene) (Beltrame *et al.*, 2017). In order to contribute with the study of the palaeobiogeographical origins of *F. hepatica* and their settlement across America, the main objective of the present study is to identify the digenean eggs found from CPD from an ancient DNA (aDNA) study.

## Material and methods

### Sample collection

In a previous study (Beltrame *et al.*, 2017), 34 coprolites were processed for palaeoparasitological purposes. Coprolites belonged to native deer identified as *P. puda* or *H. bisulcus*. Samples were collected from the CPD archaeological site, Lanín National Park, North Patagonia, Argentina (40°19'93"S, 71°20'74"W). This site is a rock shelter part of the archaeological locality named Meliquina. It is located at 964 m.a.s.l. and 50 m close to the Hermoso River. The archaeological sequence was divided into three components representing different hunter-gatherer occupation processes. The Upper Component was dated between 760 ± 60 and 580 ± 60 <sup>14</sup>C years B.P. (vegetal charcoal), the Middle Component was dated between 990 ± 60 and 900 ± 60 <sup>14</sup>C years B.P. (vegetal charcoal) and the Lower Component was dated at 2370 ± 70 <sup>14</sup>C years B.P. (vegetal charcoal). The site was occupied by hunters-gatherers and fishermen along the late Holocene (Pérez, 2010; Pérez *et al.*, 2015). The weather in the area is cold and wet, with annual precipitations around 1500–2000 mm.

Eighteen of the 34 samples were positive for digenean eggs. Positive coprolites were found in the upper, middle and lower components. The analysed eggs (Fig. 1) in this study belong to two positive samples studied in Beltrame *et al.* (2017) from the lower component. The eggs were identified under a light microscope (100× magnification) and were manually isolated by the use of a micropipette and stored in PCR tubes with phosphate-buffered saline. The isolated eggs were used for aDNA extraction, amplification and sequencing of partial regions from the cytochrome *c* oxidase subunit 1 (COI) and the nicotinamide adenine dinucleotide dehydrogenase subunit 1 (NADHI) mitochondrial genes.

### aDNA extraction and polymerase chain reaction amplification

Before DNA extraction, 30 eggs were washed three times in ultrapure water (Invitrogen by Thermo Fisher Scientific, Waltham, MA, USA). Once washed, a first disruption step was performed by five continuous cycles of freezing (immersion in liquid nitrogen for 10 s) and heating (immersion of the tube in boiling water for 2–3 s).

DNA was extracted using the ZR Fecal DNA Miniprep kit (Zymo Research, Irvine, CA, USA), following the manufacturer's instructions. PCR reactions were individually carried out for each partial fragment of the *mtDNA* genes coding for NADHI and COI. Each reaction was constituted into a final volume of 50 µL, containing 37.5 µL of ultrapure distilled water (Invitrogen), 5 µL of 10×



**Fig. 1.** Ancient trematode egg found from deer coprolites from 'Cueva Parque Diana' archaeological site, Patagonia, Argentina. Bar = 40 µm.

buffer, 4.5 µL of PCR mix containing 1 mmol of MgCl<sub>2</sub>, (2 µL) 0.2 mM of dNTPs (1 µL), 50 pmol of each primer (1 µL) and 2.5 U of *Taq* Polymerase (0.5 µL) (all reagents from Invitrogen), plus 3 µL of each template. PCR primers and temperature settings were previously described by Ichikawa and Itagaki (2012) for NADHI and COI. The primers' sequences were 5'-AAGGATG-TTGCTTTGTCGTGG-3' (forward), 5'-GGAGTACGGTTACATTCACA-3' (reverse), for NADHI, and 5'-ACGTTGGATCAT-AAGCGTGT-3' (forward), 5'-CCTCATCCAACATAACCTCT-3' (reverse), for COI. Negative controls (ultrapure water and PCR reagents) were also ran along with templates amplification.

Amplicons were visualized in 2% agarose gels stained with SybrSafe (Invitrogen) using a blue light transilluminator (Safe Imager, Invitrogen), and purified by a commercial kit (DNA Clean & Concentrator-5, Zymo Research), according to the manufacturer's instructions. Purified PCR products were elected into 10 µL and submitted to Macrogen's sequencing service (Seoul, Korea), for capillary electrophoretic sequencing using the same primers as for the PCR reactions. Once available, the sequences were submitted to the GenBank database (NCBI).

### Data analysis

Electropherograms were scored and analysed using Chromas2.01 (Technelysium, Helensvale, QLD, Australia) and aligned using CLUSTALW2 (Larkin *et al.*, 2007) in MEGA 7.0.26 (Kumar *et al.*, 2016) using default settings. Homologies were performed using the BLASTN programme from the National Center for Biotechnology Information website (<http://www.ncbi.nlm.nih.gov/BLAST>).

In order to prove the phylogenetic identity, our mitochondrial partial sequences of aDNA from COI and NADHI were compared with the corresponding mitochondrial sequences of COI and NADHI of current individuals of *F. hepatica*. These last mitochondrial sequences of both molecular markers correspond to the

following GenBank accession numbers: KR422380–KR422388, MG870561, MG870563–MG870566, MG870568–MG870570, MG987190, MG9871902, LC273097, LC273100, LC273110, LC273111 and LC273113 from COI; and LC273198–LC273202, MG972375–MG972379, MG972405–MG972409, LC076246, LC076249, LC076255, LC076259, LC076261, KR422389–KR422393 and KJ852771 from NADHI. Both COI and NADHI sequences were selected from several continental regions in order to include a broad representation of genetic diversity of these parasites characterized by a wide distribution worldwide. As out-groups we used mitochondrial sequences from two species of the family Fasciolidae (*F. gigantica* and *Fascioloides magna*); with the GenBank accession numbers AB385622, AB385621, AB385620, AB207176, AB207181, EF534996, EF534997 and EF534998 from COI; and MG972405, MG972406, MG972407, MG972408, MG972409, EF534999 and EF535000 from NADHI.

### Phylogenetic inferences

We used jModelTest (Darriba *et al.*, 2012) to infer the best-fit substitution model for both sets of data (COI and NADHI). To assess the robustness of parameter estimates, four independent chains were run with identical settings. Log-files were analysed in Tracer v1.7.1 (Rambaut *et al.*, 2018) and effective sample sizes were used to evaluate Markov chain Monte Carlo (MCMC) convergence within chains. We used the most credible substitution models in both cases: the HKY + G substitution model with four  $\gamma$  categories was performed for the COI dataset, and the HKY + I substitution model was performed for the NADHI dataset, using in each case a Yule branching rate prior, with rate variation across branches assumed to be uncorrelated and log-normally distributed (Drummond *et al.*, 2006).

We constructed a Bayesian phylogenetic tree using BEAST v2.5.2 (Bouckaert *et al.*, 2019), which employs a Bayesian MCMC approach. Each MCMC chain was run for  $10^7$  iterations (with a burn-in of 50% of the total chains), with parameters sampled every 1000 steps. For comparison, three independent MCMC runs were performed to validate the topology of each phylogenetic tree. Examination of MCMC samples using Tracer v1.7.1 (Rambaut *et al.*, 2018) suggested that the independent chains were each adequately sampling the same probability distribution and effective sample sizes for all parameters of interest were >500, conditions suggested by the authors for the proper functioning of the analysis.

A phylogenetic tree was also constructed using a maximum likelihood approach (ML) implemented in MEGA7 (Kumar *et al.*, 2016), considering the same out-groups used in the Bayesian phylogenetic inference. The programme implements simultaneous Nearest Neighbor Interchanges (NNIs) to improve a reasonable topology of the starting tree. The final run of MEGA7 considered the best substitution model inferred by jModeltest (see above; Darriba *et al.*, 2012), where both the transitions/transversions ratio and  $\gamma$  distribution parameter were empirically estimated. Consistency for internal branch and nodes was assessed using the standard bootstrapping method (sample with replacement, 1000 bootstrap replicates) implemented in MEGA7.

### Results

The morphological observations of the ancient eggs indicated that they all belonged to the same trematode. The eggs were ovoid, operculated, brown-yellowish and thin-shelled. The length ( $n = 102$ ) ranged from was  $131.7 \pm 7.82 \mu\text{m}$  (range = 120.0–147.5) and the width was  $72.8 \pm 5.96 \mu\text{m}$  (range = 62.5–87.5). Eggs were well-preserved and were identified as Class Trematoda, Subclass Digenea, Family Fasciolidae, possibly *F. hepatica*. The PCR

analysis, elicited bands of ~600 and 550 bp for NADHI and COI, in concordance with controls of *F. hepatica* and previous reports (Ichikawa and Itagaki, 2012). The sequences were assigned with the accession numbers MN207487 and MN207488 for COI and NADHI, respectively. Final alignments represented smaller fragments of 417 bp from NADHI, and 350 bp from COI, compared to the length of the PCR band. Compared to the whole mitochondrial genes (Le *et al.*, 2000), our 417 bp partial sequence included positions 99–516 of the NADHI gene, and the 350 bp partial sequence included the bases 15–365 of the COI gene. Non-indels or gaps were detected neither for NADHI nor COI when we compared our target sequence and the other sequences of this analysis.

In both cases, these genes showed 100% of identity compared to some of the GenBank's records of *F. hepatica* (NCBI nucleotide Blast: <https://blast.ncbi.nlm.nih.gov/Blast.cgi>).


The phylogenetic inferences (ML and Bayesian) strongly support the hypothesis that our target sequences of deer effectively correspond to *F. hepatica* (Fig. 2). In our phylogenetic tree, these two sequences of mitochondrial aDNA were clearly included within the clade of *F. hepatica*, showing high node supports using both phylogenetic inferences. Also, these sequences showed high genetic distances relative to *F. gigantica* and *F. magna* within the phylogenetic tree.

### Discussion

The sequences of the mitochondrial aDNA used in this study demonstrate that the trematode eggs from native deer coprolites from the CPD archaeological site belong to *F. hepatica*. The results confirm the presence of this trematode in South America from at least 2300 years B.P. This is the first report and the first aDNA study of *F. hepatica* in South America prior to the arrival of the European cattle in the 15th century. Previous palaeoparasitological studies have reported the presence of this parasite in Europe and Asia (e.g. Bouchet, 1995; Bouchet *et al.*, 2003; Dittmar and Teegen, 2003; Askari *et al.*, 2018). The first study that identifies aDNA of *Fasciola* sp. was made by S e *et al.* (2015). In this study, *Fasciola* sp. eggs were recovered from environmental samples collected at a Viking-age settlement in Viborg, Denmark, dated at 1018–1030 A.D. Ancient DNA studies also were performed from European archaeological sites such as C t  *et al.* (2016) and S e *et al.* (2018). In a recent study, Le Bailly *et al.* (2019) found one trematode egg in domestic camelids which could belong to *Fasciola* recovered from the pre-Hispanic Chim  culture site of Huanchaquito-Las Llamas, Peru. Although its diagnosis has not been confirmed yet, this would correspond to an additional evidence of the presence of the liver fluke genus *Fasciola* in pre-Hispanic times of America.

The current knowledge of the presence of *F. hepatica* on current native deer from Patagonia is limited. Some studies reported the presence of *F. hepatica* in the southern pud  (Cort s, 2006; Bravo Antilef, 2013) and in the huemul (D az and Smith-Flueck, 2000; Serret, 2001). It was also registered in the introduced red deer (*Cervus elaphus*) (Flueck and Smith-Flueck, 2012).

Our COI sequence of aDNA was mostly related with contemporary South American DNA sequences (Ecuador and Peru), and was most distant to DNA sequences from Asia, Africa and Europe. However, the NADHI sequence not only showed a great phylogenetic affinity with sequences of South America, but with Armenia and Egypt (see Fig. 2). The objective of this work was to identify digenean eggs found from deer coprolites from an aDNA study, without a deeper more conclusive evaluation of their possible palaeobiogeographical origin or colonization routes within this continent. In this sense, there are many methodological limitations that hinder the inference about a



**Fig. 2.** The evolutionary history was inferred by using the ML method (values of the nodes are given in percentage) and the Bayesian phylogenetic inference (values of the nodes are given in posterior probabilities) based on the Hasegawa–Kishino–Yano model + G for COI (A), and on the Hasegawa–Kishino–Yano model + I for NADH1 (B) (Tamura and Nei, 1993). For the ML tree, initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. The rate variation model allowed for some sites to be evolutionarily invariable (+I), 39.94% sites). Topology of the tree corresponds to that obtained from Bayesian phylogenetic inference (Drummond and Rambaut, 2007). The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. *Fascioloides magna* and *Fasciola gigantica* were selected as out-groups in both phylogenetic trees. The analysis involved 33 nucleotide sequences for COI (350 positions in the final dataset) and 29 nucleotide sequences for NADH1 (417 positions in the final dataset). Genbank accession numbers and their geographical origin by countries are given in parenthesis (Ec.: Ecuador; Pol.: Poland; Ir.: Iran; Ar.: Armenia; Pe.: Peru; Eg.: Egypt). Only the geographical origin of the *F. hepatica* sequences is shown.

possible geographical origin of these aDNA sequences reported here, beyond including contemporary DNA samples from various regions of the world in our phylogenetic analyses. In principle, knowledge about the dynamics of migration and colonization of this parasite among different continents seems to be very complex. Demographic patterns of *F. hepatica* have not only been conditioned by the regional history of domestic animal translocations performed by humans in recent decades, but also to their associations with populations of native mammal species. The latter can lead to confusing the biogeographic and demographic history of

*F. hepatica* at the continental level, especially when the ancient material is compared with current DNA samples. The need for future palaeoparasitological studies around the world is evident in order to contribute to the palaeobiogeographical origin of this trematode.

The fact that *F. hepatica* was present in America before the arrival of the Europeans in the 15th century indicates that the species was not first introduced by the European cattle during this time, but there was another alternative route of prior entry. At the moment our data do not allow us to propose a plausible

hypothesis about the possible entry of *F. hepatica* to the American continent prior to this period. Future palaeoparasitological studies are needed which should consider the different migratory routes into the American continent in pre-Hispanic times. Evolutionary relationship among *F. hepatica* and American native hosts, both definitive and intermediate, is an interesting point to be studied in future studies. The pre-Hispanic presence of *F. hepatica* in South America brings new insights to the common assumption on the palaeobiogeography and settlement of this species.

### Financial support

None.

### Conflict of interest

None.

### Ethical standards

Not applicable.

### References

- Askari Z, Mas-Coma S, Bouwman AS, Boenke N, Stöllner T, Aali A, Rezaian M and Mowlavi G (2018) *Fasciola hepatica* eggs in paleofaeces of the Persian onager *Equus hemionus onager*, a donkey from Chehrabad archaeological site, dating back to the Sassanid Empire (224–651 CE), in ancient Iran. *Infection, Genetic and Evolution* **62**, 233–243.
- Bargues MD, Gayo V, Sanchis J, Artigas P, Khoubbane M, Birriel S and Mas-Coma S (2017) DNA multigene characterization of *Fasciola hepatica* and *Lymnaea neotropica* and its fascioliasis transmission capacity in Uruguay, with historical correlation, human report review and infection risk analysis. *PLoS Neglected Tropical Diseases* **11**, e0005352.
- Beltrame MO, Tietze E, Pérez AE and Sardella NH (2017) First paleoparasitological record of digenae eggs from a native deer from Patagonia Argentina (Cueva Parque Diana archaeological site). *Veterinary Parasitology* **235**, 83–85.
- Bouchet F (1995) Recovery of helminth eggs from archaeological excavations of the Grand Louvre (Paris, France). *Journal of Parasitology* **81**, 785–787.
- Bouchet F, Harter S and Le Bailly M (2003) The state of the art of paleoparasitological research in the Old World. *Memórias Instituto Oswaldo Cruz* **98**, 95–101.
- Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones G, Kühnert D, De Maio N, Matschiner M, Mendes FK, Müller NE, Ogilvie HA, du Plessis L, Poppinga A, Rambaut A, Rasmussen D, Siveroni I, Suchard MA, Wu C, Xie D, Zhang C, Stadler T and Drummond AJ (2019) BEAST 2.5: an advanced software platform for Bayesian evolutionary analysis. *PLoS Computational Biology* **15**, e1006650.
- Bravo Antilef MJ (2013) *Probables causas de muerte y principales hallazgos en la necropsia de pudúes (Pudu puda) examinados durante 20 años en el sur de Chile*. Memoria de Título. Universidad Austral de Chile. Facultad de Ciencias Veterinarias. Instituto de Patología Animal. Chile.
- Carmona C and Tort JF (2016) Fasciolosis in South America: epidemiology and control challenges. *Journal of Helminthology* **91**, 99–109.
- Cortés M (2006) *Identificación de formas reproductivas de parásitos gastrointestinales, en mamíferos nativos presentes en el Buin Zoo, Chile*. Memoria de título. Escuela de Medicina Veterinaria, Universidad de Concepción, Chillán, Chile.
- Côté NML, Daligault J, Pruvost M, Bennett EA, Gorgé O, Guimaraes S, Capelli N, Le Bailly M, Geigl E and Grange T (2016) A new high-throughput approach to genotype ancient human gastrointestinal parasites. *PLoS ONE* **11**, e0146230.
- Darriba D, Taboada GL, Doallo R and Posada D (2012) Jmodeltest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**, 772.
- Díaz NI and Smith-Flueck JA (2000) *El Huemul patagónico: un misterioso cérvido al borde de la extinción*. Argentina: L.O.L.A., pp. 156.
- Dittmar K and Teegen WR (2003) The presence of *Fasciola hepatica* (liver-fluke) in humans and cattle from a 4500 year old archaeological site in the Saale-Unstrut-Valley, Germany. *Memórias do Instituto Oswaldo Cruz* **98**, 141–145.
- Drummond AJ and Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* **7**, 214.
- Drummond AJ, Ho SYW, Phillips MJ and Rambaut A (2006) Relaxed phylogenetics and dating with confidence. *PLoS Biology* **4**, e88.
- Ferreira LF (2014) An introduction to paleoparasitology. In Ferreira LF, Reinhard K and Araújo A (eds), *Foundations of Paleoparasitology*. Rio de Janeiro, Brazil: Fiocruz/International Federation of Tropical Medicine, pp. 27–41.
- Flueck WT and Smith-Flueck JM (2012) Diseases of red deer introduced to Patagonia and implications for native ungulates. *Animal Production Science* **52**, 766–773.
- Ichikawa M and Itagaki T (2012) Molecular analysis of aspermic *Fasciola* flukes from Korea on the basis of the nuclear ITS1 region and mitochondrial DNA markers and comparison with Japanese aspermic *Fasciola* flukes. *The Journal of Veterinary Medical Science* **74**, 899–904.
- Issia L, Petrokovsky S, Sousa-Figueiredo J, Russell Stothard J and Wisnivesky-Colli C (2009) *Fasciola hepatica* infections in livestock flock, guanacos and coypus in two wildlife reserves in Argentina. *Veterinary Parasitology* **165**, 341–344.
- Kumar S, Stecher G and Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**, 1870–1874.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ and Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics (Oxford, England)* **23**, 2947–2948.
- Le TH, Blair D, Agatsuma T, Humair PF, Campbell NJ, Iwagami M, Littlewood DT, Peacock B, Johnston DA, Bartley J, Rollinson D, Herniou EA, Zarlenga DS and McManus DP (2000) Phylogenies inferred from mitochondrial gene orders—a cautionary tale from the parasitic flatworms. *Molecular Biology and Evolution* **17**, 1123–1125.
- Le Bailly M, Goepfert N, Prieto G, Verano J and Dufour B (2019) Camelid gastrointestinal parasites from the Archaeological Site of Huanchaquito (Peru): first results. *Environmental Archaeology*. doi: 10.1080/14614103.2018.1558804.
- Mas-Coma S (2005) Epidemiology of fascioliasis in human endemic areas. *Journal of Helminthology* **79**, 207–216.
- Mas-Coma S, Valero MA and Bargues MD (2009) *Fasciola*, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Advances in Parasitology* **69**, 41–146.
- Mas-Coma S, Agramunt VH and Valero MA (2014) Neurological and ocular fascioliasis in humans. *Advances in Parasitology* **84**, 27–149.
- Pérez AE (2010) La Localidad Arqueológica Lago Meliquina, Dto. Lácara, Neuquén. El registro arqueológico del interior y borde de bosque en Norpatagonia. *Actas y Memorias del XVII Congreso Nacional de Arqueología Chilena*, Valdivia, Chile, 2006, pp. 1515–1528.
- Pérez AE, Aguirre MG and Graziano JE (2015) Imprints of cariopsis de gramíneas (Poaceae) en un fragmento de alfarería de Patagonia Noroccidental Argentina. *Revista de Antropología del Museo de Entre Ríos* **1**, 77–85.
- Rambaut A, Drummond AJ, Xie D, Baele G and Suchard MA (2018) Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**, 901–904.
- Reinhard KJ (1992) Parasitology as an interpretative tool in archaeology. *American Antiquity* **57**, 231–245.
- Serret A (2001) *El Huemul: Fantasma de la Patagonia*. Ushuaia, Argentina: Zagier and Urruty.
- Søe MJ, Nejsum P, Fredensborg BL and Kapel CMO (2015) DNA typing of ancient parasite eggs from environmental samples identifies human and animal worm infections in Viking-age settlement. *Journal of Parasitology* **101**, 57–64.
- Søe MJ, Nejsum P, Seersholm FV, Fredensborg BL, Habraken R, Haase K, Hald MM, Simonsen R, Højlund F, Blanke L and Merkyte I (2018) Ancient DNA from latrines in Northern Europe and the Middle East (500 BC±1700 AD) reveals past parasites and diet. *PLoS ONE* **13**, e0195481.
- Tamura K and Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**, 512–526.
- World Health Organization (2013) *Sustaining the Drive to Overcome the Global Impact of Neglected Tropical Diseases*. Geneva: World Health Organization, WHO Headquarters, 138 pp.