

# Insight, self-stigma and psychosocial outcomes in Schizophrenia: a structural equation modelling approach

Y.-J. Lien<sup>1</sup>, H.-A. Chang<sup>2,3</sup>, Y.-C. Kao<sup>3,4\*</sup>, N.-S. Tzeng<sup>2,3</sup>, C.-W. Lu<sup>4</sup> and C.-H. Loh<sup>5</sup>

<sup>1</sup> Department of Health Promotion and Health Education, National Taiwan Normal University, Taipei, Taiwan

<sup>2</sup> Department of Psychiatry, Tri-Service General Hospital, Taipei, Taiwan

<sup>3</sup> Department of Psychiatry, National Defense Medical Center, Taipei, Taiwan

<sup>4</sup> Department of Psychiatry, Tri-Service General Hospital Songshan Branch, Taipei, Taiwan

<sup>5</sup> School of Medicine, National Defense Medical Center, Taipei, Taiwan

**Aims.** Poor insight is prevalent in patients with schizophrenia and has been associated with acute illness severity, medication non-adherence and poor treatment outcomes. Paradoxically, high insight has been associated with various undesirable outcomes, including low self-esteem, depression and low subjective quality of life (QoL) in patients with schizophrenia. Despite the growing body of studies conducted in Western countries supporting the pernicious effects of improved insight in psychosis, which bases on the level of self-stigma, the effects are unclear in non-Western societies. The current study examined the role of self-stigma in the relationship between insight and psychosocial outcomes in a Chinese population.

**Methods.** A total of 170 outpatients with schizophrenia spectrum disorders were recruited from two general university hospitals. Sociodemographic data and clinical variables were recorded and self-report scales were employed to measure self-stigma, depression, insight, self-esteem and subjective QoL. Structural equation modelling (SEM) was used to analyse the cross-sectional data.

**Results.** High levels of self-stigma were reported by 39% of the participants ( $n = 67$ ). The influences of insight, self-stigma, self-esteem and depression on subjective QoL were confirmed by the SEM results. Our model with the closest fit to the data ( $\chi^2 = 33.28$ ;  $df = 20$ ;  $p = 0.03$ ;  $\chi^2/df = 1.66$ ; CFI = 0.98; TLI = 0.97; RMSEA = 0.06) demonstrated that self-stigma might fully mediate the association of insight with low self-esteem, depression and poor subjective QoL. High insight into illness contributed to self-stigma, which caused low self-esteem and depression and, consequently, low QoL. Notably, insight did not directly affect self-esteem, depression or QoL. Furthermore, the association of insight with poor psychosocial outcomes was not moderated by self-stigma.

**Conclusions.** Our findings support the mediating model of insight relevant to the poor psychosocial outcomes of individuals diagnosed with schizophrenia in non-Western societies, in which self-stigma plays a pivotal role. These findings elucidate the direct and indirect effects of insight on psychosocial outcomes and imply that identifying and correcting self-stigma in people with schizophrenia could be beneficial. Additional studies are required to identify whether several other neurocognitive or psychosocial variables mediate or moderate the association of insight with self-esteem, depression and QoL in patients with schizophrenia. Studies with detailed longitudinal assessments are necessary to confirm our findings.

Received 4 March 2016; Accepted 8 November 2016; First published online 15 December 2016

**Key words:** Insight, mediation, moderation, quality of life, schizophrenia, self-stigma.

## Introduction

Lack of insight is a prominent and enduring core feature of schizophrenia (Amador *et al.* 1994; Bora *et al.* 2007). Insight in psychiatric research has been regarded as a multidimensional construct that refers

to the awareness of illness-related issues, which include symptoms of the illness, the need for treatment, and consequences of the illness (Amador *et al.* 1994). Furthermore, researchers have identified that a lack of insight is of clinical concern because it is a significant contributor to disastrous consequences among patients with schizophrenia (Lysaker *et al.* 2002, 2009; Mintz *et al.* 2003; Mohamed *et al.* 2014). Consequently, improving insight is a major goal in the treatment of schizophrenia.

\* Address for correspondence: Y.-C. Kao, Department of Psychiatry, Tri-Service General Hospital Songshan Branch, 131 Jiankang Road, Songshan District, Taipei 10581, Taiwan.  
(Email: freud001@ms45.hinet.net, wheelis001@gmail.com)

Paradoxically, studies have linked higher insight with depression (Cavelti *et al.* 2014; Belvederi Murri *et al.* 2015), feelings of hopelessness (Lysaker *et al.* 2007; Hasson-Ohayon *et al.* 2009), reduced self-esteem (Lysaker *et al.* 2007; Staring *et al.* 2009) and low quality of life (QoL) (Staring *et al.* 2009; Margariti *et al.* 2015). The conflicting results regarding the correlates and consequences of insight in schizophrenia were described as the insight paradox (Lysaker *et al.* 2007). Much research has been devoted to identifying the factors that contribute to the aetiology and maintenance of the insight paradox in schizophrenia. First, as Lysaker *et al.* (2007) noted, the role of self-stigma has received considerable attention, offering support for the relationship between high insight and poor psychosocial outcomes. Specifically, they reported that self-stigma moderated the associations of insight with social functioning, hope, and self-esteem among people with schizophrenia spectrum disorders. The hypothesised moderating role of self-stigma has received further support by subsequent findings reported by Staring *et al.* (2009) and Cavelti *et al.* (2012a); the detrimental effects of improved insight were more pronounced in patients with higher levels of self-stigma than those with lower self-stigma. Nevertheless, the results reinforced the notion that higher insight might be a vulnerability factor for negative psychosocial outcomes, but not an invariant one.

A theoretically divergent proposition presented in the literature is that increased self-stigma is an effect of good insight and results in poor psychosocial outcomes. The mediation hypothesis has received further support from studies suggesting that self-stigma is an intermediate variable on the association between insight and psychological distress (Hasson-Ohayon *et al.* 2011, 2014). Similarly, using a structural equation modelling (SEM) approach, Cavelti *et al.* (2012a, b) have also discovered that the effect of insight and depressive symptoms was mediated by self-stigma among people with schizophrenia spectrum disorders. This finding may indicate that reducing self-stigma in individuals with high insight may consequently diminish demoralisation.

Researchers have suggested that further exploring the mediator and moderator variables may provide valuable information regarding early interventions in psychiatric illnesses, because these variables would address the fundamental questions of how a specific intervention produces clinical benefits (i.e. what variable(s) mediate(s) the effect of the intervention) and when and for whom a specific intervention is most effective (i.e. what variable(s) moderate the effect of the intervention) (Hopwood, 2007; Breitborde *et al.* 2010). Based on the mediator and moderator criteria proposed by Baron & Kenny (1986), clarifying the


role of self-stigma in the relationship between insight into illness and psychosocial outcomes is of practical and theoretical importance. This knowledge can facilitate developing a strategy aimed at improving the psychosocial outcomes of individuals with schizophrenia.

Furthermore, previous studies have suggested that clinical presentations and evaluations of insight (Xiang *et al.* 2012; Mohamed *et al.* 2014) and mental-illness stigma (Livingston & Boyd, 2010; Yang *et al.* 2013; Boyd *et al.* 2014) are not independent of the sociocultural environment; therefore, the results of studies conducted in Western countries may be invalid in Eastern countries. Thus, in contrast to previous studies, the present study explored whether self-stigma serves as a mediator, moderator, or both, in the relationship between insight and psychosocial outcomes among Chinese patients with schizophrenia by using a larger homogeneous sample and more rigorous statistical analyses such as SEM. We hypothesised that higher insight into illness affects people's self-stigma and impedes their self-esteem and QoL. Self-stigma and low self-esteem then directly decrease their QoL. Moreover, self-stigma and low self-esteem cause depression, further contributing to low QoL (Fig. 1). We assumed that insight has both direct and indirect (through the mediating effect of self-stigma) effects on psychosocial outcomes. We also hypothesised that the relationship between insight and psychosocial outcomes would vary according to the level of self-stigma (Fig. 2). Specifically, the associations between insight and each psychosocial outcome (i.e. self-esteem, depression, and subjective QoL) at high levels of self-stigma would be stronger than those at low levels of self-stigma.

## Methods

### Participants

A total of 190 patients with diagnosed schizophrenia or schizoaffective disorder, according to the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders, fourth edition, text revision (DSM-IV-TR) (American Psychiatric Association, 2000), were recruited consecutively in 2013 and 2014 from outpatient psychiatric clinics at two hospitals. All participants were receiving ongoing outpatient treatment and were in relatively stable clinical conditions, as defined by the absence of hospitalisation or changes in medication within 6 months of the study. Participants were excluded if they: (1) were younger than 18 or older than 65, (2) had a history of brain trauma or neurological disease, (3) displayed symptoms of alcohol or substance abuse or dependence within 12 months of participating, or (4)



**Fig. 1.** Hypothesised mediation model of insight and psychosocial outcomes (i.e. self-esteem, depression and subjective QoL) via self-stigma. QoL, Quality of Life.


were unable to follow instructions. Of the 190 individuals initially approached for this study, ten declined to participate and another ten were unable to complete the cognitive tests, yielding a final sample of 170 subjects (89% of the initial sample). Ethical approval was obtained from the Institutional Review Board at the Tri-Service General Hospital National Defense Medical Center, Taiwan (ID: SS100-05). Following a comprehensive explanation of the study, participants were asked to provide written informed consent. Participation in the present study was strictly voluntary and anonymous.

Of the 170 participants, 93 (55%) were males and 147 (86%) were unmarried or single. Additionally, most participants were unemployed ( $n = 125$ , 73.5%). The mean age of the patients was 44.4 years ( $s.d. = 10.1$ ; range: 19–65), and the mean duration of formal

education was 12.3 years ( $s.d. = 2.69$ ; range: 6–18). The mean age at illness onset was 26.4 years ( $s.d. = 7.26$  years; range: 14–50), the mean illness duration was 17.9 years ( $s.d. = 10.0$  years; range: 2–45), and the average number of previous hospitalisations among the patients was 5.2 ( $s.d. = 3.93$ ; range: 0–20).

**Study measures**

Insight into illness was assessed using both an observer-rated instrument and a self-report scale. We administered the insight and judgement item (G12) of the Positive and Negative Syndrome Scale (PANSS; Kay *et al.* 1987), which provides a rating of 1 to 7 based on the global awareness of illness symptoms, need for treatment and consequences of illness. Insight was also assessed using the Chinese version (Kao & Liu, 2010) of the Self-Appraisal of illness Questionnaire (Marks *et al.* 2000), a self-report 17-item insight scale. This scale contains three subscales: the worry, need for treatment and presence/outcome of illness. Based on the suggestions of Marks *et al.* (2000), the need for treatment and presence/outcome of illness subscales were combined into a brief screening instrument for patients possibly at risk for poor insight. Essentially, the higher the score was, the higher the level of insight. In the present study, the internal consistency of this shortened insight scale was 0.88.



**Fig. 2.** Moderation model of insight and psychosocial outcomes (i.e. self-esteem, depression and subjective QoL) via self-stigma. QoL, Quality of Life.

Self-stigma was assessed using the Chinese version (Lien *et al.* 2015) of the Internalised Stigma of Mental Illness (ISMI) scale (Ritsher *et al.* 2003), which contains five subscales: alienation, stereotype endorsement, discrimination experience, social withdrawal and stigma resistance. The stigma resistance subscale was excluded because it reportedly lacks internal consistency and correlates poorly with the other ISMI subscales (Brohan *et al.* 2010; Lien *et al.* 2015). The aggregate score of the ISMI scale was modified according to the remaining 24 items to form a general index of self-stigma (Lysaker *et al.* 2007). Previous studies on self-stigma (Ritsher *et al.* 2003; Lysaker *et al.* 2007; Brohan *et al.* 2010) have considered scores of  $\geq 2.5$  as indicators of high self-stigma. In the present study, the Cronbach's alpha of self-stigma was 0.87.

To measure each variable of psychosocial outcomes, we employed several self-rating instruments (i.e. self-esteem, depression and subjective QoL) for all participants. First, the unidimensional self-report Rosenberg Self-Esteem Scale (RSES; Rosenberg, 1979) consists of ten items that are coded using a four-point Likert scale, identical to that used in the ISMI scale. Higher scores on the RSES indicate higher self-esteem levels. In a previous study, the Chinese version of the RSES exhibited high internal consistency, high test-retest reliability and adequate convergent validity (Cheng & Hamid, 1995). The scale also exhibited high internal consistency in the present study (Cronbach's alpha = 0.92). Second, the Beck Depression Inventory Revised (BDI-II) is a 21-item, self-report scale (Beck *et al.* 1996). The items comprise four statements that are scored on a scale of 0–3; high scores indicated high symptomatic severity. Respondents are instructed to describe their feelings during the previous 2 weeks. The results are scored by summing the responses to each of the items to obtain a total depression score ranging from 0 to 63. The psychometric properties of the Chinese version of the BDI-II were reviewed by Lu *et al.* (2002). For the present study, the Cronbach's alpha for reliability was 0.87. Finally, subjective QoL was assessed using the Chinese version of the Schizophrenia Quality of Life Scale Revision Four (Kuo *et al.* 2007), which is a self-administered questionnaire of 33 items in two domains: psychosocial and vitality. All but four items are coded on a scale of 0–4 according to the frequency of occurrence during the previous 7 days (0 = always, 4 = never; the four exceptions are coded 0 = never, 4 = always). A higher score indicates higher health-related QoL. In the present study, the Cronbach's alpha for subjective QoL was 0.85.

The PANSS (Kay *et al.* 1987) is a 30-item semi-structured interview scale that was developed for comprehensively assessing the symptoms of schizophrenia.

In this study, only two of the three analytically derived components were used: positive and negative symptoms (Cheng *et al.* 1996). The components contain seven items each; overall scores range from 7 to 49.

### Statistical analysis

Descriptive statistics and correlational analyses were performed using SPSS version 15.0 (SPSS Inc., Chicago, IL, USA). Independent *t* tests were used to assess between-groups differences in self-stigma. To examine the hypothesised relationships between insight, self-stigma and the psychological outcomes, we performed a path analysis by using SEM with AMOS 21 (IBM Corp). Our model was based on two latent variables, namely insight into illness and self-stigma. SEM is a general and powerful multivariate analysis technique with a confirmatory (i.e. hypothesis-testing) approach for analysing a structural theory bearing on a given phenomenon (Berkout *et al.* 2014). ISEM focuses on theoretical constructs, which are presented by the latent variables. The relationships between the theoretical constructs are represented by their regression or path coefficients (Kline, 2011). Because the multivariate normal distribution assumption of our data was not violated, full-information, maximum-likelihood estimation was conducted using raw data as the input (Mardia & Foster, 1983). Path analysis within SEM was used to test the proposed mediation model (Fig. 1). The direct relationship of insight to psychosocial outcomes, moderated by self-stigma, was examined using multigroup SEM to determine whether this direct effect was invariant across the two groups (low *v.* high levels of self-stigma). This method is critical when measuring invariance and within-groups comparisons (Deng & Yuan, 2015). In this study, we tested the invariance (e.g. factor loadings and path coefficients) of the proposed models across both low- and high-self-stigma groups (Fig. 2). The relative chi-square ( $\chi^2/df$ ) test, comparative fit index (CFI), Tucker–Lewis fit index (TLI) and root-mean-square error of approximation (RMSEA) were used to test the goodness-of-fit of the proposed model. An adequate fit between models and data is generally indicated by four fit indices (CFI > 0.95, TLI > 0.95, RMSEA < 0.08 and  $\chi^2/df < 2$ ) (Bentler & Bonett, 1980; Kline, 2011).

## Results

### Hypothesis testing

Descriptive analysis was performed and self-stigma differences were tested. To estimate the prevalence of high self-stigma, we chose to categorise groups as 'high self-

stigma' if the mean score was >2.5. The results revealed significant between-groups differences in the scores of the psychosocial outcome variables (self-esteem:  $t = -6.96, p < 0.001$ ; self-stigma:  $t = -6.64, p < 0.001$ ; subjective QoL:  $t = 8.66, p < 0.001$ ). Mean depression scores were significantly higher, whereas mean self-esteem and subjective QoL scores were significantly lower in the participants with high self-stigma ( $n = 67$ ) compared with those of low self-stigma ( $n = 103$ ).

Table 1 displays the bivariate zero-order correlations, means, standard deviations (s.d.), skewness and kurtosis of the indicator variables used to test the hypothesis. We identified a negative correlation between subjective insight and all psychosocial outcomes and a positive relationship between subjective insight and self-stigma. The direction of the associations of objective insight with self-stigma and the psychosocial outcome variables were the same as those with subjective insight, but not as strongly. Higher self-stigma was associated with less self-esteem and subjective QoL, but was associated with lower depression (all  $p < 0.01$ ). All the sociodemographic variables (i.e. sex, age, education and marital status) and clinical variables (i.e. illness onset, illness duration and number of previous hospitalisations) were non-significantly related to the dependent and independent variables in the SEM. Neither positive nor negative symptoms showed any significant correlations with our model variables. Thus, none of these variables were included in the SEM as control variables.

**Mediation analyses**

The results of the SEM for testing our hypothesised mediation model are presented in Fig. 3. The model-fit indices suggested that this model fitted the data well ( $\chi^2 = 33.3; df = 20; p = 0.03; \chi^2/df = 1.67; CFI = 0.98; TLI = 0.97; RMSEA = 0.06$ ). Figure 3 depicts the model, with significant paths denoted by solid lines and non-significant paths denoted by dotted lines. Three pathways were non-significant: (1) insight → self-esteem ( $\beta = -0.17, p = 0.21$ ), (2) insight → depression ( $\beta = -0.004, p = 0.97$ ) and (3) insight → subjective QoL ( $\beta = -0.16, p = 0.17$ ). Moreover, the previously significant correlation between insight and each psychosocial outcomes became non-significant. In summary, our findings indicated that self-stigma might fully mediate the association of insight with the psychosocial outcomes. Insight had only an indirect, negative effect on the psychosocial outcomes.


**Moderation analysis**

Because the differences in self-esteem, depression and subjective QoL between the high- and the low-self-

**Table 1.** Intercorrelations and descriptive statistics of all indices related to stigma, insight and psychosocial outcomes ( $n = 170$ )


	1	2	3	4	5	6	7	8	9	10	11	12	Mean (s.d.)	Skew	Kurtosis
1. Self-stigma (ISMI)	1												2.36 (0.52)	-0.23	0.28
2. Stereotype (ISMI)	0.85**	1											2.26 (0.52)	-0.09	0.48
3. Alienation (ISMI)	0.89**	0.69**	1										2.47 (0.60)	-0.34	0.09
4. Social withdrawal (ISMI)	0.90**	0.69**	0.73**	1									2.39 (0.61)	0.16	0.33
5. Discrimination (ISMI)	0.88**	0.69**	0.69**	0.81**	1								2.33 (0.61)	-0.05	-0.15
6. Subjective insight (SAIQ)	-0.28*	-0.21*	-0.35*	-0.23*	-0.17*	1							38.9 (4.87)	0.64	0.80
7. Objective insight (PANSS G12)	-0.17*	-0.17*	-0.19*	-0.08	-0.13	0.22**	1						3.07 (13.3)	0.26	-0.32
8. Depression (BDI-M)	0.55**	0.49**	0.51**	0.52**	0.44**	-0.30*	-0.12	1					13.0 (0.56)	1.42	2.11
9. Self-esteem (RSES)	-0.55**	-0.49**	-0.51**	-0.50**	-0.49**	-0.24*	-0.11	-0.58**	1				13.0 (5.47)	-0.16	0.11
10. Subjective QoL (SQLS-R4)	-0.63**	-0.51**	-0.59**	-0.58**	-0.53**	-0.32*	-0.17*	-0.70**	0.62**	1			54.6 (23.6)	0.53	0.49
11. Positive symptoms (PANSS)	-0.05	-0.06	-0.05	0.01	0.02	0.01	0.48**	-0.09	-0.06	-0.16*	1		18.3 (7.15)	0.40	-0.79
12. Negative symptoms (PANSS)	0.05	-0.02	0.02	0.10	0.13	-0.02	0.40**	-0.10	0.03	-0.07	0.77**	1	19.5 (8.17)	0.24	-1.27

ISMI, the Internalised Stigma of Mental Illness Scale; SAIQ, the Self-Appraisal of Illness Questionnaire; PANSS, the Positive and Negative Symptom Scale; BDI-II, the Beck Depression Inventory Revised; RSES, the Rosenberg Self-esteem Scale; SQLS-R4, the Schizophrenia Quality of Life Scale Revision Four.  
\* $p < 0.05$ ; \*\* $p < 0.01$ .



**Fig. 3.** The mediation effects of self-stigma within the relationship of insight and psychosocial outcome variables. Notes: Structural equation model: rectangles represent indicator or manifest variables; ovals, unobserved latent variables. Solid lines indicated significant paths; dashed lines, paths that did not reach the level of statistical significance. Numbers by single-headed arrows reflect standardised regression weights. R<sup>2</sup> represented the explained variance of the endogenous variables. \**p* < 0.05; \*\**p* < 0.01. PANSS, G12 item of the Positive and Negative Symptom Scale; SAIQ, the Self-Appraisal of illness Questionnaire; ISMI-1, Stereotype endorsement subscale of the Internalised Stigma of Mental Illness scale; ISMI-2, Alienation subscale of the Internalised Stigma of Mental Illness scale; ISMI-3, Social withdrawal subscale of the Internalised Stigma of Mental Illness scale; ISMI-4, Discrimination experience of the Internalised Stigma of Mental Illness scale; QoL, Quality of Life.

stigma groups were well established, a multigroup analysis was conducted. To assess the invariance of the proposed model, two models were fitted to the data. Model 1 (M1) assumed that the factor loadings for the latent variables of insight and self-stigma were invariant, and Model 2 (M2) assumed that both the factor loadings and structural path coefficients were invariant across the two groups. According to the results, the corresponding critical values of the relative goodness-of-fit indices were met for both M1 ( $\chi^2 = 24.23$ ; *df* = 14;  $\chi^2/\text{df} = 1.73$ ; *p* = 0.04; CFI = 0.93; TLI = 0.90; RMSEA = 0.07) and M2 ( $\chi^2 = 26.87$ ; *df* = 15;  $\chi^2/\text{df} = 1.79$ , *p* = 0.03; CFI = 0.92; TLI = 0.90; RMSEA = 0.07). Because the degree of fitness did not decrease significantly (Satorra–Bentler scaled  $\chi^2$  difference test score = 2.65,  $\Delta\text{df} = 1$ , *p* > 0.05) when all path coefficients were determined as equal in both groups (Fig. 4), the invariance of the proposed model was supported. This result indicated that the detrimental effect of insight on psychosocial outcomes is invariant across stigmas.



**Fig. 4.** The moderation models and standardised path coefficients. Results of multiple-group analyses with invariant factor loadings and path coefficients (M2) across both self-stigma groups (high/low), and the explained variances of the endogenous variables (R<sup>2</sup>). PANSS, G12 item of the Positive and Negative Symptom Scale; SAIQ, the Self-Appraisal of Illness Questionnaire; QoL, Quality of Life.

## Discussion

### *Detrimental effects of good insight and self-stigma in schizophrenia*

The current study is among the first to examine the mediator and moderator effects of self-stigma on the relationship between clinical insight and psychosocial outcomes in Chinese adults with schizophrenia spectrum disorders. The results of the current study are consistent with those of previous studies, indicating that patients with higher self- and expert-rated insight into illness report significantly poorer psychosocial outcomes (Cavelti *et al.* 2012a; Belvederi Murri *et al.* 2015; Margariti *et al.* 2015). These findings suggest that insight might be a 'double-edged sword' for people with schizophrenia and might complicate their understanding of the impacts on clinical outcomes. Moreover, these findings confirm the evidence concerning the relationship between self-stigma and poor psychosocial outcomes (Yanos *et al.* 2008; Livingston & Boyd, 2010). The results from this study provide further corroborating evidence that self-stigma, which is prevalent and problematic among individuals with schizophrenia, has a substantial and direct impact on psychosocial outcomes among people with schizophrenia. By contrast, our findings do not signify that insight is not a critical variable. Insight has long been considered essential for engagement in treatment, psychotherapeutic progress and a more accurate prognosis and has therefore been identified as a critical clinical outcome measure in schizophrenia (Amador *et al.* 1994; Lysaker *et al.* 2002). However, relevant studies are characterised by conflicting results (Mintz *et al.* 2003; Karow *et al.* 2008; Lysaker *et al.* 2009).

### *Mediating role of self-stigma*

The path analysis in this study supports the mediating role of self-stigma on the relationship between insight and psychosocial outcomes among people with schizophrenia. Our findings indicate that high insight into illness indirectly influences psychosocial outcomes through self-stigma (i.e. none of the direct effects of high insight on psychosocial outcomes are due to a relationship between insight and self-stigma), and self-stigma influences the direct effect of insight on psychosocial outcomes in people with schizophrenia. The results also confirm that mental-illness stigma is a major source of stress in addition to its effects on patients' clinical outcomes. Hence, reducing self-stigma would help to ameliorate some of the deleterious effects of high insight on psychosocial outcomes among people with schizophrenia, and reducing self-stigma alone appears to be sufficient to achieve this.

Improved insight not only allows individuals to acknowledge the symptoms and consequences of their mental illness as well as the need to receive treatment, but also enables them to adopt the stigmatising views (e.g. self as dangerous, self as incompetent) of society toward individuals with mental illnesses (Mak & Wu, 2006; Lysaker *et al.* 2013a). Such individuals frequently suffer additional enhanced insight into their mental illness may be more inclined to experience a greater level of self-stigma, whereas those with poor insight may be more likely to avoid stigma-related stress and harm because they are unaware that they are being stigmatised.

However, the confirmation of the mediation model in the present study does not support the paradoxical nature of insight into mental illness (Lysaker *et al.* 2007), which refers to a series of conflicting empirical findings in which insight can cause both positive and negative outcomes. Therefore, high insight is related to worsening outcomes only in the presence of moderate to high self-stigma. Some cross-sectional studies have indicated that the relationships among insight, self-stigma and negative outcomes may be more effectively explained by using a moderator model (Lysaker *et al.* 2007; Staring *et al.* 2009) than by using a mediation model (Hasson-Ohayon *et al.* 2011, 2014; Cavelti *et al.* 2012a, b), although other cross-sectional (Yanos *et al.* 2008; Schrank *et al.* 2014) and prospective (Cavelti *et al.* 2014) studies have failed to confirm the moderation effect of self-stigma. This discrepancy in findings may be due to the high level of self-stigma in our sample as well as the differing sociocultural contexts of the samples. In this study, we found that our participants reported higher mean scores of self-stigma than those reported in other studies in Western (Lysaker *et al.* 2007; Sibitz *et al.* 2011a, b) and Chinese (Lv *et al.* 2013) societies. With the cut-off score of 2.5, our study demonstrated high levels of self-stigma (45%), a percentage twice as high as that in Lv's *et al.* (2013) study (20%) in Hong Kong and which was also higher than that previously reported (Boyd *et al.* 2014).

### *Moderating role of self-stigma*

From another perspective, the model illustrating the associations between insight and psychosocial outcomes was invariant across stigma groups, which is in contrast to the study of Staring *et al.* (2009), who discovered that patients with high insight accompanied by stigmatising beliefs had the highest risk of low QoL, negative self-esteem and depression. In comparison with the mediation model, which outlined a linear process, the moderation model was employed to determine under which situations insight caused positive

and negative outcomes. Studies on patients with mental illnesses have revealed that high insight in those with low self-stigma engendered a positive outcome. However, high insight in individuals with high self-stigma produced a negative outcome (Lysaker *et al.* 2007; Staring *et al.* 2009). Thus, the moderation model illustrates the possibility of improving insight without deteriorating psychosocial outcomes such as mood, self-esteem and QoL, whereas the mediation model does not elaborate such possibility (Hasson-Ohayon *et al.* 2014). Based on this result, the circumstances under which insight might result in positive outcomes remain unclear. Other variables such as social support and coping strategies might moderate the relation between insight and psychosocial outcomes, thus necessitating further research aimed at comparing the role of these variables relative to the present findings.

### Limitations and perspectives

First, this study was cross-sectional and should be replicated using longitudinal methods (Maxwell & Cole, 2007). Second, we used only a single PANSS item to measure insight objectively and did not consider the dimensions distinctive of clinical insight. Third, self-stigma was the only potential mediator examined in the present study, although other mediators are likely to have important roles in the association between insight and psychosocial outcomes; other variables such as social cognition or metacognition could be examined (Lysaker *et al.* 2013b). Finally, we used the BDI-II to measure depression in individuals with schizophrenia. However, the claim that the BDI is a valid instrument for assessing depressive symptoms in chronic schizophrenia requires qualification because of the overlap of negative and extrapyramidal symptoms of schizophrenia and depression (Kim *et al.* 2006; Chemerinski *et al.* 2008).

### Conclusions

Early recognition and psychoeducational programmes for patients at risk of poor insight are particularly crucial, given the findings from previous research highlighting the potential benefits of such treatments (Xia *et al.* 2011; Lysaker *et al.* 2013a; Chien & Thompson, 2014). However, it appears that interventions aimed at improving insight may be insufficient for overcoming the deleterious effects of improved insight because these interventions fail to address emotional, interpersonal and stigma-related problems (Valiente *et al.* 2011). Thus, additional interventions such as cognitive behavioural therapy (Corrigan & Calabrese, 2005; Yanos *et al.* 2008, 2010, 2011) are required to facilitate positive responses to stigma, shame and social

exclusion in people with schizophrenia. Enhancing personal empowerment or self-esteem might also reduce self-stigma (Brohan *et al.* 2010; Mittal *et al.* 2012).

### Acknowledgements

The authors would like to express their sincere thanks to Professor Ritsher, the original ISMI scale designers, for their permission to administer the ISMI scale in our study. We also thank all participants who kindly volunteered to take part in this study.

### Financial Support

This study was supported by grants from the Ministry of Science and Technology, Taiwan (grant numbers NSC 101-2314-B-003-002 and 102-2410-H-003-016). This article was subsidised by the National Taiwan Normal University (NTNU), Taiwan.

### Conflict of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

### Availability of data and materials

Due to ethical restrictions, data are only available upon request. Although our data are anonymised (medical record number), schizophrenia is an infrequent and highly stigmatising mental disorder, which is why we are concerned about placing the data into an online repository. These concerns are shared by the lead IRB of this project, in particular as the informed consent form that all patients signed does not include the possibility of an online repository of individual patient data. Interested researchers may submit requests for minimised anonymous datasets to the corresponding author.

### References

- Amador XF, Flaum M, Andreasen NC, Strauss DH, Yale SA, Clark SC, Gorman JM (1994). Awareness of illness in schizophrenia and schizoaffective and mood disorders. *Archives General Psychiatry* 51, 826–836.
- American Psychiatric Association (2000). *Diagnostic and Statistical Manual of mental Disorders*, 4th edn, Text Revision. American Psychiatric Association: Washington, DC.
- Baron RM, Kenny DA (1986). The moderator-mediator variable distinction in social psychological research:



- conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology* **51**, 1173–1182.
- Beck AT, Steer RA, Ball R, Ranieri W** (1996). Comparison of Beck Depression Inventories-IA and -II in psychiatric outpatients. *Journal of Personality Assessment* **67**, 588–597.
- Belvederi Murri M, Respino M, Innamorati M, Cervetti A, Calcagno P, Pompili M, Lamis DA, Ghio L, Amore M** (2015). Is good insight associated with depression among patients with schizophrenia? Systematic review and meta-analysis. *Schizophrenia Research* **162**, 234–247.
- Bentler PM, Bonett DG** (1980). Significance tests and goodness-of-fit in the analysis of covariance structures. *Psychological Bulletin* **88**, 588–600.
- Berkout OV, Gross AM, Young J** (2014). Why so many arrows? Introduction to structural equation modeling for the novice user. *Clinical Child and Family Psychology Review* **17**, 217–229.
- Bora E, Schitoglu G, Asiler M, Atabay I, Veznedaroglu B** (2007). Theory of mind and unawareness of illness in schizophrenia: is poor insight a mentalizing deficit? *European Archives of Psychiatry and Clinical Neuroscience* **257**, 104–111.
- Boyd JE, Adler EP, Otilingam PG, Peters T** (2014). Internalized Stigma of Mental Illness (ISMI) scale: a multinational review. *Comprehensive Psychiatry* **55**, 221–231.
- Breitborde NJ, Srihari VH, Pollard JM, Addington DN, Woods SW** (2010). Mediators and moderators in early intervention research. *Early Intervention Psychiatry* **4**, 143–152.
- Brohan E, Elgie R, Sartorius N, Thornicroft G** (2010). Self-stigma, empowerment and perceived discrimination among people with schizophrenia in 14 European countries: the GAMIAN-Europe study. *Schizophrenia Research* **122**, 232–238.
- Cavelti M, Kvrđic S, Beck EM, Rüşch N, Vauth R** (2012a). Self-stigma and its relationship with insight, demoralization, and clinical outcome among people with schizophrenia spectrum disorders. *Comprehensive Psychiatry* **53**, 468–479.
- Cavelti M, Kvrđic S, Beck EM, Kossowsky J, Vauth R** (2012b). The role of subjective illness beliefs and attitude toward recovery within the relationship of insight and demoralization among people with schizophrenia spectrum disorders. *Journal of Clinical Psychology* **68**, 462–476.
- Cavelti M, Rüşch N, Vauth R** (2014). Is living with psychosis demoralizing? Insight, self-stigma, and clinical outcome among people with schizophrenia across 1 year. *The Journal of Nervous and Mental Diseases* **202**, 521–529.
- Chemerinski E, Bowie C, Anderson H, Harvey PD** (2008). Depression in schizophrenia: methodological artifact or distinct feature of the illness? *The Journal of Neuropsychiatry and Clinical Neurosciences* **20**, 431–440.
- Cheng JJ, Ho H, Chang CJ, Lan SY** (1996). Positive and Negative Syndrome Scale (PANSS): establishment and reliability study of a Mandarin Chinese language version. *Taiwanese Journal of Psychiatry* **10**, 251–258.
- Cheng ST, Hamid PN** (1995). An error in the use of translated scales: the Rosenberg Self-Esteem Scale for Chinese. *Perceptual and Motor Skills* **81**, 431–434.
- Chien WT, Thompson DR** (2014). Effects of a mindfulness-based psychoeducation programme for Chinese patients with schizophrenia: 2-year follow-up. *British Journal of Psychiatry* **205**, 52–59.
- Corrigan PW, Calabrese JD** (2005). Strategies for assessing and diminishing self-stigma. In *On the Stigma of Mental Illness: Practical Strategies for Research and social Change* (ed. PW Corrigan), pp. 239–256. American Psychological Association: Washington, DC.
- Deng L, Yuan KH** (2015). Multi-group analysis for structural equation modeling with dependent samples. *Structural Equation Modeling: a Multidisciplinary Journal* **22**, 552–567.
- Hasson-Ohayon I, Kravetz S, Meir T, Rozencwaig S** (2009). Insight into severe mental illness, hope, and quality of life of persons with schizophrenia and schizoaffective disorders. *Psychiatry Research* **167**, 231–238.
- Hasson-Ohayon I, Levy I, Kravetz S, Vollanski-Narkis A, Roe D** (2011). Insight into mental illness, self-stigma, and the family burden of parents of persons with a severe mental illness. *Comprehensive Psychiatry* **52**, 75–80.
- Hasson-Ohayon I, Tuval-Mashiach R, Morag-Yaffe M, Gaziel M, Schapir L, Zalsman G, Shoval G** (2014). Parents of adolescents with psychiatric disorders: insight into the disorder, self-stigma and parental stress. *Comprehensive Psychiatry* **55**, 1106–1110.
- Hopwood CJ** (2007). Moderation and mediation in structural equation modeling: applications for early intervention research. *Journal of Early Intervention* **29**, 262–272.
- Kao YC, Liu YP** (2010). The clinical applicability of the Self-Appraisal of Illness Questionnaire (SAIQ) to chronic schizophrenic patients. *Psychiatric Quarterly* **81**, 215–225.
- Karow A, Pajonk FG, Reimer J, Hirdes F, Osterwald C, Naber D, Moritz S** (2008). The dilemma of insight into illness in schizophrenia: self- and expert-rated insight and quality of life. *European Archives of Psychiatry and Clinical Neuroscience* **258**, 152–159.
- Kay SR, Fiszbein A, Opler LA** (1987). The Positive and Negative Syndrome Scale (PANSS) for schizophrenia. *Schizophrenia Bulletin* **13**, 261–276.
- Kim SW, Kim SJ, Yoon BH, Kim JM, Shin IS, Hwang MY, Yoon JS** (2006). Diagnostic validity of assessment scales for depression in patients with schizophrenia. *Psychiatry Research* **144**, 57–63.
- Kline RB** (2011). *Principles and Practice of Structural Equation Modeling*, 3rd edn. The Guilford Press: New York.
- Kuo PJ, Chen-Sea MJ, Lu RB, Chung MS, Kuo CC, Huang WC** (2007). Validation of the Chinese version of the Schizophrenia Quality of Life Scale Revision 4 (SQLS-R4) in Taiwanese patients with schizophrenia. *Quality Life Research* **16**, 1533–1538.
- Lien YJ, Kao YC, Liu YP, Chang HA, Tzeng NS, Lu CW, Loh CH** (2015). Internalized stigma and stigma resistance among patients with mental illness in Han Chinese population. *Psychiatric Quarterly* **86**, 181–197.
- Livingston JD, Boyd JE** (2010). Correlates and consequences of internalized stigma for people living with mental illness stigma: a systemic review and meta-analysis. *Social Science & Medicine* **71**, 2150–2161.

- Lu ML, Che SS, Chang SW, Shen WW (2002). Reliability and validity of the Chinese version of the Beck Depression Inventory-II. *Taiwanese Journal of Psychiatry* **16**, 301–310.
- Lv Y, Wolf A, Wang X (2013). Experienced stigma and self-stigma in Chinese patients with schizophrenia. *General Hospital Psychiatry* **35**, 83–88.
- Lysaker PH, Bryson GJ, Bell MD (2002). Insight and work performance in schizophrenia. *The Journal of Nervous and Mental Diseases* **190**, 142–146.
- Lysaker PH, Roe D, Yanos PT (2007). Toward understanding the insight paradox: internalized stigma moderates the association between insight and social functioning, hope, and self-esteem among people with schizophrenia spectrum disorders. *Schizophrenia Bulletin* **33**, 192–199.
- Lysaker PH, Buck KD, Salvatore G, Popolo R, Dimaggio G (2009). Lack of awareness of illness in schizophrenia: conceptualizations, correlates and treatment approaches. *Expert Review of Neurotherapeutics* **9**, 1035–1043.
- Lysaker PH, Vohs J, Hillis JD, Kukla M, Popolo R, Salvatore G, Dimaggio G (2013a). Poor insight into schizophrenia: contributing factors, consequences and emerging treatment approaches. *Expert Review of Neurotherapeutics*, **13**, 785–793.
- Lysaker PH, Vohs J, Hasson-Ohayon I, Kukla M, Wierwille J, Dimaggio G (2013b). Depression and insight in schizophrenia: comparisons of levels of deficits in social cognition and metacognition and internalized stigma across three profiles. *Schizophrenia Research* **148**, 18–23.
- Mak WW, Wu CF (2006). Cognitive insight and causal attribution in the development of self-stigma among individuals with schizophrenia. *Psychiatric Services* **57**, 1800–1802.
- Mardia KV, Foster K (1983). Omnibus tests of multinormality based on skewness and kurtosis. *Communications in Statistics -Theory and Methods* **12**, 207–221.
- Margariti M, Ploumpidis D, Economou M, Christodoulou GN, Papadimitriou GN (2015). Quality of life in schizophrenia spectrum disorders: associations with insight and psychopathology. *Psychiatry Research* **225**, 695–701.
- Marks KA, Fastenau PS, Lysaker PH, Bond GR (2000). Self-Appraisal of Illness Questionnaire (SAIQ): relationship to researcher-rated insight and neuropsychological function in schizophrenia. *Schizophrenia Research* **45**, 203–211.
- Maxwell SE, Cole DA (2007). Bias in cross-sectional analyses of longitudinal mediation. *Psychological Methods* **12**, 23–44.
- Mintz AR, Dobson KS, Romney DM (2003). Insight in schizophrenia: a meta-analysis. *Schizophrenia Research* **61**, 75–88.
- Mittal D, Sullivan G, Chekuri L, Allee E, Corrigan PW (2012). Empirical studies of self-stigma reduction strategies: a critical review of the literature. *Psychiatric Services* **63**, 974–981.
- Mohamed S, Rosenheck R, He H, Yuping N (2014). Insight and attitudes towards medication among inpatients with chronic schizophrenia in the US and China. *Social Psychiatry & Psychiatric Epidemiology* **49**, 1063–1070.
- Ritsher J, Otilingam PG, Grajales M (2003). Internalized stigma of mental illness: psychometric properties of a new measure. *Psychiatry Research* **121**, 31–49.
- Rosenberg M (1979). *Conceiving the Self*. Basic Books: New York.
- Schrank B, Amering M, Hay AG, Weber M, Sibitz I (2014). Insight, positive and negative symptoms, hope, depression and self-stigma: a comprehensive model of mutual influences in schizophrenia spectrum disorders. *Epidemiology and Psychiatric Sciences* **23**, 271–279.
- Sibitz I, Unger A, Woppmann A, Zidek T, Amering M (2011a). Stigma resistance in patients with schizophrenia. *Schizophrenia Bulletin* **37**, 316–323.
- Sibitz I, Amering M, Unger A, Seyringer ME, Bachmann A, Schrank B, Benesch T, Schulze B, Woppmann A (2011b). The impact of the social network, stigma, and empowerment on the quality of life in patients with schizophrenia. *European Psychiatry* **26**, 28–33.
- Staring AB, Van der Gaag M, Van den Berge M, Duivenvoorden HJ, Mulder CL (2009). Stigma moderates the associations of insight with depressed mood, low self-esteem, and low quality of life in patients with schizophrenia spectrum disorders. *Schizophrenia Research* **115**, 363–369.
- Valiente C, Provencio M, Espinosa R, Chaves C, Fuentesnebro F (2011). Predictors of subjective well-being in patients with paranoid symptoms: is insight necessarily advantageous? *Psychiatry Research* **189**, 190–194.
- Xia J, Merinder LB, Belgamwar MR (2011). Psychoeducation for schizophrenia. *Cochrane Database of Systemic Reviews* **15**, CD002831.
- Xiang YT, Wang Y, Wang CY, Chiu HF, Chen Q, Chan SS, Lai KY, Lee EH, Ungvari GS (2012). Association of insight with sociodemographic and clinical factors, quality of life, and cognition in Chinese patients with schizophrenia. *Comprehensive Psychiatry* **53**, 140–144.
- Yang LH, Purdie-Vaughns V, Kotabe H, Link BG, Saw A, Wong G, Phelan JC (2013). Culture, threat, and mental illness stigma: identifying culture-specific threat among Chinese-American groups. *Social Science & Medicine* **88**, 56–67.
- Yanos PT, Roe D, Markus K, Lysaker PH (2008). Pathways between internalized stigma and outcomes related to recovery in schizophrenia spectrum disorders. *Psychiatric Services* **59**, 1437–1442.
- Yanos PT, Roe D, Lysaker PH (2010). The impact of illness identity on recovery from severe mental illness. *American Journal of Psychiatric Rehabilitation* **13**, 73–93.
- Yanos PT, Roe D, Lysaker P (2011). Narrative enhancement and cognitive therapy: a new group-based treatment for internalized stigma among persons with severe mental illness. *International Journal of Group Psychotherapy* **61**, 577–595.