

Research Article

Cite this article: Liao L-Z, Lin J-N, Ding X-S, Feng S, Yan T (2024). *Karenia mikimotoi* induced adverse impacts on abalone *Haliotis discus hannai* in Fujian coastal areas, China. *Journal of the Marine Biological Association of the United Kingdom* **104**, e47, 1–9. <https://doi.org/10.1017/S0025315424000353>

Received: 6 October 2023

Revised: 17 February 2024

Accepted: 18 March 2024

Keywords:

adverse impacts; abalone; *Haliotis discus hannai*; *Karenia mikimotoi* bloom

Corresponding author:

Tian Yan;

Email: tianyancas@163.com;

Jia-Ning Lin;

Email: linjianing@sdu.edu.cn

Karenia mikimotoi induced adverse impacts on abalone *Haliotis discus hannai* in Fujian coastal areas, China

Ling-Zhi Liao^{1,2} , Jia-Ning Lin^{1,2,3} , Xin-Shu Ding^{1,2}, Song Feng³ and Tian Yan³

¹School of Environmental Science and Engineering, Shandong University, Qingdao 266237, PR China;

²Institute of Eco-Environmental Forensics, Shandong University, Qingdao 266237, PR China and ³CAS Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

Abstract

Large-scale outbreaks of the dinoflagellate *Karenia mikimotoi* caused substantial mortality of abalone, *Haliotis discus hannai* in Fujian, China in 2012, resulting in 20 billion in economic losses to abalone industries. However, the mechanism behind the mortality, especially the reaction of abalone to this microalgal toxicity, which possibly differed significantly from the former ‘fish killer’ strain in the South China Sea (SCS). Our study revealed that *K. mikimotoi* FJ-strain exhibited a four-fold higher haemolytic toxicity than the SCS-strain during the late exponential phase. At the microalgal cell density of 3×10^7 cell L⁻¹, the FJ-strain caused abalone mortality of 67% in 48 h, with decreased granulocyte–hyalinocytes ratio and phagocytic activity by 58.96% and 75.64%, respectively, increased haemocyte viability by 4.8-fold and severe gill damage. The toxic effect only worked for the haemolytic toxicity from active algal cells, which were probably produced under the contact of algal cells and abalone gills. However, under exposure to the SCS-strain, more than 80% of individuals survived under aeration. The results indicated that FJ-strain was a new *K. mikimotoi* ecotype with stronger toxicity. It evoked severe effects, with complete abalone mortality within 24 h under the cascading effect of non-aeration (dissolved oxygen declined to 2.0 mg L⁻¹), when exposed to *K. mikimotoi* FJ-strain at the above density. Thus, apart from the microalgal toxicity, DO depletion exacerbated the mortality of abalone in the experiment. The massive abalone mortalities in Fujian were probably caused by the combination of microalgal toxic effects and oxygen depletion, leading to immunological depression and histopathological disruption.

Introduction

In the summer of 2012, an exceptionally massive bloom of *Karenia mikimotoi* occurred off the coastal waters of Fujian province, with a high biomass of 10^7 cells L⁻¹ and a long duration of 21 d (SOAC, 2013). During such outbreaks of *K. mikimotoi*, mass mortality of abalone, *Haliotis discus hannai* was observed in the farming regions and, consequently more than \$330 million was lost (SOAC, 2013), which may be one of the most serious harmful algal bloom (HAB) economic losses recorded throughout the world. The large scale and huge economic impact made further research into the bloom essential.

In fact, *K. mikimotoi* has always been one of the most prevalent HAB species along Chinese coasts in recent decades (Zhou, 2010). Previously, the *K. mikimotoi* bloom was known as a famous ‘fish killer’ in China and frequently appeared in the South China Sea (Qi *et al.*, 2004; Ou, 2006; SOAC, 2006; Li *et al.*, 2009). Most seriously, it caused almost all of the cultured fish in cages to be killed in the adjacent coastal areas near Pearl estuary and was also associated with nearly \$60 million in losses (Yang and Hodgkiss, 2004). However, this outbreak in Fujian unexpectedly killed the abundant abalone instead of fish. It was thought possible that the lethal mechanisms of *K. mikimotoi* from Fujian differed from previous blooms, in particular the nature of the microalgal toxicity between them, given their various geographical biotope around China’s coasts.

The cases of *Karenia* spp. blooms causing the death of shellfish in the wild have been long-known in Japan, South Africa, and other countries (Table 1). Except for the toxic effect of autologous algal toxicity, the dissolved oxygen (DO) depletion resulted from *Karenia* spp. blooms were also a major factor in the mortality of shellfish (Yasumoto *et al.*, 1990; Satake and Tanaka, 2005; Zou *et al.*, 2010; O’Boyle *et al.*, 2016). In Fujian, abalones were cultured densely and abundantly in offshore rafts along the coast, and these also suffered from hypoxia in the presence of *K. mikimotoi* blooms. Therefore, it was necessary to further determine whether massive mortalities of abalones were associated with the microalgal toxicity of *K. mikimotoi* or DO depletion during the blooms.

A similar type of abalone mortality, when exposed to *Karenia cristata* and *Karenia concordia* has been recorded (Botes *et al.*, 2003; Chang *et al.*, 2008), however, little is known about the reaction of abalone to *Karenia* hazards. During harmful algal blooms, the mortality of molluscs was usually due to damage to their main organs or immune system. Of these, gills are the most susceptible because of the direct contact with seawater for respiratory gas exchange.



Table 1. Records of shellfish species mortality due to *Karenia* species bloom

Algal species	Shellfish species	Location	Reference
<i>K. mikimotoi</i> (<i>Gyrodinium aureolum</i>)	<i>Littorina rudis</i> <i>Maton</i>	Maquoit Bay	Heinig and Campbell (1992)
	<i>Mytilus edulis</i>	Southern Ireland	Southgate <i>et al.</i> (1984)
	Annelids, molluscs	Norwegian	Tangen (1977);
	<i>Crassostrea gigas</i>	Scottish	Davidson <i>et al.</i> (2009)
	<i>Haliotis discus hannai</i>	China	SOAC (2013)
<i>Karenia concordia</i>	<i>Haliotis iris</i>	New Zealand	Chang <i>et al.</i> (2008)
<i>Karenia cristata</i>	<i>Haliotis midae</i>	South Africa	Botes <i>et al.</i> (2003)

ND, No data.

Harmful algae can cause some vacuolation, abnormality, and even shrinkage of mollusc gills (eg. *Mercenaria mercenaria*, *Ruditapes philippinarum*) (Basti and Segawa, 2010; Hégaret *et al.*, 2010). Additionally, haemocytes are crucial for the proper function of the immune system and are important health indicators. Exposure to HABs allows for the interaction with haemocytes found throughout the open circulatory system. The immune system is also easily destroyed by HABs through inducing haemocyte diapedesis (Haberkorn *et al.*, 2010; Medhioub *et al.*, 2012; Lassudrie *et al.*, 2014, 2015). Therefore, *K. mikimotoi* blooms in Fujian probably damaged the gills and immune systems of abalones leading to mass mortality.

To test the hypotheses above, we evaluated the differential algal character and toxicity on abalones between *K. mikimotoi* strains isolated from the South China Sea (*K. mikimotoi* – SCS strain) and Fujian coastal areas (*K. mikimotoi* – FJ strain). The immunological and histopathological changes in abalones were also investigated by *K. mikimotoi* – FJ at low bloom concentrations to determine the specific mechanisms of killing abalones.

Materials and methods

Algal species and abalone cultures

Algal cultures and juvenile abalones were used to determine the effects of exposure to *K. mikimotoi* on abalones. The dinoflagellate *K. mikimotoi* – FJ strain was isolated during the 2012 bloom in the Fujian coastal area, China, provided by the Centre for Collections of Marine Algae, Xiamen University. *K. mikimotoi* – SCS strain was isolated from the South China Sea in 2005 and provided by two National Basic Research Priority Programmes of China, CEOHAB-I and -II. The two strains were cultured in modified *f/2* medium in 5-L flasks, and prepared without the addition of silicate. The cultures were kept at 20 ± 1 °C on a 12:12 L:D cycle, with irradiance of $56 \mu\text{E m}^{-2} \text{s}^{-1}$.

Considering that a large amount of the juvenile abalone died during the algal bloom accident in Fujian, therefore, the juvenile abalone was used in this research. The juvenile abalone *H. hannai* was bought from Longwan Biotechnology Corporation, Qingdao, China. Abalones were transported to the laboratory within 2 h in small aerated tanks. Healthy abalones of similar size (initial weight 0.9 ± 0.1 g; shell length 21.0 ± 3.4 mm) were picked out in the laboratory and placed in several 100 L tanks filled with

filtered seawater and continuously aerated with air stones. The abalones were acclimated to the experimental condition for 7 days, during which they were fed daily on the macroalgae *Laminaria japonica*. All of the seawater in the tanks was completely renewed every day, and the tanks were cleaned every 2–3 days.

The natural seawater used in this study was pumped from Taipingjiao (a clean site with no known pollution history) at Qingdao and sand filtered prior to use in the laboratory. Prior to the experiments, the seawater was also subjected to filtration through a 0.45 mm pore-size cellulose nitrate membrane, boiling for sterilization. The pH of the seawater was measured using a HI991000 pH instrument. Salinity was adjusted to 31 ± 1 ‰ with distilled water as determined using an ATAGO hand-held refractometer.

Haemolytic activity

The preparation of intra- and extra-cellular lipophilic crude extracts: haemolytic activities of the two strains at different growth phases were measured by the protocol adapted from Zhou *et al.* (2011). For each strain, four batches of 3 L were cultured in a modified *f/2* medium in 5 L flasks. Initial cell numbers were estimated at 0.5×10^7 cells L^{-1} . Cells were counted every 2 days. Sampling was done at the following algal phases: pre-exponential phase, exponential phase, stationary phase, and the anaphase. At every phase, one batch for each strain was harvested and the cell density was recorded in Table 1. To detect the intra- and extra-cellular toxicity, ultrasonic ruptured cell suspensions (URCS) and cell-free culture supernatants (CFCS) were prepared from the algal culture, respectively. 3 L of culture was filtered through a 0.45 mm Whatman GF/C membrane at a negative pressure of 0.01 MPa. The filters and cells were washed with 100 mL of extraction liquid (chloroform: methanol: water = 13: 7: 5, v/v/v) and then ruptured in a sonicator for 30 min at 200 W, 0 °C, 5 : 5 s. The ruptured medium was observed by microscope to confirm that no cells existed and identified as URCS. URCS was centrifuged at $5000 \times g$ (4 °C) for 10 min. The supernatant was transferred into a separating funnel for the lipophilic liquid parts collection, which was then evaporated to dryness at 40 °C at negative pressure of 0.05 MPa. The dry, lipophilic crude extract was stored in 2 mL absolute methanol (final volume) at -20 °C till use. The filtering medium served as CFCS and was immediately filtered by SPE C18 Cartridges. The SPE was collected and eluted with 10 mL methanol with the concentration of 20%, 40%, 60%, 80%, and 100%, respectively. The final elutes were collected separately and stored at -20 °C until use.

Preparation of rabbit erythrocytes: The rabbit for the haemolytic assay was bought from Qingdao Bio-test and Test-animal Center in Qingdao, China, and acclimated in cages with water and food prior to the experiment. The erythrocytes were withdrawn by needle and syringe from the rabbit's ear and added to 10% sodium citrate to avoid clotting, which was centrifuged at $500 \times g$ (4 °C) for 10 min. After centrifugation, the supernatant was discarded. The erythrocytes were washed three times with phosphate-buffered saline (PBS) and diluted to a final concentration of 4% (v/v) in PBS for use within 24 h (Zhou *et al.*, 2011).

The hemolytic test: For the standardization of a known hemolytic agent, the standard curve between Saponin (Sigma) concentration and haemolytic activity was conducted prior to the test. The protocol set up nine saponin concentration medium, including 0.5, 1.0, 2.5, 4.5, 5.0, 6.0, 7.0, 8.0, and $9.0 \mu\text{g mL}^{-1}$. A triplicate of 150 μL saponin medium and 50 μL erythrocytes suspension were added to Clean 96-well V-bottom plates (Costar). Erythrocytes suspension in PBS buffer was performed as zero blank, and the erythrocytes with 1% Triton X-100 as

100% positive control. The 96-well plates were incubated at 30 °C for 60 min in an incubator, and then centrifuged at 1000 × g (4 °C) for 10 min. 150 µL supernatant were serially collected on other 96-well plates, and haemolysis was measured by a microplate reader at 550 nm absorbance. Haemolytic activity was calculated according to Zhou *et al.* (2011). Thus, the relationship between saponin concentration and haemolytic activity was performed.

The stocks of intra- and extra-cellular lipophilic crude extracts were diluted five-fold by PBS buffer. A triplicate of 150 µL samples and 50 µL erythrocytes suspension were added to Clean 96-welled V-bottom plates (Costar). Other steps were the same as those described above. According to the final haemolytic activity, the relative saponin concentration for each extract was calculated by the standard curve.

Acute toxicity test on abalones

Based on the bloom density occurring in the field, one high algal density (3×10^7 cells L⁻¹) and one relatively low algal density (1×10^7 cells L⁻¹) were used for each algal strain in the experiment. Algal cells were harvested at the early stationary phase. A 1 mL subsample was taken and counted under the microscope after fixation in Lugol's solution. Based on this cell count, the algae were diluted to the designated density.

The experiment was conducted in 3 L beakers. Each beaker contained 10 healthy juvenile individuals in 2.5 L of the prepared algal treatments. As a negative control, 10 individuals were cultured in 2.5 L of seawater alone. Every treatment consisted of three replicates, with 30 individuals in total. Each beaker was individually aerated using air stones and replaced with newly prepared algal treatment every day. To avoid the interaction between *K. mikimotoi* and the food macroalgae *L. japonica*, no food was offered during the experiment. The culture conditions were measured as follows: water temperature (19 ± 0.5 °C), pH value (7.5–8.2), ammonia concentration (no more than 0.1 mg L⁻¹, and dissolved oxygen (7.3–7.8 mg L⁻¹). Mortality was recorded at 12, 24, 36, and 48 h. A final mortality test was processed according to Reddy-Lopata *et al.* (2006). The abalone with no response by repeated stimulation could be judged as the dead. Dead individuals were removed immediately to avoid water contamination.

Constituent toxicity in *K. mikimotoi* was performed to further identify the algal toxicity origin. Intact cell suspension (ICS), URCS, CFCS, and methanol crude extracts (MCE) were prepared from fresh algal cells with a density of 3×10^7 cells L⁻¹, and the method was in accordance with Li *et al.* (2017). Thereafter, the prepared constituents were tested on abalones, respectively. Other conditions for the test were the same as those described above.

To identify whether the abalone mortality resulted from the combination of *K. mikimotoi* – FJ algal toxic effects and related oxygen depletion during bloom, another experiment was conducted in a non-aeration condition. No air stones were added during the experiment, with DO lower than 2.0 mg L⁻¹. Other environmental conditions and algal densities were the same as those described above.

Abalone gills and haemocyte parameters of *K. mikimotoi* FJ-strain

To evaluate the possible damage to the gill and haemocyte organs of abalones due to *K. mikimotoi* FJ-strain, another batch of healthy juvenile individuals was exposed to a low density of 1×10^7 cells L⁻¹ for 48 h under the same experimental conditions described above. After 48 h, five alive abalones from each treatment were randomly sampled for haemocyte parameters, gill configuration, and enzyme activity measurements.

Analysis of haemocyte morphology and function was done on haemolymph extracted from the prepared abalones above. Haemolymph was withdrawn by needle and syringe from the blood sinus by cutting the adductor muscle with a scalpel. During the collection, the haemolymph samples were stored temporarily on ice and immediately used for immune function measurement by flow cytometry (Accuri 6, BD Biosciences, San Diego, CA). According to Hégaret *et al.* (2003a, 2003b), we measured three main immune function parameters, including haemocyte characterization, phagocytosis activity, and haemocyte viability.

Configuration of abalone gill and the enzyme activity changes were also examined. Histological photos recorded the effects of *K. mikimotoi* on abalone gills. The abalone gill specimens were first fixed in 10% buffered formalin and then routinely processed for the preparation of histological slides, which was conducted according to Dutra *et al.* (2017). The images of the gills were captured under a microscope (Leica DM2000, Germany). Another batch of fresh abalone gills were collected for the major enzyme activity measurement involved in immune responses and energy transfer, including superoxide dismutase (SOD), Catalase (CAT), Na⁺-K⁺-ATPase, Ca²⁺-Mg²⁺-ATPase. The enzyme was measured using a commercialized kit (Jiancheng, Nanjing, China), according to the manufacturer's instructions. SOD and CAT were expressed as U/mg protein. Na⁺-K⁺-ATPase and Ca²⁺-Mg²⁺-ATPase were expressed as µmol Pi/mg protein/hour.

Data analysis

Data were analyzed using the Excel 2010, Origin 8.5, and SPSS 16.0 software packages. Due to the different cell density gradients for every alga, the statistical analysis of the data hardly corresponded to a two-way ANOVA with algal species and algal density as factors. Therefore, one-way analysis of variance (ANOVA) was applied for statistical evaluations. Prior to statistical analysis, all data were tested for normality and homogeneity (SPSS 16.0). Mean differences were considered to be significant at the 0.05 level. If the overall ANOVA results were significant, a Fisher's least significance difference (LSD) post-hoc test was performed to test among experimental combinations.

Results

Various haemolytic activity between *K. mikimotoi* FJ- and SCS-strain

The preliminary assays showed that ultrasonic ruptured cell suspensions (URCS) and cell-free culture supernatants (CFCS) from the two strains all had haemolytic activity (Figure 1). For the URCS extract, there was no significant difference during the pre- and exponential phases in the two strains ($P > 0.05$). However, during the stationary phase and the anaphase, the relative concentration of saponin became significantly higher in FJ-strain ($P < 0.05$), which exceeded 3.27, and 4.38 folds, respectively, to those in the SCS-strain. For the CFCS extract, the relative concentration of saponin remained significantly higher in the SCS-strain than those in FJ-strain ($P < 0.05$) during the pre- and exponential phase, however, which declined as the population reached the stationary phase and anaphase, with no significant difference in the two strains ($P > 0.05$). Totally, at the high bloom density *in situ* (above 3×10^7 cells L⁻¹, around stationary phase), the FJ-strain exhibited more haemolytic toxicity than SCS-strain.

Differential toxicity of *K. mikimotoi* FJ- and SCS- strain to abalones

The mortality of *H. hannai* eventually increased significantly with the increasing bloom densities of *K. mikimotoi*-FJ during




Figure 1. Haemolytic activity of *K. mikimotoi* FJ- and SCS- strain during different algal phases (URCS: ultrasonic ruptured cell suspensions ; CFCS: cell-free culture supernatants; algal phases 1, 2, 3, and 4: pre-exponential phase, exponential phase, stationary phase, and anaphase, respectively; each data point represents the mean percentage \pm standard deviation of triplicate cultures).

the aeration. The high algal density even led to the sharp fatality in a short period (e.g. mortality rate of 67% in 48 h at 3×10^7 cells L^{-1}). More seriously, all the abalones would rapidly die out at the algal density of 3×10^7 cells L^{-1} within 24 h through the cascading effect of non-aeration (DO below $2.0 \text{ mg } L^{-1}$, Figure 2). In contrast, *K. mikimotoi* SC-strain showed weaker lethal effects on abalones. Even at high bloom density (3×10^7 cells L^{-1}), over 90% of individuals still survived at the end of the experiment (up to 48 h) during the aeration (Figure 3).

To further identify the algal toxicity origin in *K. mikimotoi*-FJ, algal constituent toxicity was performed. Intact cell suspensions (ICS) showed toxic effects on abalones, with mortality rates of

66.67% at 48 h at equal algal density (3×10^7 cells L^{-1}). However, when exposed to CFCS, URCS, and methanol crude extracts (MCE) of algal cells, no abalone individual died at 48 h (Figure 4).

Specific effects on abalones by *K. mikimotoi* FJ-strain

Under the exposure to a relatively low bloom density of *K. mikimotoi* FJ-strain (1×10^7 cells L^{-1}), abalones exhibited significant histopathological and immunological changes despite high mortality rates. The normal configuration of the gill was disrupted including apparent shrinkage and deformation of the afferent and efferent branchial vessels in comparison to the control groups




Figure 2. Survival rates of abalones exposed to *K. mikimotoi*-FJ strain at bloom density during the aeration and non-aeration condition (each data point represents the mean percentage \pm standard deviation of triplicate cultures).




Figure 3. Survival rates of abalones exposed to *K. mikimotoi*-SCS strain at bloom density during the aeration and non-aeration condition (each data point represents the mean percentage \pm standard deviation of triplicate cultures).

(Figures 5A, and B). Gill filaments were also found broken (Figure 5C). All major enzymes in the gill involved in immune responses and energy transfer (SOD, CAT, $\text{Na}^+\text{-K}^+\text{-ATPase}$, $\text{Ca}^{2+}\text{-Mg}^{2+}\text{-ATPase}$) showed inhibited activities (Figure 6), which were significantly ($p < 0.05$) decreased by 21.46%, 62.99%, 46.32%, 50.34% in 48 h, respectively.

In the response of abalone haemocytes (Figure 7), granulocytes with respect to hyalinocyte type cells and phagocytosis activities declined significantly by 58.96%, and 75.64% ($p < 0.05$), respectively, with respect to control groups. The rate of haemocyte mortality was significantly increased by 4.8 folds ($p < 0.05$).

Discussion

Specific algal toxicity effects among *K. mikimotoi* strains from various biotopes

K. mikimotoi has dominated coastal waters around Europe, Japan, China, and other countries recently. (Haywood *et al.*, 2004; Taylor

et al., 2004; Steidinger *et al.*, 2008) with these populations being commonly associated with fish and marine invertebrate mortalities. In the present research, we identified the major differences in the toxicity potency and toxic effects of the FJ and SCS strains of *K. mikimotoi* on abalones. Consistent with these findings, Zou *et al.* (2010) also compared the haemolytic activity and effects on rotifers by two *K. mikimotoi* strains (SUO-1 and FUK) isolated from Japan. The SUO-1 strain (2×10^7 cells L^{-1}) showed strong haemolytic activity and high toxicity to rotifers, whereas the FUK strain (2×10^7 cells L^{-1}) was less toxic. The different effects were also observed on the brine shrimp *Artemia salina*. Sun *et al.* (2010) and Dang *et al.* (2015) reported that the SCS strain (1.5×10^7 cells L^{-1}) led to all *A. salina* dying at 96 h, however, the mortality rate was just 23% when exposed to FJ strain (3×10^7 cells L^{-1} , Li *et al.*, 2017). In our laboratory, the ITS sequences comparison also identified differences between FJ and SCS strains (Zhang QC, unpublished data). It was possible that there was a range of *K. mikimotoi* ecotypes with varying toxicities and modes of action.




Figure 4. Survival rates of abalones exposed to different algal constituents in *K. mikimotoi*-FJ strain (ICS: intact cell suspension; URCS: ultrasonic ruptured cell suspensions; CFCS: cell-free culture supernatants; MCE: methanol crude extracts; each data point represents the mean percentage \pm standard deviation of triplicate cultures).




Figure 5. The configuration of abalone gills in the control and experimental groups (A: the normal branchial vessel in control group; B: apparent shrink and deformation of afferent and efferent branchial vessel in experimental group at the algal bloom density; C: broken gill filaments in experimental group at the algal bloom density, as the arrow showed).

To test whether the characters of isolates from other geographical areas were discriminating or not, we also documented haemolytic toxin production and the adverse impacts of different *K. mikimotoi* strain worldwide on a range of marine organisms (Table 2). Although the toxic effects of *K. mikimotoi* JP isolates were just reported on rotifers, there was much comprehensive data on various marine invertebrates by other *K. mikimotoi* strains. Interestingly, many similarities were observed among the three *K. mikimotoi* strains (CH-FJ, NZ, and EU isolates) in the toxic effects. No acute effects of the three *K. mikimotoi* strains were observed for the brine shrimp *A. salina* tested, even at the algal density of $5\text{--}6 \times 10^7$ cells L^{-1} (Botes et al., 2003; Neely and Campbell, 2006; Shi et al., 2012; Li et al., 2017). However, lethal effects on abalone juveniles were all observed following exposure to any of the three *K. mikimotoi* isolates at the density of $1\text{--}3 \times 10^7$ cells L^{-1} . Due to the similarity in toxicity on marine invertebrates, one possible view is that the FJ strain may be more relative to NZ and EU isolates. Interestingly, Al-Kandari et al. (2011) have proposed the isolates from Europe and New Zealand are more closely related to each other in ITS regions

and the *rbcl* gene. However, the genotype comparison between Chinese strains and other isolates was not conducted, which is more needed to identify the differences among *K. mikimotoi* strains worldwide in future research.

For many algal strains, including *Karenia* spp., *Chattonella* spp., *Heterocapsa* spp., and *Alexandrium* spp., haemolytic toxicity was detected by haemolysis of mammalian erythrocytes (Zou et al., 2010; Basti et al., 2015; Cho et al., 2017). For example, for *K. mikimotoi* in Japan, 40% haemolysis of mammalian erythrocytes was detected in JP-Fuk strain with an algal density of 5×10^7 cells L^{-1} (Zou et al., 2010), however, 50% haemolysis was found from JP-Hiroshima strain with an algal density of 2×10^7 cells L^{-1} (Cho et al., 2017). Neely and Campbell (2006) also proposed a modified assay to determine haemolytic toxicity by the percentage of saponin. However, due to the differences in the algal density during measurement, the haemolytic toxicity was hardly compared among strains in various research. In this study, we conducted an adapted protocol to determine haemolytic toxin variability among *Karenia* clones isolated from China. Haemolytic toxicity was assessed by relative saponin concentration in each algal cell, based




Figure 6. The activity of major enzymes in the gills in the control and experimental groups (each data point represents the mean percentage \pm standard deviation of triplicate cultures).




Figure 7. The abalone haemocyte parameters after 48 h exposure (each data point represents the mean percentage \pm standard deviation of triplicate cultures).

on the standard curve between saponin (Sigma) concentration and mammalian erythrocytes haemolysis. Using this approach, the difference in the production of haemolytic compounds among various *Karenia* clones will be examined in future research.

Causes of abalone mortality resulted from *K. mikimotoi*-FJ strain

Although several similar phenomena on abalone mortality were previously reported in *Karenia* clones (eg. *K. mikimotoi*, *Karenia cristata*, and *Karenia Concordia*) in the laboratory, the lethal mechanisms have not been clarified. In this study, *K. mikimotoi* was able to kill the abalones during the aeration. The mortality rate significantly increased and lethal time sharply shortened as the dissolved oxygen-depleted during the non-aeration. These results indicated the death of abalones was caused by *K. mikimotoi* own as well as oxygen depletion secondary-developed.

In *Karenia* spp., some researchers have demonstrated that haemolytic toxins had toxic effects on fish and shellfish (Yasumoto *et al.*, 1990; Satake and Tanaka, 2005; Zou *et al.*, 2010) by primarily destroying the gill structure and immune systems (Ou, 2006). However, the lethal effect of *K. mikimotoi* only occurred in URCS. All of the abalones successfully survive in CFCs. This suggested that the haemolytic toxins killing the abalones originated from active cells of *K. mikimotoi* FJ-strain. The direct contact of active cells with gills could produce haemolytic compounds, which worked and led to the histopathological changes in gills as the previous reports (Neely and Campbell, 2006; Zou *et al.*, 2010). In fish, the haemolytic toxin of *K. mikimotoi* caused gill arch, raker, primary filament, and secondary filament (Ou, 2006). Similarly, in this experiment, the afferent and efferent branchial vessels for abalone gills become shrinking and deformed evidently. Many gill filaments were also broken. In this research, the ratio of granulocytes to hyalinocytes type

Table 2. Literature review of the toxic impacts of different *K. mikimotoi* strains on marine organisms

Taxa	Haemolytic activity	Mortality of zooplankton	Mortality of shellfish	Reference
<i>K. mikimotoi</i> (CH-FJ)	29.43 pg saponin/cell (algal density: 3×10^7 cell L^{-1})	Brine shrimp (N) rotifer (Y)	Abalone juvenile (Y)	This study; Li <i>et al.</i> (2017)
<i>K. mikimotoi</i> (NZ)	ND	Brine shrimp (N)	Abalone juvenile (Y)	Shi <i>et al.</i> (2012)
<i>K. mikimotoi</i> (EU)	48% of saponin activity	Brine shrimp (N)	Abalone juvenile (Y)	Neely and Campbell (2006) Botes <i>et al.</i> , (2003)
<i>K. mikimotoi</i> (CH-SCS)	8.99 pg saponin /cell (algal density: 3×10^7 cell L^{-1})	Rotifer (Y) brine shrimp (Y)	Scallop juvenile (N) abalone juvenile (N)	This study Sun <i>et al.</i> (2010) Dang <i>et al.</i> (2015)
<i>K. mikimotoi</i> (JP-Suo)	No effect	Rotifer (N)	ND	Zou <i>et al.</i> (2010)
<i>K. mikimotoi</i> (JP-Fuk)	40% haemolysis of mammalian erythrocytes (algal density: 5×10^7 cell L^{-1})	Rotifer (Y)	ND	Zou <i>et al.</i> (2010)
<i>K. mikimotoi</i> (JP-Hiroshima)	50% Haemolysis of mammalian erythrocytes (algal density: 2×10^7 cell L^{-1})	Rotifer (Y)	ND	Cho <i>et al.</i> (2017) Basti <i>et al.</i> (2015)

ND, No data; Y, inimical impacts found; N, no inimical impacts found.

cells and phagocytosis activity of abalones decreased, and the mortality of haemocytes significantly increased, which was constant with the previous studies. Many researchers have reported that shellfish immune systems were attacked by many HAB algal species (Haberkorn et al., 2010; Medhioub et al., 2012; Lassudrie et al., 2014, 2015). For example, some algal cells of *Alexandrium fundyense* can increase the shellfish blood cell mortality rate, and increase the proportion of shellfish apoptotic blood cells (Galimany et al., 2008). Apart from the abalones, it has been tested that *K. mikimotoi* caused harmful effects on the immune systems of other shellfish species, such as *Ruditapes philippinarum* and *Argopectens irradians* (Smolowitz and Shumway, 1997; Hégaret et al., 2007; Hégaret et al., 2011).

In addition, DO depletion was another crucial stressor, which aggravated the mortality of abalones in the experiment. Similar physiological responses were also found in other molluscs when exposed to low oxygen. Cheung et al. (2008) found that prolonged hypoxia (3.0 mg O₂ L⁻¹ and 1.5 mg O₂ L⁻¹) could reduce food consumption, respiration, growth, and reproduction in gastropod *Nassarius festivus*. DO depletion has also been documented to cause depression in the immune system. Under the 24-hour exposure of 2.05 mg O₂ L⁻¹, the abalone *Haliotis diversicolor supertexta* showed a significantly decreased total haemocyte counts and respiratory burst (Cheng et al., 2004). The green-lipped mussel *Perna viridis* also exhibited high haemocyte mortality and reduced phagocytosis at the oxygen level of 1.5 mg O₂ L⁻¹ (Wang et al., 2011). During harmful algal bloom, the decomposition and respiration of algal cells always lead to the depletion of oxygen (Wang et al., 2012). For example, laboratory simulation found that DO could drop to 2.9 mg O₂ L⁻¹ after 5 days in the decay stage of *Alexandrium catenella* (Wang et al., 2021). In the natural environment, DO levels showed a depressed tendency and could persist for weeks during the *K. mikimotoi* bloom collapse (O'Boyle et al., 2016). Thus, the extensive death of abalones was largely associated with the combination of algal toxicity and DO depletion in Fujian during *K. mikimotoi* blooms.

Apart from the oxygen, the survival, growth, and physiological metabolism of abalone could also be obviously influenced by other environmental factors such as pH and ammonia nitrogen. For example, non-ionic ammonia nitrogen in seawater with a concentration exceeding 35 µg L⁻¹, and 9.8 µg L⁻¹, could cause a significant effect on the growth of *Haliotis discus hannai*, and *Haliotis midae*, respectively (Matsuyama et al., 1999). In this research, pH, temperature, and salinity were maintained under suitable conditions. Although the total ammonia nitrogen in the water showed some fluctuation, the maximum value reached 0.084 mg L⁻¹. Whereas, according to Bower and Bidwell (1978), the maximum non-ionic ammonia nitrogen can be calculated to 2.50 µg L⁻¹. Therefore, the concentration of non-ionic ammonia nitrogen in the experimental system could not affect the survival of abalone.

Conclusion

The outbreaks of *K. mikimotoi* caused massive mortalities of abalones and serious economic losses in Fujian. Our experiments verified that *K. mikimotoi* Fujian-strain could kill the abalones through haemolytic toxicity destroying the afferent, efferent branchial vessels and filaments of abalone gill and decreasing granulocytes respect to hyalinocytes type cells, phagocytosis activity and the survival of haemocytes at the bloom density *in situ*. The toxic effect just worked for the haemolytic toxicity from active algal cells, which were probably produced under the contract of algal cells and abalone gills. Compared to the *K. mikimotoi* SCS-strain as 'fish killer', the haemolytic toxicity of Fujian strain was stronger, suggesting a new ecotype different from the former

one. Apart from the algal toxicity, DO depletion exacerbated the mortality of abalones in the experiment. Since DO could decrease during *K. mikimotoi* blooms, DO depletion cooperating with the algal toxicity of *K. mikimotoi* led to the numerous deaths of abalones in Fujian.

Data availability. All data generated or images used during this study are included in this published article.

Author contributions. Lingzhi Liao conducted paper writing. Jianing Lin made project administration and laboratory experiments. Xinshu Ding conducted data analysis. Song Feng edited the paper. Tian Yan made project administration and methodology.

Financial support. This work was supported by Open Fund of CAS Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences (grant number KLMEES201801) and the National Nature Science Foundation of China (NSFC 41476102).

Competing interests. None.

Ethical standards. All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

References

- Al-Kandari MA, Highfield AC, Hall MJ, Hayes P and Schroeder DC (2011) Molecular tools separate harmful algal bloom species, *Karenia mikimotoi*, from different geographical regions into distinct sub-groups. *Harmful Algae* 10, 636–643.
- Basti L and Segawa S (2010) Mortality of the short-neck clam *Ruditapes philippinarum* induced by the toxic dinoflagellate *Heterocapsa circularisquama*. *Fisheries Science* 76, 625–631.
- Basti L, Nagai S, Go J, Okano S, Nagai K, Watanabe R, Suzuki T and Tanaka Y (2015) Differential inimical effects of *Alexandrium* spp. and *Karenia* spp. on cleavage, hatching, and two larval stages of Japanese pearl oyster *Pinctada fucata martensii*. *Harmful Algae* 43, 1–12.
- Botes L, Smit AJ and Cook PA (2003) The potential threat of algal blooms to the abalone (*Haliotis midae*) mariculture industry situated around the South African coast. *Harmful Algae* 2, 247–259.
- Bower C and Bidwell E (1978) Ionization of ammonia in seawater: effects of temperature, pH and salinity. *Journal of the Fisheries Research Board of Canada* 35, 1012–1016.
- Chang FH, Uddstrom MJ, Pinkerton MH and Richardson KM (2008) Characterising the 2002 toxic *Karenia concordia* (Dinophyceae) outbreak and its development using satellite imagery on the north-eastern coast of New Zealand. *Harmful Algae* 7, 532–544.
- Cheng W, Li CH and Chen JC (2004) Effect of dissolved oxygen on the immune response of *Haliotis diversicolor supertexta* and its susceptibility to *Vibrio parahaemolyticus*. *Aquaculture* 232, 103–115.
- Cheung SG, Chan HY, Liu C and Shin P (2008) Effect of prolonged hypoxia on food consumption, respiration, growth and reproduction in marine scavenging gastropod *Nassarius festivus*. *Marine Pollution Bulletin* 57, 280–286.
- Cho K, Kasaoka T, Ueno M, Basti L, Yamasaki Y, Kim D and Oda T (2017) Haemolytic activity and reactive oxygen species production of four harmful algal bloom species. *European Journal of Phycology* 52, 311–319.
- Dang LX, Li Y, Liu F, Zhang Y, Yang WD, Li HY and Liu JS (2015) Chemical response of the toxic dinoflagellate *Karenia mikimotoi* against grazing by three species of zooplankton. *Journal of Eukaryotic Microbiology* 62, 470–480.
- Davidson K, Miller P, Wilding TA, Shutler J, Bresnan E, Kennington K and Swan S (2009) A large and prolonged bloom of *Karenia mikimotoi* in Scottish waters in 2006. *Harmful Algae* 8, 349–361.
- Dutra FM, Rönnau M, Sponchiado D, Forneck SC, Freire CA and Ballester E (2017) Histological alterations in gills of *Macrobrachium amazonicum* juveniles exposed to ammonia and nitrite. *Aquatic Toxicology* 187, 115–123.
- Galimany E, Sunila I, Hégaret H, Ramón M and Wikfors GH (2008) Experimental exposure of the blue mussel (*Mytilus edulis*, L.) to the toxic dinoflagellate *Alexandrium fundyense*: histopathology, immune responses, and recovery. *Harmful Algae* 7, 702–711.
- Haberkorn H, Lambert C, Le Gou'c N, Gue'guen M, Moal J, Palacios E, Lasso P and Soudant P (2010) Effects of *Alexandrium minutum* exposure upon physiological and hematological variables of diploid and triploid oysters, *Crassostrea gigas*. *Aquatic Toxicology* 97, 96–108.

- Haywood AJ, Steidinger KA, Truby EW, Bergquist PR, Adamson J and MacKenzie L (2004) Comparative morphology and molecular phylogenetic analysis of three new species of the genus *Karenia* (Dinophyceae) from New Zealand. *Journal of Phycology* **40**, 165–179.
- Hégaret H, Wikfors GH and Soudant P (2003a) Flow-cytometric analysis of haemocytes from eastern oysters, *Crassostrea virginica*, subjected to a sudden temperature elevation: I. Haemocyte types and morphology. *Journal of Experimental Marine Biology and Ecology* **293**, 237–248.
- Hégaret H, Wikfors GH and Soudant P (2003b) Flow cytometric analysis of haemocytes from eastern oysters, *Crassostrea virginica*, subjected to a sudden temperature elevation: II. Haemocyte functions: aggregation, viability, phagocytosis, and respiratory burst. *Journal of Experimental Marine Biology and Ecology* **293**, 249–265.
- Hégaret H, Da Silva PM, Wikfors GH, Lambert C, De Bettignies T, Shumway SE and Soudant P (2007) Hemocyte responses of Manila clams, *Ruditapes philippinarum*, with varying parasite, *Perkinsus olseni*, severity to toxic-algal exposures. *Aquatic Toxicology* **84**, 469–479.
- Hégaret H, Smolowitz RM, Sunila I, Shumway SE, Alix J, Dixon M and Wikfors GH (2010) Combined effects of a parasite, QPX, and the harmful-alga, *Prorocentrum minimum* on northern quahogs, *Mercenaria mercenaria*. *Marine Environmental Research* **69**, 337–344.
- Hégaret H, Da Silva PM, Wikfors GH, Haberkorn H, Shumway SE and Soudant P (2011) In vitro interactions between several species of harmful algae and haemocytes of bivalve molluscs. *Cells Biology and Toxicology* **27**, 249–266.
- Heinig CS and Campbell DE (1992) The environmental context of a *Gyrodinium aureolum* bloom and shellfish kill in Maquoit Bay, Maine, September 1988. *Journal of Shellfish Research* **11**, 111–122.
- Lassudrie M, Soudant P, Richard G, Henry N, Medhioub W, Silva PM, Donval A, Bunel M, Goïc N, Lambert C, Montaudouin X, Fabioux C and Hégaret H (2014) Physiological responses of Manila clams *Venerupis (=Ruditapes) philippinarum* with varying parasite *Perkinsus olseni* burden to toxic algal *Alexandrium ostenfeldii* exposure. *Aquatic Toxicology* **154**, 27–38.
- Lassudrie M, Wikfors GH, Sunila I, Alix JH, Dixon MS, Combout D, Soudant P, Fabioux C and Hégaret H (2015) Physiological and pathological changes in the eastern oyster *Crassostrea virginica* infested with the trematode *Bucephalus* sp. and exposed to the toxic dinoflagellate *Alexandrium fundyense*. *Journal of Invertebrate Pathology* **126**, 51–63.
- Li J, Glibert PM, Zhou MJ, Lu SH and Lu DD (2009) Relationships between nitrogen and phosphorus forms and ratios and the development of dinoflagellate blooms in the East China Sea. *Marine Ecology Progress Series* **383**, 11–26.
- Li XD, Yan T, Lin JN, Yu RC and Zhou MJ (2017) Detrimental impacts of the dinoflagellate *Karenia mikimotoi* in Fujian coastal waters on typical marine organisms. *Harmful Algae* **61**, 1–12.
- Matsuyama Y, Uchida T and Honjo T (1999) Effects of harmful dinoflagellates, *Gymnodinium mikimotoi* and *Heterocapsa circularisquama*, red-tide on filtering rate of bivalve molluscs. *Fisheries Science* **65**, 248–253.
- Medhioub W, Lassus P, Truquet P, Bardouil M, Amzil Z, Sechet V, Sibat M and Soudant P (2012) Spirolied uptake and detoxification by *Crassostrea gigas* exposed to the toxic dinoflagellate *Alexandrium ostenfeldii*. *Aquaculture* **358–359**, 108–115.
- Neely T and Campbell L (2006) A modified assay to determine hemolytic toxin variability among *Karenia* clones isolated from the Gulf of Mexico. *Harmful Algae* **5**, 592–598.
- O'Boyle S, McDermott G, Silke J and Cusack C (2016) Potential impact of an exceptional bloom of *Karenia mikimotoi* on dissolved oxygen levels in waters off western Ireland. *Harmful Algae* **53**, 77–85.
- Ou XY (2006) Histopathological effect of *Karenia mikimotoi* bloom on gills of common farmed fishes in the East China Sea (PhD thesis). Jinan University, Guangzhou, China.
- Qi YZ, Chen JF, Wang ZH, Xu N, Wang Y, Shen PP, Lu SH and Hodgkiss IJ (2004) Some observations on harmful algal bloom (HAB) events along the coast of Guangdong, southern China in 1998. *Hydrobiologia* **512**, 209–214.
- Reddy-Lopata K, Auerswald L and Cook P (2006) Ammonia toxicity and its effect on the growth of the South African abalone *Haliotis midae* Linnaeus. *Aquaculture* **261**, 678–687.
- Satake M and Tanaka Y (2005) Gymnocin-B with the largest contiguous polyether rings from the red tide dinoflagellate, *Karenia* (formerly *Gymnodinium*) *mikimotoi*. *Tetrahedron Letters* **46**, 3537–3540.
- Shi F, McNabb P, Rhodes L, Holland P, Webb S, Adamson J, Immers A, Gooneratne R and Holland J (2012) The toxic effects of three dinoflagellate species from the genus *Karenia* on invertebrate larvae and finfish. *New Zealand Journal Of Marine And Freshwater Research* **46**, 149–165.
- Smolowitz R and Shumway SE (1997) Possible cytotoxic effects of the dinoflagellate, *Gyrodinium aureolum*, on juvenile bivalve molluscs. *Aquaculture International* **5**, 291–300.
- SOAC (State oceanic administration of China) (2006) Report of marine environment quality in 2005. <http://www.mee.gov.cn/hjzl/shj/jagb/201605/P020160527557747968290.pdf>. Accessed online 27 July 2006.
- SOAC (State oceanic administration of China) (2013) Report of marine environment quality in 2012. <http://www.mee.gov.cn/hjzl/shj/jagb/201605/P020160527563756757405.pdf>. Accessed online 27 July 2013.
- Southgate T, Wilson K, Cross T and Myers A (1984) Recolonization of a rocky shore in SW Ireland following a toxic bloom of the dinoflagellate, *Gyrodinium aureolum*. *Journal of the Marine Biological Association of the United Kingdom* **64**, 485–492.
- Steidinger KA, Wolny JL and Haywood AJ (2008) Identification of Kareniaceae (Dinophyceae) in the Gulf of Mexico. *Nova Hedwigia* **133**, 269p.
- Sun K, Yan T, Zhou MJ and Ho KC (2010) Effect of *Karenia mikimotoi* on the survival of rotifer *Brachionus plicatilis*, brine shrimp *Artemia salina*, and *Neomysis awatschensis* (in Chinese). *Marine Science* **34**, 76–81.
- Tangen K (1977) Blooms of *Gyrodinium aureolum* (Dinophyceae) in North European waters, accompanied by mortality in marine organisms. *Sarsia* **63**, 123–133.
- Taylor FJR, Fukuyo Y, Larsen J and Hallegraeff GM (2004) Taxonomy of harmful dinoflagellates. In Hallegraeff GM (ed.), *Manual on Harmful Marine Microalgae, Monographs on Oceanographic Methodology*. Paris: Unesco, pp. 389–432.
- Wang YJ, Hu MH, Shin PKS and Cheung SG (2011) Immune responses to combined effect of hypoxia and high temperature in the green-lipped mussel *Perna viridis*. *Marine Pollution Bulletin* **63**, 201–208.
- Wang C, Yu RC and Zhou MJ (2012) Effects of the decomposing green macroalga *Ulva (Enteromorpha) prolifera* on the growth of four red-tide species. *Harmful Algae* **16**, 12–19.
- Wang QN, Li XD, Yan T, Song JJ, Yu RC and Zhou MJ (2021) Laboratory simulation of DO reduction and ammonia nitrogen generation in the decay stage of HAB. *Journal of Oceanology and Limnology* **39**, 500–507.
- Yang Z and Hodgkiss I (2004) Hong Kong's worst "red tide" causative factors reflected in a phytoplankton study at Port Shelter station in 1998. *Harmful Algae* **3**, 149–161.
- Yasumoto T, Uderal B and Aune T (1990) Screening for hemolytic and ichthyotoxic components of *Chrysochromulina polyplepis* and *Gyrodinium aureolum* from Norwegian coastal waters. In Graneli E (ed.), *Toxic Marine Phytoplankton*. New York: Elsevier, pp. 436–440.
- Zhou MJ (2010) Environmental settings and harmful algal blooms in the Sea area adjacent to the Changjiang river estuary. In Ishimatsu A and Lie HJ (eds), *Coastal Environmental and Ecosystem Issues of the East China Sea*. Tokyo: Terrapub, pp. 133–149.
- Zhou CX, Fernandez N, Chen HM, You YR and Yan XJ (2011) Toxicological studies of *Karlodinium micrum* (Dinophyceae) isolated from East China Sea. *Toxicon* **57**, 9–18.
- Zou Y, Yamasaki Y, Matsuyama Y, Yamaguchi K, Honjo T and Oda T (2010) Possible involvement of hemolytic activity in the contact-dependent lethal effects of the dinoflagellate *Karenia mikimotoi* on the rotifer *Brachionus plicatilis*. *Harmful Algae* **9**, 367–373.