

The potential nutrition-, physical- and health-related benefits of cow's milk for primary-school-aged children

Penny Rumbold^{1,*†}, Nicola McCulloch¹ , Ruth Boldon¹, Crystal Haskell-Ramsay² , Lewis James³, Emma Stevenson⁴ and Benjamin Green^{1,5,†}

¹Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

²Department of Psychology, Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK

³National Centre for Sport and Exercise Medicine East Midlands, School of Sport, Exercise and Health Sciences, National Centre of Sport and Exercise Medicine, Loughborough University, Loughborough, UK

⁴Human Nutrition Research Centre, Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK

⁵Danone Specialised Nutrition, Trowbridge, UK

Abstract

Cow's milk is a naturally nutrient-dense foodstuff. A significant source of many essential nutrients, its inclusion as a component of a healthy balanced diet has been long recommended. Beyond milk's nutritional value, an increasing body of evidence illustrates cow's milk may confer numerous benefits related to health. Evidence from adult populations suggests that cow's milk may have a role in overall dietary quality, appetite control, hydration and cognitive function. Although evidence is limited compared with the adult literature, these benefits may be echoed in recent paediatric studies. This article, therefore, reviews the scientific literature to provide an evidence-based evaluation of the associated health benefits of cow's milk consumption in primary-school-aged children (4–11 years). We focus on seven key areas related to nutrition and health comprising nutritional status, hydration, dental and bone health, physical stature, cognitive function, and appetite control. The evidence consistently demonstrates cow's milk (plain and flavoured) improves nutritional status in primary-school-aged children. With some confidence, cow's milk also appears beneficial for hydration, dental and bone health and beneficial to neutral concerning physical stature and appetite. Due to conflicting studies, reaching a conclusion has proven difficult concerning cow's milk and cognitive function; therefore, a level of caution should be exercised when interpreting these results. All areas, however, would benefit from further robust investigation, especially in free-living school settings, to verify conclusions. Nonetheless, when the nutritional-, physical- and health-related impact of cow's milk avoidance is considered, the evidence highlights the importance of increasing cow's milk consumption.

Keywords: Milk: Dairy: Children: Nutrition: Health: Paediatrics

(Received 20 September 2020; revised 25 March 2021; accepted 5 April 2021; accepted manuscript published online 27 April 2021)

Introduction

Commonly cited as nature's perfect food⁽¹⁾, milk is the foundation of life for all mammalian neonates^(2,3). Representing a complex and unique liquid, milk (which is produced by the mammary glands⁽⁴⁾) is the only foodstuff designed by nature to serve as a complete food, at least during the pre-weaning period. In this sense, milk contains numerous biological constituents and an assortment of nutrients necessary for immunological protection and initial growth^(2,5). Shortly after weaning, most mammals stop consuming milk⁽⁶⁾. With domestication and

milking of various animals, however, some humans continue to drink milk throughout life⁽⁷⁾, mostly cow's milk. The value of cow's milk in the human diet has been heavily debated for many years⁽⁸⁾ and has been portrayed by some as 'white poison', a health hazard and promoter of Western chronic diseases⁽⁹⁾. This position is based on numerous hypotheses including that cow's milk contains blood, pus, hormones and antibiotics, causes acne and cancer and has a high cholesterol and fat/saturated fat content collectively contributing to cardiovascular disease risk, weight gain and obesity. Nevertheless, such percep-

* **Corresponding Author:** Penny Rumbold, Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK. E-mail: Penny.rumbold@northumbria.ac.uk

† These authors contributed equally to this work.

tions of cow's milk being harmful to health are not supported by evidence. Conversely, cow's milk has often been signified as the white elixir⁽⁶⁾.

Childhood is a key stage for the development of healthy eating patterns⁽¹⁰⁾, and the school environment provides a valuable setting to develop such behaviours. This is particularly true considering children spend much of their time at school. Indeed, dietary habits shaped throughout the childhood years might carry forth and track into adulthood. Between the ages of 4 and 11 years (primary-school age), children grow and mature at a rapid rate preceding the onset of puberty⁽¹¹⁾. Childhood is therefore a critical transition period preceding adolescence, characterised by growing independence and marked physical development. Good nutrition during the childhood period is therefore particularly important⁽¹²⁾. Not only may good nutrition support proper growth and maturation, but it may therefore act as a base for immediate and lasting health, wellbeing and disease prevention. This is best achieved by consuming a balanced and varied diet that provides all the nutrients needed. The inclusion of cow's milk as a staple component of a healthy balanced diet has been long recognised and is central in most public dietary recommendations⁽¹³⁾. Milk is naturally nutrient-dense and is a significant source of many essential macro and micronutrients. Indeed, cow's milk may confer nutrition-, physical- and health-related benefits beyond that of helping children to simply meet nutrient targets. Based on the available literature, it appears that many of these effects are a product of milk's nutritional composition. It therefore seems relevant to provide readers with a brief description of the nutritional composition of cow's milk.

Nutritional Composition of Cow's Milk and Patterns of Consumption

The nutritional composition of cow's milk is influenced by factors including genes (species and breed), physical state (age and stage of lactation) and environment (available nutrition and climatic conditions)⁽¹⁴⁾. The composition of whole cow's milk is approximately 87% water and 13% solids⁽¹⁾. The percentage split of water and solids of cow's milk, as well as energy, is primarily determined by the amount of fat⁽¹⁵⁾ but is also influenced by added sugars and sweeteners. Cow's milk is approximately 4.9% carbohydrate. The primary carbohydrate portion of milk is lactose, a disaccharide comprising glucose and galactose. The lipid component of cow's milk and other milk-based dairy foods contributes numerous properties including the provision of fat-soluble micronutrients and essential fatty acids, as well as influencing flavour, texture and appearance⁽¹⁵⁾. Of the lipid content within cow's milk, roughly 64% is saturated fatty acids, with a considerable amount (~26%) from monounsaturated fatty acids and a small contribution from trans- and polyunsaturated fatty acids (both ~3%)⁽¹⁶⁾. Additionally, cow's milk provides high-quality proteins, namely casein and whey. Casein and whey constitute approximately 82% and 18% of the total protein found in cow's milk and provide an abundance of essential amino acids⁽¹⁷⁾. Aside from the macronutrient content, cow's milk contains essential micronutrients that contribute to dietary quality and overall nutritional status. As presented in Table 1, with the exception of vitamin C (which is

broken down during pasteurisation) and vitamin D (unless fortified), cow's milk is a good source of all vitamins⁽¹⁸⁾. Calcium and phosphorus, crucial to healthy growth and maturation, as well as other biological processes, are the most prominent minerals present in cow's milk. Cow's milk also makes significant contributions to intakes of other major minerals⁽¹⁹⁾. In this sense, cow's milk makes a substantial contribution to, and is the main dietary source of, calcium (26%), iodine (37%), riboflavin (25%), magnesium (10%) and potassium (14%) in the diets of primary-school-aged children⁽²⁰⁾. While volume-specific data are limited, in the UK, the National Diet and Nutrition Survey (NDNS) provides a nationally representative assessment concerning dietary habits of individuals, aged 1.5 years and older, living in private households and remains the only surveillance programme to do so. Temporal and age-related trends of cow's milk to average daily total energy and macro and micronutrient intake from 2009/10 through to 2015/16 are presented in Table 2 for children (4–10 years) and adolescents (11–18 years). Despite the clear value of cow's milk in the everyday diet, it is clear that intakes steadily decline as children age. This trend is not only restricted to the UK; it is also true in the USA^(21,22), Australia⁽²³⁾ and other European countries⁽²⁴⁾. This is of great concern among children, especially as dietary habits may track into adulthood, as milk avoidance may have detrimental implications over time, leaving populations vulnerable to micronutrient deficiencies (e.g. calcium, iodine and riboflavin deficiencies to name a few⁽²⁵⁾) and lasting nutrition- and health-related complications (e.g. cardiovascular disease⁽²⁶⁾, metabolic syndrome⁽²⁷⁾, hypertension^(27,28), poor weight management and bone health^(25,29)).

Scope and Methodology of the Narrative Review

Evidence from adult populations suggests that adequate cow's milk consumption is associated with a reduced risk of cardiovascular disease⁽³⁰⁾, metabolic syndrome^(28,31) and obesity^(32–34). Emerging data also suggest that cow's milk may have a role in overall dietary quality, appetite control, hydration and cognitive function. Although the evidence to date is limited compared with the adult literature, these benefits appear echoed in recent paediatric studies. There is a need to review the literature to assess whether there is sufficient evidence of a beneficial or detrimental effect of cow's milk in the diet and health of children. The aim of this narrative review is therefore to summarise and appraise the scientific literature to form an accurate, evidence-based evaluation of the associated nutrition-, physical- and health-related benefits of cow's milk consumption in primary-school-aged children (4–11 years). Based on recent publications, it is not the intention of this narrative review to focus on body weight and body composition, as it is known with some confidence that cow's milk consumption is inversely (or not) associated with body weight and body composition in children and adolescents⁽³⁵⁾. Where possible, this review focuses solely on literature in children aged 4–11 years; however, where there is no applicable literature, data from general child studies will be reviewed (and possibly some adult studies). This is particularly true for mechanistic considerations, as much of this evidence comes from adult studies; therefore, any adult-specific data should be interpreted with some caution as the observations cited may not always be replicated in

Table 1. Nutritional composition of cow's milk

Per 100 g	Whole milk	Semi-skimmed milk	Skimmed milk	Flavoured milk*
Energy (kJ)	274	195	144	270
Protein (g)	3.3	3.5	3.5	3.6
Carbohydrate (g)	4.6	4.7	4.8	9.6
of which sugars (g)	4.6	4.7	4.8	8.9
Fat (g)	3.9	1.7	0.3	1.5
Saturates	2.5	1.1	0.1	1
Monounsaturates	1	0.4	0.1	0.3
Polyunsaturates	0.1	Trace	Trace	0.1
Trans fatty acids	0.1	0.1	Trace	Trace
Thiamin (mg)	0.03	0.03	0.03	0.03
Niacin (mg)	0.2	0.1	0.1	0.1
Niacin from tryptophan (mg)	0.6	0.6	0.7	0.8
Calcium (mg)	118	120	125	120
Riboflavin (mg)	0.23	0.24	0.22	0.17
Vitamin B6 (mg)	0.06	0.06	0.06	0.03
Vitamin B12 (µg)	0.6	0.9	0.8	0.1
Folate (µg)	8	9	9	2
Vitamin D (µg)	Trace	Trace	Trace	Trace
Biotin (µg)	2.5	3.0	2.5	2.2
Pantothenate (mg)	0.6	0.7	0.5	0.3
Vitamin C (mg)	2	2	1	Trace
Retinol (µg)	30	19	1	20
Carotene (µg)	19	9	Trace	8
Sodium (mg)	43	43	44	52
Potassium (mg)	155	156	162	168
Magnesium (mg)	11	11	11	12
Phosphorus (mg)	93	94	96	102
Zinc (mg)	0.4	0.4	0.5	0.4
Copper (mg)	Trace	Trace	Trace	Trace
Selenium (µg)	1	1	1	N/A
Manganese (mg)	Trace	Trace	Trace	Trace
Iodine (µg)	31	30	30	N/A

Note: N/A = values not available for this food; Trace = nutrient is present in less than 0.1 g per 100 g. *Data taken from a sample of strawberry- and banana-flavoured sugar-sweetened milk. Adapted from the Dairy UK. Available at: <http://www.milk.co.uk/publications/default.aspx>

Table 2. Percentage contribution of cow's milk to average daily total energy and macro- and micro-nutrient intake over time by sex and age

	NDNS Rolling Programme Years 1–2; 2009/10				NDNS Rolling Programme Years 3–4; 2011/12				NDNS Rolling Programme Years 5–6; 2013/14				NDNS Rolling Programme Years 7–8; 2015/16	
	4–10 years		11–18 years		4–10 years		11–18 years		4–10 years		11–18 years		4–10 years	11–18 years
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	Combined	Combined
Energy	8	8	4	4	7	7	5	4	8	7	4	4	7	4
Protein	14	12	8	7	13	12	9	8	13	13	10	7	11	8
Carbohydrate	5	5	4	3	5	5	4	3	5	5	4	3	5	4
Fat	9	10	6	5	9	10	6	4	10	11	6	4	9	6
Saturated fat	17	16	8	9	15	14	9	7	16	17	9	8	14	9
Trans fatty acids	14	12	8	8	19	19	14	11	20	21	14	12	17	13
Vitamin A	11	10	7	7	10	10	9	4	12	12	9	7	11	9
Calcium	31	29	21	22	29	27	22	19	29	29	23	20	26	22
Riboflavin	32	29	22	21	28	29	23	20	29	31	23	18	25	19
Vitamin D	0	0	0	0	1	0	0	0	0	1	0	0	1	1
Sodium	5	4	3	4	6	5	3	2	6	6	3	4	6	4
Potassium	16	15	9	9	14	13	11	10	15	16	11	8	14	9
Magnesium	12	10	8	8	11	10	8	7	11	12	8	7	10	7
Zinc	14	13	8	9	13	13	9	8	13	13	10	8	12	8
Iron	1	1	0	1	1	0	1	0	1	1	1	1	1	1
Selenium	6	5	3	3	5	5	3	3	5	5	3	3	4	3
Folate	9	8	4	5	8	7	5	5	8	9	6	4	7	5
Iodine	40	38	30	30	37	37	32	28	39	28	34	27	37	30

Note: Percentage contribution information taken from National Diet and Nutrition Survey (NDNS) rolling programme. Percentage contributions for vitamin B₆, vitamin B₁₂, vitamin C and phosphorus are not reported in the NDNS. *The NDNS rolling programme for 2015/16 (<https://www.gov.uk/government/statistics/ndns-results-from-years-7-and-8-combined>) stopped providing sex-specific intake data, so the data are presented as boys and girls combined.

children. The narrative review will also highlight current knowledge gaps in this field and suggest directions for future research.

To identify research articles to facilitate this narrative review, PubMed (US National Library of Medicine National Institutes of Health) and Web of Science (Thomson Reuters, UK), were searched using various combinations of keywords relevant to the scope of the review up to August 2020. Search terms included: humans, child, milk, flavoured milk, animal milk, cow milk, infant, adolescent, preschool, primary-school, paediatric, diet, energy intake, nutritional status, diet quality, obesity, school-age children, food habits, dairy products, milk beverages, height, stature, anthropometry, calcium, health association, health benefits, dental health, caries, decay, bone health, appetite, cognition, memory, mental performance, school milk, insulin, glucose, hydration, snack. Studies were considered eligible for inclusion in the review if (i) they were conducted in children 4–11 years; (ii) participants received cow's milk, and not supplemental forms of dairy minerals (unless a milk group was included); and (iii) the study had been peer-reviewed. Results from studies that grouped data over a wider age range were also considered where this was deemed appropriate. This was especially true for prospective and intervention studies, yet the mean age of participants had to fall between the stipulated age range. Studies were excluded if (i) human participants were not used; (ii) participants lay directly outside of the stipulated age range; (iii) studies used supplements only; (iv) studies used different dairy foods (i.e. yogurt, cheese, dairy desserts); (v) the milk consumed included different sources of animal milk; or (vi) data reporting was poor and/or not published in English.

In this narrative review, we have summarised the evidence base, which we identified by nutrition-, physical- and health-related related benefits and provide a summary for the studies identified. As this is not a systematic review, we did not perform a quality assessment of each study (based on design and implementation) or conduct a meta-analysis by study design but summarise the findings for each study and sub-section in Tables 3 (nutrition-related benefits), 4 (physical-related benefits) and 5 (health-related benefits). Symbols \uparrow and \downarrow indicate a statistically significant positive and negative effect/relationship, respectively, between cow's milk and related subsections, with \leftrightarrow indicating no statistically significant effect/relationship or a neutral effect. This approach has successfully been used in a recent narrative review in this journal⁽³⁵⁾, and therefore readers should be directed to this paper for justifications and interpretations for this approach. In brief, however, a tabulated summary allowed us to draw overall conclusions for this narrative review, although we did not undertake a detailed assessment of quality of individual studies.

Nutrition-Related Benefits of Cow's Milk

Nutritional Status

The importance of regular cow's milk consumption in children has been recognised for over 40 years⁽³⁶⁾. As described, cow's milk is a naturally nutrient-dense foodstuff, providing a rich source of many essential nutrients. While the potential of cow's milk to improve nutritional status may be unsurprising, the

effect of this in primary-school-aged children has only been investigated by numerous observational studies^(37–42) and three intervention-based studies^(43–45) (Table 3). All reported improved nutritional status with increased cow's milk consumption, yet, because of the study designs utilised, findings must be interpreted with caution. In the few intervention-based studies^(43–45), which ranged from 16 weeks to 6 months, primary school children were provided with flavoured milk or no drink (control)⁽⁴⁴⁾, or 200 ml⁽⁴³⁾ and 250 ml⁽⁴⁵⁾ of multi-micronutrient-fortified milk or unfortified milk, to establish whether unfortified and fortified cow's milk influences micronutrient status in primary-school-aged children. In those consuming fortified versus unfortified cow's milk, significant increases in blood riboflavin and vitamin B₁₂ were observed, with all other analytes (selenium, ferritin and vitamin D) comparable between drinks⁽⁴³⁾. Additionally, significant increases in serum ferritin and zinc have been observed after 6 months with both fortified versus unfortified cow's milk compared with a control drink⁽⁴⁵⁾. With respect to nutritional status, dietary patterns characterised by high cow's milk consumption (both plain and flavoured milk) resulted in greater intakes of energy, protein, phosphorus, magnesium, calcium, potassium, vitamin A, zinc, riboflavin (vitamin B₂) and niacin compared with children who seldom consumed cow's milk. Notably, diets high in cow's milk may also limit intakes of foods and beverages high in fat and sugar. In studies of Dutch⁽³⁷⁾ and American primary-school-aged children⁽⁴⁶⁾, for example, cow's milk intake was inversely related with the intake of sugar-sweetened beverages. In these studies, children with low cow's milk intakes had lower protein, fibre, calcium, magnesium, potassium and phosphorus intakes. Low cow's milk consumption has significant implications for intakes of several key nutrients that are of importance in childhood. Based on the nutritional contribution to dietary intakes, it is important to note that children who drink cow's milk regularly are more likely to meet dietary recommendations for many nutrients, and thus have a better nutritional status⁽¹³⁾. It could therefore be argued that milk intake might be a marker for healthier eating habits⁽³⁷⁾.

With the above in mind, in the UK, it should be noted that dietary intakes of vitamin D among primary-school-aged children are low. Considering UK cows' milk is generally not a good source of vitamin D because it is not fortified, as it is in other countries, this narrative review may provide justification for UK policy makers to reconsider widespread fortification. There is evidence (albeit from a theoretical modelling perspective) from Northern Ireland that cows' milk can be used as a successful vehicle for vitamin D fortification⁽⁴⁷⁾. Fortification of cows' milks with 1 μg , 1.5 μg and 2.0 $\mu\text{g}/100\text{ g}$ theoretically increased median vitamin D intakes from 2.0 $\mu\text{g}/\text{d}$ to 4.2 $\mu\text{g}/\text{d}$, 5.1 $\mu\text{g}/\text{d}$ and 5.9 $\mu\text{g}/\text{d}$, respectively and may therefore provide strong evidence for the efficacy of widespread fortification. This modelling appears to translate to human studies. In support of this, a recent review comprising 20 studies showed positive associations between the consumption of vitamin D fortified milk and 25(OH)D status in different population groups⁽⁴⁸⁾. Furthermore, in Finland, Canada and the USA, who exercise a national vitamin D fortification policy (covering various fluid milk products), milk products contributed 28–63% to vitamin D intake, while in countries without a fortification policy, or where

Table 3. Studies in primary-school-aged children that measured nutritional status and hydration with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Nutritional status					
Cook <i>et al.</i> (1975) ⁽³⁸⁾	<i>n</i> = 312 (159 boys) Age: 8–11 years United Kingdom	Observational	7 d diet record	Nutrient intake was higher in children drinking school milk every day than in non-consumers Boys: energy (<i>P</i> < 0.05), protein (<i>P</i> < 0.01), fat (<i>P</i> < 0.05), calcium (<i>P</i> < 0.001), thiamin (<i>P</i> < 0.01) and riboflavin (<i>P</i> < 0.001) intakes higher than in non-consumers Girls: energy (<i>P</i> < 0.001), protein (<i>P</i> < 0.001), fat (<i>P</i> < 0.01), calcium (<i>P</i> < 0.001), thiamin (<i>P</i> < 0.001) and riboflavin (<i>P</i> < 0.001) intakes higher than in non-consumers	↑ ↑ ↑
LaRowe <i>et al.</i> (2007) ⁽³⁹⁾	<i>n</i> = 793 (410 boys) Age: 6–11 years United States	Observational	24 h dietary recalls collected during NHANES 2001–2002.	General linear models showed children clustered as high-fat milk consumers had higher (<i>P</i> < 0.05 for all) intakes of energy, protein, riboflavin, folate, vitamin A, vitamin C and calcium compared with clusters of water, SSB, soda and mix/light drinks	↑
Albala <i>et al.</i> (2008) ⁽⁴⁴⁾	<i>n</i> = 98 (56 boys) Age: 8–10 years Chile	Randomised, control trial	16 week intervention. Children drank 3 servings/d of flavoured milk or control (no drink)	Protein and calcium intakes increased (<i>P</i> = 0.0001) and energy intake decreased (<i>P</i> = 0.009) with milk compared with controls	↑
Murphy <i>et al.</i> (2008) ⁽⁴⁰⁾	<i>n</i> = 2097 (1061 boys) Age: 6–11 years United States	Observational	24 h dietary recalls collected during NHANES 1999–2002. Participants grouped by age (2–5 years; 6–11 years; 12–18 years).	Calcium, phosphorus, magnesium, potassium and vitamin A intakes comparable between plain and flavoured milk consumers Calcium, phosphorus, magnesium, potassium and vitamin A intakes significantly higher (<i>P</i> < 0.05 for all) with milk consumption (plain and flavoured) compared with non-consumers	↔ ↑
Lien <i>et al.</i> (2009) ⁽⁴⁵⁾	<i>n</i> = 454 (217 boys) Age: 7–8 years Vietnam	Double-blind, intervention	Volume-matched drink (250 ml) served twice daily (morning), 6 d per week for 6 months: (a) multi-micronutrient fortified milk (b) unfortified milk (c) control (no drink)	Intakes of energy, protein, fat, sugar and vitamin A increased (<i>P</i> < 0.05 for all) after 6 months in conditions (a) and (b) and compared with condition (c) Serum ferritin and zinc increased (<i>P</i> < 0.05 for all) after 6 months in conditions (a) and (b) and compared with condition (c) Zinc levels, however, also increased in condition (c) (<i>P</i> < 0.05) FM drinkers had a higher mean intake for calcium (930 versus 837 mg; <i>P</i> = 0.010), vitamin D (6.9 versus 5.9 µg; <i>P</i> = 0.021) and total sugar (99 g versus 90 g, <i>P</i> = 0.015) than non-consumers	↑ ↑
Wang <i>et al.</i> (2012) ⁽⁴²⁾	<i>n</i> = 632 Age: 8–10 years Canada	Observational	Three 24 h diet recalls	FM drinkers had a higher mean intake for calcium (930 versus 837 mg; <i>P</i> = 0.010), vitamin D (6.9 versus 5.9 µg; <i>P</i> = 0.021) and total sugar (99 g versus 90 g, <i>P</i> = 0.015) than non-consumers	↑
Rangan <i>et al.</i> (2013) ⁽⁴¹⁾	<i>n</i> = 222 (121 boys) Age: 8–10 years Australia	Observational	Three non-consecutive 24 h diet recalls	Milk (and dairy) intake significantly associated with increased intake of energy (<i>P</i> < 0.001), protein (<i>P</i> = 0.02), calcium (<i>P</i> < 0.001), phosphorus (<i>P</i> < 0.001), magnesium (<i>P</i> < 0.001), potassium (<i>P</i> = 0.009), zinc (<i>P</i> = 0.019), vitamin A (<i>P</i> < 0.001), riboflavin (<i>P</i> < 0.001) and niacin (<i>P</i> = 0.03)	↑
Campmans-Kuijpers <i>et al.</i> (2016) ⁽³⁷⁾	<i>n</i> = 1007 (504 boys) Age: 7–13 years Netherlands	Observational	Two non-consecutive 24 h dietary recalls with an interval of 2–6 weeks	Higher milk consumption was associated with significantly higher intakes of energy (<i>P</i> = 0.003), protein (<i>P</i> < 0.0001), fat (<i>P</i> = 0.03), fibre (<i>P</i> = 0.02), calcium (<i>P</i> < 0.0001), folate (<i>P</i> < 0.0001), iodine (<i>P</i> < 0.0001), potassium (<i>P</i> < 0.0001), magnesium (<i>P</i> < 0.0001), phosphorus (<i>P</i> < 0.0001), selenium (<i>P</i> = 0.002), zinc (<i>P</i> < 0.0001) and vitamins B ₁ (<i>P</i> < 0.0001), B ₂ (<i>P</i> < 0.0001) and B ₁₂ (<i>P</i> < 0.0001)	↑
Kuriyan <i>et al.</i> (2016) ⁽⁴³⁾	<i>n</i> = 225 (52 boys) Age: 7–10 years India	Double-blind, randomised placebo-controlled	Volume-matched drink (200 ml) served twice daily (morning and afternoon), 6 d per week for 5 months: (a) multi-micronutrient fortified milk (b) unfortified milk	At the end of the intervention, levels of blood Vitamin B ₁₂ , and riboflavin were significantly different (<i>P</i> < 0.001) between the study groups, in favour of the fortified milk group Vitamin D, selenium and body iron showed no difference with either group	↑ ↔

Table 3. (Continued)

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Hydration Montenegro-Bethancourt et al. (2013) ⁽⁶⁷⁾	n = 442 (309 boys) Age: 4–10 years, Germany	Observational	DONALD Study 3 d weighed dietary records	Milk consumption significantly increased FWR ($P < 0.05$) by 25 ml in boys and 33 ml in girls per 100 g of intake	↑
Volterman et al. (2014) ⁽⁶⁸⁾	n = 38 (19 boys) Age: 7–11 years and 14–17 years, Canada	Randomised, repeated-measures cross-over	Three exercise sessions in 34.5°C (2 × 20-min cycling bouts at 60 % peak oxygen uptake) followed by consumption of: (a) plain water (W) (b) a carbohydrate–electrolyte solution (CES) (c) skimmed milk (SM)	Fraction of ingested beverage retained at 2 h of recovery was greater with SM (74 % ± 18 %) than W (47 % ± 26 %) and CES (59 % ± 20 %, $P < 0.001$ for both), and greater in CES than water ($p < 0.001$)	↑
Volterman et al. (2016) ⁽⁶⁸⁾	n = 15 (7 boys) Age: 10–12 Canada	Randomised, counter-balanced crossover design	45 min of alternating running and cycling exercise in 34.5°C followed by consumption (150% of body mass loss) of isoenergetic and electrolyte-matched beverages containing: (a) 0 g (control), (b) 0.76 g (Lo-PRO) (c) 1.5 g (Hi-PRO) of milk protein/100 ml	Less negative fluid balance at 2 h of recovery after consumption of Hi-PRO versus CONT ($P = 0.01$) Compared with CONT, beverage retention was enhanced by Hi-PRO at 2 h ($P < 0.05$)	↑ ↑

Abbreviations in order of appearance: FM = flavoured milk; NHANES = National Health and Nutrition Examination Survey; SSB = sugar-sweetened beverage; FWR = free water reserve; W = water; CES = carbohydrate/electrolyte solution; SM = skimmed milk; Lo-PRO = low protein; Hi-PRO = high protein.

the fortification covered only some dairy products (Sweden, Norway), the contribution was much lower or negligible⁽⁴⁸⁾. Based on the above, widespread fortification of milk seems to help bolster vitamin D intakes and could be adopted by the UK to increase vitamin D intakes in primary-school children.

Concerns about added sugars in foods and beverages have increased recently. Accordingly, many schools have limited access to foods and beverages high in added sugars, including flavoured cow's milk. In some studies, removal of flavoured cow's milk from the school environment reduces energy and sugar intake, but negatively impacts essential nutrient intake and even reduced the overall intake of non-flavoured cow's milk^(49,50). In this sense, total milk consumption (both plain and flavoured) decreased by 12.3 %⁽⁵⁰⁾. Contrariwise, children who drink more flavoured cow's milk generally have higher non-flavoured milk intakes and display nutritional intakes similar to children who only consume non-flavoured milk⁽⁵¹⁾. While greater cow's milk consumption (both plain and flavoured) is generally associated with higher daily energy intake, we know with some confidence that this does not impact body mass or composition⁽⁵⁵⁾. If daily energy intake is of concern for children predisposed to overweight and obesity, results of a recent study suggest that replacement of whole cow's milk or semi-skimmed cow's milk with skimmed cow's milk may help reduce total energy intake, without impacting nutrient provision⁽⁵²⁾.

When taken together, these results suggest that cow's milk consumption (both plain and flavoured) might serve as a useful strategy to boost nutritional status in primary-school-age children and may act as a surrogate marker of diet quality. There may be a need for widespread vitamin D fortification of cow's milk, however, and based on the above seems justified and should be reviewed by UK policy makers. Cow's milk is a readily available, accessible and affordable means of providing valuable essential nutrients to the diets of primary-school children. The nutritional implications of cow's milk provision and/or removal in the school environment must be considered and is especially relevant in children suffering from nutritional inadequacies. The data are, however, primarily limited to observational investigations and require verification in more controlled intervention-based studies covering differing populations.

Hydration

Whole cow's milk is approximately 87 % water⁽¹⁾ and may therefore be a beneficial choice for hydration. For children, maintenance of euhydration is important for good health⁽⁵³⁾ but may also increase concentration and mental performance (cognitive function)⁽⁵⁴⁾ while helping reduce instances of headaches, constipation and other disorders⁽⁵³⁾. This is important as children are at a greater risk of dehydration compared with adults, having relatively greater fluid losses at rest and during exercise and being less able to recognise thirst⁽⁵⁴⁾. Given that water is continually being lost from the body water pool, it is important to constantly replenish fluid losses to prevent dehydration. Establishing beverages that encourage longer-term fluid retention and maintenance of euhydration has real clinical and practical implications⁽⁵⁵⁾. This is particularly true in situations where free access to fluids is limited or when frequent breaks for urination



are not desirable⁽⁵⁵⁾, such as in the school setting where many children arrive at school in an already hypo-hydrated state⁽⁵⁶⁾. There is little research concerning the impact of cow's milk on hydration in primary-school-aged children (Table 3). To date, three studies have been conducted^(57–59). Two of these were exercise-based intervention studies conducted by the same research group in Canada^(58,59), and the remaining study was cross-sectional⁽⁵⁷⁾. Nonetheless, all reported improved hydration status with cow's milk in comparison with water or alternative beverages. While these findings show promise, additional controlled-intervention studies should be performed in different settings to examine repeatability of these effects.

In the two exercise-based intervention studies, researchers examined the influence of post-exercise cow's milk consumption on rehydration in 7–11-year-old⁽⁵⁹⁾ and 10–12-year-old children⁽⁵⁸⁾. The rehydration potential of cow's milk (or milk protein) was compared with water or a carbohydrate–electrolyte drink or zero or low cow's milk protein beverage, given at 100 % and 150 % of the children's body mass losses following exercise, respectively. In both studies, children exercised in the heat (~34.5°C) and consumed the test beverages immediately following exercise. The children were subsequently observed for a period of 2 h⁽⁵⁹⁾ and 4 h⁽⁵⁸⁾, respectively, during which measures of hydration status were collected (urine output and fluid balance). Findings over 2 h showed cow's milk was more effective than both water and the carbohydrate–electrolyte drink at replacing fluid loss during exercise⁽⁵⁹⁾. This was similar over 4 h. The authors suggested that the protein in cow's milk may be a factor responsible for its rehydrating properties⁽⁵⁸⁾.

Montenegro-Bethancourt and colleagues⁽⁵⁷⁾ conducted a cross-sectional study designed initially to establish the contribution of fruit and vegetable intake on hydration status. Children (4–10 years old) recorded all food and beverages consumed over 3 d using a weighed food diary alongside 24 h urine samples. Regular intake of fruit and vegetables made a substantial contribution to hydration status, but notably, cow's milk consumption was also a strong dietary predictor of hydration status. In particular, cow's milk increased free water reserve by 25 ml in boys and 33 ml in girls per 100 g of intake. These findings corroborate studies in adult populations^(60–62).

Based solely on adult studies, there are a number of potential mechanisms explaining the greater fluid retention (and thus hydration potential) with cow's milk⁽⁶³⁾. Firstly, milk contains modest amounts of sodium (~20 mmol/L, similar to most commercial sports drinks) and large amounts of potassium (~40 mmol/L). Sodium, as the main cation in the extracellular fluid, plays a major role in fluid retention⁽⁶⁴⁾, whilst potassium may exert some beneficial effects⁽⁶⁵⁾, although this is not a consistent finding. Secondly, the protein content of cow's milk may help facilitate greater fluid retention^(66,67). While water and alternative beverages meet the basic intentions of rehydration, they do not offer the abundance of nutrients present in cow's milk that will also aid in normal growth and maturation. Nevertheless, it should be considered that, while these findings suggest that cow's milk helps improve the hydration status of children, there remains room for further studies to clarify the role of cow's milk in hydration, especially in a free-living school setting. While the

intervention-based studies show promise, many primary-school children do not exercise in such conditions.

Physical-Related Benefits of Cow's Milk

Dental Health

Milk contains multiple nutrients that may offer anticariogenic properties, protecting against the development of dental caries, and thus supporting dental health in children^(68,69). The nutrients principally believed to play a role in dental health include calcium, phosphorus and protein^(68–70). As reported earlier, cow's milk intake has been shown to be inversely related with the intake of sugar-sweetened beverages^(37,46). Although speculative, one could argue that increased cow's milk intake might indirectly improve dental health. Calcium is required for bone and tooth formation, whereas phosphorus ions work alongside calcium to maintain tooth strength⁽⁷⁰⁾. Cow's milk proteins (particularly α_{s1} -, α_{s2} - and β -casein) may act to prevent tooth enamel erosion and demineralisation of the tooth surface by producing casein phosphopeptides⁽⁷¹⁾. Based on the available evidence, inverse associations between cow's milk intake and the incidence of tooth decay have frequently been reported^(72–75) (Table 4). The evidence, however, has been derived from cross-sectional research, so causal conclusions cannot be justified and caution must be exercised.

The findings of the cited studies were generated from recall methods which have obvious shortcomings. Nevertheless, all studies (following visual examinations accompanied with food frequency questionnaires) reported inverse associations with cow's milk and the development of dental caries. Interestingly, though sugar is suggested to possess acidogenic and cariogenic potential⁽⁷⁶⁾, one study⁽⁷⁴⁾ reported that the association between cow's milk consumption and protection against dental caries was stronger for children with diets high in sucrose. This might suggest that cow's milk offers protection against the harmful effects of sugar, though this is speculative. To this end, cow's milk intake during primary-school years has been reported as a predictor of incidences of caries later in childhood⁽⁷²⁾. In this sense, greater cow's milk intake is inversely associated with indices of dental caries. Collectively, these studies support the suggestion that dietary habits established and maintained during primary-school years could have longer-term effects on health outcomes.

Although no controlled trials have been conducted in primary-school children, the available literature appears to suggest that cow's milk could help reduce the incidence of dental caries and contribute to dental health in children. To reduce the occurrence of tooth decay, it is recommended that primary-school children limit their consumption of sugary beverages (especially when not consumed with a meal) and increase consumption of cow's milk. The exact mechanism by which cow's milk reduces the incidence of dental caries remains uncertain, though calcium, phosphate and casein phosphopeptides may all play a role^(68–70). Casein phosphopeptides are phosphorylated casein-derived peptides produced by tryptic digestion of casein in the duodenum⁽⁷¹⁾. The anticariogenic activity of casein phosphopeptides is due to their ability to stabilise high levels of

amorphous calcium phosphate on tooth surface, preventing demineralisation and enhancing remineralisation of enamel caries⁽⁷⁷⁾. In addition, milk fat could be adsorbed onto the enamel surface and may have a protective role. Thirdly, milk enzymes may have a role in reducing the growth of acidogenic plaque bacteria⁽⁷⁸⁾. Where prior observational research provides a solid foundation, any future work should seek to implement robust randomised clinical trials (RCT) to confirm any causal relationships between cow's milk consumption and dental health in primary-school-aged children.

Bone Health and Physical Stature

Childhood is a critical time for bone growth and lasting bone health^(79,80), and the nutritional composition of cow's milk has evolved to stimulate and support this^(2,5). The scientific opinion that regular cow's milk consumption is associated with greater physical stature has a long history, dating back to the 1920s^(81,82). Cow's milk contains multiple nutrients that may support childhood growth⁽⁸⁰⁾ and lasting bone health⁽⁷⁹⁾. The beneficial effects of cow's milk consumption on bone health in children may include increased bone mineral content (BMC) and bone mineral density (BMD), characteristics necessary for the prevention of bone-related diseases later in life (osteoporosis). Evidence of these benefits, however, is equivocal, showing a beneficial-to-neutral effect of cow's milk on these constructs^(25,29,83–88). In this sense, of the cited studies, two reported greater total body BMC^(85,88), while one reported no effect. One study reported no effect on total body BMD⁽⁸⁷⁾, yet three studies reported increased regional BMC^(29,83,86), and another two studies reported increased regional BMD^(25,84) (Table 4). The mixed findings reported throughout these studies may be due to a lack of consistency in methodological approaches, length of study, location and measures of bone health, all of which confound comparisons and prevent a clear conclusion. Furthermore, age and sex differences must be considered, especially as puberty and bone mineralisation typically occurs earlier in girls compared with boys⁽⁸⁹⁾.

In two recent studies^(85,88), school milk interventions increased total body and regional (forearm) BMC and BMD compared with children who seldom drank cow's milk. In these studies, the beneficial effects of cow's milk on bone health were observed in $n = 435$ – 757 children (mean age 11 years) following daily school milk intake for 1–2 years.

On a comparative basis with other animal sources, whole cow's milk is the richest source of calcium and represents the biggest contributor of dietary calcium during childhood⁽⁹⁰⁾. In addition, considering beef and eggs for example, cow's milk is the cheapest source of protein, calcium, phosphorus and vitamin D⁽⁹¹⁾. During childhood, the beneficial effects of cow's milk on bone health in children are commonly attributed to calcium⁽⁹²⁾. However, many other nutrients, including phosphorus and protein, are needed for normal growth and lasting bone health⁽⁹³⁾. In 5–11-year-old children ($n = 99$), for example, calcium supplementation for 10 months did not influence total body or regional BMC⁽⁸⁷⁾. In contrast, in an earlier study where cow's milk (and dairy) was supplemented daily for 1 year (distributed to deliver 1200 mg calcium daily), lumbar (lower back) BMD increased compared with children who maintained their usual eating habits⁽⁸⁴⁾. This may illustrate that calcium and other nutrients work together for bone growth and, thus,

lasting bone health⁽⁹³⁾. It is important to note, however, that children in the calcium supplementation study were already consuming near daily recommended amounts, which may illustrate that intakes exceeding calcium recommendations (from either supplemental calcium or cow's milk) offer no further benefit to bone health in children.

In several studies, it has been observed that children who avoid consuming cow's milk characteristically exhibit low calcium intakes, short statures, increased fatness and lower BMC (and thus exhibit reduced bone health) compared with their cow's milk-drinking counterparts^(25,86). In children (mean age 8 years) who previously avoided cow's milk, the introduction of cow's milk to the diet increased not only habitual cow's milk consumption but also increased total body BMC⁽²⁹⁾. This may suggest it is never too late for children to introduce cow's milk into their diet for bone health benefits.

From a stature perspective, available data appear to suggest that cow's milk consumption almost certainly has a positive effect on growth in children. To date, 14 studies in primary-school-aged children have evaluated this aspect of cow's milk consumption^(25,29,36,44,81–83,85,94–99). Seven were intervention-based, four were observational and the remaining three were prospective designs. Based on these studies, the evidence suggests that cow's milk intake positively influences physical stature in primary-school-aged children. All six intervention studies showed increased stature with increased cow's milk (one study included milk calcium). In addition, all prospective studies reported increased stature with increased cow's milk. With regard to the cross-sectional studies, two illustrated that cow's milk avoiders displayed stunted growth compared with cow's milk drinkers, while the remaining study showed increased adult stature with increased cow's milk consumption in childhood. Although trends are consistent across studies, there remains a need for evidence from robust controlled-intervention trials in primary-school-aged children to verify causality.

Beneficial effects on physical stature were observed with both whole- and reduced-fat cow's milk, distributed at a range of 190–568 ml daily. In several of these studies, it was reported that childhood cow's milk intake was associated with higher skeletal development (BMD of the hip and the forearm), bone growth and periosteal bone expansion. These were likely established earlier during growth periods and maintained into late adolescence and young adulthood^(96–99), supporting the notion that dietary habits established and maintained during the primary- and secondary-school years may not only induce short-term effects but offer lasting benefits. Indeed, in children with prolonged cow's milk avoidance, stunted growth and physical stature is observed compared with children who habitually consume cow's milk, and this is maintained into adulthood^(25,29,86). During a pubertal growth spurt, about 37% of the entire skeletal mass is accumulated⁽¹⁰⁰⁾. Therefore, inadequate calcium intake during this period may compromise volumetric bone density and overall stature attained. Notably, in a study that explored both cow's milk and supplemental calcium, those children in the cow's milk (and dairy) group were 3 cm taller (166 cm) compared with the supplemental calcium (163 cm) and placebo (163 cm) group⁽⁹⁶⁾. This may indicate that cow's milk has more of a

Table 4. Studies in primary-school-aged children that measured dental health, bone health and physical stature with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Dental health					
Petti <i>et al.</i> (1997) ⁽⁷⁴⁾	<i>n</i> = 439 (217 boys) Age: 6–11 years Italy	Descriptive cross-sectional	24 h diet diary, oral examination using Plaque Index and number of decayed, extracted or filled teeth	Milk consumption significantly reduced the probability of carries (<i>P</i> < 0.05); greater significance in high-sucrose (<4/day) consuming children (<i>P</i> < 0.01)	↑
Levine <i>et al.</i> (2007) ⁽⁷²⁾	<i>n</i> = 315 Age: 7–11 years England	Prospective cohort/cross-sectional	3 d dietary diary, tooth brushing habits, dental examination using BASCD survey	Significantly (<i>P</i> < 0.05) less caries associated with moderate consumption (1–2/d) of milk (and dairy) products by children aged 11–15 years	↑
Llena & Forner (2008) ⁽⁷³⁾	<i>n</i> = 369 (220 boys) Age: 6–10 years Spain	Descriptive cross-sectional	Food frequency questionnaire, one dental examination for caries and fillings	No significant impact of milk on the incidence of caries	↔
Curtis <i>et al.</i> (2018) ⁽⁷⁵⁾	<i>n</i> = 392 (183 boys) Age: 0–19 years United States	Longitudinal, observational	Dietary questionnaires at 6-month intervals, cavities assessed age 5, 9, 13 and 17 years	Higher milk intake associated with lower expected adjusted decayed and filled surface increments; however, was not significant	↑
Bone health and physical stature					
Cook <i>et al.</i> (1979) ⁽³⁶⁾	<i>n</i> = 1210 (581 boys) Age: 6–7 years United Kingdom	Prospective cohort	Free school milk consumed (190 ml/d at school) versus control; height gain	Significantly greater height gain in the milk group in Scottish females (<i>P</i> < 0.05) No difference in males or any groups in England	↑ ↔
Rona & Chinn (1989) ⁽⁹⁸⁾	<i>n</i> = 670 Age: 5–11 years United Kingdom	Prospective cohort	Free school availability (190 ml/d at school) versus unavailable; change in standardised height score	Significantly greater increments in standardised height scores in 'milk available' group after 2 years in Scotland (<i>P</i> < 0.05) No difference in England	↑ ↔
Baker <i>et al.</i> (1980) ⁽⁹⁴⁾	<i>n</i> = 581 (267 boys) Age: 7–8 years United Kingdom	Randomised, control trial	12 month intervention; 190 ml/d of milk at school versus control; height gain	Significantly greater height gain in milk group than control (<i>P</i> < 0.05)	↑
Chan <i>et al.</i> (1995) ⁽⁸⁴⁾	<i>n</i> = 48 females Age: 9–13 years USA	Randomised control trial	12 month intervention; ≥ 1200 mg/d calcium from dairy versus control; DXA of total body bone mineral content	Greater gain in total body bone mineral content in dairy calcium group (<i>P</i> < 0.001)	↑
Bonjour <i>et al.</i> (1997) ⁽⁸³⁾	<i>n</i> = 149 females Age: 6–10 years Switzerland	Randomised control trial	12 month intervention; 850 mg/d milk-extracted calcium versus placebo; DXA of regional bone mineral content, height gain and change in lumbar spine length	Greater increment in six-site mean bone mineral content in supplementation group (<i>P</i> < 0.05). At 1 year follow-up, greater increment in femoral shaft bone mineral content (<i>P</i> < 0.02) No difference in height, though significance approached (<i>P</i> ≤ 0.08) favouring supplement group Significant difference in lumbar spine length following intervention and 1 year follow-up (<i>P</i> ≤ 0.05)	↑ ↔ ↑
Black <i>et al.</i> (2002) ⁽²⁵⁾	<i>n</i> = 250 (120 boys) Age: 3–10 years New Zealand	Cross-sectional	Food frequency questionnaire, and DXA of total body and regional bone mineral content and height	Milk avoiders had significantly lower total body bone mineral content than age-, sex-matched controls (<i>P</i> < 0.01) Milk avoiders significantly shorter than control children of the same age and sex (<i>P</i> < 0.01)	↑ ↑
Du <i>et al.</i> (2004) ⁽⁸⁵⁾	<i>n</i> = 757 (girls) Age: 10–12 years China	Randomised control trial	24 month intervention; 330 ml fortified milk (560 mg calcium)/school day versus 300 ml fortified milk (560 mg calcium) and 5 µg or 8 µg cholecalciferol/school day versus control group. DXA of regional and total body bone mineral content with stature assessed pre-, mid- and post-trial	Greater increase in total-body bone mineral content (> 1.2%) and bone mineral density (> 3.2%) in milk groups compared to control Significantly greater % change in height in milk supplement groups compared with control group (<i>P</i> < 0.005)	↑ ↑



Table 4. (Continued)

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Goulding <i>et al.</i> (2004) ⁽⁸⁶⁾	<i>n</i> = 50 (20 boys) Age: 3–10 years New Zealand	Observational longitudinal	Milk avoiders versus birth cohort population, DXA of total body and regional bone mineral content, history of bone injuries, estimation of calcium intake	Greater number of children reporting fractures and increased total fractures in milk-avoidance group compared with control (<i>P</i> < 0.001)	↑
Rockell <i>et al.</i> (2005) ⁽²⁹⁾	<i>n</i> = 46 (18 boys) Age: 5–12 years New Zealand	Cross-sectional, longitudinal	Food frequency questionnaire, 4 d food diary and DXA of total body bone mineral content and stature	Increase in total body bone mineral content in prior milk avoiders when milk consumption had increased (<i>P</i> < 0.05) but remained lower than in non milk avoiders	↑
Wiley (2005) ⁽⁹⁹⁾	<i>n</i> = 2592 Age: 5–11 year United States	Cross-sectional	24 h recall, milk frequency in past 30 d; height and adult height	Milk intake not associated with height at age 5–11 years (<i>P</i> = 0.385)	↔
Iuliano-Burns <i>et al.</i> (2006) ⁽⁸⁷⁾	<i>n</i> = 99 (58 boys) Age 5–11 years Austria	Randomised control trial	10 month intervention; 800 mg/d of calcium from CaCO ₃ versus 800 mg/d of calcium from milk versus placebo, DXA of total and regional bone mineral content	Greater gain in pelvic bone mineral content in pre-pubertal children versus controls at 10 months in the milk mineral group (<i>P</i> < 0.02) but not biologically meaningful	↔
Albala <i>et al.</i> (2008) ⁽⁴⁴⁾	<i>n</i> = 98 (56 boys) Age: 8–10 years Chile	Randomised, control trial	16 week intervention; Children drank 3 servings/d of flavoured milk or control (no drink)	Significantly greater increase (<i>P</i> = 0.01) Smaller increase in height in females in intervention group versus control (<i>P</i> = 0.10)	↑ ↔
Zhou <i>et al.</i> (2011) ⁽⁸⁸⁾	<i>n</i> = 435 (188 boys) Age 10–12 years China	Retrospective cohort	School milk group (1 year, 4/week; or 3 year, 1–3/week) versus 'seldom milk' group, ulna bone mineral content	Greater ulna bone mineral content in school milk group (<i>P</i> < 0.001)	↑
Guo <i>et al.</i> (2020) ⁽⁹⁵⁾	<i>n</i> = 41 439 (19 618 boys) Age: 6–17 years China	Cross-sectional	Validated questionnaire assessing milk intake classified as low, medium and high; stature	Milk intake not associated with stature (<i>P</i> > 0.05)	↔

Abbreviations in order of appearance: DXA = dual energy X-ray absorptiometry; CaCO₃ = calcium carbonate.

Table 5. Studies in primary-school-aged children that measured appetite and cognitive function with cow's milk consumption

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Appetite regulation					
Rumbold <i>et al.</i> (2013) ⁽¹¹¹⁾	<i>n</i> = 25 (11 boys) Age: 9–10 years United Kingdom	Randomised controlled cross-over	Energy-matched (0.3 MJ) preloads during school mid-morning break: (a) 160 ml of semi-skimmed milk (b) 153 g of apple	Boys and girls felt less hunger and expressed a lower desire to eat when apple was consumed compared with semi-skimmed milk (<i>P</i> = 0.02) Energy intake comparable at lunch between semi-skimmed milk and apple (<i>P</i> > 0.05)	↓ ↔
Mehrabani <i>et al.</i> (2014) ⁽¹¹⁰⁾	<i>n</i> = 34 (obese boys) Age: 10–12 years Iran	Three-way repeated measure randomised controlled cross-over	Volume-matched preloads with breakfast (all 240 ml): (a) low fat milk (1.5 %) (b) apple juice (isoenergetic control) (c) water (control)	Energy intake lower at lunch following low fat milk consumption compared to water (<i>P</i> < 0.005) and apple juice (<i>P</i> < 0.05)	↑
Vien <i>et al.</i> (2014) ⁽¹¹²⁾	Normal and overweight children Mean age: 11.5 years Average age: 11.5 years Canada	Not stated	Volume (250 ml) and energy-matched (543.9 kJ) with the exception of water given pre- and within meals: (a) 1 % fat (1 g/100 g) flavoured milk (b) 2 % fat (2 g/100 g) milk (c) 1.5 % fat (1.5 g/100 g) yogurt drink (d) fruit punch (e) water	<i>Experiment 1</i> Energy intake lower after chocolate milk and yogurt (<i>P</i> < 0.01) compared with water Post-meal SA was lower after milk than yogurt (<i>P</i> < 0.01) Post-meal SA was lower after milk than fruit punch (<i>P</i> < 0.01) <i>Experiment 2</i> GLP-1 AUC was higher after milk than fruit punch and in OWOB compared with NW children (<i>P</i> < 0.03). Post-prandial drop in ghrelin was greater after milk than fruit punch in OWOB children (–24 versus –14%) but was not significant (<i>P</i> = 0.06) Energy intake, insulin and glucose AUC were comparable between all preloads (<i>P</i> > 0.05)	↑ ↑ ↑ ↔ ↔
Mehrabani <i>et al.</i> (2016) ⁽¹⁰⁹⁾	<i>n</i> = 34 obese boys Age: 10–12 years Iran	Three-way repeated measure randomised controlled cross-over	Volume-matched preloads with breakfast (all 240 ml): (a) low fat milk (1.5 %) (b) apple juice (isoenergetic control) (c) water (control)	Higher satiety scores after drinking low-fat milk with breakfast compared with water and apple juice (<i>P</i> < 0.05) Energy intake lower at lunch following low-fat milk compared with water (<i>P</i> < 0.001) and apple juice (<i>P</i> = 0.03)	↑ ↑
Cognitive function					
Rahmani <i>et al.</i> (2011) ⁽¹²⁴⁾	<i>n</i> = 469 (230 boys) Average age: 8 years Iran	Case-control population-based intervention	Drink at morning school break: (a) tetra-pack sterilised and homogenised milk (250 ml) (b) control (no milk supplementation)	Grade-point average increased from pre- to post-test in girls with milk (<i>P</i> < 0.05) No change for the control group, nor for either of the groups of boys Mean IQ after milk was better in boys compared with boys at post-test in the control (<i>P</i> < 0.05)	↑ ↔ ↑
Brindal <i>et al.</i> (2013) ⁽¹²³⁾	<i>n</i> = 40 (19 boys) Age: 10–12 years, Australia	Double-blind, randomised three-way repeated measures crossover	Volume-matched (455 ml) morning drink following ≥ 8 h fasting: (a) glucose beverage (b) full milk beverage (c) half milk/glucose beverage	No effect of beverage type on speed of processing, working memory, short-term memory, attention switching, perceptual speed and inspection time No interactions between beverage type and timing of the cognitive testing (60, 120 and 180 min post-drink) In the conditions with milk, girls recalled 0.7–0.8 more words, but in the glucose condition they recalled 0.5 fewer words (<i>P</i> = 0.014) For boys, no difference between beverages was found (<i>P</i> > 0.09) Sex differences identified for change in word recall after full milk (<i>P</i> < 0.001) and half milk/glucose (<i>P</i> < 0.001) conditions: girls showed an increase, and boys showed a decrease	↔ ↔ ↑ ↔ ↔



Table 5. (Continued)

Reference	Details	Design	Methodological approach	Results and conclusion	Effect
Kuriyan <i>et al.</i> (2016) ⁽⁴³⁾	<i>n</i> = 225 (52 boys) Age: 7–10 years India	Double-blind, randomised placebo-controlled design	Volume-matched drink (200 ml) served twice daily (morning and afternoon), 6 d per week for 5 months: (a) multi-micronutrient fortified milk (b) unfortified milk	No group × time interaction for short-term memory and executive functions from baseline	↔
Zahrou <i>et al.</i> (2016) ⁽¹³²⁾	<i>n</i> = 200 Age: 7–9 years Morocco	Double-blind, controlled design	Volume-matched (200 ml) drink served daily at school for 9 months, including weekends: (a) fortified milk; added potassium iodide and other micronutrients (b) non-fortified milk	Fortified milk group performed significantly better than the non-fortified milk group (<i>P</i> = 0.02) on a dynamic testing procedure designed to assess children's learning potential	↔
Faught <i>et al.</i> (2017) ⁽¹²⁶⁾	<i>n</i> = 1595 (732 boys) Age: 10–11 years Canada	Observational	Analysis of secondary data (food frequency questionnaire) from a 2012 population-based survey using a stratified random sampling design	Boys meeting the nutrition recommendations for milk and alternatives (3–4 servings per d) scored 3.45 % better on an average measure of Math and Language Arts standardised provincial achievement tests than those not meeting the recommendations (<i>P</i> = 0.02)	↑
Petrova <i>et al.</i> (2019) ⁽¹²⁵⁾	<i>n</i> = 103 (52 boys) Age: 8–14 years Spain	Double-blind, randomised controlled design	Volume-matched (200 ml) drink served three times per d for 5 months: (a) fortified milk beverage; added micronutrients and essential fatty acids (b) regular full milk	Greater baseline to post-intervention increases in working memory capacity in the fortified milk group (32 % increase) compared with the regular milk group (13 % increase) (<i>P</i> = 0.027)	↔
Sheng <i>et al.</i> (2019) ⁽¹³³⁾	<i>n</i> = 75 (42 boys) Age: 5–6 years China	Double-blind, multicentre randomised controlled parallel crossover	Volume-matched (150 ml) drinks consumed twice daily following meals for 5 d: (a) conventional milk, containing A1 and A2 β-casein (b) milk containing only A2 β-casein	Subtle Cognitive Impairment Test improved following both conventional milk (<i>P</i> < 0.014) and milk containing only A2 β-casein (<i>P</i> < 0.002) No difference in Subtle Cognitive Impairment Test between drinks Consumers of conventional milk in phase 1 and milk containing only A2 β-casein in phase 2, no effect on error rates in phase 1 (<i>P</i> = 0.101) but a decrease in error rates in phase 2 (<i>P</i> < 0.001) In consumers of milk containing only A2 β-casein in phase 1 and conventional milk in phase 2, there was a decrease in the error rate in phase 1 which continued through the washout period (<i>P</i> < 0.001), but error rates then increased in phase 2 (<i>P</i> < 0.001)	↔ ↔ ↔ ↓
Anderson <i>et al.</i> (2020) ⁽¹²⁷⁾	<i>n</i> = 84 (39 boys) Age: 8–12 years, United States	Randomised counterbalanced crossover	Volume-matched (237 ml) morning drink following ≥ 8 h fasting: (a) 1 % fat milk (b) apple juice	<i>Inhibitory control</i> Reaction time significantly faster after milk compared to apple juice (<i>P</i> < 0.05) No effect of beverage on accuracy <i>Speeded working memory</i> No difference between the beverages on reaction time (<i>P</i> = 0.45) Sex × beverage interaction for accuracy (<i>P</i> = 0.003); compared with apple juice, milk decreased performance accuracy for females whereas non-significant increase in accuracy for males <i>Sustained attention</i> Sex × beverage interaction (<i>P</i> = 0.02); compared with apple juice, milk significantly improved reaction time for females, whereas reaction time decreased in males (though not significantly)	↑ ↔ ↔ ↑↔ ↑↔

Benefits of milk for primary-school children

Abbreviations in order of appearance: SA = subjective appetite; GLP-1 = glucagon like peptide-1; OWOB = overweight and obese; NW = normal weight; AUC = area under the curve; IQ = intelligence quotient.

beneficial effect on physical stature and growth than single cow's milk constituents (i.e. calcium).

The precise mechanisms or nutrients in cow's milk responsible for stimulating growth and lasting bone health are not yet clear. Evidence suggests that maintaining adequate calcium intake during childhood might be advantageous for the attainment of peak bone mass, which may be crucial in reducing the risk of bone-related diseases later in life⁽¹⁰¹⁾. Interestingly, it appears that whole foods may offer greater benefits than the equivalent amount of calcium in supplemental form. It has also been suggested that the growth-stimulating effect of cow's milk is likely attributed to hormonal effects that can be influenced by ingested cow's milk proteins (predominantly whey protein and the release of amino acids during digestion but also casein), micronutrients and also energy^(102–104). Observations from child studies show that these nutrients stimulate the secretion of insulin-like growth factor-1 and insulin, both of which are anabolic hormones that play an essential role in the regulation of growth and accrual of bone mass during childhood^(102–104), though there is some controversy that cow's milk intake upregulates insulin and insulin-like growth factor-1 signalling and thus promotes chronic diseases such as cancer (prostate, breast and colorectal) and cardiovascular disease⁽⁹⁾, though these perceptions are hypothetical at present and not supported by evidence.

Nonetheless, when taken together, it appears that cow's milk consumption promotes health and may increase physical stature in primary-school-aged children. This suggests that cow's milk consumption during childhood might be important to ensure full growth potential is achieved. While there is some suggestion about the improved bone health with increased cow's milk consumption and the mechanisms responsible for the growth-stimulating properties of cow's milk, more intervention studies are needed to elucidate the components responsible for these effects and to prove and/or disprove the chronic disease hypotheses. This is especially prudent when considering bone health, as the methodological approaches previously employed have been diverse in nature.

Health-Related Benefits of Cow's Milk

Appetite Regulation

Appetite comprises numerous regulatory processes associated with the initiation and termination of eating and the selection and amount of food consumed. The regulation of appetite and eating behaviour depend on the detection and integration of signals relaying nutritional status, and their interaction with signals associated with food palatability and gastrointestinal handling, in addition to circadian, social, emotional, habitual and other situational influences⁽¹⁰⁵⁾. Consequently, appetite and the regulation of eating behaviour are complex processes, regulated by homeostatic and non-homeostatic influences⁽¹⁰⁶⁾. Cow's milk contains a host of nutrients that might exert a favourable effect on appetite and eating behaviour⁽¹⁰⁷⁾, yet in primary-school children there is limited evidence concerning the short-term effect of cow's milk on these measures. There are also no data on the moderate- and longer-term effects of daily cow's milk consumption on these outcomes in primary-school children.

From the available studies^(108–112), cow's milk has principally been given as a mid-morning snack or alongside breakfast, with the volume of cow's milk offered to children ranging from 160 ml to 250 ml. Based on these studies, the evidence concerning cow's milk and appetite regulation is inconclusive. Three of these studies found a decrease in energy intake after cow's milk consumption, yet three reported no effect. Two studies reported that cow's milk consumption reduced subjective appetite, and one reported increased subjective appetite compared with a fruit-based snack, while two studies did not measure subjective appetite. Only one study measured hormonal indicators of appetite and reported that cow's milk consumption stimulated the secretion of glucagon-like peptide-1.

In two studies of 34 overweight and obese boys (mean age 11 years)^(109,110), when compared with volume- and energy-matched servings of water or fruit juice, 240 ml of low-fat cow's milk with breakfast reduced energy intake at an *ad libitum* lunchtime meal. In another study comprising 48 obese children (mean age 11 years), girls reported higher satiety scores 4 h after drinking whole cow's milk compared with skimmed milk, and low-fat cow's milk significantly reduced appetite compared with apple juice 2 h after consumption. These differences, however, did not translate to changes in energy intake at an *ad libitum* lunchtime meal across all conditions in girls, boys and the group as a whole⁽¹⁰⁸⁾. As mentioned, only one investigation (comprising two experiments) has sought to establish the effect of cow's milk (and other dairy products) on appetite and feeding behaviour in normal weight and overweight/obese children (mean age 11.5 years), where subjective appetite and appetite-related analytes were measured⁽¹¹²⁾. In both experiments, preloads (experiment 1: 1 % fat (1 g/100 g) chocolate cow's milk, 2 % (2 g/100 g) fat cow's milk, 1.5 % (1.5 g/100 g) fat yogurt drink, fruit punch or a water drink; experiment 2: 2 % (2 g/100 g) fat cow's milk or a fruit punch) were provided 60 min preceding and during an *ad libitum* pizza meal. All preloads were matched for volume (250 ml) and energy content (543.9 kJ; except water in experiment 1). The first experiment comprised measures of subjective appetite, whereas the second experiment included measures of subjective appetite together with appetite-related analytes (serum glucose, insulin and plasma GLP-1 and peptide YY). Reduced energy intake was observed following chocolate cow's milk and yogurt consumption compared with a water drink in the first experiment. Consistent with a reduction in energy intake, subjective appetite (combined appetite score) was significantly lower following 2 % (2 g/100 g) fat cow's milk compared with the yogurt drink only. No additional effects were observed concerning energy intake following the consumption of 2 % (2 g/100 g) fat cow's milk and fruit punch or on subjective measures of appetite after 1 % fat chocolate cow's milk, 1.5 % (1.5 g/100 g) fat yogurt drink, fruit punch or water. Compared with the fruit punch preload, cow's milk consumption resulted in a significantly greater GLP-1 area under the curve. Nonetheless, *ad libitum* energy intake, insulin and glucose AUC were comparable between trials. Considering all aforementioned studies, it is important to consider that, in these studies, energy intake was principally assessed via *ad libitum* assessments which may not be reflective of free-living eating behaviour.

The mechanism(s) by which cow's milk consumption might influence eating behaviour are not fully understood, but there are



several plausible suggestions from the adult literature and constituents of cow's milk that may act synergistically to explain possible actions. In the studies where cow's milk reduced appetite and subsequent eating behaviour, it is probably unsurprising that cow's milk consumption suppressed energy intake at *ad libitum* assessment meals, considering the macronutrient composition of cow's milk compared with fruit-juice drinks and water. Cow's milk contains considerably more protein than fruit-juice drinks and water. Although it is not a universal finding, it is widely recognised that dietary proteins are more satiating than energetic equivalents of carbohydrate and fat under most conditions, commonly suppressing eating behaviour at the next meal^(113–115). Consequently, cow's milk proteins (whey and casein, and their products of digestion) may act to potentiate peptides of gastrointestinal, pancreatic and adipose tissue origin, increasing perceptions of satiety and acutely reducing energy intake (anorexigenic behaviours)⁽¹¹³⁾. Moreover, medium-chain triglycerides, conjugated linoleic acid (CLA) and lactose may also be implicated in the reduction of energy intake after cow's milk intake⁽¹⁰⁷⁾. Medium-chain triglycerides are absorbed directly into the portal circulation and transported to the liver for rapid oxidation. A combined action of increased energy expenditure, decreased fat deposition and increased satiety may reduce of energy intake via pre-absorptive signals, post-absorptive changes in metabolites and appetite-related analytes⁽¹⁰⁷⁾, and similar appetite-related analyte responses have been observed with lactose⁽¹¹⁶⁾. When considering CLA intake and its potential implications in appetite regulation, cow's milk (and dairy) and cattle meat (cows and lamb) represent the almost exclusive dietary sources⁽¹¹⁷⁾ where the predominant isomer of CLA (accounting for more than 90 % of the total CLA intake) is *cis*-9,*trans*-11-CLA. It is, however, strongly proposed that other isomers, such as *trans*-10, *cis*-12-CLA, may influence body-weight and fat changes⁽¹¹⁸⁾. In agreement with an earlier narrative review⁽³⁴⁾, it remains unknown whether cow's milk (and dairy), when providing physiological doses of CLA, can elicit any meaningful impact on appetite and body-weight regulation in humans. This is especially prudent when considering experimental design, age, sex, energy intake and CLA metabolism of the participants, and the dose and chemical form of the CLA isomer administered, as differences may arise solely from these methodological differences⁽¹¹⁹⁾. From a child perspective, potential underlying mechanisms of CLA on appetite regulation are poorly understood, though evidence from adult studies suggests that CLA can inhibit fatty acid synthase and stearoyl-CoA desaturase-1⁽¹¹⁸⁾, enhance fat oxidation and thermogenesis and reduce lipogenesis and preadipocyte differentiation and proliferation⁽¹²⁰⁾.

In summary, evidence suggests that cow's milk may have a unique potential to influence elements of energy balance. Macro- and micronutrients and other bioactive constituents might act individually, though probably synergistically, to impart beneficial effects on energy balance and body mass regulation through actions related to appetite, eating behaviour and metabolism. However, there is little mechanistic exploration of cow's milk consumption and appetite regulation from a paediatric perspective. Understanding the relationship between cow's milk consumption and appetite regulation could provide novel nutritional interventions to contribute toward the fight against childhood overweight and obesity⁽¹²¹⁾, whilst bolstering nutritional status and improving elements of cognitive function and

hydration. Controlled intervention studies are necessary to determine the best possible timings to administer cow's milk and establish whether consumption delivers these benefits when administered during the school day.

Collectively, the effects of cow's milk intake on appetite regulation in primary-school-aged children are unclear and could be dependent on BMI. The studies summarised suggest that mid-morning milk consumption influences eating behaviour at the next meal in overweight and obese children, showing that cow's milk could be beneficial for reducing body mass. In lean children, the evidence suggests there is no effect of cow's milk consumption on appetite and eating behaviour, but cow's milk may boost the nutritional quality of the diet. There is much scope for further studies to clarify the role of cow's milk on appetite and eating behaviour, especially in a free-living school environment where methodological approaches are more reflective of habitual eating behaviours. In addition, it will be important to fully distinguish the effects of cow's milk on appetite- and on metabolism-related peptides and subsequent eating behaviour.

Cognitive Function

Compared with children with better dietary quality, those with nutritional inadequacies demonstrate decreased attention and academic performance⁽¹²²⁾. Aside from improving nutritional status and dietary quality, emerging evidence illustrates that cow's milk may positively influence cognitive function in primary-school children^(43,123,124). Varying in duration from 2 h to 9 months, studies of an intervention-based nature generally report that cow's milk consumption increases elements of cognitive function, though some of the specific outcome measures demonstrate no effect or, in some cases, a negative effect of cow's milk. These studies highlight that consideration should be given to potential moderators such as sex and to the use of cow's milk as a non-stigmatised method for providing nutrients through fortification⁽¹²⁵⁾. One non-interventional study considered the adherence of $n = 1595$ children to Canadian nutrition recommendations in Grade 5 (10–11 years) and in relation to academic achievement in the provincial achievement tests taken approximately 1 year later⁽¹²⁶⁾. A positive association was identified for boys, with those who met the nutrition recommendations for milk and alternatives (at the time of the study: 3–4 servings/d) scoring 3.45 % better on an average measure of Math and Language Arts tests than those who did not meet the recommendations⁽¹²⁶⁾. An account of the intervention studies follows, but overall, the varying methodological approaches used suggest caution is needed in making firm conclusions about potential links between cow's milk consumption and improved cognitive function.

In a study⁽¹²⁴⁾ involving $n = 469$ boys and girls (mean age 8 years), evaluating the effects of daily mid-morning cow's milk consumption on physical, mental and school performance, researchers found that a school feeding scheme focusing on daily cow's milk intake had beneficial effects on school performance for girls. In this study, children received a serving of cow's milk (250 ml) daily for 12 weeks. Assessments of physical, mental and school performance were conducted prior to and at the end of the 12-week supplementation period. Similarly, in a smaller study⁽¹²³⁾ involving 40 children (mean age 11 years), the effects of a carbohydrate drink, a cow's milk drink or a

combination of both on subsequent cognitive function (processing speed, memory, attention and perceptual speed) were assessed over a 3 h period⁽¹²³⁾. Findings showed cow's milk consumption improved short-term memory. Children were able to recall 0.7–0.8 more words compared with 0.5 fewer words after the carbohydrate drink. However, this effect was only observed in girls and not boys^(123,124). There were slightly more mixed findings from a crossover study in which 84 children (mean age 10 years) were given 237 ml of cow's milk or apple juice⁽¹²⁷⁾. While the beverages were not standardised for temperature, participants were asked to complete computerised tasks of inhibitory control, speeded working memory and sustained attention at baseline and 30 min, 90 min and 120 min following beverage consumption. Although the significant results following cow's milk compared with juice consumption were, again, only apparent in girls, there was a negative effect (decreased working memory accuracy) alongside the positive one (improved reaction time on the sustained attention task). There were non-significant trends in the opposite direction for boys. No significant effects of the beverages were observed for on-task behaviour during the testing.

The mechanisms responsible for the beneficial effects of cow's milk on improved cognitive function are unclear. One suggestion is a sustained blood glucose response following consumption^(128,129). Findings from Anderson *et al.*⁽¹²⁷⁾, an adult-based study, suggest that glucoregulation may play a role, as participants with higher fasting glucose levels demonstrated faster reaction times on an inhibition task following cow's milk compared with juice. There were, however, no sex differences in initial glucose elevation or change in glucose levels over time to explain the apparent sex differences in cognition. Such findings suggest a likelihood that factors other than glucoregulation explain why cow's milk differentially affects the cognition and behaviour of boys and girls. The micronutrient content of cow's milk is another potential mechanism for the observed effects⁽⁴³⁾. Low intakes of vitamin B₁, folate and vitamin B₁₂ affect short-term memory and impair learning, causing low cognitive scores and development in primary-school-age children⁽¹³⁰⁾. In addition, low iron and riboflavin intake may adversely affect motor skill development and psychomotor performance⁽¹³¹⁾. All of these micronutrients are heavily present in cow's milk. In one of the longer intervention-based studies available (5 months), Kuriyan and colleagues⁽⁴³⁾ attempted to establish if fortification of cow's milk with multiple micronutrients influenced the mental and physical performance of children (7–10 years) compared with an unfortified cow's milk drink. The children were randomised into groups and were provided with cow's milk (2 × 200 ml) 6 d per week for 5 months, with assessments of attention and executive function conducted at baseline and 5 months. The findings showed improved cognitive and physical performance in both groups, illustrating that further fortification of cow's milk provided no additional benefits to cognitive and physical performance but did improve some elements of nutritional status⁽³⁹⁾. Finally, in a double-blind RCT (9 months), the learning potential of 7–9-year-olds (31.9 % with moderate or severe iodine deficiency at baseline) improved to a greater degree following 200 ml daily of fortified cow's milk [fortified with 45 µg iodine

(given as potassium iodide) and other micronutrients] than following non-fortified milk (20.8 µg iodine)⁽¹³²⁾.

Some of the components of cow's milk may be detrimental for the cognitive performance of lactase-deficient children. In a study involving children of 5–6 years (85 % lactase deficient), information processing efficiency was assessed after 5 d consumption of 150 ml twice daily of conventional cow's milk (containing A1 and A2 β-casein) or cow's milk containing A2 β-casein only⁽¹³³⁾. While post-intervention response times were significantly improved from baseline for both cow's milk products, error rates decreased in the A2 β-casein only condition. Furthermore, consuming conventional cow's milk in the second phase of the crossover study appeared to undo the positive effects on error rate obtained from the A2 β-casein only milk in the first phase, which were maintained through the 9 d wash-out period.

It is prudent to highlight that none of the studies of cognition in primary-school children have compared cow's milk with a control beverage. It is therefore difficult to ascertain whether these studies simply show less detrimental effects of cow's milk than comparator beverages. Furthermore, where no comparators have been employed, this could reflect practice effects. Nonetheless, based on the published studies, and bearing in mind the varying methodological approaches employed, it is difficult to ascertain the role of cow's milk in cognitive function for primary-school-aged children, but it may at the very least be a useful conduit for nutrient fortification. This could be particularly meaningful within the school environment. The intervention studies did not measure academic achievement directly, but their findings have relevant scholastic implications that warrant further investigation using RCTs to establish if increased milk consumption influences academic achievement in primary-school-aged children, especially given the identification of a positive association for boys between academic achievement and meeting the recommendations for consumption of milk and alternatives⁽¹²⁶⁾.

Future Directions and Conclusions

The aim of this narrative review was to evaluate evidence for the potential nutritional-, physical- and health-related benefits of cow's milk consumption for primary-school-aged children (4–11 years). Cow's milk consumption (both plain and flavoured) improves nutritional status without adversely impacting body mass and body composition⁽³⁵⁾. With some confidence, cow's milk also appears beneficial for hydration, dental and bone health and to have a beneficial-to-neutral effect on physical stature and appetite. Due to conflicting studies, reaching a conclusion has proven difficult concerning cow's milk and cognitive function; therefore, a level of caution should be exercised when interpreting these results. All areas, however, would benefit from further robust investigation, especially in free-living school settings, to verify conclusions. Improving elements of cognitive function, hydration and appetite regulation could have important long-term health and scholastic implications that should certainly be explored further. Indeed, recent research involving adolescent populations illustrates that high intakes of cow's milk are positively associated with academic performance⁽¹³⁴⁾ and increased motivation for learning⁽¹³⁴⁾ and



impact favourably on eating behaviour following acute and chronic consumption⁽¹³⁵⁾.

Despite a growing body of scientific literature exploring the potential benefits of cow's milk consumption, there are several pertinent knowledge gaps that would benefit from further investigation. This is particularly true for cognitive function, hydration status and appetite regulation, where there is little research available, especially in free-living school settings. Further research, especially of a robust and methodologically sound nature (such as double-blind randomised controlled trials), should seek to explore the mechanisms responsible for any effects observed. At present, few studies on this population have attempted to establish mechanisms. Most purported mechanisms given throughout this narrative review come from adult-specific data and should therefore be interpreted with some caution, as the observations cited may not always be replicated in children. When working with child populations, there are various considerations that must be accounted for. The methodological approaches deemed most appropriate for the study of cognitive function, hydration status and appetite regulation in children will differ according to the objective of the study, type of data required and resources available⁽¹³⁶⁾. In children, it is of great importance to adopt approaches that are non-invasive with a low level of participant stress. Current techniques available to assess cognitive function, hydration status and appetite are arguably invasive, elicit a moderate level of participant stress and have increased ethical risk. This might explain the current lack of studies and mechanistic insight from a child perspective. In recent years, however, research groups have been pursuing non-invasive techniques that offer an opportunity to conduct comprehensive mechanistic work in vulnerable populations. For example, developments in appetite and metabolism research have identified fingertip-capillary blood sampling as an efficacious, comparable and reproducible alternative to antecubital-venous blood sampling for the quantification of appetite-related peptides^(137,138). These developments will certainly help provide a better understanding of mechanisms that influence appetite and eating behaviour in younger populations, where traditional methods of venous blood sampling might be contraindicated. Furthermore, fingertip-capillary blood sampling offers many advantages, including simplistic application, cost/time effectiveness and reduced volume of blood required for analysis⁽¹³⁹⁾.

Considering the nutritional-, physical- and health-related impact of cow's milk avoidance, the evidence begins to highlight the importance of increasing cow's milk consumption. Cow's milk is a naturally nutrient-dense foodstuff, providing a significant contribution of several essential nutrients and bioactive constituents that potentially impact human health favourably. Establishing and shaping healthy eating behaviours during the primary-school years is vital. Dietary behaviours shaped throughout the childhood years progress through to adolescence and adulthood⁽¹⁴⁰⁾, making healthy eating environments crucial. For primary-school-aged children, the school setting may be an ideal environment to promote cow's milk consumption, and school milk schemes are a great place to start developing healthy eating behaviours, given 35–40 % of children's daily nutritional needs are met at school⁽¹⁴¹⁾. Cow's milk is a readily available, accessible

and affordable means of providing valuable essential nutrients to the diets of primary-school children. Based on the evidence presented in this manuscript, there appears no reason for primary-school children to limit cow's milk consumption, and there may, in fact, be many potential benefits to milk consumption.

Author Contributions

Benjamin Green and Penny Rumbold conceived and planned this narrative review. Penny Rumbold, Nicola McCulloch, Ruth Boldon and Benjamin Green sourced the relevant literature. All authors contributed to the writing and critically reviewed and approved the final manuscript.

Financial Support

This work was supported by funding from Cool Milk Ltd. The funding sponsors had no role in the design, collection, analysis or study interpretation, in the writing of the manuscript, and in the decision to publish the results.

Conflicts of Interest

Penny Rumbold has previously received funding from The Dairy Council (UK), Nourishmenow and Cool Milk Ltd. Benjamin Green has previously been supported by The Dairy Council (UK) in the award of a PhD studentship and has received funding from Cool Milk Ltd and is presently an employee of Danone Specialised Nutrition. Lewis James has previously received funding from Volac International Ltd. Emma Stevenson has previously received funding from The Dairy Council (UK) and Arla Food Ingredients. Collectively, the above (with the exception of Cool Milk Ltd) was not related in any way to the work presented in this article. Nicola McCulloch, Ruth Boldon and Crystal Haskell-Ramsay declare no conflicts of interest.

References

1. Nagpal R, Behare PV, Kumar M, *et al.* (2012) Milk, milk products, and disease free health: an updated overview. *Critical Reviews in Food Science and Nutrition* **52**(4):321–333.
2. Ebringer L, Ferenčík M, Krajčovič J. (2008) Beneficial health effects of milk and fermented dairy products – Review. *Folia Microbiologica* **53**(5):378–394.
3. Nongonierma AB, FitzGerald RJ. (2015) The scientific evidence for the role of milk protein-derived bioactive peptides in humans: A review. *Journal of Functional Foods* **17**: 640–656.
4. Ferreira AM, Bislev SL, Bendixen E, *et al.* (2013) The mammary gland in domestic ruminants: A systems biology perspective. *Journal of Proteomics* **94**:110–123.
5. Rezaei R, Wu Z, Hou Y, *et al.* (2016) Amino acids and mammary gland development: nutritional implications for milk production and neonatal growth. *Journal of Animal Science and Biotechnology* **7**(1):20.
6. Givens DI, Livingstone KM, Pickering JE, *et al.* (2014) Milk: White elixir or white poison? An examination of the associations between dairy consumption and disease in human subjects. *Animal Frontiers* **4**(2):8–15.
7. Velten H. (2010) Milk: A global history: Reaktion Books.

8. Ludwig DS, Willett WC. (2013) Three daily servings of reduced-fat milk: An evidence-based recommendation? *JAMA Pediatrics* **167**(9):788–789.
9. Melnik BC. (2009) Milk – The promoter of chronic Western diseases. *Medical Hypotheses* **72**(6):631–639.
10. Morgan PJ, Warren JM, Lubans DR, *et al.* (2010) The impact of nutrition education with and without a school garden on knowledge, vegetable intake and preferences and quality of school life among primary-school students. *Public Health Nutrition* **13**(11):1931–1940.
11. Patton GC, Viner R. (2007) Pubertal transitions in health. *The Lancet* **369**(9567):1130–1139.
12. Wilson T, Temple NJ (2016) How beverages impact health and nutrition. In: Wilson T, Temple NJ, editors. *Beverage Impacts on Health and Nutrition: Second Edition*. Cham: Springer International Publishing. p. 3–9.
13. Rozenberg S, Body J-J, Bruyère O, *et al.* (2016) Effects of dairy products consumption on health: benefits and beliefs – A commentary from the Belgian Bone Club and the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases. *Calcified Tissue International* **98**(1):1–17.
14. Clare DA, Swaisgood HE. (2000) Bioactive milk peptides: A prospectus. *Journal of Dairy Science* **83**(6):1187–1195.
15. Parodi PW. (2004) Milk fat in human nutrition. *Australian Journal of Dairy Technology* **59**(1):3.
16. MacGibbon AKH, Taylor MW (2006) Composition and structure of bovine milk lipids. In: Fox PF, McSweeney PLH, editors. *Advanced Dairy Chemistry Volume 2 Lipids*. Boston, MA: Springer US. p. 1–42.
17. Wilkinson SB, Tarnopolsky MA, MacDonald MJ, *et al.* (2007) Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *The American Journal of Clinical Nutrition* **85**(4):1031–1040.
18. Tunick MH, Van Hekken DL. (2015) Dairy products and health: Recent insights. *Journal of Agricultural and Food Chemistry* **63**(43):9381–9388.
19. Bates B, Lennox A, Prentice A, *et al.* (2014) National diet and nutrition survey: Results from years 1, 2, 3 and 4 (combined) of the rolling programme (2008/2009–2011/2012): A survey carried out on behalf of public health England and the food standards agency: Public Health England.
20. Roberts C, Steer T, Maplethorpe N, *et al.* (2018) National Diet and Nutrition Survey: Results from years 7 and 8 (combined) of the rolling programme (2014/2015–2015/2016).
21. Cavadini C, Siega-Riz AM, Popkin BM. (2000) US adolescent food intake trends from 1965 to 1996. *Archives of Disease in Childhood* **83**(1):18–24.
22. Fiorito LM, Mitchell DC, Smiciklas-Wright H, *et al.* (2006) Dairy and dairy-related nutrient intake during middle childhood. *Journal of the American Dietetic Association* **106**(4):534–542.
23. Baird DL, Syrette J, Hendrie GA, *et al.* (2012) Dairy food intake of Australian children and adolescents 2–16 years of age: 2007 Australian National Children's Nutrition and Physical Activity Survey. *Public Health Nutrition* **15**(11):2060–2073.
24. van Rossum CT, Fransen HP, Verkaik-Kloosterman J, *et al.* (2011) Dutch National Food Consumption Survey 2007–2010: Diet of children and adults aged 7 to 69 years. RIVM report 350050006/2011:1–143.
25. Black RE, Williams SM, Jones IE, *et al.* (2002) Children who avoid drinking cow milk have low dietary calcium intakes and poor bone health. *The American Journal of Clinical Nutrition* **76**(3):675–680.
26. Appel LJ, Moore TJ, Obarzanek E, *et al.* (1997) A clinical trial of the effects of dietary patterns on blood pressure. *The New England Journal of Medicine* **336**(16):1117–1124.
27. Pereira MA, Jacobs J, David R., Van Horn L, *et al.* (2002) Dairy consumption, obesity, and the insulin resistance syndrome in young adults: The CARDIA Study. *JAMA* **287**(16):2081–2089.
28. Elwood PC, Pickering JE, Fehily AM. (2007) Milk and dairy consumption, diabetes and the metabolic syndrome: The Caerphilly prospective study. *Journal of Epidemiology and Community Health* **61**(8):695–698.
29. Rockell JEP, Williams SM, Taylor RW, *et al.* (2005) Two-year changes in bone and body composition in young children with a history of prolonged milk avoidance. *Osteoporosis International* **16**(9):1016–1023.
30. Soedamah-Muthu SS, de Goede J. (2018) Dairy consumption and cardiometabolic diseases: Systematic review and updated meta-analyses of prospective cohort studies. *Current Nutrition Reports* **7**(4):171–182.
31. Crichton GE, Bryan J, Buckley J, *et al.* (2011) Dairy consumption and metabolic syndrome: a systematic review of findings and methodological issues. *Obesity Reviews* **12**(5):e190–e201.
32. Abargouei AS, Janghorbani M, Salehi-Marzjafari M, *et al.* (2012) Effect of dairy consumption on weight and body composition in adults: a systematic review and meta-analysis of randomized controlled clinical trials. *International Journal of Obesity* **36**(12):1485–1493.
33. Barba G, Russo P. (2006) Dairy foods, dietary calcium and obesity: A short review of the evidence. *Nutrition, Metabolism and Cardiovascular Diseases* **16**(6):445–451.
34. Dougkas A, Reynolds CK, Givens ID, *et al.* (2011) Associations between dairy consumption and body weight: a review of the evidence and underlying mechanisms. *Nutrition Research Reviews* **24**(1):72–95.
35. Dougkas A, Barr S, Reddy S, *et al.* (2019) A critical review of the role of milk and other dairy products in the development of obesity in children and adolescents. *Nutrition Research Reviews* **32**(1):106–127.
36. Cook J, Irwig LM, Chinn S, *et al.* (1979) The influence of availability of free school milk on the height of children in England and Scotland. *Journal of Epidemiology and Community Health* **33**(3):171–176.
37. Campmans-Kuijpers MJE, Singh-Povel C, Steijns J, *et al.* (2016) The association of dairy intake of children and adolescents with different food and nutrient intakes in the Netherlands. *BMC Pediatrics* **16**(1):2.
38. Cook J, Altman DG, Jacoby A, *et al.* (1975) The contribution made by school milk to the nutrition of primary schoolchildren. *British Journal of Nutrition* **34**(1):91–103.
39. LaRowe TL, Moeller SM, Adams AK. (2007) Beverage patterns, diet quality, and body mass index of US preschool and school-aged children. *Journal of the American Dietetic Association* **107**(7):1124–1133.
40. Murphy MM, Douglass JS, Johnson RK, *et al.* (2008) Drinking flavored or plain milk is positively associated with nutrient intake and is not associated with adverse effects on weight status in US children and adolescents. *Journal of the American Dietetic Association* **108**(4):631–639.
41. Rangan AM, Flood VM, Denyer G, *et al.* (2012) Dairy consumption and diet quality in a sample of Australian children. *Journal of the American College of Nutrition* **31**(3):185–193.
42. Wang J, Shang L, Johnson-Down L, *et al.* (2012) Contribution of flavoured milk on dietary intakes and its association with body mass index percentile in children at high risk of overweight. *The FASEB Journal* **26**(S1):lb389–lb389.
43. Kuriyan R, Thankachan P, Selvam S, *et al.* (2016) The effects of regular consumption of a multiple micronutrient fortified milk

- beverage on the micronutrient status of school children and on their mental and physical performance. *Clinical Nutrition* **35**(1):190–198.
44. Albala C, Ebbeling CB, Cifuentes M, *et al.* (2008) Effects of replacing the habitual consumption of sugar-sweetened beverages with milk in Chilean children. *The American Journal of Clinical Nutrition* **88**(3):605–611.
 45. Lien DTK, Nhung BT, Khan NC, *et al.* (2009) Impact of milk consumption on performance and health of primary school children in rural Vietnam. *Asia Pacific Journal of Clinical Nutrition* **18**:326–334.
 46. Fiorito LM, Marini M, Mitchell DC, *et al.* (2010) Girls' early sweetened carbonated beverage intake predicts different patterns of beverage and nutrient intake across childhood and adolescence. *Journal of the American Dietetic Association* **110**(4):543–550.
 47. Weir RR, Johnston M, Lewis C, *et al.* (2020) Vitamin D3 content of cows' milk produced in Northern Ireland and its efficacy as a vehicle for vitamin D fortification: A UK model. *International Journal of Food Sciences and Nutrition*: 1–9.
 48. Itonen ST, Erkkola M, Lamberg-Allardt CJE. (2018) Vitamin D fortification of fluid milk products and their contribution to vitamin D intake and vitamin D status in observational studies – A review. *Nutrients* **10**(8):1054.
 49. Hanks AS, Just DR, Wansink B. (2014) Chocolate milk consequences: A pilot study evaluating the consequences of banning chocolate milk in school cafeterias. *PLoS One* **9**(4): e91022.
 50. Henry C, Whiting SJ, Phillips T, *et al.* (2015) Impact of the removal of chocolate milk from school milk programs for children in Saskatoon, Canada. *Applied Physiology, Nutrition, and Metabolism* **40**(3):245–250.
 51. Fayet-Moore F. (2015) Effect of flavored milk vs plain milk on total milk intake and nutrient provision in children. *Nutrition Reviews* **74**(1):1–17.
 52. Rehm CD, Drownowski A, Monsivais P. (2015) Potential population-level nutritional impact of replacing whole and reduced-fat milk with low-fat and skim milk among US children aged 2–19 years. *Journal of Nutrition Education and Behavior* **47**(1):61–68.e61.
 53. Popkin BM, D'Anci KE, Rosenberg IH. (2010) Water, hydration, and health. *Nutrition Reviews* **68**(8):439–458.
 54. D'Anci KE, Constant F, Rosenberg IH. (2006) Hydration and cognitive function in children. *Nutrition Reviews* **64**(10): 457–464.
 55. Maughan RJ, Watson P, Cordery PA, *et al.* (2015) A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index. *The American Journal of Clinical Nutrition* **103**(3):717–723.
 56. Bonnet F, Lepicard EM, Cathrin L, *et al.* (2012) French children start their school day with a hydration deficit. *Annals of Nutrition and Metabolism* **60**(4):257–263.
 57. Montenegro-Bethancourt G, Johner SA, Remer T. (2013) Contribution of fruit and vegetable intake to hydration status in schoolchildren. *The American Journal of Clinical Nutrition* **98**(4):1103–1112.
 58. Volterman K, Moore D, Obeid J, *et al.* (2016) The effect of postexercise milk protein intake on rehydration of children. *Pediatric Exercise Science* **28**(2):286.
 59. Volterman KA, Obeid J, Wilk B, *et al.* (2014) Effect of milk consumption on rehydration in youth following exercise in the heat. *Applied Physiology, Nutrition, and Metabolism* **39**(11):1257–1264.
 60. James L. (2012) Milk ingestion in athletes and physically active individuals. *Nutrition Bulletin* **37**(3):257–261.
 61. Pegoretti C, Antunes AEC, Manchado-Gobatto FdB, *et al.* (2015) Milk: An alternative beverage for hydration? *Food and Nutrition Sciences* **6**(6):8.
 62. Shirreffs SM, Watson P, Maughan RJ. (2007) Milk as an effective post-exercise rehydration drink. *British Journal of Nutrition* **98**(1):173–180.
 63. James LJ, Stevenson EJ, Rumbold PLS, *et al.* (2019) Cow's milk as a post-exercise recovery drink: implications for performance and health. *European Journal of Sport Science* **19**(1):40–48.
 64. Maughan RJ, Leiper JB. (1995) Sodium intake and post-exercise rehydration in man. *European Journal of Applied Physiology and Occupational Physiology* **71**(4):311–319.
 65. Maughan RJ, Owen JH, Shirreffs SM, *et al.* (1994) Post-exercise rehydration in man: Effects of electrolyte addition to ingested fluids. *European Journal of Applied Physiology and Occupational Physiology* **69**(3):209–215.
 66. James LJ, Clayton D, Evans GH. (2011) Effect of milk protein addition to a carbohydrate–electrolyte rehydration solution ingested after exercise in the heat. *British Journal of Nutrition* **105**(3):393–399.
 67. James LJ, Evans GH, Madin J, *et al.* (2013) Effect of varying the concentrations of carbohydrate and milk protein in rehydration solutions ingested after exercise in the heat. *British Journal of Nutrition* **110**(7):1285–1291.
 68. Aimutis WR. (2004) Bioactive properties of milk proteins with particular focus on anticariogenesis. *The Journal of Nutrition* **134**(4):989S–995S.
 69. Levine RS. (2001) Milk, flavoured milk products and caries. *British Dental Journal* **191**(1):20–20.
 70. Moynihan P. (2000) Foods and factors that protect against dental caries. *Nutrition Bulletin* **25**(4):281–286.
 71. Ferrazzano G, Cantile T, Quarto M, *et al.* (2008) Protective effect of yogurt extract on dental enamel demineralization in vitro. *Australian Dental Journal* **53**(4):314–319.
 72. Levine RS, Nugent ZJ, Rudolf MC, *et al.* (2007) Dietary patterns, toothbrushing habits and caries experience of schoolchildren in West Yorkshire, England. *Community Dental Health* **24**(2):82–87.
 73. Llana C, Forner L. (2008) Dietary habits in a child population in relation to caries experience. *Caries Research* **42**(5):387–393.
 74. Petti S, Simonetti R, D'Arca AS. (1997) The effect of milk and sucrose consumption on caries in 6-to-11-year-old Italian schoolchildren. *European Journal of Epidemiology* **13**(6):659–664.
 75. Curtis AM, VanBuren J, Cavanaugh JE, *et al.* (2018) Longitudinal associations between dental caries increment and risk factors in late childhood and adolescence. *Journal of Public Health and Dentistry* **78**(4):321–328.
 76. Giacaman RA, Campos P, Muñoz-Sandoval C, *et al.* (2013) Cariogenic potential of commercial sweeteners in an experimental biofilm caries model on enamel. *Archives of Oral Biology* **58**(9):1116–1122.
 77. Cross KJ, Huq NL, Palamara JE, *et al.* (2005) Physicochemical characterization of casein phosphopeptide-amorphous calcium phosphate nanocomplexes. *Journal of Biological Chemistry* **280**(15):15362–15369.
 78. McDougall WA. (1977) Effect of milk on enamel demineralization and remineralization in vitro. *Caries Research* **11**(3):166–172.
 79. Weaver CM. (2014) Milk consumption and bone health. *JAMA Pediatrics* **168**(1):12–13.
 80. Wiley AS. (2010) Dairy and milk consumption and child growth: Is BMI involved? *An analysis of NHANES 1999–2004. American Journal of Human Biology* **22**(4):517–525.

81. Leighton G, Clark ML. (1929) Milk consumption and the growth of school children: Second preliminary report on tests to the Scottish Board of Health. *British Medical Journal* **1**(3548):23–25.
82. Orr JB. (1928) Influence of amount of milk consumption on the rate of growth of school children. *British Medical Journal* **1**(3499):140–141.
83. Bonjour JP, Carrie AL, Ferrari S, *et al.* (1997) Calcium-enriched foods and bone mass growth in prepubertal girls: A randomized, double-blind, placebo-controlled trial. *The Journal of Clinical Investigation* **99**(6):1287–1294.
84. Chan GM, Hoffman K, McMurry M. (1995) Effects of dairy products on bone and body composition in pubertal girls. *The Journal of Pediatrics* **126**(4):551–556.
85. Du X, Zhu K, Trube A, *et al.* (2004) School-milk intervention trial enhances growth and bone mineral accretion in Chinese girls aged 10–12 years in Beijing. *British Journal of Nutrition* **92**(1):159–168.
86. Goulding A, Rockell JEP, Black RE, *et al.* (2004) Children who avoid drinking cow's milk are at increased risk for prepubertal bone fractures. *Journal of the American Dietetic Association* **104**(2):250–253.
87. Iuliano-Burns S, Wang XF, Evans A, *et al.* (2006) Skeletal benefits from calcium supplementation are limited in children with calcium intakes near 800 mg daily. *Osteoporosis International* **17**(12):1794–1800.
88. Zhou Y, Hu Y, Ma Z, *et al.* (2011) Relation of long-term consumption of school milk to bone density in children. *Journal of Hygiene Research* **40**(1):65–67.
89. Feskanich D, Bischoff-Ferrari HA, Frazier AL, *et al.* (2014) Milk consumption during teenage years and risk of hip fractures in older adults. *JAMA Pediatrics* **168**(1):54–60.
90. Sunycz JA. (2008) The use of calcium and vitamin D in the management of osteoporosis. *Therapeutics and Clinical Risk Management* **4**(4):827–836.
91. Gaucheron F. (2011) Milk and dairy products: A unique micro-nutrient combination. *Journal of the American College of Nutrition* **30**(sup5):400S–409S.
92. Caroli A, Poli A, Ricotta D, *et al.* (2011) Invited review: Dairy intake and bone health: A viewpoint from the state of the art. *Journal of Dairy Science* **94**(11):5249–5262.
93. Rizzoli R. (2014) Dairy products, yogurts, and bone health. *The American Journal of Clinical Nutrition* **99**(5):1256S–1262S.
94. Baker IA, Elwood PC, Hughes J, *et al.* (1980) A randomised controlled trial of the effect of the provision of free school milk on the growth of children. *Journal of Epidemiology and Community Health* **34**(1):31–34.
95. Guo Q, Wang B, Cao S, *et al.* (2020) Association between milk intake and childhood growth: results from a nationwide cross-sectional survey. *International Journal of Obesity*.
96. Matkovic V, Landoll JD, Badenhop-Stevens NE, *et al.* (2004) Nutrition influences skeletal development from childhood to adulthood: A study of hip, spine, and forearm in adolescent females. *The Journal of Nutrition* **134**(3):701S–705S.
97. Okada T. (2004) Effect of cow milk consumption on longitudinal height gain in children. *American Journal of Clinical Nutrition* **80**(4):1088–1089.
98. Rona RJ, Chinn S. (1989) School meals, school milk and height of primary school children in England and Scotland in the eighties. *Journal of Epidemiology and Community Health* **43**(1):66–71.
99. Wiley AS. (2005) Does milk make children grow? *Relationships between milk consumption and height in NHANES 1999–2002*. *American Journal of Human Biology* **17**(4):425–441.
100. Matkovic V, Jelic T, Wardlaw GM, *et al.* (1994) Timing of peak bone mass in Caucasian females and its implication for the prevention of osteoporosis. Inference from a cross-sectional model. *The Journal of Clinical Investigation* **93**(2):799–808.
101. Luiza Loro M, Sayre J, Roe TF, *et al.* (2000) Early identification of children predisposed to low peak bone mass and osteoporosis later in life. *The Journal of Clinical Endocrinology and Metabolism* **85**(10):3908–3918.
102. Hoppe C, Mølgaard C, Vaag A, *et al.* (2005) High intakes of milk, but not meat, increase s-insulin and insulin resistance in 8-year-old boys. *European Journal of Clinical Nutrition* **59**(3):393–398.
103. Hoppe C, Udam TR, Lauritzen L, *et al.* (2004) Animal protein intake, serum insulin-like growth factor I, and growth in healthy 2.5-y-old Danish children. *American Journal of Clinical Nutrition* **80**(2):447–452.
104. Rogers I, Emmett P, Gunnell D, *et al.* (2006) Milk as a food for growth? The insulin-like growth factors link. *Public Health Nutrition* **9**(3):359–368.
105. Woods SC, Lutz TA, Geary N, *et al.* (2006) Pancreatic signals controlling food intake; insulin, glucagon and amylin. *Philosophical Transactions of the Royal Society B* **361**(1471):1219–1235.
106. Berthoud H-R. (2006) Homeostatic and non-homeostatic pathways involved in the control of food intake and energy balance. *Obesity* **14**(S8):197S–200S.
107. Aziz A, Anderson GH. (2007) The effects of dairy components on food intake and satiety. In: Saarela M, editor. *Functional Dairy Products*: Woodhead Publishing. p. 19–45.
108. Kavezade S, Mozaffari-Khosravi H, Aflatoonian M, *et al.* (2018) The effects of whole milk compared to skim milk and apple juice consumption in breakfast on appetite and energy intake in obese children: A three-way randomized crossover clinical trial. *BMC Nutrition* **4**(1):44.
109. Mehrabani S, Safavi SM, Mehrabani S, *et al.* (2016) Effects of low-fat milk consumption at breakfast on satiety and short-term energy intake in 10- to 12-year-old obese boys. *European Journal of Nutrition* **55**(4):1389–1396.
110. Mehrabani S, Salehi-Abargouei A, Asemi M, *et al.* (2014) Effect of low-fat milk consumption compared to apple juice and water on the energy intake among 10-12-year-old obese boys: A three-way cross-over clinical trial. *International Journal of Preventive Medicine* **5**(11):1405–1411.
111. Rumbold PLS, Dodd-Reynolds CJ, Stevenson EJ. (2013) Informing primary school nutritional policy: Effects of mid-morning snacks on appetite and energy control. *Food and Nutrition Sciences* **4**(5):529–537.
112. Vien S, Patel B, Panahi S, *et al.* (2014) The effect of fluid dairy products on food intake, glycemic and appetite hormone responses in children (120.3). *The FASEB Journal* **28**(S1):120–123.
113. Anderson GH, Moore SE. (2004) Dietary proteins in the regulation of food intake and body weight in humans. *The Journal of Nutrition* **134**(4):974S–979S.
114. Astrup A. (2005) The satiating power of protein – a key to obesity prevention? *The American Journal of Clinical Nutrition* **82**(1):1–2.
115. Rolls BJ, Hetherington M, Burley VJ. (1988) The specificity of satiety: The influence of foods of different macronutrient content on the development of satiety. *Physiology & Behavior* **43**(2):145–153.
116. Bowen J, Noakes M, Trenergy C, *et al.* (2006) Energy intake, ghrelin, and cholecystokinin after different carbohydrate and protein preloads in overweight men. *The Journal of Clinical Endocrinology and Metabolism* **91**(4):1477–1483.



117. Terpstra AH. (2004) Effect of conjugated linoleic acid on body composition and plasma lipids in humans: An overview of the literature. *American Journal of Clinical Nutrition* **79**(3):352–361.
118. Li J-J, Huang CJ, Xie D. (2008) Anti-obesity effects of conjugated linoleic acid, docosahexaenoic acid, and eicosapentaenoic acid. *Molecular Nutrition and Food Research* **52**(6):631–645.
119. Plourde M, Jew S, Cunnane SC, *et al.* (2008) Conjugated linoleic acids: Why the discrepancy between animal and human studies? *Nutrition Reviews* **66**(7):415–421.
120. Wang YW, Jones PJH. (2004) Conjugated linoleic acid and obesity control: Efficacy and mechanisms. *International Journal of Obesity* **28**(8):941–955.
121. Jo H. (2008) Understanding and addressing the epidemic of obesity: An energy balance perspective. *Journal of Cardiopulmonary Rehabilitation and Prevention* **28**(5):345.
122. Florence MD, Asbridge M, Veugelers PJ. (2008) Diet quality and academic performance. *78*(4):209–215.
123. Brindal E, Baird D, Slater A, *et al.* (2013) The effect of beverages varying in glycaemic load on postprandial glucose responses, appetite and cognition in 10–12-year-old school children. *British Journal of Nutrition* **110**(3):529–537.
124. Rahmani K, Djazayeri A, Habibi MI, *et al.* (2011) Effects of daily milk supplementation on improving the physical and mental function as well as school performance among children: Results from a school feeding program. *Journal of Research in Medical Sciences* **16**(4):469–476.
125. Petrova D, Bernabeu Litrán MA, García-Mármol E, *et al.* (2019) Effects of fortified milk on cognitive abilities in school-aged children: Results from a randomized-controlled trial. *European Journal of Nutrition* **58**(5):1863–1872.
126. Faught EL, Montemurro G, Storey KE, *et al.* (2017) Adherence to dietary recommendations supports Canadian children's academic achievement. *Canadian Journal of Dietetic Practice and Research* **78**(3):102–108.
127. Anderson JR, Gunstad J, Updegraff J, *et al.* (2020) Biological sex and glucoregulation modulate postprandial cognition following dairy milk and fruit juice in healthy school-age children. *Nutritional Neuroscience* **23**(5):374–383.
128. Amiel SA, Pottinger RC, Archibald HR, *et al.* (1991) Effect of antecedent glucose control on cerebral function during hypoglycemia. *Diabetes Care* **14**(2):109–118.
129. Gold AE, MacLeod KM, Thomson KJ, *et al.* (1995) Cognitive function during insulin-induced hypoglycemia in humans: Short-term cerebral adaptation does not occur. *Psychopharmacology* **119**(3):325–333.
130. Vaz M, Pauline M, Unni US, *et al.* (2011) Micronutrient supplementation improves physical performance measures in Asian Indian school-age children. *The Journal of Nutrition* **141**(11):2017–2023.
131. Swaminathan S, Edward BS, Kurpad AV. (2013) Micronutrient deficiency and cognitive and physical performance in Indian children. *European Journal of Clinical Nutrition* **67**(5):467–474.
132. Zahrou FE, Azlaf M, El Menchawy I, *et al.* (2016) Fortified iodine milk improves iodine status and cognitive abilities in schoolchildren aged 7–9 years living in a rural mountainous area of Morocco. *Journal of Nutrition and Metabolism* **2016**:8468594.
133. Sheng X, Li Z, Ni J, *et al.* (2019) Effects of conventional milk versus milk containing only A2 β -casein on digestion in Chinese children: A randomized study. *Journal of Pediatric Gastroenterology and Nutrition* **69**(3):375–382.
134. Kim SH, Kim WK, Kang M-H. (2016) Relationships between milk consumption and academic performance, learning motivation and strategy, and personality in Korean adolescents. *Nutrition Research and Practice* **10**(2):198–205.
135. Green BP, Stevenson EJ, Rumbold PLS. (2017) Metabolic, endocrine and appetite-related responses to acute and daily milk snack consumption in healthy, adolescent males. *Appetite* **108**:93–103.
136. Gibbons C, Finlayson G, Dalton M, *et al.* (2014) Metabolic phenotyping guidelines: Studying eating behaviour in humans. *Journal of Endocrinology* **222**(2):G1.
137. Allsop S, Rumbold PLS, Green BP. (2016) The between-day reproducibility of fasting, satiety-related analytes, in 8 to 11 year-old boys. *Physiology & Behavior* **164**:207–213.
138. Green BP, Gonzalez JT, Thomas K, *et al.* (2014) Agreement between fingertip-capillary and antecubital-venous appetite-related peptides. *Endocrine Connections* **3**(4):233.
139. Dayre McNally J, Matheson LA, Sankaran K, *et al.* (2008) Capillary blood sampling as an alternative to venipuncture in the assessment of serum 25 hydroxyvitamin D levels. *The Journal of Steroid Biochemistry and Molecular Biology* **112**(1):164–168.
140. Lake AA, Mathers JC, Rugg-Gunn AJ, *et al.* (2006) Longitudinal change in food habits between adolescence (11–12 years) and adulthood (32–33 years): the ASH30 Study. *Journal of Public Health* **28**(1):10–16.
141. Gortmaker SL, Peterson K, Wiecha J, *et al.* (1999) Reducing obesity via a school-based interdisciplinary intervention among youth: Planet Health. *Archives of Pediatrics & Adolescent Medicine* **153**(4):409–418.