

Involvement of PLA₂, COX and LOX in *Rhinella arenarum* oocyte maturation

Maria Eugenia Ortiz², Marta Inés Bühler³ and Liliana Isabel Zelarayán¹

Centro de Referencias para Lactobacilos (CERELA)-CONICET, Tucumán, Argentina; and Instituto de Biología, Facultad de Bioqca, Qca. y Farmacia, INSIBIO-UNT, Tucumán, Argentina

Date submitted: 14.08.2012. Date accepted: 28.10.2012

Summary

In *Rhinella arenarum*, progesterone is the physiological nuclear maturation inducer that interacts with the oocyte surface and starts a cascade of events that leads to germinal vesicle breakdown (GVBD). Polyunsaturated fatty acids and their metabolites produced through cyclooxygenase (COX) and lipoxygenase (LOX) pathways play an important role in reproductive processes. In amphibians, to date, the role of arachidonic acid (AA) metabolites in progesterone (P₄)-induced oocyte maturation has not been clarified. In this work we studied the participation of three enzymes involved in AA metabolism – phospholipase A₂ (PLA₂), COX and LOX in *Rhinella arenarum* oocyte maturation. PLA₂ activation induced maturation in *Rhinella arenarum* oocytes in a dose-dependent manner. Oocytes when treated with 0.08 μM melittin showed the highest response (78 ± 6% GVBD). In follicles, PLA₂ activation did not significantly induce maturation at the assayed doses (12 ± 3% GVBD). PLA₂ inhibition with quinacrine prevented melittin-induced GVBD in a dose-dependent manner, however PLA₂ inactivation did not affect P₄-induced maturation. This finding suggests that PLA₂ is not the only phospholipase involved in P₄-induced maturation in this species. P₄-induced oocyte maturation was inhibited by the COX inhibitors indomethacin and rofecoxib (65 ± 3% and 63 ± 3% GVBD, respectively), although COX activity was never blocked by their addition. Follicles showed a similar response following the addition of these inhibitors. Participation of LOX metabolites in maturation seems to be correlated with seasonal variation in ovarian response to P₄. During the February to June period (low P₄ response), LOX inhibition by nordihydroguaiaretic acid or lysine clonixinate increased maturation by up to 70%. In contrast, during the July to January period (high P₄ response), LOX inhibition had no effect on hormone-induced maturation.

Keywords: Cyclooxygenase, Lipoxygenase, Oocyte maturation, Phospholipase A₂, *Rhinella arenarum*

Introduction

Amphibian oocytes are arrested in prophase I of meiosis until ovulation. At this time, follicular cells release progesterone (P₄) to induce meiosis resumption, however mechanisms involved in this process have not been fully elucidated. P₄ interacts

with the oocyte surface and starts a cascade of events that leads to meiosis resumption or germinal vesicle breakdown (GVBD) (Sánchez Toranzo *et al.*, 2006; Zelarayán *et al.*, 2012).

Although lipids have traditionally been considered as storage molecules, their involvement in gonadal function regulation is being increasingly recognized (Sorbera *et al.*, 2001). Seasonal variations in polyunsaturated fatty acids (PUFAs) composition have been linked to reproductive performance because of their effect on pituitary and gonadal hormone levels (Cerdá *et al.*, 1995, 1997; Navas *et al.*, 1997, 1998). Moreover, PUFAs and their metabolites produced through cyclooxygenase (COX) and lipoxygenase (LOX) pathways may have modulatory effects on the gonadal steroid metabolism of mammals and birds (Lin, 1985; Johnson *et al.*, 1991).

¹All correspondence to: Liliana Isabel Zelarayán. Instituto de Biología, Facultad de Bioqca, Qca. y Farmacia, INSIBIO-UNT, Chacabuco 461, 4000. San Miguel de Tucumán, Argentina. Tel: +54 381 4247752 Ext. 7093. Fax: +54 381 4247752 Ext. 7004. E-mail: lzelarayan@fbqf.unt.edu.ar

²Centro de Referencias para Lactobacilos (CERELA)-CONICET, Chacabuco 145, Tucumán, Argentina.

³Instituto de Biología, Facultad de Bioqca, Qca. y Farmacia, INSIBIO-UNT, Chacabuco 461, Tucumán, Argentina.

The role of PUFAs in oocyte maturation was examined in certain species. In starfish, arachidonic acid (AA) and eicosapentaenoic acid were capable of inducing maturation. Moreover, LOX inhibition prevented meiosis resumption. PUFAs appear to be involved in the mechanism by which methyladenine induces oocyte maturation (Meijer *et al.*, 1986). In zebrafish, Lister & Van Der Kraak (2008) showed that follicles expressed enzymes that were involved in AA release, such as phospholipase A₂ (PLA₂) and phospholipase C_{γ1}, or enzymes that can metabolize AA such as COX₁, COX₂ and prostaglandin synthase-2. In agreement with this finding, it was shown in *Rhinella arenarum* that P₄-induced maturation caused significant changes in the amount and composition of phospholipids and neutral lipids in both fully grown ovarian oocytes and plasma membrane-enriched fractions (Caldironi & Alonso, 1996; Mata *et al.*, 2000; Zelarayán *et al.*, 1999). These changes may occur due to lipolytic enzymes activation, especially PLA₂, that release AA. In some mammalian species, AA has been proposed to be a prostaglandins (PGs) source. PGs are produced by COX under gonadotrophin control in granulosa cells. It is well known that PGs are the major paracrine agents in ovarian physiology during ovulation. To our knowledge, there are no previous works that explore the role of AA and its metabolites in oocyte maturation in *Rhinella arenarum*.

In this work we evaluated the role of AA metabolites during *Rhinella arenarum* oocyte maturation using *in vitro* approaches; we focussed on the participation of enzymes involved in AA metabolism – PLA₂, COX and LOX.

Materials and methods

Adult specimens of *Rhinella arenarum* were collected in northwestern Argentina throughout the year and kept at 15°C until use, up to 15 days after collection. Follicles and denuded oocytes were obtained in accordance with Zelarayán *et al.* (2000) and kept in amphibian Ringer solution (AR) until use. *In vitro* cultures were carried out using multiwell culture dishes (Costar 3524, Cambridge, MA, USA). Randomized samples of 20 oocytes or follicles were distributed into separate wells that contained 2 ml of AR. Duplicates were routinely run in each experimental group. Oocyte maturation was assessed by detection of germinal vesicle breakdown (GVBD) 18–20 h after inducer addition. All reagents were purchased from Sigma or Merck.

In order to obtain a dose–response curve of melittin (0.02–0.08 μM), denuded oocytes or follicles were incubated in the presence of melittin for 60 min, then transferred to AR where incubation was completed

and GVBD was assessed. Continuous exposure or high doses of melittin were deleterious for oocytes (lysis signs and irregular pigment distribution).

Inhibition experiments were conducted by pre-incubation of samples for 60 min in the presence of inhibitors and then the inducer was added. The inhibitors used were: quinacrine (Quin): 10–20 μM, indomethacin (Indo): 5–100 μM, rofecoxib: 50–400 μM, lysine clonixinate (LC): 25–100 μM, and nordi-hydroguaiaretic acid (NDGA): 2.5–30 μM. In all cases, P₄ (3 μM) was used as the positive maturation control.

Results are expressed as means ± standard deviation (SD). Comparisons among different treatments were carried out using Student's *t*-test. A value of *P* < 0.05 was considered to be statistically significant.

Results and Discussion

Given that PLA₂-mediated hydrolysis of phospholipids results in AA release, we analysed whether this enzyme was involved in the maturation process. Denuded oocytes and follicles were treated with melittin (0.02–0.08 μM), an enzyme activator. In denuded oocytes, PLA₂ activation induced meiosis resumption in a dose-dependent manner; highest response (78 ± 6% GVBD) was scored with 0.08 μM (Fig. 1). In follicles, PLA₂ activation did not significantly induce meiosis resumption at the assayed doses (12 ± 3% GVBD).

The direct role of PLA₂ activation in GVBD was demonstrated using quinacrine, a specific enzyme inhibitor. As PLA₂ activation had no effect on follicles, they were not treated with quinacrine. Denuded oocytes were pre-incubated for 1 h in the presence of different quinacrine doses (0–20 μM) before induction of maturation with melittin (0.08 μM). PLA₂ inactivation prevented melittin-induced GVBD in a dose-dependent manner (Fig. 2). Highest quinacrine dose (20 μM) significantly reduced maturation to 15 ± 3% GVBD, while the control reached 78 ± 6% GVBD. This result suggests that PLA₂ could participate in mechanisms related to maturation. Enzyme inhibition by quinacrine had a limited effect on P₄-induced maturation.

In order to analyse the involvement of PLA₂ in P₄-induced maturation, denuded oocytes and follicles were pre-incubated in the presence of different quinacrine doses (0–20 μM) for 1 h before induction of maturation with P₄. PLA₂ inhibition did not significantly affect P₄-induced maturation in oocytes or follicles (Fig. 3). This result suggests that other phospholipases could be involved in P₄-induced maturation. One of these could be phospholipase C (PLC), which hydrolyzes membrane phosphatidylinositol 4,5-bisphosphate (PIP₂) into inositol 1,4,5-trisphosphate




Figure 1 Melittin dose–response curve. Denuded oocytes and follicles were incubated in the presence of different melittin doses (0.02–0.08 μM) for 60 min and then samples were transferred to amphibian Ringer solution (AR) in which incubation was completed at 18–20 h. After this procedure, meiosis resumption was assessed, determined as a percentage of germinal vesicle breakdown (GVBD). Progesterone (P_4) 3 μM was used as a control. Values represent the mean \pm standard deviation (SD) of four experiments performed in triplicate on different animals.




Figure 2 Effect of quinacrine on melittin-induced maturation. Denuded oocytes were incubated for 1 h with different quinacrine doses (0–20 μM) and then maturation was induced with melittin (0.08 μM). After incubation for 20 h, germinal vesicle breakdown (GVBD) was assessed. Values represent the mean \pm standard deviation (SD) of six experiments performed in triplicate on different animals.

(IP_3) and diacylglycerol (DAG). It was reported that these messengers would act as potential maturation inducers in *Rhinella arenarum* oocytes treated with P_4




Figure 3 Effect of quinacrine on progesterone (P_4)-induced maturation. Denuded oocytes and complete follicles were pre-incubated with different quinacrine doses (10–20 μM) for 60 min before the addition of P_4 (1 $\mu\text{g}/\text{ml}$). The samples were incubated for 20 h and fixed to assess germinal vesicle breakdown (GVBD). Values represent the mean \pm standard deviation (SD) of three experiments performed in triplicate on different animals.

(Zelarayán *et al.*, 2000). However, results obtained in fish (European sea bass) indicate that PLA_2 blockade with quinacrine significantly inhibited human chorionic gonadotropin (hCG)-induced maturation (Sorbera *et al.*, 2001) in follicles.

PLA_2 has been mainly associated with ovulation rather than maturation. In rodents, expression and activity of PLA_2 in granulosa cells were induced by gonadotropins (Kurusu *et al.*, 1998). In ruminants, an increase in PLA_2 expression after preovulatory gonadotropin surge has been reported, so that this enzyme could be responsible for the AA mobilization necessary for PGs synthesis during ovulation (Diouf *et al.*, 2006).

Arachidonic acid is usually metabolized by an oxidation process in which COX and/or LOX participate. It has been suggested that AA metabolites produced by these enzymes were involved in fish oocyte maturation (Sorbera *et al.*, 2001). The participation of COX, a key enzyme in PGs synthesis, was studied in P_4 -induced maturation. Two COX inhibitors were assayed: indomethacin (5–100 μM), which inhibits both isoforms (COX₁ and COX₂), and rofecoxib (50–400 μM), a selective COX₂ inhibitor. Highest indomethacin dose (100 μM) inhibited P_4 -induced maturation by 50% (Fig. 4). Denuded oocytes and follicles showed a similar response. Maturation was never blocked completely. Indomethacin had a stronger effect than rofecoxib on maturation inhibition in oocytes and follicles. In fact, oocytes reached




Figure 4 Effect of indomethacin or rofecoxib on progesterone (P₄)-induced maturation. Denuded oocytes and follicles were pre-incubated with different doses of each inhibitor (indomethacin: 5–100 μM; rofecoxib: 50–400 μM) for 1 h before the addition of P₄. At the end of incubation (20 h) germinal vesicle breakdown (GVBD) was assessed. Values represent the mean ± standard deviation (SD) of eight experiments performed in triplicate on different animals.

63 ± 3% GVBD with the highest assayed dose (400 μM) (Fig. 4), a finding that suggests that COX would participate in the mechanism by which P₄ induces maturation. Inhibition caused by these nonsteroidal antiinflammatory drugs was partial; other enzymes would be involved in this process (Fig. 4).

Although in *Rhinella arenarum* oocytes P₄-induced maturation takes place throughout the year, oocytes showed a different P₄ response capacity dependent upon the period in which animals had been captured (Zelarayán *et al.*, 2009). Moreover, Medina *et al.* (2004) reported that *Rhinella arenarum* females revealed seasonal variations in the serum levels of steroid hormones associated with reproductive biology.

The participation of the AA cascade in the reproductive process of *Rhinella arenarum* seems to be also correlated with seasonal variations in the ovarian response to P₄. During the February to June period, P₄-induced maturation was low in both oocytes and follicles (56 ± 5% and 40 ± 5% GVBD, respectively) (period of low P₄ response capacity). During this period, LOX inhibition by NDGA (5–30 μM) or LC (25–50 μM) had a stimulatory effect on P₄-induced maturation. Interestingly, oocytes and follicles treated with P₄ plus NDGA 30 μM reached higher maturation values (80 ± 5% and 90 ± 4% GVBD, respectively) than controls (Fig. 5). Samples incubated in the presence of NDGA or LC alone did not show maturation signs.

During the July to January period, when oocytes and follicles showed maximum P₄-induced maturation values, LOX inhibition had no effect on hormone-induced maturation at the assayed doses. Only a slight decrease in GVBD percentage was observed at NDGA 30 μM and LC 100 μM (Fig. 6). During all seasons of the year, higher doses resulted in increased lysis of samples treated with the inhibitors.

Montelukast, an antagonist for leukotriene Cyst-LT₁-receptor, was also evaluated. The addition of this antagonist did not affect hormone-induced maturation and oocytes reached GVBD values similar to the one reached with P₄ (results not shown).

Arachidonic acid can be metabolized through either the COX or LOX pathways. In *Rhinella arenarum* oocytes, it is probable that inhibition of LOX results in metabolism of available AA through the COX pathway only, and thereby leads to an increase in PG synthesis that could be positively correlated with the maturation process. LOX participation in oocyte maturation in fish showed different results; LOX inhibition by NDGA inhibited hCG-induced maturation, a finding that suggested that products of this enzyme could be involved in hormone-induced maturation (Patiño *et al.*, 2003).

In summary, these results suggest that AA metabolites that result from COX or LOX pathways could be involved in P₄-induced *Rhinella arenarum* oocytes maturation.




Figure 5 Effect of nordihydroguaiaretic acid (NDGA) or lysine clonixinate (LC) on progesterone (P_4)-induced maturation during the February to June period. Denuded oocytes and follicles were incubated for 60 min in the presence of different doses of NDGA (5–30 μ M) or LC (25–50 μ M), after which P_4 (3 μ M) was added. The incubations were performed in triplicate at a controlled constant temperature of 26°C. At 20 h of incubation germinal vesicle breakdown (GVBD) was assessed. Values represent the mean \pm standard deviation (SD) of six experiments performed on different animals in triplicate.




Figure 6 Effect of nordihydroguaiaretic acid (NDGA) or lysine clonixinate (LC) on progesterone (P_4)-induced maturation during the July to January period. Maturation was considered as the percentage of germinal vesicle breakdown (GVBD) after 20 h of incubation of denuded oocytes and follicles. They were pre-incubated for 60 min with different doses of NDGA (2.5–30 μ M) or LC (25–100 μ M) before the addition of P_4 . Values represent the mean \pm standard deviation (SD) of five experiments performed on different animals in triplicate.

References

- Caldironi, H.A. & Alonso, T.S. (1996). Lipid characterization of full-grown amphibian oocytes and their plasma membrane-enriched fractions. *Lipids* **31**, 651–6.
- Cerdá, J., Zanuy, S., Carrillo, M., Ramos, J. & Serrano, R. (1995). Short- and long-term dietary effects on female

sea bass (*Dicentrarchus labrax*): seasonal changes in plasma profiles of lipids and sex steroids in relation to reproduction. *Comp. Biochem. Physiol. Part C: Pharmacology, Toxicology and Endocrinology* **111**, 83–91.

- Cerdá, J., Zanuy, S. & Carrillo, M. (1997). Evidence for dietary effects on plasma levels of sexual steroids

- during spermatogenesis in the sea bass. *Aquacult. Int.* **5**, 473–7.
- Diouf, M.N., Sayasith, K., Lefebvre, R., Silversides, D.W., Sirois, J. & Lussier, J.G. (2006). Expression of phospholipase A2 group IVA (PLA₂G4A) is upregulated by human chorionic gonadotropin in bovine granulosa cells of ovulatory follicles. *Biol. Reprod.* **74**, 1096–103.
- Johnson, A.L., Tilly, J.L. & LeVorse, J.M. (1991). Possible role for arachidonic acid in the control of steroidogenesis in hen theca. *Biol. Reprod.* **44**, 338–44.
- Kurusu, S., Iwao, M., Kawaminami, M. & Hashimoto, I. (1998). Involvement of cytosolic phospholipase A2 in the ovulatory process in gonadotropin-primed immature rats. *Prostaglandins Leukot. Essent. Fatty Acids* **58**, 405–11.
- Lin, T. (1985). Mechanism of action of gonadotropin-releasing hormone stimulated Leydig cell steroidogenesis III. The role of arachidonic acid and calcium/phospholipid dependent protein kinase. *Life Sci.* **36**, 1255–64.
- Lister, A.L. & Van Der Kraak, G. (2008). An investigation into the role of prostaglandins in zebrafish oocyte maturation and ovulation. *Gen. Comp. Endocrinol.* **159**, 46–57.
- Mata, A., Zelarayán, L.I., Sánchez Toranzo, G., Alonso, T. & Bühler, M.I. (2000). Neutral lipid characterization of *Bufo arenarum* oocyte capable and not capable of undergoing spontaneous maturation. Abstract of XVI Annual Scientific Meeting, Tucuman Biology Society. *Biocell* **24**, 134.
- Medina, M., Ramos, I., Crespo, C.A., González-Calvar, S. & Fernández, S.N. (2004). Changes in serum sex steroid levels throughout the reproductive cycle of *Bufo arenarum* females. *Gen. Comp. Endocrinol.* **136**, 143–51.
- Meijer, L., Brash, A., Bryant, R., Kwokei, N.G., Maclouff, J. & Sprecher, H. (1986). Stereospecific induction of starfish oocyte maturation by (8R)-hydroxyeicosatetraenoic acid. *J. Biol. Chem.* **261**, 17040–7.
- Navas, J.M., Bruce, M., Thrush, M., Farnedale, B.N., Bromage, N., Zanuy, S., Carrillo, M., Bell, J. & Ramos, J. (1997). The impact of seasonal alteration in the lipid composition of broodstock diets on egg quality in the European sea bass (*Dicentrarchus labrax*). *J. Fish Biol.* **51**, 760–73.
- Navas, J.M., Mañanos, E., Thrush, M., Ramos, J., Zanuy, S., Carrillo, M., Zohar, Y. & Bromage, N. (1998). Effect of dietary lipid composition on vitellogenin, 17 β -estradiol and gonadotropin plasma levels and spawning performance in captive sea bass (*Dicentrarchus labrax*). *Aquaculture* **165**, 65–79.
- Patiño, R., Yoshizaki, G., Bolamba, D. & Thomas, P. (2003). Role of arachidonic acid and protein kinase C during maturation-inducing hormone-dependent meiotic resumption and ovulation in ovarian follicles of Atlantic croaker. *Biol. Reprod.* **68**, 516–23.
- Sánchez Toranzo, G., Bonilla, F., Zelarayán, L., Oterino, J. & Bühler, M.I. (2006). Activation of maturation promoting factor in *Bufo arenarum* oocytes: injection of mature cytoplasm and germinal vesicle contents. *Zygote* **14**, 305–16.
- Sorbera, L.A., Asturiano, J.F., Carrillo, M. & Zanuy, S. (2001). Effects of polyunsaturated fatty acids and prostaglandins on oocyte maturation in a marine teleost, the European Sea Bass (*Dicentrarchus labrax*). *Biol. Reprod.* **64**, 382–9.
- Zelarayán, L.I., Alonso, T., Mata, A., Sánchez Toranzo, G., Oterino, J. & Bühler, M.I. (1999). Evaluation of parameters related with the *Bufo arenarum* oocyte competence to mature spontaneously. Abstract of XV Annual Scientific Meeting, Tucuman Biology Society. *Biocell* **23**, 84.
- Zelarayán, L.I., Sánchez Toranzo, G., Oterino, J. & Bühler, M. (2000). Involvement of purines and phosphoinositides on spontaneous and progesterone induced maturation in *Bufo arenarum* oocytes. *J. Exp. Zool.* **287**, 151–7.
- Zelarayán, L.I., Ortiz, M.E., Unías, L., Ajmat, M.T., Bonilla, F., Sánchez Toranzo, G. & Bühler, M.I. (2009). Prostaglandins effects on *Rhinella arenarum* oocyte maturation. Abstract of XXV Annual Scientific Meeting, Tucuman Biology Association. *Biocell* **33**, 161.
- Zelarayán, L.I., Ajmat, M.T., Bonilla, F. & Bühler, M.I. (2012). Involvement of G protein and purines in *Rhinella arenarum* oocyte maturation. *Zygote* **2**, 1–10.