

RESEARCH ARTICLE

Unitary groups, K -theory, and traces

Pawel Sarkowicz 

Department of Pure Mathematics, University of Waterloo, Waterloo, ON, N2L 3G1, Canada
Email: psarkowi@uwaterloo.ca

Received: 29 May 2023; **Revised:** 13 November 2023; **Accepted:** 20 November 2023;

First published online: 15 December 2023

Keywords: K -theory; traces; unitary groups; C^* -algebras

2020 Mathematics Subject Classification: *Primary* - 46L35, 46L80

Abstract

We show that continuous group homomorphisms between unitary groups of unital C^* -algebras induce maps between spaces of continuous real-valued affine functions on the trace simplices. Under certain K -theoretic regularity conditions, these maps can be seen to commute with the pairing between K_0 and traces. If the homomorphism is contractive and sends the unit circle to the unit circle, the map between spaces of continuous real-valued affine functions can further be shown to be unital and positive (up to a minus sign).

1. Introduction

Unitary groups of C^* -algebras have been long studied, and for many classes of operator algebras they form a complete invariant. In [14], Dye studied the unitary group isomorphism problem between nonatomic W^* -algebras, with the assumption of *weak bicontinuity* of the isomorphism. He later showed that the unitary group, this time as an algebraic object, determined the type of a factor [15] (except for type I_{2n}). He showed that such group isomorphisms were the restrictions of a $*$ -isomorphism or a conjugate linear $*$ -isomorphism multiplied by a possibly discontinuous character ([3, Appendix A] gives exposition). Sakai generalized Dye's results to show that any uniformly continuous unitary group isomorphism between AW^* -factors comes from a $*$ -isomorphism or conjugate-linear $*$ -isomorphism [35] (see also [40] for general AW^* -algebras which have no component of type I_n).

Dye's method was generalized to large classes of real rank zero C^* -algebras by Al-Rawashdeh, Booth and Giordano in [2], where they applied the method to obtain induced maps between K -theory, with a general linear variant being done by Giordano and Sierakowski in [22]. The stably finite and purely infinite cases were handled separately. The unital, simple AH-algebras of slow dimension growth and of real rank zero were classified by the topological group isomorphism class of their unitary groups (or general linear groups), and the unital, simple, purely infinite UCT algebras were classified via the algebraic isomorphism classes of their unitary groups (or general linear groups). These results made use of the abundance of projections in real rank zero C^* -algebras (at least to show there were isomorphic K_0 -groups) and made use of the Dadarlat-Elliott-Gong [10, 20] and Kirchberg-Phillips [31] classification theorems, respectively (see Theorems 3.3.1 and 8.4.1 of [34] for each respective case).

In [28], it was proven by Paterson that two unital C^* -algebras are isomorphic if and only there is an isometric isomorphism of the unitary groups, which acts as the identity on the circle. In a similar vein, the metric structure of the unitary group has also played a role in determining the Jordan $*$ -algebra structure on C^* -algebras. In [24], Hatori and Molnár showed that two unital C^* -algebras are Jordan $*$ -isomorphic if and only if their unitary groups are isometric as metric spaces, not taking into account any algebraic structure.

Chand and Robert have shown in [9] that if A and B are prime traceless C^* -algebras with full square zero elements such that $U^0(A)$, the subgroup of unitaries, which are path connected to the identity, is algebraically isomorphic to $U^0(B)$, then A is either isomorphic or anti-isomorphic to B . In fact, the group

isomorphism is the restriction of a $*$ -isomorphism or anti- $*$ -isomorphism which follows from the fact that unitary groups associated to these C^* -algebras have certain automatic continuity properties that allow one to use characterizations of *commutativity preserving maps* [4] (see [1]). Chand and Robert also show that if A is a unital separable C^* -algebra with at least one tracial state, then $U^0(A)$ admits discontinuous automorphisms. Thus, the existence of traces is an obstruction to classification via algebraic structure on the unitary groups—at least an obstruction to unitary group homomorphisms being the restrictions of $*$ -homomorphisms or anti- $*$ -homomorphisms.

In this paper, we show that uniformly continuous unitary group homomorphisms yield maps between traces which have several desirable K -theoretic properties—especially under stricter continuity assumptions. Namely, that the homomorphism sends the circle to the circle and is contractive, which would be automatic if it had a lift to a $*$ -homomorphism or conjugate-linear $*$ -homomorphism.

We state our main results. Recall that if A is a unital C^* -algebra, $T(A)$ denotes the simplex of tracial states, and $\text{Aff } T(A)$ is the real Banach space of continuous affine functions $T(A) \rightarrow \mathbb{R}$.

Theorem A (Corollary 3.6). *Let A, B be unital C^* -algebras. If $\theta : U^0(A) \rightarrow U^0(B)$ is a uniformly continuous group homomorphism, then there exists a bounded \mathbb{R} -linear map $\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ such that*

$$\begin{CD}
 \pi_1(U^0(A)) @>\tilde{\Delta}_A^1>> \text{Aff } T(A) \\
 @V\pi_1(\theta)VV @VV\Lambda_\theta V \\
 \pi_1(U^0(B)) @>\tilde{\Delta}_B^1>> \text{Aff } T(B)
 \end{CD} \tag{1.1}$$

commutes.

Here $\pi_1(\theta)$ is the map between fundamental groups induced by θ , and, for a C^* -algebra A , $\tilde{\Delta}_A^1$ is the *pre-determinant* (used in the definition of the de la Harpe-Skandalis determinant associated to the universal trace) that takes a piece-wise smooth path in $U^0(A)$ beginning at the unit to an element of $\text{Aff } T(A)$. See Section 2.2 for details.

Recall that the K_0 -group of a unital C^* -algebra can be identified with the fundamental group $\pi_1(U_\infty^0(A))$. Restricting to C^* -algebras with sufficient K_0 -regularity—by this we mean C^* -algebras whose K_0 -group can be realized as loops in the connected component of its unitary group—we get a map between K_0 -groups and a map between spaces of continuous real-valued affine functions on the trace simplex which commute with the pairing.

Corollary B (Corollary 3.6). *Let A, B be unital C^* -algebras such that the canonical maps*

$$\pi_1(U^0(A)) \rightarrow K_0(A) \text{ and } \pi_1(U^0(B)) \rightarrow K_0(B) \tag{1.2}$$

are isomorphisms. If $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous group homomorphism, then there exists a bounded linear map $\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ such that

$$\begin{CD}
 K_0(A) @>\rho_A>> \text{Aff } T(A) \\
 @V K_0(\theta) VV @VV \Lambda_\theta V \\
 K_0(B) @>\rho_B>> \text{Aff } T(B)
 \end{CD} \tag{1.3}$$

commutes, where $K_0(\theta)$ is the map induced by $\pi_1(\theta)$ together with the isomorphisms of (1.2).

C^* -algebras satisfying the above hypothesis are quite common—for example, C^* -algebras having stable rank one [32] or that are \mathcal{Z} -stable [25] have this property. Viewing $\text{Aff } T(A)$ and $\text{Aff } T(B)$ as partially

ordered real Banach spaces (under the uniform norm) with order units, it is not, however, true that Λ_θ is unital or positive (see Example 3.4). This is remedied by adding stricter continuity assumptions on the homomorphism θ (and possibly by replacing Λ_θ with $-\Lambda_\theta$).

When $\theta : U(A) \rightarrow U(B)$ is contractive, injective, and sends the circle to the circle, then we show (Lemma 4.3) that either Λ_θ or $-\Lambda_\theta$ is unital and positive, and therefore, θ induces a map between K -theory and traces in such a manner that respects the pairing (which in turn gives a map between Elliott invariants for certain simple C^* -algebras). As a consequence, we can identify certain unitary subgroups with C^* -subalgebras by using K -theoretic classification of embeddings [6].

Theorem C (Corollary 4.13). *Let A be a unital, separable, simple, nuclear C^* -algebra satisfying the UCT which is either \mathcal{Z} -stable or has stable rank one and B be a unital, separable, simple, nuclear \mathcal{Z} -stable C^* -algebra. If there is a contractive injective group homomorphism $U(A) \rightarrow U(B)$ that maps the circle to the circle, then there is a unital embedding $A \hookrightarrow B$.*

This paper is structured as follows. In Section 3, we use a continuous unitary group homomorphism to construct a map between spaces of continuous affine functions on the trace simplices and use the de la Harpe–Skandalis determinant to show that this map has several desirable properties with respect to the map induced on the fundamental groups of the unitary groups. In Section 4, we discuss how our map between spaces of affine functions respects or flips the order under certain continuity assumptions on the unitary group homomorphism. In Section 5, we discuss a slight general linear variant. We finish in Section 6 with some open questions.

2. Preliminaries and notation

2.1. Notation

For a group G , we denote by DG the derived subgroup of G , that is

$$DG := \langle ghg^{-1}h^{-1} \mid g, h \in G \rangle. \tag{2.1}$$

If G has an underlying topology, we denote by CG the closure of DG and G^0 the connected component of the identity.

For a unital C^* -algebra A , $U(A)$ denotes the unitary group of A , while $U^0(A)$ denotes the connected component of the identity in $U(A)$. For $n \in \mathbb{N}$, we write $U_n(A) := U(M_n(A))$, $U_n^0(A) := U^0(M_n(A))$, and we set

$$U_\infty(A) := \varinjlim U_n(A), \tag{2.2}$$

to be the inductive limit with connecting maps $U_n(A) \ni u \mapsto u \oplus 1 \in U_{n+1}(A)$. This makes $U_\infty(A)$ both a topological space (with the inductive limit topology) and a group.¹ We have general linear analogs by replacing U with GL , where $GL(A)$ denotes the group of invertible elements of A . Similarly, we define $M_\infty(A) = \varinjlim M_n(A)$ (as an algebraic direct limit) with connecting maps $x \mapsto x \oplus 0$. If E is real Banach space and $\tau : A_{sa} \rightarrow E$ is a linear map that is tracial (i.e., $\tau(a^*a) = \tau(aa^*)$ for all $a \in A$), we extend this canonically to $(M_\infty(A))_{sa}$ by setting $\tau((a_{ij})) := \sum_i \tau(a_{ii})$ for $(a_{ij}) \in (M_n(A))_{sa}$.

We write $\pi_0(X)$ for the space of connected components of a topological space X , and $\pi_1(X)$ for the fundamental group of X with distinguished base point. In our case, we will usually have $X = U_n(A)$ or $X = U_n^0(A)$, for $n \in \mathbb{N} \cup \{\infty\}$, with the base point being the unit.

For a unital C^* -algebra A , we let $K_0(A), K_1(A)$ be the topological K -groups of A . We will often use the identification of $K_0(A)$ with the fundamental group $\pi_1(U_\infty^0(A))$ (see e.g., [33, Chapter 11.4]). The set of tracial states on A will be denoted $T(A)$, which is a Choquet simplex [36, Theorem 3.1.18], and

¹As pointed out in footnote 58 of [6], $U_\infty(A)$ is not in general a topological group since multiplication is not in general jointly continuous in this topology.

we denote by $\text{Aff } T(A)$ the set of continuous affine functions $T(A) \rightarrow \mathbb{R}$, which is an interpolation group with order unit (see [21, Chapter 11]). For unital A , the pairing map $\rho_A : K_0(A) \rightarrow \text{Aff } T(A)$ is defined as follows: if $x \in K_0(A)$, we can write $x = [p] - [q]$ where $p, q \in M_n(A)$ are projections, and then

$$\rho_A(x)(\tau) := \tau(p - q), \quad \tau \in T(A). \tag{2.3}$$

2.2. The de la Harpe–Skandalis determinant and Thomsen’s variant

We recall the definition of the unitary variant of the de la Harpe–Skandalis determinant [12] (see [11] for a more in-depth exposition). By a bounded trace, we mean a bounded linear map $\tau : A_{sa} \rightarrow E$, where E is a real Banach space, such that $\tau(a^*a) = \tau(aa^*)$ for all $a \in A$. For $n \in \mathbb{N} \cup \{\infty\}$, a bounded trace $\tau : A_{sa} \rightarrow E$, and a piece-wise smooth path $\xi : [0, 1] \rightarrow U_n(A)$, set

$$\tilde{\Delta}_\tau^n(\xi) := \int_0^1 \tau \left(\frac{1}{2\pi i} \xi'(t)\xi(t)^{-1} \right) dt, \tag{2.4}$$

where this integral is just the Riemann integral in E .² We state the unitary variant of [12, Lemme 1].

Proposition 2.1. *Let $\tau : A_{sa} \rightarrow E$ be a bounded trace and $n \in \mathbb{N} \cup \{\infty\}$. The map $\tilde{\Delta}_\tau^n$, which takes a piece-wise smooth path in $U_n^0(A)$ to an element in E , has the following four properties:*

- 1. *it takes pointwise products to sums: if ξ_1, ξ_2 are two piece-wise smooth paths, then*

$$\tilde{\Delta}_\tau^n(\xi_1 \xi_2) = \tilde{\Delta}_\tau^n(\xi_1) + \tilde{\Delta}_\tau^n(\xi_2), \tag{2.5}$$

where $\xi_1 \xi_2$ is the piece-wise smooth path $t \mapsto \xi_1(t)\xi_2(t)$ from $\xi_1(0)\xi_2(0)$ to $\xi_1(1)\xi_2(1)$;

- 2. *if $\|\xi(t) - 1\| < 1$ for all $t \in [0, 1]$, then*

$$2\pi i \tilde{\Delta}_\tau^n(\xi) = \tau(\log \xi(1) - \log \xi(0)); \tag{2.6}$$

- 3. *$\tilde{\Delta}_\tau^n(\xi)$ depends only on the continuous homotopy class of ξ ;*
- 4. *if $p \in M_n(A)$ is a projection, then the path $\xi_p : [0, 1] \rightarrow U_n^0(A)$ given by $\xi_p(t) := pe^{2\pi it} + (1 - p)$ satisfies $\tilde{\Delta}_\tau^n(p) = \tau(p)$.*

The de la Harpe–Skandalis determinant associated to τ (at the n^{th} level) is then the map

$$\Delta_\tau^n : U_\infty^0(A) \rightarrow E / \tilde{\Delta}_\tau^n(\pi_1(U_n^0(A))) \tag{2.7}$$

given by $\Delta_\tau^n(x) := \left[\tilde{\Delta}_\tau^n(\xi_x) \right]$ where ξ_x is any piece-wise smooth path $\xi_x : [0, 1] \rightarrow U_n^0(A)$ from 1 to x . This is a well-defined group homomorphism (using Proposition 2.1) to an abelian group and therefore factors through the derived group, i.e., $DU_n^0(A) \subseteq \ker \Delta_\tau^n$. For the $n = \infty$ case, we just write $\tilde{\Delta}_\tau$ and Δ_τ for $\tilde{\Delta}_\tau^\infty$ and Δ_τ^∞ , respectively.

We will often be interested in the *universal trace* $\text{Tr}_A : A_{sa} \rightarrow \text{Aff } T(A)$, which is given by $\text{Tr}_A(a) := \hat{a}$, where $\hat{a} \in \text{Aff } T(A)$ is the function given by $\hat{a}(\tau) := \tau(a)$ for $\tau \in T(A)$. We note that in this case, for $[x] \in K_0(A)$, we have that $\text{Tr}(x) = \rho_A([x])$. When considering the universal trace, we will write Δ^n and Δ for Δ_{Tr}^n and $\Delta_{\text{Tr}}^\infty$, respectively. If the C^* -algebra needs to be specified, we write Δ_A^n or Δ_A .

Proposition 2.2. *Let $n \in \mathbb{N} \cup \{\infty\}$. Every continuous path $\xi : [0, 1] \rightarrow U_n(A)$ is homotopic to a piece-wise smooth path (even a piece-wise smooth exponential path if we are in $U_n^0(A)$). Moreover, there exists $a \in A_{sa}$ such that $\tilde{\Delta}_\omega^n(\xi) = \omega(a)$ whenever $\omega : A_{sa} \rightarrow E$ is a bounded trace.*

In particular, as $\tilde{\Delta}^n$ is homotopy invariant, it makes sense to apply $\tilde{\Delta}^n$ to any continuous path.

²Note that $\xi'(t)\xi(t)$ is skew-adjoint by [30, Proposition 1.4] so that $\frac{1}{2\pi i}\xi'(t)\xi(t)^{-1}$ is self-adjoint.

Proof. This is essentially [12, Lemme 3]. Take a continuous path $\xi : [0, 1] \rightarrow U_n(A)$ and choose k such that

$$\left\| \xi \left(\frac{j-1}{k} \right)^{-1} \xi \left(\frac{j}{k} \right) - 1 \right\| < 1 \text{ for all } j = 1, \dots, k. \tag{2.8}$$

Then taking

$$a_j := \frac{1}{2\pi i} \log \left(\xi \left(\frac{j-1}{k} \right)^{-1} \xi \left(\frac{j}{k} \right) \right), j = 1, \dots, k, \tag{2.9}$$

ξ will be homotopic to the path

$$\eta(t) = \xi \left(\frac{j-1}{k} \right) e^{2\pi i(kt-j+1)a_j}, t \in \left[\frac{j-1}{k}, \frac{j}{k} \right], j = 1, \dots, k. \tag{2.10}$$

We note that $\tilde{\Delta}_\tau^n(\eta) = \sum_{j=1}^k \tau(a_j)$. Indeed, for simplicity denote by

$$X_j := \xi \left(\frac{j-1}{k} \right) \text{ and } Y_j := e^{2\pi i(kt-j+1)a_j}. \tag{2.11}$$

Then

$$\begin{aligned} \tilde{\Delta}_\omega^n(\xi) &= \tilde{\Delta}_\omega^n(\eta) \\ &= \sum_{j=1}^k \int_{\frac{j-1}{k}}^{\frac{j}{k}} \tau \left(\frac{1}{2\pi i} X_j 2\pi i k a_j Y_j Y_j^* X_j^* \right) dt \\ &= \sum_{j=1}^k \int_{\frac{j-1}{k}}^{\frac{j}{k}} k \tau(a_j) dt \\ &= \sum_{j=1}^k \tau(a_j). \end{aligned} \tag{2.12}$$

If we take $a := \text{tr}(a')$, where $a' = \sum_{j=1}^k a_j$ (here $\text{tr} : M_\infty(A) \rightarrow A$ is the unnormalized trace), we see that $\tilde{\Delta}_\omega^n(\xi) = \omega(a)$. □

Let A_0 consist of elements $a \in A_{sa}$ satisfying $\tau(a) = 0$ for all $\tau \in T(A)$. This is a norm-closed real subspace of A_{sa} such that $A_0 \subseteq \overline{[A, A]}$, and there is an isometric identification $A_{sa}/A_0 \simeq \text{Aff } T(A)$ sending an element $[a]$ to \hat{a} . Indeed, it is not hard to see that the map $A_{sa}/A_0 \rightarrow \text{Aff } T(A)$ given by $[a] \mapsto \hat{a}$ is a well-defined \mathbb{R} -linear map. Moreover [8, Theorem 2.9], together with a convexity argument, gives that this is isometric identification. To see that we have all the real-valued affine functions, we note that the image of this map contains constant functions and separates points, so [21, Corollary 7.4] gives that the image is dense and therefore all of $\text{Aff } T(A)$ (since this is an isometry). We freely identify A_{sa}/A_0 with $\text{Aff } T(A)$.

2.3. Thomsen’s variant

Thomsen’s variant of the de la Harpe–Skandalis determinant is the Hausdorffized version, taking into account the closure of the image of the homotopy groups. For a bounded trace $\tau : A_{sa} \rightarrow E$, we consider the map

$$\bar{\Delta}_\tau^n : U_n^0(A) \rightarrow E / \overline{\tilde{\Delta}_\tau^n(\pi_1(U_n^0(A)))}, \tag{2.13}$$

given by $\bar{\Delta}_\tau^n(x) := \left[\tilde{\Delta}_\tau^n(\xi_x) \right]$ where $\xi_x : [0, 1] \rightarrow U_n^0(A)$ is any piece-wise smooth path from 1 to $x \in U_n^0(A)$. This is similar to the map Δ_τ^n , except the codomain is now the quotient by the closure of the

image of the fundamental group under the predeterminant (i.e., the Hausdorffization of the codomain). When considering the universal trace, we just write $\overline{\Delta}^n$ for $\overline{\Delta}_{\text{Tr}}^n$ and $\overline{\Delta}$ for $\overline{\Delta}_{\text{Tr}}$. If the C^* -algebra needs to be specified, we write $\overline{\Delta}_A^n$ or $\overline{\Delta}_A$.

If one considers the universal trace, the kernel of $\overline{\Delta}^n$ can be identified with $CU_n^0(A)$ (where the closure is taken with respect to the inductive limit topology in the $n = \infty$ case).

Lemma 2.3 (Lemma 3.1, [39]). *Let A be a unital C^* -algebra. Then*

$$\ker \overline{\Delta}^n = CU_n^0(A). \tag{2.14}$$

It is not in general true that the kernel of Δ^n can be identified with the derived group $DU_n^0(A)$, although there are several positive results [13, 26, 27, 38].

It immediately follows that the quotient of $U_n^0(A)$ by the closure of the commutator subgroup (under the inductive limit topology in the $n = \infty$ case) can be identified with a quotient of $\text{Aff } T(A)$.

Theorem 2.4 (Theorem 3.2, [39]). *$\overline{\Delta}^n$ gives a homeomorphic group isomorphism*

$$U_n^0(A)/CU_n^0(A) \simeq \text{Aff } T(A)/\overline{\Delta^n(\pi_1(U_n^0(A)))} \tag{2.15}$$

for every $n \in \mathbb{N} \cup \{\infty\}$. In particular,

$$U_\infty^0(A)/CU_\infty^0(A) \simeq \text{Aff } T(A)/\overline{\rho_A(K_0(A))}. \tag{2.16}$$

2.4. The KT_u invariant

Following [6], for a unital C^* -algebra, we let

$$KT_u(A) := (K_0(A), [1_A]_0, K_1(A), \rho_A, \text{Aff } T(A)) \tag{2.17}$$

be the invariant consisting of the K_0 group, the position of the unit in K_0 , the K_1 group, the pairing between K_0 and traces, and the continuous real-valued affine functions on the trace simplex (viewed as a partially ordered Banach space with order unit). For two unital C^* -algebras A, B , a KT_u morphism

$$(\alpha_0, \alpha_1, \gamma) : KT_u(A) \rightarrow KT_u(B), \tag{2.18}$$

will be a triple $(\alpha_0, \alpha_1, \gamma)$ consisting of $\alpha_0 : K_0(A) \rightarrow K_0(B)$ a group homomorphism such that $\alpha_0([1_A]_0) = [1_B]_0$, $\alpha_1 : K_1(A) \rightarrow K_1(B)$ a group homomorphism, and $\gamma : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ is a unital positive map such that

$$\begin{array}{ccc} K_0(A) & \xrightarrow{\rho_A} & \text{Aff } T(A) \\ \alpha_0 \downarrow & & \downarrow \gamma \\ K_0(B) & \xrightarrow{\rho_B} & \text{Aff } T(B) \end{array} \tag{2.19}$$

commutes.

We note that for large classes of unital simple C^* -algebras—for example, the class of unital, simple, separable, nuclear \mathcal{Z} -stable C^* -algebras satisfying the UCT— $KT_u(\cdot)$ recovers the Elliott invariant.

3. Continuous unitary group homomorphisms and traces

Throughout, A and B will be unital C^* -algebras with nonempty trace simplices, and $\theta : U^0(A) \rightarrow U^0(B)$ will denote a uniformly continuous group homomorphism between the connected components of the identities in the respective unitary groups. We will specify any additional assumptions as we go along. As

θ is a continuous group homomorphism, it send commutators to commutators and limits of commutators to limits of commutators. Thus, there are induced group homomorphisms

$$\begin{aligned} U^0(A)/CU^0(A) &\rightarrow U^0(B)/CU^0(B) \text{ and} \\ U^0(A)/DU^0(A) &\rightarrow U^0(B)/DU^0(B). \end{aligned} \tag{3.1}$$

Thomsen’s isomorphism [39, Theorem 3] then brings about maps between quotients of $\text{Aff } T(A)$ and $\text{Aff } T(B)$:

$$\begin{array}{ccc} U^0(A)/CU^0(A) & \xrightarrow{\simeq} & \text{Aff } T(A)/\overline{\tilde{\Delta}_A^1(\pi_1(U^0(A)))} \\ \downarrow & & \downarrow \\ U^0(B)/CU^0(B) & \xrightarrow{\simeq} & \text{Aff } T(B)/\overline{\tilde{\Delta}_B^1(\pi_1(U^0(B)))}. \end{array} \tag{3.2}$$

In a similar vein, when $DU^0(A) = \ker \Delta_A^1$ and $DU^0(B) = \ker \Delta_B^1$, there is a purely algebraic variant of the above diagram:

$$\begin{array}{ccc} U^0(A)/DU^0(A) & \xrightarrow{\simeq} & \text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) \\ \downarrow & & \downarrow \\ U^0(B)/DU^0(B) & \xrightarrow{\simeq} & \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))). \end{array} \tag{3.3}$$

In fact, there is always a diagram as above with $DU^0(A)$ and $DU^0(B)$ replaced with $\ker \Delta_A^1$ and $\ker \Delta_B^1$, respectively, by Proposition 3.5(3). That is, whether or not the kernel of the determinant agrees with the commutator subgroup of the connected component of the identity, we have:

$$\begin{array}{ccc} U^0(A)/\ker \Delta_A^1 & \xrightarrow{\simeq} & \text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) \\ \downarrow & & \downarrow \\ U^0(B)/\ker \Delta_B^1 & \xrightarrow{\simeq} & \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))). \end{array} \tag{3.4}$$

The diagram in (3.3) is just a special case of (3.4).

In the setting where $\pi_1(U^0(A)) \rightarrow K_0(A)$ and $\pi_1(U^0(B)) \rightarrow K_0(B)$ are surjections, we have induced maps between quotients

$$\begin{aligned} \text{Aff } T(A)/\overline{\rho_A(K_0(A))} &\rightarrow \text{Aff } T(B)/\overline{\rho_B(K_0(B))}, \\ \text{Aff } T(A)/\rho_A(K_0(A)) &\rightarrow \text{Aff } T(B)/\rho_B(K_0(B)) \end{aligned} \tag{3.5}$$

in the respective Hausdorffized and non-Hausdorffized settings.

One question to be answered is whether or not we can lift the maps on the right of (3.2) and (3.4) to maps $\text{Aff } T(A) \rightarrow \text{Aff } T(B)$. These spaces have further structure as dimension groups with order units [21, Chapter 7], so we would like to be able to alter the lift to get a map which is unital and positive. We show that we can always lift this map, and altering it to be unital and positive is possible under a certain continuity assumption on θ .

If we further assume that $K_0(A) \simeq \pi_1(U^0(A))$ and $K_0(B) \simeq \pi_1(U^0(B))$ in the canonical way (which is true in the presence of \mathcal{Z} -stability by [25] or stable rank one [32]), we would like this map to be compatible with the group homomorphism

$$K_0(\theta) : K_0(A) \rightarrow K_0(B) \tag{3.6}$$

arising from the diagram

$$\begin{CD}
 K_0(A) @>\simeq>> \pi_1(U^0(A)) \\
 @V K_0(\theta) VV @VV \pi_1(\theta) V \\
 K_0(B) @>\simeq>> \pi_1(U^0(B)).
 \end{CD} \tag{3.7}$$

By compatible, we mean that

$$\begin{CD}
 K_0(A) @>\rho_A>> \text{Aff } T(A) \\
 @V \pi_1(\theta) VV @VV V \\
 K_0(B) @>\rho_B>> \text{Aff } T(B)
 \end{CD} \tag{3.8}$$

commutes, where the map on the right is the lift coming from maps induced by (3.2) and (3.4). If our map between spaces of affine continuous functions is not unital and positive, but we can alter it accordingly, we must do the same to our map between K_0 . We would still have a commuting diagram as above, but it would give that maps induced on $K_0(\cdot)$ and $\text{Aff } T(\cdot)$ respect the pairing.

Stone’s theorem [7, Section X.5] allows one to recover from a strongly continuous one parameter family $U(t)$ of unitaries a (possibly unbounded) self-adjoint operator X such that $U(t) = e^{itX}$ for all $t \in \mathbb{R}$. If it is a norm-continuous one parameter family of unitaries, one can recover a bounded self-adjoint operator X , and X will lie in the C^* -algebra generated by the unitaries. The use of Stone’s theorem to deduce that continuous group homomorphisms between unitary groups send exponentials to exponentials is not new. Sakai used it in the 1950s in order to show that a norm-continuous group isomorphism between unitary groups of AW^* -algebras are induced by a $*$ -isomorphism or conjugate-linear $*$ -isomorphism between the algebras themselves [35]. More recently, this sort of idea has been used to understand how the metric structure of the unitary groups can be related to the Jordan $*$ -algebra structure of the algebras [24].

Lemma 3.1. *Let A, B be unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ be a continuous group homomorphism. Suppose that $a \in A_{sa}$ and represent $B \subseteq \mathcal{B}(\mathcal{H})$ faithfully. Then $(\theta(e^{2\pi ita}))_{t \in \mathbb{R}}$ is a one-parameter norm-continuous family of unitaries, and consequently is of the form $(e^{2\pi itb})_{t \in \mathbb{R}}$ for a unique $b \in B_{sa}$.*

Proof. Using the fact that θ is a norm-continuous homomorphism, $t \mapsto \theta(e^{2\pi ita})$ is a norm-continuous one-parameter family of unitaries. Stone’s theorem gives that there is a unique self-adjoint $b \in \mathcal{B}(\mathcal{H})$ such that $\theta(e^{2\pi ita}) = e^{2\pi itb}$ for all $t \in \mathbb{R}$. The boundedness of b follows from norm-continuity, and the uniqueness follows from the fact that $b = \frac{1}{2\pi it_0} \log \theta(e^{2\pi it_0 a})$ for sufficiently small $t_0 > 0$. This also gives that $b \in B$ by functional calculus. □

Let $S_\theta : A_{sa} \rightarrow B_{sa}$ be defined via the correspondence given above:

$$\theta(e^{2\pi ita}) = e^{2\pi itS_\theta(a)} \text{ for all } t \in \mathbb{R}. \tag{3.9}$$

Then S_θ is a bounded \mathbb{R} -linear map (see [24, 35], or note that it is easy to see that its kernel is closed). It is also easily checked to respect commutation, and that its canonical extension to a map from A to B actually sends commutators to commutators, although we will not explicitly use this. Recall that for a C^* -algebra A , A_0 denotes the set of self-adjoint elements that vanish on every tracial state.

Lemma 3.2. *If $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous group homomorphism, then S_θ is a bounded linear map and the following hold.*

1. *If θ is injective, then S_θ is injective.*
2. *If θ is a homeomorphism, then S_θ is bijective.*

Proof. As already remarked, S_θ is bounded. Assuming that θ is injective, suppose that $S_\theta(a) = S_\theta(b)$. Then

$$\theta(e^{2\pi ita}) = \theta(e^{2\pi itb}) \tag{3.10}$$

for all $t \in \mathbb{R}$. Injectivity of θ gives that $e^{2\pi ita} = e^{2\pi itb}$ for all $t \in \mathbb{R}$. But this implies that $a = b$ since we can take t appropriately close to 0 and take logarithms.

Now if we further assume that θ is a homeomorphism, then $(\theta^{-1}(e^{2\pi itb}))_{t \in \mathbb{R}} \subseteq U^0(A)$ is a norm-continuous one-parameter family of unitaries which we can write as $(e^{2\pi ita})_{t \in \mathbb{R}}$ for some $a \in A_{sa}$. But then

$$\theta(e^{2\pi ita}) = \theta \circ \theta^{-1}(e^{\pi itb}) = e^{2\pi itb}. \tag{3.11}$$

Thus, $S_\theta(a) = b$ by the uniqueness in Lemma 3.1. □

Recall that a linear map $\tau : A_{sa} \rightarrow E$, where E is a real Banach space, is a bounded trace if it is a bounded \mathbb{R} -linear map such that $\tau(a^*a) = \tau(aa^*)$ for all $a \in A$. Note that this is equivalent to $\tau \circ \text{Ad}_u = \tau$ for all $u \in U(A)$ – to see this, follow the steps in [33, Exercise 3.6] together with a complexification argument. This is further equivalent to $\tau \circ \text{Ad}_u = \tau$ for all $u \in U^0(A)$ (as one can write every element in a C^* -algebra as the linear combination of four unitaries, each of which can be made to not have full spectrum).

Proposition 3.3. *Let $\theta : U^0(A) \rightarrow U^0(B)$ be a continuous group homomorphism, E be a real Banach space and $\tau : B_{sa} \rightarrow E$ a bounded trace. Then $\tau \circ S_\theta : A_{sa} \rightarrow E$ is a bounded trace. In particular, $S_\theta(A_0) \subseteq B_0$ and S_θ induce a bounded \mathbb{R} -linear map*

$$\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B). \tag{3.12}$$

Proof. Observe that, for $a \in A_{sa}$ and $u \in U^0(A)$, we have

$$\begin{aligned} e^{2\pi itS_\theta(uau^*)} &= \theta(e^{2\pi ituau^*}) \\ &= \theta(ue^{2\pi ita}u^*) \\ &= \theta(u)e^{2\pi itS_\theta(a)}\theta(u)^* \\ &= e^{2\pi it\theta(u)S_\theta(a)\theta(u)^*} \end{aligned} \tag{3.13}$$

for all $t \in \mathbb{R}$. Therefore, $S_\theta(uau^*) = \theta(u)S_\theta(a)\theta(u)^*$, and applying τ yields

$$\begin{aligned} \tau \circ S_\theta(uau^*) &= \tau(\theta(u)S_\theta(a)\theta(u)^*) \\ &= \tau(S_\theta(a)), \end{aligned} \tag{3.14}$$

i.e., $\tau \circ S_\theta$ is tracial.

Thus, if $a \in A_0$, it vanishes on every tracial state (hence on every tracial functional), and so $\tau \circ S_\theta(a) = 0$ for all $\tau \in T(B)$. Therefore, $S_\theta(A_0) \subseteq B_0$ and so S_θ factors through a map

$$\Lambda_\theta : \text{Aff } T(A) \simeq A_{sa}/A_0 \rightarrow B_{sa}/B_0 \simeq \text{Aff } T(B), \tag{3.15}$$

where we identify $A_{sa}/A_0 \simeq \text{Aff } T(A)$ and $B_{sa}/B_0 \simeq \text{Aff } T(B)$. □

One cannot expect Λ_θ (or S_θ) to be unital or positive, as the following examples show.

Example 3.4.

1. Consider a continuous homomorphism $\theta : \mathbb{T} \rightarrow \mathbb{T} = U^0(\mathbb{C}) = U(\mathbb{C})$. By Pontryagin duality, $\theta(z) = z^n$ for some $n \in \mathbb{Z}$. We then have that $S_\theta : \mathbb{R} \rightarrow \mathbb{R}$ is given by $S_\theta(x) = nx$. If $n \neq 1$, clearly S_θ is not unital. If $n < 0$, then S_θ is not positive since it sends 1 to $n < 0$. An important observation, however, is that if $n < 0$, $-S_\theta : \mathbb{R} \rightarrow \mathbb{R}$ is positive, and $-\frac{1}{n}S$ is unital and positive.

2. Consider $\theta : \mathbb{T}^3 \rightarrow \mathbb{T}$ given by $\theta(z, w, v) = \bar{z}wv$. The corresponding map $S_\theta : \mathbb{R}^3 \rightarrow \mathbb{R}$ is given by

$$S_\theta(a, b, c) = -a + b + c. \tag{3.16}$$

Clearly $(1, 0, 0) \in \mathbb{C}^3$ is a positive element; however, $S_\theta(1, 0, 0) = -1 < 0$. This map is, however, unital.

3. Let $\theta : U_2 \rightarrow \mathbb{T}$ be defined by $\theta(u) = \det(u)$. Then $S_\theta : (M_2)_{sa} \rightarrow \mathbb{R}$ is defined by $S_\theta(A) = \text{tr}A$, where tr is the unnormalized trace. Clearly this map is not unital, but it is positive.
4. Let $\theta : U_2 \rightarrow U_3$ be defined by $\theta(u) = u \oplus 1$. Then $S_\theta : (M_2)_{sa} \rightarrow (M_3)_{sa}$ is given by $S_\theta(A) = A \oplus 0$, which is again not unital, but is positive. The induced map $\Lambda_\theta : \mathbb{R} \rightarrow \mathbb{R}$ is given by $\Lambda_\theta(x) = \frac{2}{3}x$ for $x \in \mathbb{R}$.
5. Let $\theta : \mathbb{T} \hookrightarrow U_2$ be defined by

$$\theta(\lambda) = \begin{pmatrix} \lambda & \\ & \lambda \end{pmatrix}. \tag{3.17}$$

Then S_θ is a unital, positive isometry and Λ_θ gives rise the identity map

$$\mathbb{R} = \text{Aff } T(\mathbb{C}) \rightarrow \text{Aff } T(M_2) = \mathbb{R}. \tag{3.18}$$

6. Let $\theta : \mathbb{T} \hookrightarrow U_2$ be defined by

$$\theta(\lambda) = \begin{pmatrix} \lambda & \\ & \bar{\lambda} \end{pmatrix}. \tag{3.19}$$

Then $S_\theta : \mathbb{R} \rightarrow (M_2)_{sa}$ is defined by

$$S_\theta(x) = \begin{pmatrix} x & \\ & -x \end{pmatrix} \tag{3.20}$$

and Λ_θ is identically zero.

The above examples are important. If $\theta(\mathbb{T}) \subseteq \mathbb{T}$, which is a moderate assumption (e.g., if θ was the restriction of a unital *-homomorphism or an anti-*-homomorphism), we can restrict the homomorphism to the circle to get a continuous group homomorphism $\mathbb{T} \rightarrow \mathbb{T}$. We understand such homomorphisms by Pontryagin duality [17, Chapter 4].

We now use (pre-)determinant techniques in order to show desirable relationships between our maps.

Proposition 3.5. *Let A, B be unital C^* -algebras, $\theta : U^0(A) \rightarrow U^0(B)$ be a uniformly continuous homomorphism, E a real Banach space, and $\tau : B_{sa} \rightarrow E$ a bounded trace.*

1. Let $\xi : [0, 1] \rightarrow U^0(A)$ be a piece-wise smooth path with $\xi(0) = 1$. Then

$$\tilde{\Delta}_{\tau \circ S_\theta}^1(\xi) = \tilde{\Delta}_\tau^1(\theta \circ \xi). \tag{3.21}$$

In particular, $\tilde{\Delta}_{\tau \circ S_\theta}^1(\pi_1(U^0(A))) \subseteq \tilde{\Delta}_\tau^1(\pi_1(U^0(B)))$.

2. The following diagram commutes:

$$\begin{array}{ccc} \pi_1(U^0(A)) & \xrightarrow{\tilde{\Delta}_{\tau \circ S_\theta}^1} & E \\ \pi_1(\theta) \downarrow & & \downarrow id \\ \pi_1(U^0(B)) & \xrightarrow{\tilde{\Delta}_\tau^1} & E. \end{array} \tag{3.22}$$

3. The following diagram commutes:

$$\begin{CD}
 U^0(A) @>\Delta_{\tau \circ S_\theta}^1>> E/\tilde{\Delta}_{\tau \circ S_\theta}^1(\pi_1(U^0(A))) \\
 @V\theta VV @VVV \\
 U^0(B) @>\Delta_\tau^1>> E/\tilde{\Delta}_\tau^1(\pi_1(U^0(B))),
 \end{CD} \tag{3.23}$$

where the map on the right is the canonical map induced from the inclusion $\tilde{\Delta}_{\tau \circ S_\theta}^1(\pi_1(U^0(A))) \subseteq \tilde{\Delta}_\tau^1(\pi_1(U^0(B)))$ coming from (1). In particular, $\theta(\ker \Delta_{\tau \circ S_\theta}^1) \subseteq \ker \Delta_\tau^1$. The analogous diagram commutes if we consider Thomsen’s variant of the de la Harpe–Skandalis determinant associated to τ and $\tau \circ S_\theta$ in (3.23).

Proof. By the proof of Proposition 2.2, we can find $k \in \mathbb{N}$ and $a_1, \dots, a_k \in (M_n(A))_{sa}$ such that ξ is homotopic to the path $\eta : [0, 1] \rightarrow U(A)$ given by

$$\eta(t) = \xi \left(\frac{j-1}{k} \right) e^{2\pi i(kt-j+1)a_j}, t \in \left[\frac{j-1}{k}, \frac{j}{k} \right] \tag{3.24}$$

and we have that

$$\tilde{\Delta}_\omega^n(\xi) = \tilde{\Delta}_\omega^n(\eta) = \sum_{j=1}^k \omega(a_j) \tag{3.25}$$

whenever $\omega : A_{sa} \rightarrow F$ is a bounded trace to a real Banach space F .

Now $\theta \circ \xi$ is homotopic to $\theta \circ \eta$, which has the following form, for $j = 1, \dots, k$ and $t \in [\frac{j-1}{k}, \frac{j}{k}]$:

$$\begin{aligned}
 \theta \circ \eta(t) &= \theta \circ \xi \left(\frac{j-1}{k} \right) \theta \left(e^{2\pi i(kt-j+1)a_j} \right) \\
 &= \theta \circ \xi \left(\frac{j-1}{k} \right) e^{2\pi i(kt-j+1)S_\theta(a_j)}.
 \end{aligned} \tag{3.26}$$

By taking $X_j := \theta \circ \xi \left(\frac{j-1}{k} \right)$ and $Y_j := e^{2\pi i(kt-j+1)S_\theta(a_j)}$ in (2.12), we see that

$$\tilde{\Delta}_\tau^1(\theta \circ \xi) = \tilde{\Delta}_\tau^1(\theta \circ \eta) = \sum_{j=1}^k \tau(S_\theta(a_j)), \tag{3.27}$$

and this last quantity is just $\tilde{\Delta}_{\tau \circ S_\theta}^1(\xi)$ by (3.25).

As for parts (2) and (3), they follow from (1). Indeed, (2) is obvious and to see (3) we let

$$q : E/\tilde{\Delta}_{\tau \circ S_\theta}^1(\pi(U^0(A))) \rightarrow E/\tilde{\Delta}_\tau^1(\pi(U^0(B))) \tag{3.28}$$

be the canonical surjection. If $\xi_u : [0, 1] \rightarrow U^0(A)$ is a path from 1 to u , then we have that

$$\begin{aligned}
 q(\Delta_{\tau \circ S_\theta}^1(u)) &= q\left(\tilde{\Delta}_{\tau \circ S_\theta}^1(\xi_u) + \tilde{\Delta}_{\tau \circ S_\theta}^1(\pi_1(U^0(A)))\right) \\
 &= \tilde{\Delta}_{\tau \circ S_\theta}^1(\xi_u) + \tilde{\Delta}_\tau^1(\pi_1(U^0(B))) \\
 &\stackrel{(1)}{=} \tilde{\Delta}_\tau^1(\theta \circ \xi_u) + \tilde{\Delta}_\tau^1(\pi_1(U^0(B))) \\
 &= \Delta_\tau^1(\theta(u)).
 \end{aligned} \tag{3.29}$$

The remark about Thomsen’s variant is obvious. □

Corollary 3.6. *The following diagram commutes:*

$$\begin{array}{ccc}
 \pi_1(U^0(A)) & \xrightarrow{\tilde{\Delta}_A^1} & \text{Aff } T(A) \\
 \pi_1(\theta) \downarrow & & \downarrow \Lambda_\theta \\
 \pi_1(U^0(B)) & \xrightarrow{\tilde{\Delta}_B^1} & \text{Aff } T(B).
 \end{array} \tag{3.30}$$

In particular, when the canonical maps

$$\pi_1(U^0(A)) \rightarrow K_0(A) \text{ and } \pi_1(U^0(B)) \rightarrow K_0(B) \tag{3.31}$$

are isomorphisms, we have that

$$\begin{array}{ccc}
 K_0(A) & \xrightarrow{\rho_A} & \text{Aff } T(A) \\
 K_0(\theta) \downarrow & & \downarrow \Lambda_\theta \\
 K_0(B) & \xrightarrow{\rho_B} & \text{Aff } T(B)
 \end{array} \tag{3.32}$$

commutes, where $K_0(\theta) : K_0(A) \rightarrow K_0(B)$ is the map induced from the diagram

$$\begin{array}{ccc}
 K_0(A) & \xrightarrow{\cong} & \pi_1(U^0(A)) \\
 K_0(\theta) \downarrow & & \downarrow \pi_1(\theta) \\
 K_0(B) & \xrightarrow{\cong} & \pi_1(U^0(B)).
 \end{array} \tag{3.33}$$

Proof. By definition, we have that $\Lambda_\theta \circ \text{Tr}_A = S_\theta \circ \text{Tr}_B$ and so if $\xi : [0, 1] \rightarrow U^0(A)$ is a (piece-wise smooth) path, we have³

$$\begin{aligned}
 \Lambda_\theta \left(\tilde{\Delta}_{\text{Tr}_A}^1(\xi) \right) &= \Lambda_\theta \left(\int_0^1 \text{Tr}_A \left(\xi'(t) \xi(t)^{-1} \right) dt \right) \\
 &= \int_0^1 \Lambda_\theta \circ \text{Tr}_A \left(\xi'(t) \xi(t)^{-1} \right) dt \\
 &= \int_0^1 \text{Tr}_B \circ S_\theta \left(\xi'(t) \xi(t)^{-1} \right) dt \\
 &= \tilde{\Delta}_{\text{Tr}_B \circ S_\theta}^1(\xi).
 \end{aligned} \tag{3.34}$$

Now we note that the diagram

$$\begin{array}{ccc}
 & \text{Aff } T(A) & \\
 \tilde{\Delta}_A^1 \nearrow & & \searrow \Lambda_\theta \\
 \pi_1(U^0(A)) & \xrightarrow{\tilde{\Delta}_{\text{Tr}_B \circ S_\theta}^1} & \text{Aff } T(B) \\
 \pi_1(\theta) \downarrow & & \downarrow \text{id} \\
 \pi_1(U^0(B)) & \xrightarrow{\tilde{\Delta}_B^1} & \text{Aff } T(B)
 \end{array} \tag{3.35}$$

³ If $\mathfrak{X}, \mathfrak{Y}$ are Banach spaces over the same base field, $f : [0, 1] \rightarrow \mathfrak{X}$ is a Riemann integrable function and $T : \mathfrak{X} \rightarrow \mathfrak{Y}$ is a bounded linear map, then $T \circ f : [0, 1] \rightarrow \mathfrak{Y}$ is Riemann integrable with $T \left(\int_0^1 f(t) dt \right) = \int_0^1 T \circ f(t) dt$.

commutes, where the top triangle commutes by the (3.34) and the bottom square commutes by Proposition 3.5(2). This gives that (3.30) commutes.

The second part follows since if ξ is a (piece-wise smooth) path in $U_n(A)$ and $m > n$, then

$$\tilde{\Delta}_\tau^m (\xi \oplus \mathbf{1}_{m-n}) = \tilde{\Delta}_\tau^n (\xi). \tag{3.36}$$

□

Proposition 3.7. *Let A, B be unital C^* -algebras. Then the map Λ_θ is a lift of the map*

$$\text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) \rightarrow \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))) \tag{3.37}$$

as described in (3.4).

Proof. Let us label the maps in the diagram (3.4):

$$\begin{array}{ccc} U^0(A)/\ker \Delta_A^1 & \xrightarrow{\delta_A^1} & \text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) \\ \tilde{\theta} \downarrow & & \downarrow P \\ U^0(B)/\ker \Delta_B^1 & \xrightarrow{\delta_B^1} & \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))) \end{array} \tag{3.38}$$

where $\tilde{\theta}([u]) := [\theta(u)]$, $\delta_A^1([e^{2\pi ia}]) := [\hat{a}]$, δ_B^1 is defined similarly, and

$$P = \delta_B^1 \circ \tilde{\theta} \circ (\delta_A^1)^{-1}. \tag{3.39}$$

But then we have that

$$\begin{aligned} P([a]) &= \delta_B^1 \circ \tilde{\theta} \circ (\delta_A^1)^{-1}([\hat{a}]) \\ &= \delta_B^1 \circ \tilde{\theta}([e^{2\pi ia}]) \\ &= \delta_B^1([\theta(e^{2\pi ia})]) \\ &= \delta_B^1([e^{2\pi iS_\theta(a)}]) \\ &= [\widehat{S_\theta(a)}] \\ &= [\Lambda_\theta(\hat{a})] \end{aligned} \tag{3.40}$$

In particular, the diagram

$$\begin{array}{ccc} \text{Aff } T(A) & \xrightarrow{q_A^1} & \text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) \\ \Lambda_\theta \downarrow & & \downarrow P \\ \text{Aff } T(B) & \xrightarrow{q_B^1} & \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))) \end{array} \tag{3.41}$$

commutes, where q_A^1 and q_B^1 are the respective quotient maps. □

Proposition 3.8. *Let A, B be unital C^* -algebras and $\theta:U^0(A) \rightarrow U^0(B)$ be a continuous group homomorphism. Then, $\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ is a lift of the map*

$$\text{Aff } T(A)/\overline{\tilde{\Delta}_A^1(\pi_1(U^0(A)))} \rightarrow \text{Aff } T(B)/\overline{\tilde{\Delta}_B^1(\pi_1(U^0(B)))} \tag{3.42}$$

as described in (3.2).

Proof. One can mimic the proof above or apply the above result and appeal to the commuting diagram

$$\begin{array}{ccc}
 \text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A))) & \longrightarrow & \overline{\text{Aff } T(A)/\tilde{\Delta}_A^1(\pi_1(U^0(A)))} \\
 \downarrow & & \downarrow \\
 \text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B))) & \longrightarrow & \overline{\text{Aff } T(B)/\tilde{\Delta}_B^1(\pi_1(U^0(B)))},
 \end{array} \tag{3.43}$$

where the vertical maps are defined via the diagrams from (3.4) and (3.2), respectively, and the horizontal maps are the canonical surjections. □

In particular, assuming some K_0 -regularity gives that Λ_θ is a lift of a map between quotients of spaces of continuous real-valued affine functions by images of K_0 .

Corollary 3.9. *Let A, B be unital C^* -algebras such that the canonical maps*

$$\pi_1(U^0(A)) \rightarrow K_0(A) \text{ and } \pi_1(U^0(B)) \rightarrow K_0(B) \tag{3.44}$$

are surjections. If $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous homomorphism, then Λ_θ is a lift of the maps on the right of the following two commutative diagrams:

$$\begin{array}{ccc}
 U^0(A)/CU^0(A) & \xrightarrow{\cong} & \overline{\text{Aff } T(A)/\rho_A(K_0(A))} \\
 \downarrow & & \downarrow \\
 U^0(B)/CU^0(B) & \xrightarrow{\cong} & \overline{\text{Aff } T(B)/\rho_B(K_0(B))}
 \end{array} \tag{3.45}$$

and

$$\begin{array}{ccc}
 U^0(A)/\ker \Delta_A^1 & \xrightarrow{\cong} & \text{Aff } T(A)/\rho_A(K_0(A)) \\
 \downarrow & & \downarrow \\
 U^0(B)/\ker \Delta_B^1 & \xrightarrow{\cong} & \text{Aff } T(B)/\rho_B(K_0(B)).
 \end{array} \tag{3.46}$$

Further, if $\ker \Delta_A^1 = DU^0(A)$ and $\ker \Delta_B^1 = DU^0(B)$, then Λ_θ is a lift of the map induced by the diagram

$$\begin{array}{ccc}
 U^0(A)/DU^0(A) & \xrightarrow{\cong} & \overline{\text{Aff } T(A)/\rho_A(K_0(A))} \\
 \downarrow & & \downarrow \\
 U^0(B)/DU^0(B) & \xrightarrow{\cong} & \overline{\text{Aff } T(B)/\rho_B(K_0(B))}.
 \end{array} \tag{3.47}$$

C^* -algebras satisfying the last condition arise naturally—for example, unital, separable, simple, pure C^* -algebras of stable rank 1 such that every 2-quasitracial state on A is a trace have this property [27].

4. The order on $\text{Aff } T(\cdot)$

We now examine when the map induced on $\text{Aff } T(\cdot)$ is positive in order to compare K -theory, traces, and the pairing. As we saw in Example 3.4, the map we get between spaces of affine functions on the

trace simplices need not be positive nor unital in general. In this section, we will be able to use the map Λ_θ to construct a unital positive map, under some extra assumptions on θ .

We record the following results as they give us necessary and sufficient conditions for the Λ_θ to be positive. We use the C^* -algebra-valued analog of the fact that any unital, contractive linear functional is positive, along with the fact that completely positive maps are (completely) bounded with the norm determined by the image of the unit. Recall that an operator system is a self-adjoint unital subspace of a C^* -algebra. The following is a combination of Proposition 2.11, Theorem 3.9, and Proposition 3.6 in [29].

Proposition 4.1. *Let S be an operator system and B a unital C^* -algebra.*

1. *If $\phi : S \rightarrow B$ is a unital contraction, then ϕ is positive.*
2. *If $B = C(X)$ and $\phi : S \rightarrow B$ is positive, then it is bounded with $\|\phi\| = \|\phi(1)\|$.*

Lemma 4.2. *Suppose A, B are unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ be a continuous group homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. If $\theta|_{\mathbb{T}}$ is injective, then $S_\theta(1) \in \{1, -1\}$.*

Proof. The restriction $\theta|_{\mathbb{T}} : \mathbb{T} \rightarrow \mathbb{T}$ is a continuous group homomorphism, hence by Pontryagin duality is of the form $\theta(z) = z^n$ for some n . Injectivity implies that $n \in \{1, -1\}$. We then have that

$$e^{2\pi i S_\theta(1)t} = \theta(e^{2\pi it}) = e^{2\pi int} \tag{4.1}$$

for all $t \in \mathbb{R}$. This implies that $S_\theta(1) = n \cdot 1 \in \{1, -1\}$. □

In the sequel, we will be interested in the case that $T(A) \neq \emptyset$. The map S_θ will still descend to a map $\Lambda_\theta : A_{sa}/A_0 \rightarrow B_{sa}/B_0$ regardless of their being tracial states on either C^* -algebra, but these quotients are zero if there are no traces. If we take the zero Banach space to be partially ordered with order unit trivially, then one can speak of unital and positive maps $\text{Aff } T(A) \rightarrow \text{Aff } T(B)$, regardless of whether $T(B)$ is empty or not (any map from a partially ordered Banach space with order unit to the zero Banach space will be unital and positive). However, if $T(A) = \emptyset$ and $T(B) \neq \emptyset$, then our map Λ_θ would have no chance of being unital. We identify the zero Banach space with $\text{Aff } (\emptyset) = C_{\mathbb{R}}(\emptyset)$ and with the complex analogs as well. However, the following lemma is true regardless of whether $T(A)$ is empty or not.

Lemma 4.3. *Suppose A, B are unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous group homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. If θ is injective, the following are equivalent:*

1. *one of Λ_θ or $-\Lambda_\theta$ is positive;*
2. *Λ_θ is contractive.*

Proof. First suppose that $T(A), T(B) \neq \emptyset$. By Lemma 4.2, we know that $S_\theta(1) \in \{1, -1\}$ and consequently $\Lambda_\theta(\hat{1}) \in \{1, -1\}$ (where we recall that, for $a \in A_{sa}$, $\hat{a} \in \text{Aff } T(A)$ is the affine function $\hat{a}(\tau) = \tau(a)$). By replacing Λ_θ with $-\Lambda_\theta$, we can without loss of generality assume that Λ_θ is unital. Using the fact that $\text{Aff } T(A) + i\text{Aff } T(A) \subseteq C(T(A))$ is an operator system and the canonical extension

$$\Lambda_\theta^{\mathbb{C}} : \text{Aff } T(A) + i\text{Aff } T(A) \rightarrow \text{Aff } T(B) + i\text{Aff } T(B) \subseteq C(T(B)) \tag{4.2}$$

is a unital linear map with abelian target algebra, this is an easy consequence of the two parts of Proposition 4.1.

Finally, if $T(A) = \emptyset$, then any map from $\text{Aff } T(A)$ is both contractive and positive trivially. The same is true of any map with codomain $\text{Aff } T(B)$ if $T(B) = \emptyset$. □

Lemma 4.4. *Suppose A, B are unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous group homomorphism. Suppose that $T(A) \neq \emptyset$.*

1. $\|\Lambda_\theta\| \leq \|S_\theta\|$.
2. If $K > 0$ is such that $\|\theta(u) - \theta(v)\| \leq K\|u - v\|$ for all $u, v \in U^0(A)$, then $\|S_\theta\| \leq K$ and $\|\Lambda_\theta\| \leq K$.
3. If θ is a homeomorphism, then Λ_θ is bijective with bounded inverse being $\Lambda_{\theta^{-1}}$.
4. If θ is isometric, then so is S_θ . If θ is a surjective isometry, then Λ_θ is a surjective isometry.

Proof. As $S_\theta(A_0) \subseteq B_0$, we have

$$\begin{aligned} \|\Lambda_\theta(\hat{a})\|_{\text{Aff } T(B)} &= \|\widehat{S_\theta(a)}\|_{\text{Aff } T(B)} \\ &= \inf_{b \in B_0} \|S_\theta(a) + b\| \\ &\leq \inf_{b \in \theta(A_0)} \|S_\theta(a) + b\| \\ &= \inf_{a' \in A_0} \|S_\theta(a) + S_\theta(a')\| \\ &\leq \|S_\theta\| \inf_{a' \in A_0} \|a + a'\| \\ &= \|S_\theta\| \|\hat{a}\|_{\text{Aff } T(A)} \end{aligned} \tag{4.3}$$

whenever $a \in A_{sa}$. This gives that $\|\Lambda_\theta\| \leq \|S_\theta\|$.

The fact that S_θ is an isometry whenever θ is an isometry follows from an argument in [24]; we exemplify said argument to show the bound condition. We use the observation that

$$\frac{e^{2\pi ita} - 1}{t} \rightarrow 2\pi ia \tag{4.4}$$

as $t \rightarrow 0$. Since

$$\|e^{2\pi itS_\theta(a)} - 1\| \leq K\|e^{2\pi ita} - 1\| \tag{4.5}$$

for all $t \in \mathbb{R}$, we can divide both sides by $\frac{1}{2\pi}|t|$ and take $t \rightarrow 0$ to get that

$$\|S_\theta(a)\| \leq K\|a\|. \tag{4.6}$$

Thus, $\|S_\theta\| \leq K$. It then follows from (1) that $\|\Lambda_\theta\| \leq K$ as well.

If θ is a homeomorphism, S_θ and $S_{\theta^{-1}}$ are both defined and its clear that $S_\theta^{-1} = S_{\theta^{-1}}$.

The surjectivity of θ implies the surjectivity of S_θ and thus if $b \in B_{sa}$, we can find $a \in A_{sa}$ with $S_\theta(a) = b$. But then $\Lambda_\theta(\hat{a}) = \widehat{S_\theta(a)} = \hat{b}$. In particular, Λ_θ is surjective. Now if θ is a surjective isometry, we identify $\text{Aff } T(A) \simeq A_{sa}/A_0$ and $\text{Aff } T(B) \simeq B_{sa}/B_0$, noting that $S_\theta(A_0) = B_0$, and that Λ_θ will preserve the quotient norms. □

Corollary 4.5. *Suppose A, B are unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous group homomorphism. Suppose that $T(A) \neq \emptyset$. If $S_\theta(1) = n$ and $\|S_\theta\| = |n|$, then $\frac{1}{n}S_\theta$ is a unital contraction, hence positive. In particular, if $\theta(\mathbb{T}) = \mathbb{T}$ and $\theta|_{\mathbb{T}}$ is an injection, then either Λ_θ or $-\Lambda_\theta$ is unital and positive.*

Proof. The first part follows from the above lemma. If θ is an injection with $\theta(\mathbb{T}) = \mathbb{T}$, we have that $S_\theta(\hat{1}) \in \{\hat{1}, -\hat{1}\}$ and that Λ_θ is contractive, so one of Λ_θ or $-\Lambda_\theta$ is a unital contraction, hence positive by part (1) of Proposition 4.1. □

Theorem 4.6. *Suppose A, B are unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ is a contractive injective homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. Suppose that $T(A) \neq \emptyset$. Then there is a continuous affine map $T_\theta : T(B) \rightarrow T(A)$ such that $\Lambda_\theta(f) = f \circ T_\theta$ or $-\Lambda_\theta(f) = f \circ T_\theta$, depending on whether Λ_θ or $-\Lambda_\theta$ is positive.*

Proof. This follows from the fact that the induced map $\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ will have the property that Λ_θ or $-\Lambda_\theta$ will be a unital positive map. Therefore, by contravariant identification of compact

convex sets (of locally convex Hausdorff linear spaces) with the state space of the space of continuous real-valued affine valued functions on them [21, Chapter 7], there exists a continuous affine map $T_\theta : T(B) \rightarrow T(A)$. \square

Theorem 4.7. *Suppose A, B be unital C^* -algebras and $\theta : U^0(A) \rightarrow U^0(B)$ is a contractive topological group isomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. Suppose that $T(A) \neq \emptyset$. Then the map $T_\theta : T(B) \rightarrow T(A)$ induced by Λ_θ is an affine homeomorphism.*

Proof. As $\theta(\mathbb{T}) = \mathbb{T}$, $S_\theta(1) \in \{-1, 1\}$. Let $\pm\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ be either Λ_θ or $-\Lambda_\theta$, depending on which is unital, positive, contractive and surjective by combining Lemmas 4.4, 4.3 and 3.2(2). By the duality of (compact) simplices and continuous affine functions on them, the map $T_\theta : T(B) \rightarrow T(A)$ is an affine homeomorphism. \square

Theorem 4.8. *Let A, B be unital C^* -algebras and $\theta : U(A) \rightarrow U(B)$ be a contractive injective homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. Suppose that $T(A) \neq \emptyset$. If*

$$\pi_i(U(A)) \simeq K_{i-1}(A) \text{ and } \pi_i(U(B)) \simeq K_{i-1}(B) \text{ for } i = 0, 1,^4 \tag{4.7}$$

via the canonical maps, then there is an induced map

$$KT_u(\theta) : KT_u(A) \rightarrow KT_u(B). \tag{4.8}$$

Proof. Let

- $\Lambda := \Lambda_{\theta|_{U^0(A)}} : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$,
- $\theta_0 : \pi_1(U^0(A)) \rightarrow \pi_1(U^0(B))$ be the map induced on fundamental groups by $\theta|_{U^0(A)}$,
- $K_0(\theta)$ be the map induced on K_0 by θ_0 together with (4.7) for $i = 1$,
- $\theta_1 : \pi_0(U(A)) \rightarrow \pi_0(U(B))$ be the map induced by θ on connected components (so that $\theta_1([u]_{\sim_n}) = [\theta_1(u)]_{\sim_n}$) and
- $K_1(\theta)$ be the map induced by θ_1 together with (4.7) for $i = 0$.

Then

$$(\pm K_0(\theta), K_1(\theta), \pm\Lambda) : KT_u(A) \rightarrow KT_u(B) \tag{4.9}$$

is a KT_u -morphism, where $\pm\Lambda$ is either Λ or $-\Lambda$ depending on which one is unital and positive, and $\pm K_0(\theta)$ is either $K_0(\theta)$ if Λ is positive or $-K_0(\theta)$ if $-\Lambda$ is positive. Indeed, $\pm K_0(\theta), \theta_1, \pm\Lambda$ are all appropriate morphisms, and we have that

$$\begin{array}{ccc} K_0(A) & \xrightarrow{\rho_A} & \text{Aff } T(A) \\ \pm K_0(\theta) \downarrow & & \downarrow \pm\Lambda \\ K_0(B) & \xrightarrow{\rho_B} & \text{Aff } T(B) \end{array} \tag{4.10}$$

commutes⁵ by Corollary 3.6. \square

Corollary 4.9. *Let A, B be unital C^* -algebras, $\theta : U^0(A) \rightarrow U^0(B)$ is a contractive topological group isomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$, and suppose that $T(A) \neq \emptyset$. If*

$$\pi_i(U(A)) \simeq K_{i-1}(A) \text{ and } \pi_i(U(B)) \simeq K_{i-1}(B) \text{ for } i = 0, 1, \tag{4.11}$$

via the canonical maps, then $KT_u(A) \simeq KT_u(B)$.

⁴Note that $\pi_1(U(A)) = \pi_1(U^0(A))$ and $\pi_1(U(B)) = \pi_1(U^0(B))$ since we are taking our base point to be the identity in each case.
⁵The map $-K_0(\theta)$ will take a piece-wise smooth loop ξ to the loop $-\theta \circ \xi$ defined by $(-\theta \circ \xi)(t) = \theta(\xi(-t))$. From here its obvious that the diagram commutes.

Proof. By Corollary 4.8, we have an induced KT_u -morphism. This map is necessarily an isomorphism since θ is. □

Corollary 4.10. *Let A, B be unital C^* -algebras which are either \mathcal{Z} -stable or of stable rank one and suppose that $T(A) \neq \emptyset$. Let $\theta : U(A) \rightarrow U(B)$ be a contractive injective homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$. Then there is an induced map*

$$KT_u(\theta) : KT_u(A) \rightarrow KT_u(B).$$

Proof. C^* -algebras which are \mathcal{Z} -stable or have stable rank one satisfy the hypotheses of Theorem 4.8 by [25] and [32], respectively. So Theorem 4.8 applies. □

Remark 4.11. *The strict ordering on $\text{Aff } T(A)$ is given by $f \gg g$ if $f(\tau) > g(\tau)$ for all $\tau \in T(A)$. If A, B are unital with $T(A) \neq \emptyset$ and $\theta : U^0(A) \rightarrow U^0(B)$ is a contractive injective homomorphism such that $\theta(\mathbb{T}) = \mathbb{T}$, then $\pm\Lambda_\theta : \text{Aff } T(A) \rightarrow \text{Aff } T(B)$ is a unital positive contraction by Lemma 4.3 (again $\pm\Lambda_\theta$ is Λ_θ or $-\Lambda_\theta$ depending on which is positive). We moreover have that*

$$\pm\Lambda_\theta(f) \gg \pm\Lambda_\theta(g) \iff f \gg g. \tag{4.12}$$

Indeed, let us show that $f \gg 0$ if and only if its image is $\gg 0$. As $\pm\Lambda_\theta$ has the form $\pm\Lambda_\theta(\hat{a}) = \widehat{\pm S_\theta(a)}$, it suffices to show that if $\sigma(a) > 0$ for all $\sigma \in T(A)$, then $\tau(\pm S_\theta(a)) > 0$ for all $\tau \in T(B)$. But this is trivial because $\tau \circ \pm S_\theta : A_{sa} \rightarrow \mathbb{R}$ extends canonically to a tracial state $A \rightarrow \mathbb{C}$ by Proposition 3.3, so evaluating it against a must give that it is strictly positive.

The above says the following: for certain C^* -algebras, we can read off positivity in K_0 , thinking of it as the fundamental group of the unitary group, from the strict positivity of the predeterminant applied a the loop. Precisely, a nonzero element $x \in K_0(A)$, where A is a unital, simple C^* -algebra with strict comparison, is in the positive cone if and only if the corresponding loop ξ_x satisfies $\Delta_\tau(\xi_x) > 0$ for all $\tau \in T(A)$.

Although the following is known, for example, by very strong results in [1, Chapter 6] pertaining to certain prime C^* -algebras, we give the following as a corollary by using K -theoretic classification results.

Corollary 4.12. *Let A, B be unital, separable, simple, nuclear \mathcal{Z} -stable C^* -algebras satisfying the UCT. Then $A \simeq B$ if and only if there is a contractive isomorphism $U(A) \simeq U(B)$.*

Proof. Its clear that two isomorphic C^* -algebras have isomorphic unitary groups. On the other hand, if $U(A) \simeq U(B)$, then since these C^* -algebras are \mathcal{Z} -stable, Corollary 4.9 applies. As $KT_u(\cdot)$ recovers the Elliott invariant, which is a complete invariant for the C^* -algebras as in the statement of the theorem (by [5, Corollary D], [16, 18, 19] and the references therein), $A \simeq B$. □

Using the state-of-the art classification of embeddings [6], there is an enlarged invariant of $KT_u(\cdot)$, which is able to classify morphisms between certain C^* -algebras. Any KT_u -morphisms automatically has a lift to this larger invariant [6, Theorem 3.9], and so under the assumption that the KT_u -morphism is faithful (i.e., the map $T(B) \rightarrow T(A)$ induced by the map $\text{Aff } T(A) \rightarrow \text{Aff } T(B)$ sends traces on B to faithful traces on A), there is a $*$ -homomorphism witnessing the KT_u -morphism. Therefore, as a corollary of their main theorem, we have that for an abundance of C^* -algebras, there is an (contractive) embedding of unitary groups if and only if there is an embedding of C^* -algebras.

Corollary 4.13. *Let A be a unital, separable, simple nuclear C^* -algebra satisfying the UCT which is either \mathcal{Z} -stable or of stable rank one and B be a unital, separable, simple, nuclear \mathcal{Z} -stable C^* -algebra.*

If there is a contractive injective homomorphism $\theta : U(A) \rightarrow U(B)$ such that $\theta(\mathbb{T}) = \mathbb{T}$, then there is an embedding $A \hookrightarrow B$.

Proof. Assuming such a θ exists, it gives rise to a KT_u -morphism

$$KT_u(\theta) : KT_u(A) \rightarrow KT_u(B) \tag{4.13}$$

by Corollary 4.10. As A, B are simple, the map $T_\theta : T(B) \rightarrow T(A)$ necessarily maps traces on B to faithful traces on A , and thus the KT_u -morphism $KT_u(\theta)$ is “faithful.” Therefore, $KT_u(\theta)$ induces an embedding $A \hookrightarrow B$ by [6, Theorem B]. \square

5. A slight general linear variant

Here, we briefly describe a slight general linear variant of some of the results above. Unfortunately, the maps we get at the level of A, B and complex-valued continuous affine functions are not \mathbb{C} -linear in general (see Example 5.3). In the presence of a continuous homomorphism $\theta : GL^0(A) \rightarrow GL^0(B)$, we have corresponding maps

$$\begin{array}{ccc} GL^0(A)/CGL^0(A) & \xrightarrow{\cong} & \left(A / \overline{[A, A]} \right) / \overline{\tilde{\Delta}_A^1(\pi_1(U^0(A)))} \\ \downarrow & & \downarrow \\ GL^0(B)/CGL^0(B) & \xrightarrow{\cong} & \left(A / \overline{[A, A]} \right) / \overline{\tilde{\Delta}_B^1(\pi_1(U^0(B)))}. \end{array} \tag{5.1}$$

Again, by modding out by algebraic commutator subgroups when $DGL^0(A) = \ker \Delta_A^1$ and $DGL^0(B) = \ker \Delta_B^1$ (both with respect to the general linear variant of the de la Harpe–Skandalis determinant, as originally introduced in [12]), instead of closures of derived groups, there is a purely algebraic variant of the above diagram:

$$\begin{array}{ccc} GL^0(A)/DGL^0(A) & \xrightarrow{\cong} & \left(A / \overline{[A, A]} \right) / \tilde{\Delta}_A^1(\pi_1(GL^0(A))) \\ \downarrow & & \downarrow \\ GL^0(B)/DGL^0(B) & \xrightarrow{\cong} & \left(A / \overline{[B, B]} \right) / \tilde{\Delta}_B^1(\pi_1(GL^0(B))). \end{array} \tag{5.2}$$

Thinking of $K_0(A)$ as the Grothendieck group of the semigroup of equivalence classes of idempotents and $K_0(A) \simeq \pi_1(GL_\infty^0(A))$, we would like to lift the maps on the right of (5.1) and (5.2) to a map

$$A / \overline{[A, A]} \rightarrow B / \overline{[B, B]} \tag{5.3}$$

(the latter holding true when A, B are C^* -algebras whose determinant has appropriate kernel).

We can always achieve a bounded \mathbb{R} -linear map.

Proposition 5.1. *Let A, B be unital C^* -algebras and $\theta : GL^0(A) \rightarrow GL^0(B)$ be a continuous group homomorphism. Then there is a continuous \mathbb{R} -linear map*

$$\tilde{G}_\theta : A / \overline{[A, A]} \rightarrow B / \overline{[B, B]} \tag{5.4}$$

which lifts the maps on the right of (5.1) and (5.2) (the latter holding when $DGL^0(A) = \ker \Delta_A^1$ and $DGL^0(B) = \ker \Delta_B^1$).

Proof. We define $G_\theta : A \rightarrow B$ given by

$$G_\theta(a) := \lim_n \frac{n}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{n}} \right). \tag{5.5}$$

We note that the sequence on the right is eventually constant: choose N large enough such that $n \geq N$ implies that

$$\|\theta \left(e^{2\pi i \frac{a}{n}} \right) - 1\| < 1. \tag{5.6}$$

We then have for $n \geq N$,

$$\begin{aligned} \frac{n}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{n}} \right) &= \frac{n}{2\pi i} \log \theta \left(e^{2\pi i \frac{Na}{Nn}} \right) \\ &= \frac{n}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{Nn}} \right)^N \\ &= \frac{Nn}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{Nn}} \right) \\ &= \frac{N}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{Nn}} \right)^n \\ &= \frac{N}{2\pi i} \log \theta \left(e^{2\pi i \frac{na}{Nn}} \right) \\ &= \frac{N}{2\pi i} \log \theta \left(e^{2\pi i \frac{a}{N}} \right). \end{aligned} \tag{5.7}$$

To see that the map is additive, one can use the Lie product formula

$$e^{a+b} = \lim_k \left(e^{\frac{a}{k}} e^{\frac{b}{k}} \right)^k \tag{5.8}$$

(see for example [23, Theorem 2.11]), along with the fact that θ is a continuous homomorphism. From here, it is clear that G_θ is continuous and \mathbb{Q} -linear, hence \mathbb{R} -linear. Moreover, one can use the formula

$$e^{[a,b]} = \lim_k \left(e^{-\frac{a}{k}} e^{-\frac{b}{k}} e^{\frac{a}{k}} e^{\frac{b}{k}} \right)^{k^2} \tag{5.9}$$

(a variation of the argument given in the proof of [23, Theorem 2.11] will give the desired formula), together with θ being a continuous group homomorphism, to show that it respects commutation. Note that the same proof indeed shows that S_θ respects commutation, although we never explicitly used this. From here, $G_\theta([A, A]) \subseteq [B, B]$, and consequently, $G_\theta(\overline{[A, A]}) \subseteq \overline{[B, B]}$ by continuity. Thus, there is an induced \mathbb{R} -linear map

$$\tilde{G}_\theta : A/\overline{[A, A]} \rightarrow B/\overline{[B, B]}. \tag{5.10}$$

The fact that \tilde{G}_θ is a lift of the maps on the right of (5.1) and (5.2) follows from the same arguments as in Propositions 3.7 and 3.8. □

Remark 5.2. We note that G_θ could have also been defined in the same manner as S_θ . In particular, there is a correspondence

$$\theta \left(e^{2\pi i t a} \right) = e^{2\pi i t G_\theta(a)}, t \in \mathbb{R}. \tag{5.11}$$

One can use this to show that $\theta(\ker \Delta_A^1) \subseteq \ker \Delta_B^1$ as in Proposition 3.5 by using G_θ in place of S_θ , along with the fact that $G_\theta([A, A]) \subseteq [B, B]$. So there will always be a general linear variant of the commuting diagram (3.4).

Let us use (5.11) to show that the maps G_θ and \tilde{G}_θ in Proposition 5.1 are not always \mathbb{C} -linear.

Example 5.3. Consider $A = B = \mathbb{C}$ and $\theta : \mathbb{C}^\times \rightarrow \mathbb{C}^\times$ given by $\theta(z) = |z|^{\alpha+\beta i} z^n$ where $\alpha, \beta \in \mathbb{R}$ and $n \in \mathbb{Z}$. It is easy to see that θ is a continuous group homomorphism. However, the map G_θ is not \mathbb{C} -linear. Indeed, we have that

$$\begin{aligned} \theta(e^{2\pi i t(a+bi)}) &= |e^{2\pi i t(a+bi)}|^{\alpha+\beta i} e^{2\pi i t(a+bi)} \\ &= (e^{-2\pi t b})^{\alpha+\beta i} e^{2\pi i t(a+bi)} \\ &= e^{-2\pi t b \alpha} e^{-2\pi i t \beta b} e^{2\pi i t(a+bi)} \\ &= e^{2\pi i t((n\alpha-\alpha b)+i(n-\beta)b)}. \end{aligned} \tag{5.12}$$

In particular, thinking of \mathbb{C} as \mathbb{R}^2 with 1 and i corresponding to the basis vectors $(1, 0)$ and $(0, 1)$, respectively, we have that $G_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the map

$$G_\theta \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} n & -\alpha \\ 0 & n - \beta \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}. \tag{5.13}$$

In this example, the map G_θ is \mathbb{C} -linear if and only if $\alpha = \beta = 0$. We do note, however, that θ sends unitaries to unitaries and $\theta|_{\mathbb{T}}(z) = z^n$.

In general, it is clear that if $\theta : GL(A) \rightarrow GL(B)$ sends unitaries to unitaries, then we can use techniques in Section 4 to get maps between spaces of continuous affine functions on the trace simplices. If one had that θ was the restriction of a $*$ -homomorphism or a conjugate-linear $*$ -homomorphism, then this would be true.

6. Final remarks and open questions

An alternate way to construct the map Λ_θ , using duality of traces, is as follows. Denote by $\mathfrak{T}_s(A)$ the set of all tracial functionals on A . Suppose that A, B are unital C^* -algebras with $T(A), T(B) \neq \emptyset$, and $\theta : U^0(A) \rightarrow U^0(B)$ is a continuous homomorphism. Define, for $a \in A_{sa}$ and $\tau \in \mathfrak{T}_s(B)$,

$$F(\tau)(a) := \lim_{n \rightarrow \infty} \tau \left(\frac{n}{2\pi i} \log \theta(e^{2\pi i \frac{a}{n}}) \right). \tag{6.1}$$

Proposition 6.1. For $\tau \in \mathfrak{T}_s(B)$, the map $F(\tau) : A_{sa} \rightarrow \mathbb{R}$ is a well-defined, bounded, self-adjoint, tracial functional. Moreover, $F : \mathfrak{T}_s(B) \rightarrow \mathfrak{T}_s(A)$ given by $\tau \mapsto F(\tau)$ is a bounded \mathbb{R} -linear map.

Proof. Using the same arguments as in Proposition 5.1, it is clear the formula (6.1) is well defined and gives rise to a bounded \mathbb{R} -linear map $F : \mathfrak{T}_s(B) \rightarrow \mathfrak{T}_s(A)$. □

One can identify $(A_{sa}/A_0)^* \simeq \mathfrak{T}_s(A)$ (see, e.g., [8]), and so we can use duality to get a map $\tilde{\Lambda}_\theta := F^* : \mathfrak{T}_s(A)^* \rightarrow \mathfrak{T}_s(B)^*$ and restrict it to the dense set A_{sa}/A_0 . One can check that the image lies in B_{sa}/B_0 and that the restriction is just the map Λ_θ that we got before.

We finish by listing some open problems.

1. There are classes where topological isomorphisms between $U(A)$ and $U(B)$ (or even $U^0(A)$ and $U^0(B)$) come from $*$ -isomorphisms or anti- $*$ -isomorphisms. For example, if A, B are prime traceless C^* -algebras containing full square zero elements, this is true by results in [9].

If A is a unital, separable, nuclear C^* -algebra satisfying the UCT and B is a unital simple separable nuclear \mathcal{Z} -stable C^* -algebra, then unital embeddings $A \hookrightarrow B$ are classified by an invariant $KT_u(\cdot)$, which is an enlargement of KT_u [6]. Thus, any isometric unitary group homomorphism $U(A) \rightarrow U(B)$ will give a KT_u -morphism $KT_u(\theta)$ and therefore there will be an embedding

$\phi : A \hookrightarrow B$ such that $KT_u(\phi) = KT_u(\theta)$. However, it is not clear that ϕ satisfies $\phi|_{U(A)} = \theta$. More generally though – in the tracial setting – are there continuous group homomorphisms which do not have lifts to $*$ -homomorphisms or anti- $*$ -homomorphisms?

Note that in [1, Chapter 6], Lie isomorphisms between certain C^* -algebras are shown to be the sum of a Jordan $*$ -isomorphism and a center-valued trace. Is there a result for general (injective) Lie homomorphisms between certain classes of C^* -algebras?

2. This enlargement of KT_u discussed in [6] contains K -theory with coefficients (along with appropriate pairing maps—the Bosckstein maps discussed in [37]). So we ask: do continuous group homomorphisms induce maps between K -theory with coefficients?
3. For a general continuous homomorphism $\theta : U^0(A) \rightarrow U^0(B)$, does the norm $\|S_\theta\|$ determine a Lipschitz constant for θ ? We clearly have that

$$\|S_\theta\| \leq \inf\{K \mid \theta \text{ is } K\text{-Lipschitz}\} \quad (6.2)$$

by Lemma 4.4. Is this equality?

4. For A simple (or prime), is it true that any continuous injective homomorphism $\theta : U^0(A) \rightarrow U^0(B)$ is isometric? Contractive? What if B is simple (or prime)?
5. In the initial draft of this paper, we claimed that any continuous group homomorphism in Proposition 5.1 gave rise to a \mathbb{C} -linear \tilde{G}_θ . This is clearly false by Example 5.3. Is there a way to guarantee that the map \tilde{G}_θ is \mathbb{C} -linear? Or can one alter it accordingly for this to happen? Or alter it to get a map between unitary groups, which would then allow one to use the results in Section 4? Maybe if one starts with an injective, contractive group homomorphism $GL^0(A) \rightarrow GL^0(B)$ which sends C^\times to C^\times , one can say something.

Acknowledgments. Many thanks to my PhD supervisors Thierry Giordano and Aaron Tikuisis for many helpful discussions. Thanks to the authors of [6] for sharing a draft of their paper. Finally, thanks to the referee for asking for clarification on the general linear variant, which led me to consider a counterexample to \mathbb{C} -linearity.

References

- [1] P. Ara and M. Mathieu, *Local multipliers of C^* -algebras*, Springer Monographs in Mathematics (Springer, London, 2003).
- [2] A. Al-Rawashdeh, A. Booth and T. Giordano, Unitary groups as a complete invariant, *J. Funct. Anal.* **262**(11) (2012), 4711–4730.
- [3] A. Booth, The unitary group as a complete invariant for simple unital AF algebras, Master’s thesis (University of Ottawa, Canada, 1998).
- [4] M. Brešar, Commuting traces of biadditive mappings, commutativity-preserving mappings and Lie mappings, *Trans. Am. Math. Soc.* **335**(2) (1993), 525–546.
- [5] J. Castillejos, S. Evington, A. Tikuisis, S. White and W. Winter, Nuclear dimension of simple C^* -algebras, *Invent. Math.* **224**(1) (2021), 245–290.
- [6] J. R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis and S. White, Classifying $*$ -homomorphisms I: Unital simple nuclear C^* -algebras, *arXiv:2307.06480* (2023).
- [7] J. B. Conway, *A course in functional analysis*, Graduate Texts in Mathematics, vol. **96** (Springer, New York, 1985).
- [8] J. Cuntz and G. K. Pedersen, Equivalence and traces on C^* -algebras, *J. Funct. Anal.* **33**(2) (1979), 135–164.
- [9] A. Chand and L. Robert, Simplicity, bounded normal generation, and automatic continuity of groups of unitaries, *Adv. Math.* **415** (2023), PaperNo.108894,52.
- [10] M. Dadarlat, Reduction to dimension three of local spectra of real rank zero C^* -algebras, *J. Reine Angew. Math.* **460** (1995), 189–212.
- [11] P. de la Harpe, Fuglede–Kadison determinant: theme and variations, *Proc. Natl. Acad. Sci. USA* **110**(40) (2013), 15864–15877.
- [12] P. de la Harpe and G. Skandalis, Déterminant associé à une trace sur une algèbre de Banach, *Ann. Inst. Fourier (Grenoble)* **34**(1) (1984), 241–260.
- [13] P. de la Harpe and G. Skandalis, Produits finis de commutateurs dans les C^* -algèbres, *Ann. Inst. Fourier (Grenoble)* **34**(4) (1984), 169–202.
- [14] H. A. Dye, The unitary structure in finite rings of operators, *Duke Math. J.* **20** (1953), 55–69.
- [15] H. A. Dye, On the geometry of projections in certain operator algebras, *Ann. Math. (2)* **61** (1955), 61–89.

- [16] G. A. Elliott, H. L. Guihua Gong and Z. Niu, On the classification of simple amenable C^* -algebras with finite decomposition rank, II, *arXiv:1507.03437* (2015).
- [17] G. B. Folland, *A course in abstract harmonic analysis*, Textbooks in Mathematics, 2nd edition (CRC Press, Boca Raton, FL, 2016).
- [18] G. Gong, H. Lin and Z. Niu, A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, I: C^* -algebras with generalized tracial rank one, *C. R. Math. Acad. Sci. Soc. R. Can.* **42**(3) (2020), 63–450.
- [19] G. Gong, H. Lin and Z. Niu, A classification of finite simple amenable \mathcal{Z} -stable C^* -algebras, II: C^* -algebras with rational generalized tracial rank one, *C. R. Math. Acad. Sci. Soc. R. Can.* **42**(4) (2020), 451–539.
- [20] G. Gong, On inductive limits of matrix algebras over higher-dimensional spaces. I, II, *Math. Scand.* **80**(1) (1997), 41–55,56–100.
- [21] K. R. Goodearl, Partially ordered abelian groups with interpolation, *Mathematical Surveys and Monographs*, vol. **20** (American Mathematical Society, Providence, RI, 1986).
- [22] T. Giordano and A. Sierakowski, The general linear group as a complete invariant for C^* -algebras, *J. Oper. Theory* **76**(2) (2016), 249–269.
- [23] B. Hall, An elementary introduction, in *Lie groups, Lie algebras, and representations*, Graduate Texts in Mathematics, vol. **222**, 2nd edition (Springer, Cham, 2015).
- [24] O. Hatori and L. Molnár, Isometries of the unitary groups and Thompson isometries of the spaces of invertible positive elements in C^* -algebras, *J. Math. Anal. Appl.* **409**(1) (2014), 158–167.
- [25] X. Jiang, Nonstable K -theory for \mathcal{Z} -stable C^* -algebras, *arXiv preprint math/9707228* (1997).
- [26] P. W. Ng, The kernel of the determinant map on certain simple C^* -algebras, *J. Oper. Theory* **71**(2) (2014), 341–379.
- [27] P. W. Ng and L. Robert, The kernel of the determinant map on pure C^* -algebras, *Houston J. Math.* **43**(1) (2017), 139–168.
- [28] A. L. T. Paterson, Harmonic analysis on unitary groups, *J. Funct. Anal.* **53**(3) (1983), 203–223.
- [29] V. Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics, vol. **78** (Cambridge University Press, Cambridge, 2002).
- [30] N. C. Phillips, The rectifiable metric on the space of projections in a C^* -algebra, *Int. J. Math.* **3**(5) (1992), 679–698.
- [31] N. C. Phillips, A classification theorem for nuclear purely infinite simple C^* -algebras, *Doc. Math.* **5** (2000), 49–114.
- [32] M. A. Rieffel, The homotopy groups of the unitary groups of noncommutative tori, *J. Oper. Theory* **17**(2) (1987), 237–254.
- [33] M. Rørdam, F. Larsen and N. J. Laustsen, *An introduction to K -theory for C^* -algebras*, London Mathematical Society Student Texts, vol. **49** (Cambridge University Press, Cambridge, 2000).
- [34] M. Rørdam, Classification of nuclear, simple C^* -algebras, in *Classification of nuclear C^* -algebras. Entropy in operator algebras*, Encyclopaedia Math. Sci., vol. **126** (Springer, Berlin, 2002), 1–145.
- [35] S. Sakai, On the group isomorphism of unitary groups in AW^* -algebras, *Tohoku Math. J. (2)* **7**(1955), 87–95.
- [36] S. Sakai, *C^* -algebras and W^* -algebras* (Springer, Berlin, 2012).
- [37] C. Schochet, Topological methods for C^* -algebras. IV. Mod p homology, *Pac. J. Math.* **114**(2) (1984), 447–468.
- [38] K. Thomsen, Finite sums and products of commutators in inductive limit C^* -algebras, *Ann. Inst. Fourier (Grenoble)* **43**(1) (1993), 225–249.
- [39] K. Thomsen, Traces, unitary characters and crossed products by \mathbb{Z} , *Publ. Res. Inst. Math. Sci.* **31**(6) (1995), 1011–1029.
- [40] T. Yen, Isomorphism of unitary groups in AW^* -algebras, *Tohoku Math. J. (2)* **8** (1956), 275–280.