

# Oxidative stress damage in the protozoan parasite *Trypanosoma cruzi* is inhibited by Cyclosporin A

PATRICIA L. BUSTOS<sup>1</sup>, ALINA E. PERRONE<sup>1</sup>, NATALIA MILDUBERGER<sup>1,2</sup>, MIRIAM POSTAN<sup>1</sup> and JACQUELINE BUA<sup>1,2\*</sup>

<sup>1</sup> Instituto Nacional de Parasitología “Dr Mario Fatała Chabén”, A.N.L.I.S. Malbrán, Av. Paseo Colón 568, C1063AC S, Buenos Aires, Argentina

<sup>2</sup> CAECIHS, Universidad Abierta Interamericana, Av. Montes de Oca 745, 2° piso, C1270AAH, Buenos Aires, Argentina

(Received 18 October 2014; revised 20 January 2015; accepted 6 February 2015; first published online 31 March 2015)

## SUMMARY

Cyclosporin A (CsA) specifically inhibits the mitochondrial permeability transition pore (mPTP). Opening of the mPTP, which is triggered by high levels of matrix  $[Ca^{2+}]$  and/or oxidative stress, leads to mitochondrial dysfunction and thus to cell death by either apoptosis or necrosis. In the present study, we analysed the response of *Trypanosoma cruzi* epimastigote parasites to oxidative stress with 5 mM  $H_2O_2$ , by studying several features related to programmed cell death and the effects of pre-incubation with 1  $\mu$ M of CsA. We evaluated *Tc*PARP cleavage, DNA integrity, cytochrome *c* translocation, Annexin V/propidium iodide staining, reactive oxygen species production. CsA prevented parasite oxidative stress damage as it significantly inhibited DNA degradation, cytochrome *c* translocation to cytosol and *Tc*PARP cleavage. The calcein-AM/CoCl<sub>2</sub> assay, used as a selective indicator of mPTP opening in mammals, was also performed in *T. cruzi* parasites.  $H_2O_2$  treatment decreased calcein fluorescence, but this decline was partially inhibited by pre-incubation with CsA. Our results encourage further studies to investigate if there is a mPTP-like pore and a mitochondrial cyclophilin involved in this protozoan parasite.

Key words: *Trypanosoma cruzi*, cyclosporin A, programmed cell death, oxidative stress, cyclophilin.

## INTRODUCTION

Mitochondria contain a structure that forms a large and non-specific pore, permeable to molecules smaller than 1.5 kDa, which opens in the inner mitochondrial membrane and is called the mitochondrial permeability transition pore (mPTP). Opening of mPTP, which is greatly enhanced under stressful conditions such as  $Ca^{2+}$  overload, nucleotide starvation or oxidative stress, allows the release of apoptosis-related factors such as apoptosis-inducing factor, cytochrome *c* and endonuclease G into the cytosol (Crompton *et al.* 1999; Vaseva *et al.* 2012) and leads to programmed cell death. Although the identity of the pore-forming proteins remains under debate, recent works propose that dimers of the  $F_1F_0$  ATP synthase display the key features of the mPTP. A well-characterized protein regulator of the PTP is cyclophilin D (CyPD), which binds the  $F_1F_0$  ATP synthase (complex V) (Giorgio *et al.* 2013) and is the only component that has been genetically proven to be indispensable for this process (Kroemer *et al.* 2007). CyPD can be displaced from the lateral stalk of  $F_1F_0$  ATP synthase

by Cyclosporin A (CsA), which is a mPTP opening inhibitor. It is proposed that when CsA binds CyPD, its displacing from the  $F_1F_0$  ATP synthase results in enzyme reactivation (Giorgio *et al.* 2013).

However, other proteins have been extensively studied and are proposed to have a role in mPTP opening regulation, such as voltage-dependent anion channel (VDAC) and adenine nucleotide translocase (ANT) (Halestrap and Davidson, 1990; Crompton *et al.* 1999; Halestrap, 2014).

*Trypanosoma cruzi* is the protozoan parasite that causes Chagas disease or American trypanosomiasis. This is a potentially life-threatening illness and it is estimated that about 7 million to 8 million people are infected, mostly in Latin America, where Chagas disease is endemic, but also in other parts of the world through migration from endemic areas (WHO, 2014).

We have previously described the CyP gene family in *T. cruzi* which consists of 15 members (Potenza *et al.* 2006; Bua *et al.* 2008). Analyses of clusters formed by the *T. cruzi* cyclophilins with others encoded in various genomes revealed that 8 of them (*Tc*CyP19, *Tc*CyP21, *Tc*CyP22, *Tc*CyP24, *Tc*CyP35, *Tc*CyP40, *Tc*CyP42 and *Tc*CyP110) have orthologues in many different genome. Besides, 4 CyPs of the family (of 19, 22, 28 and 40 kDa, respectively) have been isolated with a

\* Corresponding author. Instituto Nacional de Parasitología “Dr Mario Fatała Chabén”, A.N.L.I.S. Malbrán, Av. Paseo Colón 568, C1063ACS, Ciudad Autónoma de Buenos Aires, Argentina. E-mail: [jacbua@yahoo.com](mailto:jacbua@yahoo.com)

CsA-affinity column from parasites lysate. They present PPIase activity, which could be inhibited by CsA *in vitro* (Potenza *et al.* 2006).

Programmed cell death has been described in mammalian cells but has also been suggested for unicellular organisms, including Trypanosomatid parasites affecting humans, such as *Leishmania* spp. (Das *et al.* 2001; Gannavaram *et al.* 2008) and *Trypanosoma* spp. (Figarella *et al.* 2005; Piacenza *et al.* 2007). Some of the main features observed in mammalian cell apoptosis, namely the loss of mitochondrial membrane potential, release of cytochrome *c*, condensation of nuclear chromatin, fragmentation of genomic DNA and surface binding of Annexin V, have been described in these unicellular organisms in response to a variety of stress inducers and parasitocidal drugs (Ameisen *et al.* 1995; Duszenko *et al.* 2006). However, the protozoan cell death pathways and the effector molecules involved in these processes remain to be identified. Several articles have reported these events as apoptotic (Jiménez-Ruiz *et al.* 2010) *et al.* or incidental (Proto *et al.* 2013). Furthermore, the existence of a mPTP-like pore in protozoan parasites has not been previously described.

In the present work, we studied the effects of oxidative stress on *T. cruzi* epimastigotes by analysing several apoptotic features described for mammals and the protective effect exerted by the cyclophilin (CyP) inhibitor CsA on the parasite.

## MATERIALS AND METHODS

### Chemical compounds

Protease inhibitors, MOPS and EDTA were obtained from Sigma-Aldrich (Sigma Co., USA). The *In Situ* Cell Death detection kit was purchased from Roche Diagnostics (Manheim, Germany). Anti-tubulin, anti-cytochrome *c* were from Santa Cruz Biotechnology, Inc. (TX, USA). The Annexin V: FITC Apoptosis Detection Kit was from BD Biosciences (Becton Dickinson & Co., NJ, USA). The anti-rabbit immunoglobulin G conjugated with horseradish peroxidase was from Jackson ImmunoResearch Labs Inc. (PA, USA). The ECL Western Blotting Detection kit was from GE Healthcare (IL, USA). MitoTracker Deep Red, dichloro-dihydro-fluorescein diacetate (DCFH-DA) and Calcein-AM were from Molecular Probes (NY, USA). CsA, used as a CyP inhibitor, was a gift from Novartis (Basel, Switzerland).

### Parasite culture

*Trypanosoma cruzi* CL Brener clone parasites were cultured in Brain Heart Infusion medium (BHI) (Difco Michigan, USA) 33 g L<sup>-1</sup>, tryptose 3 g L<sup>-1</sup>, Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O 4 g L<sup>-1</sup>, ClK 0.4 g L<sup>-1</sup>, glucose 0.3

g L<sup>-1</sup>, hemin 0.02 g L<sup>-1</sup>) supplemented with 10% fetal bovine serum-heat inactivated (FBS). Epimastigotes were collected at 5 days of culture (late log phase) at 28 °C (Potenza *et al.* 2006).

### Induction of oxidative stress

Oxidative stress was triggered by incubating 1 × 10<sup>8</sup> epimastigotes in exponential phase of growth with 5 mM H<sub>2</sub>O<sub>2</sub> (in some assays 1 mM H<sub>2</sub>O<sub>2</sub> incubation was performed) for different periods of time (30 to 180 min, as indicated for each assay) at room temperature. Parasites were centrifuged at 2300 rpm for 10 min and washed once in PBS-3% glucose. Pellets were resuspended in PBS-3% glucose for subsequent studies. To inhibit CyP, parasites were incubated with 1 μM CsA for 1 h at room temperature prior to H<sub>2</sub>O<sub>2</sub> incubation.

### Detection of Annexin V binding

A common method to detect phosphatidylserine (PS) on a cell surface is the use of Annexin V, a protein which binds PS with high affinity (Van Heerde *et al.* 1994). Oxidative stress was induced with 5 mM H<sub>2</sub>O<sub>2</sub> and samples were collected at different timepoints (30, 60 or 180 min). When inhibitory conditions were required, pre-incubation with 1 μM CsA for 1 h was performed. Annexin V binding was detected on the external surface of the plasma membrane of treated parasites using the Annexin V: FITC Apoptosis Detection Kit according to the manufacturer's protocol. Co-staining of the parasites with propidium iodide was performed, to evaluate the integrity of plasma membrane during the treatments. Fluorescence was detected in a FACSCalibur equipment (Becton Dickinson & Co., NJ, USA). Data were analysed using Cyflogic software.

### Reactive oxygen species (ROS) production

To evaluate ROS production, the fluorescent dye DCFH-DA was used. DCFH-DA is a nonpolar dye that is converted into the polar derivative non-fluorescent DCFH by cellular esterases, and can be switched into the highly fluorescent DCF when oxidized by intracellular ROS and other peroxides. Epimastigotes (1 × 10<sup>7</sup> in log phase) were collected, washed once in PBS-3% glucose and resuspended in PBS-3% glucose. Parasites were incubated with 1 μM CsA for 1 h at room temperature prior to H<sub>2</sub>O<sub>2</sub> incubation. To induce oxidative stress, two different concentrations of H<sub>2</sub>O<sub>2</sub> (1 and 5 mM) were tested for 1 h at room temperature. After treatment, parasites were loaded with 10 μM of the fluorescent dye DCFH-DA for 30 min at 28 °C. Parasites were washed once in PBS-3% glucose and resuspended in PBS for flow cytometry. Fluorescence

was detected in a FACSCalibur equipment (Becton Dickinson & Co., NJ, USA). Data were analysed using Cyflog software.

#### Subcellular fractionation

Parasites were fractionated following a described protocol (Gannavaram *et al.* 2008). Briefly,  $1 \times 10^8$  epimastigotes (in log phase) were collected and washed three times in 15 mL MES buffer (20 mM MOPS, pH 7.0, 250 mM sucrose, 3 mM EDTA). Parasites were resuspended in 0.2 mL MES buffer containing  $4 \text{ mg mL}^{-1}$  digitonin and protease inhibitors: 1 mM pepstatin A, 1 mM phenylmethylsulfonyl fluoride and 0.1 mM Na-ptyosyl-L-lysine chloro-methyl ketone. The suspension was incubated at room temperature for 5 min and centrifuged at  $10\,000 \text{ g}$  for 5 min. The resulting supernatant was collected as the cytosolic fraction and the membrane pellet as the mitochondria-enriched fraction. This fraction was resuspended in phosphate buffer (20 mM sodium phosphate, pH 7.0, 3 mM EDTA) and citrate synthase, a mitochondrial marker enzyme, was assayed as previously described (Cannata and Cazzulo, 1984), to confirm a successful procedure.

#### Cytochrome *c* release

To study cytochrome *c* localization after oxidative stress induction,  $1 \times 10^8$  epimastigotes (in log phase) were collected, washed once in PBS-3% glucose and resuspended in PBS-3% glucose. Parasites were incubated with  $1 \mu\text{M}$  CsA for 1 h at room temperature prior to  $\text{H}_2\text{O}_2$  incubation. To induce oxidative stress, parasites were incubated with 5 mM  $\text{H}_2\text{O}_2$  for 3 h at room temperature. After treatment, cytosolic and mitochondria-enriched fractions were obtained as described before. Fractions were separated by SDS-PAGE, carried out as described (Laemmli, 1970). Proteins were electrotransferred from 13.5% polyacrylamide gels to nitrocellulose membranes, which were blocked with a 5% (W/V) non-fat milk suspension for 1 h at room temperature. After incubation for 2 h with a rabbit polyclonal antibody against cytochrome *c* (1:200) and a 1:1500 dilution of the goat anti-rabbit immunoglobulin G conjugated with horseradish peroxidase. The *T. cruzi* cytochrome *c*, size  $\sim 12 \text{ kDa}$  protein (TriTrypsDB annotated as TcCLB.506949.50) was visualized by chemiluminescence with an ECL Western Blotting Detection kit.

#### In situ labelling of DNA fragments

To evaluate DNA integrity after oxidative stress treatment,  $1 \times 10^7$  epimastigotes (in log phase) were collected, washed once and resuspended in PBS-3% glucose. Parasites were incubated with  $1$

$\mu\text{M}$  CsA for 1 h at room temperature prior to  $\text{H}_2\text{O}_2$  incubation. To induce oxidative stress, parasites were incubated with 5 mM  $\text{H}_2\text{O}_2$  for 3 h at room temperature.

*In situ* detection of DNA fragments by TUNEL was performed using the *In Situ* Cell Death detection kit (Roche). Epimastigotes were harvested, fixed in 4% formaldehyde and coated onto poly-(L-lysine) covered slides. Permeabilization was done with 0.1% (v/v) Triton X-100/PBS for 5 min at room temperature, followed by incubation with TdT buffer containing nucleotide mix (50 mM fluorescein-12-dUTP, 100 mM dATP, 10 mM Tris-HCl, 1 mM EDTA, pH 7.6) for 1 h at 37 °C. The samples were counterstained with  $10 \text{ mg mL}^{-1}$  propidium iodide and visualized under a fluorescence microscope. The percentage of parasites showing a clearly visible nuclear staining was determined in treated and control parasites by counting 100 cells in triplicate (blind-coded samples).

#### DNA fragmentation assay by agarose gel electrophoresis

DNA integrity after treatments was also evaluated by electrophoresis in agarose gel. Briefly,  $1 \times 10^7$  epimastigotes (in log phase) were collected, washed once and resuspended in PBS-3% glucose. Parasites were incubated with  $1 \mu\text{M}$  CsA for 1 h at room temperature prior to  $\text{H}_2\text{O}_2$  incubation. To induce oxidative stress, parasites were incubated with 5 mM  $\text{H}_2\text{O}_2$  for 3 h at room temperature. Total cellular DNA was isolated by a usual procedure and analysed by agarose gel electrophoresis. Pellets of  $10^7$  epimastigotes were treated with sarkosyl detergent lysis buffer (50 mM Tris, 10 mM EDTA, 0.5% w/v sodium-N-lauryl sarcosine, pH 7.5) and 7 mM proteinase K, vortexed and allowed to digest overnight at 50 °C. RNase A (10 mM) treatment followed for 1 h at 37 °C. The lysates were then extracted with phenol/chloroform/isoamyl alcohol (25:24:1) and centrifuged at  $16\,000 \text{ g}$  for 5 min. The upper phase was treated with 3 M sodium acetate and 100% ethanol overnight at  $-20 \text{ }^\circ\text{C}$ . The sample was centrifuged at  $16\,000 \text{ g}$  for 10 min and the pellet was washed with 0.5 ml of 70% ethanol. DNA was solubilized in Tris/EDTA (10/1 mM) buffer and spectrophotometrically quantitated at 260/280 nm. Total DNA was mixed with tracking dye and loaded on 1% agarose gels containing ethidium bromide. Gels were run for 2.5 h at 50 V.

#### mPTP-like opening

Calcein is a fluorescent dye of 1 kDa, which enables the study of mPTP. Thus, the calcein quenching method is a highly selective indicator of sustained mPTP opening *in situ* in mammals (Petronilli *et al.* 1998). In the presence of  $\text{CoCl}_2$ , fluorescence of

calcein is quenched from the cytosol and nuclear compartments, remaining in mitochondria. When the mPTP opens, calcein is lost from mitochondria.

To perform this assay with *T. cruzi* epimastigotes, a protocol adapted from a previous report was followed (Petronilli *et al.* 1998). Briefly,  $1 \times 10^6$  epimastigotes were loaded with  $2 \mu\text{M}$  of Calcein-AM and  $1 \text{ mM}$   $\text{CoCl}_2$  at  $37^\circ\text{C}$  for 30 min. After staining, parasites were incubated with  $1 \mu\text{M}$  CsA (when CyP inhibition was assayed) or with a combination of the other inhibitors of mPTP components ( $1 \text{ mM}$  ADP and  $10 \mu\text{M}$  Bongkrekic acid) for 1 h at room temperature. Oxidative stress was induced with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  for 45 min at room temperature and washed with PBS-3% glucose to remove the excess of Calcein-AM and  $\text{CoCl}_2$ . Samples were analysed using a FACSCalibur equipment as mentioned above, with appropriate excitation and emission filters for fluorescein. The decrease in fluorescence intensity indicated mPTP - like opening.

### Statistical analysis

Experiments were performed in duplicate or triplicate, as previously indicated, in two to four independent experiments. Statistical analysis of the *in vitro* parasite experiments was performed using Student's *t* test, for paired values. The values presented are the means  $\pm$  s.d. of 2 or more independent experiments. Values of  $P < 0.05$  were considered statistically significant.

## RESULTS

### CsA decreases Annexin V binding induced by $\text{H}_2\text{O}_2$

One of the first phenotypic changes observed during metazoan programmed cell death is the exposure of PS in the external surface of plasma membrane, which occurs after the cell has received the injure signal and will start to develop the cell death program. This event can be tested with the protein Annexin V, which binds PS with high affinity.

To investigate if PS exposure also occurs in *T. cruzi* after oxidative stress induction, parasites were treated with hydrogen peroxide ( $\text{H}_2\text{O}_2$ ). CsA was included to evaluate this event in the presence of a CyP inhibitor.

Epimastigotes in exponential growth phase were incubated with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  for different periods of time. Annexin V binding was detected after 30 min incubation with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  and increased up to 180 min (Fig. 1A), compared to the Control (untreated) parasites. When incubation with  $1 \mu\text{M}$  CsA for 1 h was done prior to oxidative stress induction, Annexin V binding was significantly decreased after 30 and 60 min of  $\text{H}_2\text{O}_2$  treatment. The absence of propidium iodide staining in the elapsed time indicated membrane integrity and ensured only the




Fig. 1. The CyP inhibitor CsA decreases Annexin V binding in *T. cruzi* epimastigotes after oxidative stress. (A) Untreated epimastigotes (control), parasites treated with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  for different periods of time or pre-incubated with  $1 \mu\text{M}$  CsA were stained with Annexin V-FITC to detect phosphatidylserine exposure in the external surface of the plasma membrane by flow cytometry. Means  $\pm$  s.d.,  $n = 4$  are shown, \*\*\*( $\text{H}_2\text{O}_2$ -30 min/ $\text{H}_2\text{O}_2$ -30 min+CsA), \*\*( $\text{H}_2\text{O}_2$ -60 min/ $\text{H}_2\text{O}_2$ -60 min+CsA),  $P < 0.05$ , Student's *t*-test). (B) Propidium iodide was also measured in parasites from (A). Absence of significant propidium iodide staining during the experiment indicates plasma membrane integrity, meaning that Annexin V-FITC is only binding to the exposed phosphatidylserine. Means  $\pm$  s.d.,  $n = 4$  are shown, \*\*\*( $\text{H}_2\text{O}_2$ -30 min/ $\text{H}_2\text{O}_2$ -30 min+CsA), \*\*( $\text{H}_2\text{O}_2$ -60 min/ $\text{H}_2\text{O}_2$ -60 min+CsA),  $P < 0.05$ , Student's *t*-test).

detection of Annexin V binding in the external membrane surface (Fig. 1B).

In this oxidative stress condition, *T. cruzi* epimastigotes responded by exposing PS in their external surface of the plasma membrane, as an early indicator of programmed cell death in this protozoan parasite. Pre-incubation with CsA significantly decreased the Annexin V signal, indicating that a CyP may be involved in the cascade of events that




Fig. 2. CsA decreases ROS production in *T. cruzi* epimastigotes after oxidative stress. Parasites were incubated with two different concentrations of H<sub>2</sub>O<sub>2</sub> (1 or 5 mM). Pre-incubation with 1 μM CsA was performed when inhibitory conditions were required. After oxidative stress induction, parasites were loaded with the fluorescent dye DCFH-DA. ROS were quantified by flow cytometry (means ± s.d., n = 3, P < 0.05, Student's *t*-test).

lead to PS exposure. However, after longer exposures with the inducer (180 min), the protective effect of CsA disappeared.

*Reactive oxygen species (ROS) production after treatment with H<sub>2</sub>O<sub>2</sub>*

In mammals, mPTP opening is triggered by a synergic combination of increased ROS production together with high levels of Ca<sup>2+</sup> in the mitochondrial matrix (Figueira *et al.* 2013).

To detect oxidative species in parasites after H<sub>2</sub>O<sub>2</sub> treatment and in the presence of CsA, the DCFH-DA dye was used, as detailed in the M&M section. Epimastigotes were treated with two different concentrations of H<sub>2</sub>O<sub>2</sub> for 1 h. Incubation with 1 μM CsA for 1 h was performed prior to H<sub>2</sub>O<sub>2</sub> treatment. Pre-incubation with CsA significantly decreased ROS production when parasites were incubated with both 1 mM H<sub>2</sub>O<sub>2</sub> and 5 mM H<sub>2</sub>O<sub>2</sub> (Fig. 2), showing that when the CyP inhibitor is present there is a protection against the damaging effects of H<sub>2</sub>O<sub>2</sub>. It is important to highlight that the concentration of CsA used in control parasites did not yield a significant amount of ROS *per se*.

*Cytochrome c translocation into the cytosol after oxidative stress is prevented by CsA*

Translocation of proteins such as cytochrome *c*, apoptosis-inducing factor and endonuclease G from the mitochondrial matrix into the cytoplasm is a critical event during metazoan programmed cell death. This change in localization of apoptosis-




Fig. 3. CsA prevents cytochrome *c* translocation into the cytosol of *T. cruzi* epimastigotes during oxidative stress. Oxidative stress was induced with 5 mM H<sub>2</sub>O<sub>2</sub> for 3 h at room temperature. Pre-incubation with 1 μM CsA for 1 h at the same temperature was performed when inhibitory conditions were required. Anti-cytochrome *c* antibodies were used in Western blots with the mitochondria-enriched subcellular fraction (M) and the cytosolic subcellular fraction (C). Membranes were stripped and re-incubated with antibodies against the mitochondrial protein VDAC to confirm the purity of the mitochondria-enriched fractions.

related proteins occurs as a consequence of the change in mitochondrial membrane permeability caused by increased ROS and Ca<sup>2+</sup> overload.

Having established that the induction with H<sub>2</sub>O<sub>2</sub> caused an increase in oxidative species in *T. cruzi*, we wanted to investigate if this would also result in translocation of these mitochondrial proteins in the parasite.

To evaluate cytochrome *c* localization after oxidative stress induction, parasites were either treated with 5 mM H<sub>2</sub>O<sub>2</sub> or not treated (control), or incubated with 1 μM CsA prior to H<sub>2</sub>O<sub>2</sub>. After treatment, parasites were fractionated using digitonin to obtain a mitochondria-enriched fraction and a cytosolic fraction, and their proteins were separated by SDS-PAGE and immuno-blotted with antibodies against cytochrome *c*. After oxidative stress induction with H<sub>2</sub>O<sub>2</sub>, cytochrome *c* was released from mitochondria and detected in the cytosolic fraction (Fig. 3, H<sub>2</sub>O<sub>2</sub>). However, when parasites were pre-treated with CsA, cytochrome *c* remained mostly in the mitochondria-enriched fraction (Fig. 3, H<sub>2</sub>O<sub>2</sub> CsA). To confirm the purity of the mitochondria-enriched fractions, these blots were stripped and re-probed for VDAC, protein size ~30 kDa (TriTrypsDB annotated as TcCLB.509141.40) which is located to the outer mitochondrial membrane. These results show that the presence of CsA prevented the release of cytochrome *c* from mitochondria in response to oxidative stress, in agreement with the results obtained for ROS production, suggesting that a CyP is involved in these mitochondria-related events.

*CsA inhibits DNA degradation induced by H<sub>2</sub>O<sub>2</sub>*

A later step in the cascade of events leading to programmed cell death is marked by changes to the




Fig. 4. CsA decreases DNA damage in *T. cruzi* epimastigotes during oxidative stress. (A) *T. cruzi* epimastigotes were subjected to oxidative stress with 5 mM H<sub>2</sub>O<sub>2</sub>. Parasites with positive nucleus staining were counted as TUNEL positive. The percentage is expressed as TUNEL-positive/total parasites. Significant differences were observed when parasites were treated with CsA prior to incubation with 5 mM H<sub>2</sub>O<sub>2</sub> (means  $\pm$  s.d.,  $n = 3$ ,  $P < 0.05$ , Student's *t*-test). (B) After inducing oxidative stress with 5 mM H<sub>2</sub>O<sub>2</sub> genomic DNA was extracted with phenol/chloroform, electrophoresed in 1% agarose gel and visualized by ethidium bromide. DNA degradation observed in 5 mM H<sub>2</sub>O<sub>2</sub>-treated parasites (lane 3) was prevented by pre-incubation with 1  $\mu$ M CsA (lane 4), compared to untreated parasites (lane 1) and 1  $\mu$ M CsA-treated parasites (lane 2).

nucleus. DNA degradation, chromatin condensation and changes in nuclear morphology are often considered the best indicators of an apoptotic process, being DNA degradation probably the most frequent marker of apoptotic death used in metazoans. DNA fragmentation revealed by the presence of strand breaks is considered to be the gold standard for identification of apoptotic cells (Jiménez-Ruiz *et al.* 2010).

To further characterize the changes occurring in H<sub>2</sub>O<sub>2</sub>-treated *T. cruzi* epimastigotes, *in situ* detection of DNA fragments by TUNEL was performed to detect DNA 3' free ends after breakage. Figure 4A shows the percentage of cells with positive staining in control (C and CsA) or treatment (H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>+CsA) conditions. Epimastigotes treated with 5 mM H<sub>2</sub>O<sub>2</sub> showed 67% of TUNEL-positive cells vs 42% when parasites were pre-incubated with CsA.

To evaluate genomic DNA degradation, we used electrophoresis in agarose gel of total epimastigotes DNA. A massive degradation of genomic DNA was visualized in 5 mM H<sub>2</sub>O<sub>2</sub>-treated epimastigotes (Fig. 4B, lane 3). This pattern was not observed when parasites were incubated with CsA prior to oxidative stress induction with H<sub>2</sub>O<sub>2</sub> (Fig. 4B, lane 4).

The degraded DNA pattern observed for trypanosomatids showed some degree of smearing and although it is not as clear as the typical expected DNA ladder from metazoan DNA, it is consistent with other reports in protozoan parasites (Jiménez-Ruiz *et al.* 2010).

These results taken together showed that, when CsA is present prior to the addition of the oxidative stress inducer, the genetic material of the parasites can be partially protected from the damaging effect of H<sub>2</sub>O<sub>2</sub>. Again, this is another feature that can be partially inhibited by CsA, showing indirectly that a CyP is involved in programmed cell death in the parasite.

#### *mPTP-like opening occurs in Trypanosoma cruzi and responds to mammalian mPTP blockers*

The mPTP is a crucial component of cell death events in mammals. Although its molecular identity remains under debate, it is well established that it is the mPTP opening in mitochondrial membrane which leads to the release of apoptotic-related proteins and in consequence, the completion of cell death program.

To investigate whether the same mechanism occurs in trypanosomatids, we used the highly selective indicator of sustained mPTP opening *in situ*, the fluorescent dye calcein. When Calcein-AM is co-loaded with CoCl<sub>2</sub>, calcein fluorescence is quenched in both cytosolic and nuclear compartments, allowing the detection of calcein fluorescence in mitochondria. As we found no previous reports about the calcein quenching method in protozoan parasites, we first attempted to establish the correct loading dye in parasites. *T. cruzi* epimastigotes were loaded with calcein and the quencher CoCl<sub>2</sub>. The proper localization of calcein in the parasite mitochondrion was confirmed by fluorescence microscopy of live cells (data not shown). After finding that epimastigotes could be loaded with the fluorescent probe, we evaluated if calcein was released from parasite mitochondria in response to oxidative stress induction. After 15 min in the presence of 5 mM H<sub>2</sub>O<sub>2</sub>, almost 50% of the calcein was released from parasite mitochondria, and remained constant up to 90 min in oxidative stress conditions (Fig. 5). Having established the appropriate conditions to study calcein release from parasite mitochondria, we evaluated the effect of CsA and other inhibitors described for mammalian mPTP




Fig. 5. To evaluate mPTP-like opening in *T. cruzi* epimastigotes, the calcein quenching method was used. Parasites were loaded with  $2 \mu\text{M}$  calcein-AM and  $1 \text{ mM}$   $\text{CoCl}_2$  to detect the fluorescent dye retained within the mitochondria and treated with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  for 90 min. At the indicated time points, fluorescence was measured by flow cytometry. Data were analysed with Cyflogic software. Results are representative of two independent experiments.

components: Bongkreikic acid and ADP, which inhibit ANT. Figure 6 shows the fluorescent calcein retained from parasites treated with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  for 45 min (H) or pre-incubated either with CsA (H CsA) or a combination of CsA + Bongkreikic acid + ADP (H CsA-ADP-BKA) compared to untreated parasites (control). When parasites were treated with  $\text{H}_2\text{O}_2$ ,  $\sim 45\%$  of calcein fluorescence was retained in mitochondria (Fig. 6, column H). However, parasites pre-treated with CsA showed increased calcein retention (60%). When the combination of mPTP inhibitors was assayed, parasites were able to retain a higher amount (86%, Fig. 6).

These results suggest the existence of a mPTP-like pore in *T. cruzi* parasites, which responds to inhibitors described for mammalian mPTP components. In particular, *T. cruzi* mPTP-like appears to respond to CsA, which is one of the best characterized mPTP inhibitors in mammals by binding the mitochondrial CyPD.

## DISCUSSION

In this study, we report several programmed cell death features in the protozoan parasite *Trypanosoma cruzi* that were sensitive to the inhibitors of the mammalian mPTP. It has been described that mPTP opening in mammals leads to cell death (Vaseva *et al.* 2012), and whether cell death in protozoan parasites is programmed (Ameisen *et al.* 1995) or incidental (Proto *et al.* 2013) the molecular mechanisms and effectors remain unknown. Several cell death inducers have been tested in different protozoan parasites: chloroquine and staurosporine were assayed in *Plasmodium falciparum* (Meslin *et al.*




Fig. 6. mPTP-like opening occurs in *T. cruzi* and responds to mammalian mPTP blockers. After having established the proper conditions for the calcein quenching method in our model, the effect of CsA or a combination of known mPTP blockers were evaluated. Parasites were untreated (control) or treated with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$  (H). When inhibition of mPTP was assayed, pre-incubation with CsA (H CsA) or with a combination of the inhibitors CsA + ADP + Bongkreikic acid (H CsA-ADP-BKA) was performed. (means  $\pm$  s.d.,  $n = 3$ , \*\*( $\text{H}_2\text{O}_2/\text{H}_2\text{O}_2$  CsA), \*\*\*( $\text{H}_2\text{O}_2/\text{H}_2\text{O}_2$  CsA-BKA-ADP),  $P < 0.05$ , Student's *t*-test).

2007), prostaglandins in *T. brucei* (Figarella *et al.* 2005, 2006), miltefosine in *Leishmania donovani* (Paris *et al.* 2004) and fresh human serum (Piacenza *et al.* 2001) and starvation (Jimenez *et al.* 2008) in *T. cruzi*.

Oxidative stress with  $\text{H}_2\text{O}_2$  has also been used by other research groups and, in this work, in the same range of concentrations, from 1 to 5 mM, as an inducer of cell death in *T. cruzi* (Das *et al.* 2001; Fernández Villamil *et al.* 2008; Jiménez-Ruiz *et al.* 2010).

PS exposure, which is an early event in programmed cell death in mammals, had been already reported to occur in *T. cruzi* (Piacenza *et al.* 2007; Jimenez *et al.* 2008) and in *T. brucei* (Figarella *et al.* 2005) as shown by Annexin V binding. Although there may be other PS-containing phospholipids, Annexin V binds PS with high affinity in *T. cruzi*. After oxidative stress induction with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$ , we demonstrated that Annexin V binding occurred in *T. cruzi* epimastigotes after 30 min exposure to  $\text{H}_2\text{O}_2$  and continued up to 180 min. Interestingly, pre-incubation with CsA partially decreased Annexin V binding in  $\text{H}_2\text{O}_2$ -treated parasites.

Regarding mitochondrial function, cytochrome *c* was detected in parasite cytosolic fractions after 30 min treatment with  $5 \text{ mM}$   $\text{H}_2\text{O}_2$ , consistent with a

previous report (Piacenza *et al.* 2007). In our model, pre-incubation for 60 min with 1  $\mu\text{M}$  CsA partially prevented the translocation of cytochrome *c* into the parasite cytosolic fraction.

PARP proteolytic processing is another feature of programmed cell death frequently evidenced in metazoans. PARP is a family of abundant nuclear proteins, some of which are involved in the DNA base excision repair system. In metazoans, the specific cleavage of PARP-1 has been used extensively as a biochemical marker of apoptosis (Jiménez-Ruiz *et al.* 2010). In *T. cruzi*, the counterpart *Tc*PARP has been characterized (Fernández Villamil *et al.* 2008). After inducing oxidative stress with 5 mM  $\text{H}_2\text{O}_2$ , we were also able to detect that *Tc*PARP was cleaved. This cleavage was decreased by pre-incubation with CsA (data not shown).

During apoptosis, as repair mechanisms are inhibited, DNA degradation will occur by activated nucleases that migrate to the nucleus to degrade DNA.

In *T. cruzi* epimastigotes, DNA degradation had been reported after cell death induction with 20% (V/V) fresh human serum. We confirmed by both TUNEL and electrophoresis in agarose gel that degradation occurred after treatment with 5 mM  $\text{H}_2\text{O}_2$ . This degradation pattern was not observed when epimastigotes were pre-incubated with 1  $\mu\text{M}$  CsA, probably due to the inhibition of previous steps of the cell death cascade. This phenomenon has also been reported in other protozoan parasites as *Leishmania donovani* under oxidative stress induction with 3 mM  $\text{H}_2\text{O}_2$  (Das *et al.* 2001). It is important to highlight that, although an internucleosomal ladder pattern is the common feature in metazoan programmed cell death, this is not the case in protozoan parasites (Jiménez-Ruiz *et al.* 2010), where a smear pattern is usually observed.

It is important to point out that, in all the experiments performed in this study, a protective effect was achieved with pre-incubation with 1  $\mu\text{M}$  CsA. When we added some other inhibitors such as ADP and Bongkreikic acid, which block the mammalian mPTP related protein ANT, we observed an even higher protective effect against cell death in the parasite. This potential effect was observed using the calcein/ $\text{CoCl}_2$  quenching method, which is a highly selective indicator of mPTP opening in mammals (Petronilli *et al.* 1998; Kroemer *et al.* 2007; Vaseva *et al.* 2012). After 15 min of oxidative stress induction in *T. cruzi* epimastigotes, most calcein was released from mitochondria, but pre-incubation with CsA retained up to ~60% calcein inside the parasite. When inhibitors of other mammalian mPTP components were included (1 mM ADP+10  $\mu\text{M}$  Bongkreikic acid), the calcein retained in parasite mitochondria increased up to 86%,

evidencing a drug additive effect in inhibiting a probable mPTP-like opening present in *T. cruzi*.

When we searched similarities with proteins that participate in the formation of the mammalian mPTP in the *T. cruzi* genome databases, we found some putative predicted proteins in the parasite with high similarity to mammalian ANT and VDAC, among others. The mPTP protein components have not been fully described in mammals and, to our knowledge, in any protozoan parasite.

Although CsA may exert some inhibitory effect on other Cyclophilins, it is well known that it has a specific cell death protective effect as a mPTP blocker by binding the mitochondrial CyPD (Connern and Halestrap, 1994; Friberg *et al.* 1998). CsA has thus become the usual molecular tool to analyse the role of CyPD in the mPTP in different models (Vaseva *et al.* 2012; Kim *et al.* 2014), while more specific CyPD inhibitors are still under development (Malouitre *et al.* 2009).

We have previously described the CyP gene family in *T. cruzi* which consists of 15 members, while four CyPs of 19, 22, 28 and 40 kDa have been isolated as CsA-binding proteins (Potenza *et al.* 2006; Bua *et al.* 2008).

As *T. cruzi* has a CyPD orthologue in its genome, current efforts are being made to confirm the role of a mitochondrial parasite CyP in these mPTP-related events. Ongoing experiments are expected to produce parasites overexpressing a mitochondrial CyP, which will allow to specifically address its responsibility in the programmed cell death features described in this work.

To our knowledge, the results presented in this article describe for the first time a mPTP-like opening that would be involved in a unicellular parasite cell death. Our findings also encourage further biochemical studies to determine the specific role of a *T. cruzi* mitochondrial CyP homologous to mammalian CyPD, considering that it has not been described in any unicellular parasite yet.

#### ACKNOWLEDGEMENTS

We are grateful for the valuable help of Dr Silvia Moreno and her committed mentoring to P.L.B. at the CTEGD, University of Georgia. P.L.B.'s research visit was financed by US National Institutes of Health Grant D43TW00888. We thank Dr Alicia Graciela Fuchs and CAECIHS for spectrofluorometer facilities and Claudia Nose for technical support.

#### FINANCIAL SUPPORT

This work was supported by ANLIS Carlos G. Malbrán and Focanlis 2009, PICTO-ANLIS 00136/11, by US National Institutes of Health Grant D43TW00888 and by National Scientific and Technical Research Council (CONICET) Grant PIP 0317/10.



## REFERENCES

- Ameisen, J. C., Idziorek, T., Billaut-Mulot, O., Loyens, M., Tissier, J. P. and Potentier, A. (1995). Apoptosis in a unicellular eukaryote (*Trypanosoma cruzi*): implications for the evolutionary origin and role of programmed cell death in the control of cell proliferation, differentiation and survival. *Cell Death and Differentiation* **2**, 285–300.
- Bua, J., Fichera, L. E., Fuchs, A. G., Potenza, M., Dubin, M., Wenger, R. O., Moretti, G., Scabone, C. M. and Ruiz, A. M. (2008). Anti-*Trypanosoma cruzi* effects of cyclosporin A derivatives: possible role of a P-glycoprotein and parasite cyclophilins. *Parasitology* **135**, 217–228.
- Cannata, J. J. and Cazzulo, J. J. (1984). Glycosomal and mitochondrial malate dehydrogenases in epimastigotes of *Trypanosoma cruzi*. *Molecular and Biochemical Parasitology* **11**, 37–49.
- Connern, C. P. and Halestrap, A. P. (1994). Recruitment of mitochondrial cyclophilin to the mitochondrial inner membrane under conditions of oxidative stress that enhance the opening of a calcium-sensitive non-specific channel. *Biochemical Journal* **302**, 321–324.
- Crompton, M., Virji, S., Doyle, V., Johnson, N. and Ward, J. M. (1999). The mitochondrial permeability transition pore. *Biochemical Society Symposia* **66**, 167–179.
- Das, M., Mukherjee, S. B. and Shaha, C. (2001). Hydrogen peroxide induces apoptosis-like death in *Leishmania donovani* promastigotes. *Journal of Cell Science* **114**(Pt 13), 2461–2469.
- Duszenko, M., Figarella, K., Macleod, E. T. and Welburn, S. C. (2006). Death of a trypanosome: a selfish altruism. *Trends in Parasitology* **22**, 536–542.
- Fernández Villamil, S. H., Baltanás, R., Alonso, G. D., Vilchez Larrea, S. C., Torres, H. N. and Flawiá, M. M. (2008). TcPARP: a DNA damage-dependent poly(ADP-ribose) polymerase from *Trypanosoma cruzi*. *International Journal for Parasitology* **38**, 277–287.
- Figarella, K., Rawer, M., Uzcategui, N. L., Kubata, B. K., Lauber, K. and Madeo, F. (2005). Prostaglandin D2 induces programmed cell death in *Trypanosoma brucei* bloodstream form. *Cell Death and Differentiation* **12**, 335–346.
- Figarella, K., Uzcategui, N. L., Beck, A., Schoenfeld, C., Kubata, B. K. and Lang, F. (2006). Prostaglandin-induced programmed cell death in *Trypanosoma brucei* involves oxidative stress. *Cell Death and Differentiation* **13**, 1802–1814.
- Figueira, T. R., Barros, M. H., Camargo, A. A., Castilho, R. F., Ferreira, J. C., Kowaltowski, A. J., Sluse, F. E., Souza-Pinto, N. C. and Vercesi, A. E. (2013). Mitochondria as a source of reactive oxygen and nitrogen species: from molecular mechanisms to human health. *Antioxidants and Redox Signaling* **18**, 2029–2074.
- Friberg, H., Ferrand-Drake, M., Bengtsson, F., Halestrap, A. P. and Wieloch, T. (1998). Cyclosporin A, but not FK 506, protects mitochondria and neurons against hypoglycemic damage and implicates the mitochondrial permeability transition in cell death. *Journal of Neuroscience* **18**, 5151–5159.
- Gannavaram, S., Vedvyas, C. and Debrabant, A. (2008). Conservation of the pro-apoptotic nuclease activity of endonuclease G in unicellular trypanosomatid parasites. *Journal of Cell Science* **121**, 99–109.
- Giorgio, V., von Stockum, S., Antoniel, M., Fabbro, A., Fogolari, F., Forte, M., Glick, G. D., Petronilli, V., Zoratti, M., Szabó, I., Lippe, G. and Bernardi, P. (2013). Dimers of mitochondrial ATP synthase form the permeability transition pore. *Proceedings of the National Academy of Sciences* **110**, 5887–5892.
- Halestrap, A. P. (2014). The C ring of the F1Fo ATP synthase forms the mitochondrial permeability transition pore: a critical appraisal. *Frontiers in Oncology* **25**, 234.
- Halestrap, A. P. and Davidson, A. M. (1990). Inhibition of Ca<sup>2+</sup>-induced large-amplitude swelling of liver and heart mitochondria by cyclosporin is probably caused by the inhibitor binding to mitochondrial-matrix peptidyl-prolyl cis-trans isomerase and preventing it interacting with the adenine nucleotide translocase. *The Biochemical Journal* **268**, 153–160.
- Jimenez, V., Paredes, R., Sosa, M. A. and Galanti, N. (2008). Natural programmed cell death in *T. cruzi* epimastigotes maintained in axenic cultures. *Journal of Cellular Biochemistry* **105**, 688–698.
- Jiménez-Ruiz, A., Alzate, J. F., Macleod, E. T., Lüder, C. G., Fasel, N. and Hurd, H. (2010). Apoptotic markers in protozoan parasites. *Parasites and Vectors* **3**, 104.
- Kim, S. Y., Shim, M. S., Kim, K. Y., Weinreb, R. N., Wheeler, L. A. and Ju, W. K. (2014). Inhibition of cyclophilin D by cyclosporin A promotes retinal ganglion cell survival by preventing mitochondrial alteration in ischemic injury. *Cell Death and Disease* **5**, e1105.
- Kroemer, G., Galluzzi, L. and Brenner, C. (2007). Mitochondrial membrane permeabilization in cell death. *Physiological Reviews* **87**, 99–163.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Malouitre, S., Dube, H., Selwood, D. and Crompton, M. (2009). Mitochondrial targeting of cyclosporin A enables selective inhibition of cyclophilin-D and enhanced cytoprotection after glucose and oxygen deprivation. *Biochemical Journal* **425**, 137–148.
- Meslin, B., Barnadas, C., Boni, V., Latour, C., De Monbrison, F. and Kaiser, K. (2007). Features of apoptosis in *Plasmodium falciparum* erythrocytic stage through a putative role of PfMCA1 metacaspase-like protein. *Journal of Infectious Diseases* **195**, 1852–1859.
- Paris, C., Loiseau, P. M., Bories, C. and Bréard, J. (2004). Miltefosine induces apoptosis-like death in *Leishmania donovani* promastigotes. *Antimicrobial Agents and Chemotherapy* **48**, 852–859.
- Petronilli, V., Miotto, G., Canton, M., Colonna, R., Bernardi, P. and Di Lisa, F. (1998). Imaging the mitochondrial permeability transition pore in intact cells. *Biofactors* **8**, 263–272.
- Piacenza, L., Peluffo, G. and Radi, R. (2001). L-arginine-dependent suppression of apoptosis in *Trypanosoma cruzi*: contribution of the nitric oxide and polyamine pathways. *Proceedings of the National Academy of Sciences of USA* **98**, 7301–7306.
- Piacenza, L., Irigoín, F., Alvarez, M. N., Peluffo, G., Taylor, M. C. and Kelly, J. M. (2007). Mitochondrial superoxide radicals mediate programmed cell death in *Trypanosoma cruzi*: cytoprotective action of mitochondrial iron superoxide dismutase overexpression. *Biochemical Journal* **403**, 323–334.
- Potenza, M., Galat, A., Minning, T. A., Ruiz, A. M., Duran, R., Tarleton, R. L. and Bua, J. (2006). Analysis of the *Trypanosoma cruzi* cyclophilin gene family and identification of Cyclosporin A binding proteins. *Parasitology* **132**, 867–882.
- Proto, W. R., Coombs, G. H. and Mottram, J. C. (2013). Cell death in parasitic protozoa: regulated or incidental? *Nature Reviews Microbiology* **11**, 58–66.
- Van Heerde, W. L., Poort, S., Van't Veer, C., Reutelingsperger, C. P. and de Groot, P. G. (1994). Binding of recombinant annexin V to endothelial cells: effect of annexin V binding on endothelial-cell-mediated thrombin formation. *Biochemical Journal* **302**(Pt 1), 305–312.
- Vaseva, A. V., Marchenko, N. D., Ji, K., Tsirka, S. E., Holzmann, S. and Moll, U. M. (2012). p53 opens the mitochondrial permeability transition pore to trigger necrosis. *Cell* **149**, 1536–1548.
- WHO (2014) <http://www.who.int/mediacentre/factsheets/fs340/en/>.