

Main Article

Dr D Chaukar takes responsibility for the integrity of the content of the paper

Cite this article: Sharma S, Chaukar DA, Bal M, D'Cruz AK. Is routine neck dissection warranted at salvage laryngectomy? *J Laryngol Otol* 2021;**135**:785–790. <https://doi.org/10.1017/S0022215121001808>

Accepted: 4 January 2021
First published online: 9 July 2021

Key words:

Laryngectomy; Salvage Therapy; Neck Dissection; Neoplasm, Residual

Author for correspondence:

Dr Devendra A Chaukar,
Department of Head Neck Services,
Tata Memorial Hospital, Dr E Borges Marg,
Parel, Mumbai 400012, India
E-mail: dchaukar@gmail.com

Is routine neck dissection warranted at salvage laryngectomy?

S Sharma¹, D A Chaukar², M Bal³ and A K D'Cruz⁴

¹Department of Head Neck Surgical Oncology, Narayana Superspeciality Hospital, Gurugram, Departments of ²Head Neck Services, and ³Pathology, Tata Memorial Hospital, Mumbai and ⁴Department of Oncology, Apollo Hospital, Mumbai, India

Abstract

Background. There is controversy regarding management of the neck at salvage laryngectomy. The aim of this study was to perform an analysis to determine the incidence of occult node positivity in this group and analyse factors affecting it.

Method. A retrospective analysis of 171 patients who underwent salvage total laryngectomy between 2000 and 2015 for recurrent or residual disease following definitive non-surgical treatment and were clinico-radiologically node negative at the time salvage laryngectomy was carried out.

Results. A total of 171 patients with laryngeal or hypopharyngeal cancers underwent concurrent neck dissection at laryngectomy. There were 162 patients (94.7 per cent) who underwent bilateral neck dissection, and 9 patients (5.3 per cent) who underwent ipsilateral neck dissection. The occult lateral nodal metastasis rate was 10.5 per cent. Of various factors, initial node positive disease was the only factor predicting occult metastasis on univariable and multivariable analysis ($p = 0.001$).

Conclusion. Risk of occult metastasis is high in patients who have node positive disease before starting radiotherapy. This group should be offered elective neck dissection.

Introduction

The Veterans Affairs, European Organization for Research and Treatment of Cancer and Radiation Therapy Oncology Group 91-11 trials established non-surgical organ preservation with chemo-radiotherapy (RT) as the standard of care for locally advanced laryngeal or hypopharyngeal cancers, with surgical salvage reserved for those with recurrent or residual disease.^{1–3} Management of the clinico-radiologically node negative neck at salvage laryngectomy has been a matter of debate. Some advocate elective neck dissection, whereas others favour a wait and watch policy.^{4–7} This is primarily because neck dissection in salvage settings is associated with increased post-operative morbidity.^{8–11} On the other hand, a wait and watch approach can result in recurrences becoming unsalvageable. The risks must be balanced against potential benefits when a decision is made to address the neck in the salvage setting. The literature on this is sparse, and most studies have a small sample size and are heterogeneous. A recent systemic review addressing this issue failed to come up with concrete recommendations because of several limitations.¹² The National Comprehensive Cancer Network and other commonly used guidelines are also ambiguous on this topic.^{13,14}

We retrospectively analysed the occult metastasis rate and attempted to identify risk factors for occult metastasis in patients who were clinico-radiologically node negative following initial non-surgical organ preservation so that the decision for neck dissection could be made at the time of salvage laryngectomy.

Materials and methods

We retrospectively analysed patients who underwent salvage total laryngectomy following definitive RT or chemo-RT between September 2000 and October 2015 at a tertiary care centre. Of 1130 total laryngectomy procedures performed during the study period, 236 patients underwent salvage total laryngectomy. Of these 236 patients, 203 patients were clinico-radiologically node negative, and 33 patients had node positive disease.

Inclusion criteria were: (1) patients requiring salvage total laryngectomy for biopsy-proven recurrent or residual squamous cancer; (2) patients who were clinico-radiologically node negative at the time of surgery; and (3) patients treated with ipsilateral or bilateral neck dissection.

Patients who underwent salvage laryngectomy for dysfunctional larynx ($n = 7$), non-squamous pathology ($n = 2$) or who had a non-viable tumour on final histopathology ($n = 5$), and patients where the neck was not dissected ($n = 18$) were excluded from analysis. Our study cohort consisted of 171 patients.

Patient data were extracted from the hospital information system and electronic medical records system, which is updated at each follow-up visit ensuring effective and

authentic data capture. The demographical, clinical, surgical, histological, post-operative and follow-up details were recorded.

Radiotherapy

As per the institutional policy, patients with T₁ or T₂ node negative glottic disease received narrow field radiation. Glottic tumours that were stage T₃ or T₄ or node positive and supraglottic or hypopharyngeal primaries received irradiation to primary as well as to bilateral nodal basins.

Imaging

All patients underwent routine radiological screening for the neck using contrast-enhanced computed tomography (CT) scans or positron-emission tomography (PET)-CT. The majority of the patients operated on before 2009 underwent CT scans. Patients operated on thereafter were imaged with either CT or PET-CT.

Surgery

All patients underwent wide field total laryngectomy for residual or recurrent disease. Primary closure of the neopharynx was performed if unstretched remnant mucosa was more than 2.5 cm. The decision to put pectoralis major myofascial flap over the suture line was taken by the treating surgeon. The technique of harvesting pectoralis major myofascial flap was described in our previous publication.¹⁵

If the remnant unstretched pharyngeal mucosa was found to be less than 2.5 cm, it was augmented with pectoralis major myocutaneous flap or free flap. All patients underwent unilateral or bilateral elective lateral neck dissection encompassing levels II–IV. Any patient with occult metastasis, which was confirmed by intra-operative frozen section, underwent level V dissection. Post-operative complication details were captured from the charts and electronic medical records.

Histopathology

Lymph nodes from each lymph node level were labelled and submitted in separate containers to the pathology department. All specimens were evaluated by specialist head and neck pathologists, employing a standardised grossing and reporting protocol. Briefly, all lymph nodes retrieved from each level were submitted separately and examined microscopically. Lymph nodes less than 1 cm were embedded *in toto*, and a single section was studied. Nodes ranging from 1 to 2.5 cm in size were bisected along the long axis, and both halves were submitted for examination. Nodes larger than 2.5 cm were sectioned serially, and a minimum of 2 sections were studied.

Follow up

Patients were followed up at three monthly intervals for first two years, six monthly intervals until five years and annually thereafter. Patients were assessed clinically during each follow-up visit. Imaging was ordered in cases of clinical suspicion of disease. Recurrence was confirmed clinico-radiologically, and a histological diagnosis was established. Second primary or recurrence during a follow-up visit was captured and labelled as local, regional, loco-regional or distant.

Statistical analysis

Statistical analysis was performed using SPSS® statistical software (version 24). Qualitative data was analysed using the chi-square test and Fisher's exact test for univariable analysis. Regression analysis using the binary logistic forward regression method was performed for multivariable analysis. Survival analysis was done using the Kaplan–Meier method. Comparisons in survival were analysed using log rank test. Two-sided *p*-values less than 0.05 were considered significant.

Results

The study cohort comprised 171 patients. There were 158 men and 13 women with a median age of 57 years (range, 26–89 years). At the initial presentation, 112 patients (65.5 per cent) had glottic disease, 16 patients (9.4 per cent) had supraglottic disease and 43 patients (25.1 per cent) had hypopharyngeal disease.

About half of the patients (95 of 171; 55.6 per cent) were early stage (T₁ or T₂), 72 patients (42.1 per cent) were stage T₃ and 4 patients were stage T₄. A total of 150 patients were N₀, 6 patients (3.5 per cent) were N₁ and 6 patients (3.5 per cent) were N₂. Information on nodal stage was missing in nine patients.

A total of 100 patients (58.5 per cent) received RT alone, and 71 patients (41.5 per cent) were treated with chemo-RT (Table 1). All patients who were included in this study were clinico-radiologically node negative at the time of salvage total laryngectomy. Primary pharyngeal closure was achieved in 130 patients. In 31 of these patients, the suture line was reinforced with onlay pectoralis major myofascial flap. Thirty-two patients (18.7 per cent) required pharyngeal augmentation using patch pectoralis major myocutaneous flap, free flap reconstruction or gastric pull up. In 9 patients (5.3 per cent), skin of the neck had to be excised and reconstructed with spiral pectoralis major myocutaneous flap. Surgical details are summarised in Table 2.

Neck dissection details

A total of 162 patients (94.7 per cent) underwent bilateral neck dissection, and 9 patients (5.3 per cent) underwent ipsilateral neck dissection. All patients underwent level II–IV lymph node clearance and level V was addressed in 10 patients (5.8 per cent) because of metastatic disease in the lateral neck as reported on frozen section.

Occult node positivity

A total of 18 patients (10.5 per cent) had occult nodal disease in the lateral neck on histopathology examination. Bilateral metastatic spread to the neck was present in only 1 patient (0.5 per cent). This patient had extensive local disease involving bilateral transglottic areas, both pyriform sinuses along with soft tissue extension to thyroid gland.

The incidence of nodal metastatic disease was highest for hypopharyngeal lesions (6 of 43; 13.9 per cent) followed by glottic lesions with transglottic spread (11 of 112; 9.8 per cent) and supraglottic lesions (1 of 16; 6.3 per cent); however, this difference was not statistically significant (*p* = 0.6). Early T-stage at recurrence (T₁ or T₂) had lower incidence of occult metastasis when compared with advanced T-stage at recurrence (T₃ or T₄), with 5.2 per cent versus 12 per cent, respectively, although this difference was not statistically significant (*p* = 0.3).

Table 1. Clinical and demographic details

Parameter	Value*
Age (median; years)	57
Gender (n (%))	
– Male	158 (92.4)
– Female	13 (7.6)
Prior treatment type (n (%))	
– Radiotherapy	100 (58.5)
– Chemoradiotherapy	71 (41.5)
Site (n (%))	
– Glottis	112 (65.5)
– Supraglottis	16 (9.4)
– Hypopharynx	43 (25.1)
Presence of tracheostomy at recurrence (n (%))	
– Yes	48 (28.1)
– No	123 (71.9)
Tumour stage at initial presentation (n (%))	
– Early (T ₁ or T ₂)	95 (55.6)
– Advanced (T ₃ or T ₄)	76 (44.4)
Nodal stage at initial presentation [†] (n (%))	
– N ₀	150 (87.7)
– N ₁	6 (3.5)
– N ₂	6 (3.5)
Imaging at recurrence (n (%))	
– Computed tomography scan	115 (67.3)
– Positron emission tomography-computed tomography	56 (32.7)
Tumour stage at recurrence (n (%))	
– Early (T ₁ or T ₂)	38 (22.2)
– Advanced (T ₃ or T ₄)	133 (77.8)

*n = 171; [†]data missing in nine patients

We had 12 patients who were node positive at initial treatment (RT or chemo-RT). On comparing the occult node positivity rate in these 12 patients with remaining patients who were N₀ at the start of initial treatment, we found that the incidence of occult node positivity was significantly higher in these 12 patients (42 per cent versus 8 per cent, respectively; $p = 0.004$). This difference was significant on both univariable and multivariable analysis.

We analysed the following factors to predict occult node positivity: age, tumour stage before definitive treatment (RT or chemo-RT) and at recurrence, nodal status before initial treatment, tumour site, type of treatment received (RT versus chemo-radiation), time to recurrence, imaging type, and the presence of tracheostomy at salvage laryngectomy (Table 3). All these factors were analysed by cox regression analysis as covariates. Of these, node-positive disease prior to definitive treatment was the only significant factor for occult metastasis on univariable ($p = 0.004$) and multivariable analysis ($p = 0.001$).

Accuracy of imaging

A total of 115 patients (67.3 per cent) were imaged using contrast-enhanced CT, and 56 patients (32.7 per cent)

Table 2. Surgical details

Parameter	Value* (n (%))
Surgery type	
– Salvage total laryngectomy	99 (57.9)
– Salvage total laryngectomy + pectoralis major myofascial flap	31 (18.1)
– Salvage total laryngectomy + spiral pectoralis major myocutaneous flap	9 (5.3)
– Salvage total laryngectomy + pharyngeal reconstruction	32 (18.7)
Neck dissection	
– Unilateral	9 (5.3)
– Bilateral	162 (94.7)
Neck dissection levels	
– II–IV	161 (94.2)
– II–V	10 (5.8)
Occult node positive rate on histopathology	
– Node negative	153 (89.5)
– Node positive	18 (10.5)

*n = 171

underwent PET-CT. The negative predictive value of PET-CT and contrast-enhanced CT in detecting occult neck metastasis was 91 per cent and 88.7 per cent, respectively.

Post-operative complications

A total of 84 patients (49.1 per cent) had complications. The most common complication was pharyngocutaneous fistula which was present in 74 patients (43.3 per cent). The majority of the patients had minor fistula which healed with conservative management. Fourteen patients (8.2 per cent) required surgical intervention to close the fistula.

Recurrence and survival

Forty patients (23.3 per cent) developed recurrence, and 6 patients (3.5 per cent) had second primaries over a median follow up of 24 months. Twenty-one patients (12.3 per cent) developed isolated local recurrences. Three patients (1.7 per cent) developed isolated regional failure. Two patients (1.2 per cent) experienced simultaneous loco-regional relapse. Fourteen patients (8.1 per cent) had metastatic disease either alone or along with local or regional recurrences or both.

Seventy per cent of patients were recurrence free at the end of five years (Figure 1). We compared recurrence-free survival in patients who had occult neck positive disease on histopathology versus those who did not. There was no difference in the recurrence-free survival rate at 5 years between the 2 groups (70.6 per cent versus 68.2 per cent, respectively; log rank test $p = 0.5$; Figure 2).


Discussion

The role of elective neck dissection at salvage laryngectomy for recurrent laryngeal cancer remains controversial. Salvage laryngectomy itself is a technically challenging procedure and is associated with increased post-operative complications,

Table 3. Factors predicting occult nodal metastasis (univariable analysis)

Risk factor	Pathological node negative* (n (%))	Pathological node positive† (n (%))	P-value
Tumour stage at initial presentation			
– T ₁ or T ₂	87 (56.9)	8 (44.4)	0.3
– T ₃ or T ₄	66 (43.1)	10 (55.6)	
Tumour stage at recurrence			
– T ₁ or T ₂	36 (23.5)	2 (11.1)	0.3
– T ₃ or T ₄	117 (76.5)	16 (88.9)	
Nodal stage before starting radiotherapy or chemoradiotherapy‡			
– Node negative	138 (95.2)	12 (70.6)	0.004
– Node positive	7 (4.8)	5 (29.4)	
Initial treatment type			
– Radiotherapy	92 (60.1)	8 (44.4)	0.2
– Chemoradiotherapy	61 (39.9)	10 (55.6)	
Tumour site at initial presentation			
– Glottis	101 (66.0)	11 (61.1)	0.6
– Supraglottis	15 (9.8)	1 (5.6)	
– Hypopharynx	37 (24.2)	6 (33.3)	
Presence of tracheostomy at recurrence			
– Yes	44 (28.8)	4 (22.2)	0.5
– No	109 (71.2)	13 (77.8)	
Imaging type			
– Computed tomography scan	102 (66.7)	13 (72.2)	0.6
– Positron emission tomography-computed tomography	51 (33.3)	5 (27.8)	

*n = 153; †n = 18; ‡data missing in nine patients

**Fig. 1.** Kaplan–Meier curve showing recurrence-free survival for the study cohort.

with the incidence varying between 5 per cent and 92 per cent in literature.^{16–18}

Concurrent neck dissection has been shown to increase the risk of post-operative complications in salvage settings.^{8–10}

A meta-analysis of factors affecting pharyngocutaneous fistula by Paydarfar and Birkmeyer demonstrated that neck dissection in salvage settings increases the risk of pharyngocutaneous fistula.¹¹ As per a Radiation Therapy Oncology Group analysis, neck dissection following RT increases the risk of both acute and late toxicity.¹⁹ This raises the concern about the safety of elective neck dissection in a salvage setting. Few studies have addressed this issue, and there have been conflicting results. The actual benefit of performing this procedure can only be shown after factoring in the rate of occult positivity, post-operative morbidity and regional failure.⁴

The incidence of occult metastasis following salvage laryngectomy varies from 3 per cent to 28 per cent.^{4–7,20–23} In our study, it was 10.6 per cent. This wide variation in occult positivity rate results from variability in characteristics of the study population, such as patients recruited at different time periods, differences in treatment protocols, variations in the stage of the disease, initial node positivity, site of involvement, type of imaging method used and so on. Koss *et al.* reported the highest rate of occult positivity to the extent of 28.3 per cent.²¹ In their study, N₀ status was determined mainly by clinical examination, and imaging was used when available. Additionally, they reported the occult metastasis rate in lateral and paratracheal basins. These factors probably resulted in higher neck positivity.

In our study, all patients were clinico-radiologically node negative as assessed by either CT scan or PET-CT scan, and we analysed occult metastasis rate in lateral neck alone. We routinely dissect paratracheal lymph node basins as laryngeal




Fig. 2. Kaplan-Meier curve for recurrence-free survival comparing neck dissection with pathological node positive versus pathological node negative patients. Five-year survival was 70.6 per cent and 68.2 per cent for pathological node positive and pathological node negative patients, respectively. Log rank test: $p=0.5$. pN = pathological node; pN₀ = pathological node negative; pN+ = pathological node positive

cancers may have central node positivity. These stringent criteria could have resulted in a low occult metastasis rate in our study. Studies which have used imaging methods such as CT scan to determine N₀ status have found a lower incidence of occult metastasis as was the case in our study.^{6,20} This is probably because of greater accuracy of CT scan in determining neck node status as compared with clinical examination alone. CT scans have a high negative predictive value (94 per cent to 97 per cent) in detecting occult nodal metastasis in salvage settings.^{20,24} Recently PET-CT has become the standard of care in planning neck dissection following chemo-RT in patients with advanced nodal disease.²⁵ In our study, the negative predictive value of PET-CT was 91 per cent and for CT scan was 88 per cent. Based on our results, we hypothesise that PET-CT is a useful imaging tool to plan neck dissection in salvage laryngectomy settings.

On analysing the factors predicting occult metastasis, we found that patients who had positive lymph node status before initiating definitive treatment (RT or chemo-RT) had microscopic disease in 42 per cent of cases. If we eliminate this group (i.e. cases who were node positive before starting RT) and study the occult metastasis rate in patients who were node negative at the time of RT, then incidence of occult metastasis is further reduced to 8 per cent. This is an important observation that may be helpful in selecting cases for elective neck dissection at the time of salvage laryngectomy. The same observation has been reported by Amit *et al.*⁵

There is less clarity regarding unilateral versus bilateral neck dissection during salvage laryngectomy. Yao *et al.* recommended bilateral neck dissection, based on two patients who had contralateral or bilateral pathological disease.²³ It must be kept in mind that the type of imaging that was used for these patients was not specified. Wax and Touma

recommended ipsilateral dissection for glottis cancers and bilateral neck dissection for supraglottic cancers in view of a 15 per cent bilateral neck metastasis rate.⁴ A recent meta-analysis by Lin *et al.* tried to find out whether unilateral neck dissection was as efficacious as bilateral neck dissection but did not reach any consensus due to lack of data in this regard.²⁶ Neck dissection, particularly bilateral neck dissection, is considered a risk factor for post-operative complications.^{6,11,27} The majority of patients (94.7 per cent) in our study had bilateral neck dissection, which could be a reason for the high incidence of pharyngocutaneous fistula (41.3 per cent). This is in accordance with literature where patients undergoing neck dissection had high post-operative complications when compared with patients who did not undergo neck dissection.^{6,27,28} In our study, only one patient had bilateral neck disease. Others have also found either no or very low rates of contralateral node positivity.^{5,6,20,28} In view of a bilateral metastatic rate of 0.5 per cent, we do not recommend bilateral neck dissection unless the tumour grossly involves both sides of the larynx.

Some authors have found advanced stage to be an important determinant for neck dissection and have recommended neck dissection in this group.^{4,23,29} We did not find stage as a determinant for positive occult metastasis rate.

The proponents of elective neck dissection argue that the procedure eliminates occult metastasis and the resulting regional failure.⁴ Hilly *et al.* devised a decision analysis model to determine the need for elective neck dissection following salvage laryngectomy.³⁰ Based on their model, they recommended neck dissection only when the isolated regional recurrence rate exceeds 20 per cent. However, the isolated regional failure rate following salvage laryngectomy varies from 0–8 per cent and is much lower than the rate of observed occult metastasis.^{4,20,29}

Local recurrence remains the most common type of relapse following salvage laryngectomy. In a study by Deganello *et al.*, the incidence of local recurrence was 31 per cent, and only 2.8 per cent of patients had regional relapse.²⁷ As per the existing literature, the regional failure rate in the dissected neck is either the same or worse than in the non-dissected neck. Elective neck dissection during salvage laryngectomy is not effective in reducing the rate of regional relapse.^{6,27–29}

- Management of the node negative neck during salvage laryngectomy is debatable
- Neck dissection in salvage settings increases the risk of complications
- Incidence of occult metastasis following salvage laryngectomy is low
- Patients who are node negative at initial treatment and remain clinico-radiologically node negative at salvage laryngectomy can be observed
- Patients who are node positive before radiotherapy should be offered elective neck dissection even if they are node negative at the time of salvage laryngectomy
- Unilateral neck dissection is adequate unless the tumour grossly involves both sides

The majority of the studies have found no survival advantage with the addition of elective neck dissection in salvage laryngectomy.^{21,23,28} Some have even reported poorer outcomes following elective neck dissection. In a large study by Li *et al.*, the wait and watch group had better overall survival than the patients who received any kind of treatment to the neck, confirming the safety of the wait and watch approach.³¹ We compared the outcomes between pathological node positive and pathological node negative patients. Presence of occult

metastasis was not associated with inferior survival outcome in our study. Pezier *et al.* have reported similar findings.⁷

The limitation of our study is that it is retrospective in nature. Since the policy of our institution is to do bilateral neck dissection, we did not have a comparative group where neck dissection was not performed. However, it overcomes many limitations mentioned in a recent systemic review.¹² Our study is a single institution study with a large number of patients, with robust data collection. Unlike many studies, all patients underwent either CT or PET-CT before being labelled as node negative. We investigated the role of several factors in predicting occult metastasis, which was not possible in a recently published systemic review and meta-analysis due to inclusion of studies with small patient numbers with considerable heterogeneity.¹²

Conclusion

Patients who are clinico-radiologically node negative at the time of salvage laryngectomy have a low rate of occult metastasis. Risk of occult metastasis is high in patients who have node positive disease before starting RT. This group should be offered elective neck dissection. Unilateral neck dissection is adequate unless there is gross bilateral disease. With the use of excellent modern imaging modalities like PET-CT, it is possible to identify patients who are at low risk of harbouring occult neck node metastasis. Morbidity of neck dissection can be avoided in a select group of patients who are node negative before starting the RT or chemo-RT and remain node negative at the time of salvage laryngectomy. Our study is a useful addition to the existing literature to provide insight into the management of the neck during salvage laryngectomy.

Competing interests. None declared

References

- Department of Veterans Affairs Laryngeal Cancer Study Group, Wolf GT, Fisher SG, Hong WK, Hillman R, Spaulding M *et al.* Induction chemotherapy plus radiation compared with surgery plus radiation in patients with advanced laryngeal cancer. *N Engl J Med* 1991;**324**:1685–90
- Lefebvre JL, Chevalier D, Luboinski B, Kirkpatrick A, Collette L, Sahnoud T. Larynx preservation in pyriform sinus cancer: preliminary results of a European Organization for Research and Treatment of Cancer phase III trial. EORTC Head and Neck Cancer Cooperative Group. *J Natl Cancer Inst* 1996;**88**:890–9
- Forastiere AA, Goepfert H, Maor M, Pajak TF, Weber R, Morrison W *et al.* Concurrent chemotherapy and radiotherapy for organ preservation in advanced laryngeal cancer. *N Engl J Med* 2003;**349**:2091–8
- Wax MK, Touma BJ. Management of the N0 neck during salvage laryngectomy. *Laryngoscope* 1999;**109**:4–7
- Amit M, Hilly O, Leider-Trejo L, Popovtzer A, Gutfeld O, Shvero J *et al.* The role of elective neck dissection in patients undergoing salvage laryngectomy. *Head Neck* 2013;**35**:1392–6
- Basheeth N, O'Leary G, Sheahan P. Elective neck dissection for no neck during salvage total laryngectomy: findings, complications, and oncological outcome. *JAMA Otolaryngol Head Neck Surg* 2013;**139**:790–6
- Pezier TF, Nixon IJ, Scotton W, oshi A, Guerrero-Urbano T, Oakley R *et al.* Should elective neck dissection be routinely performed in patients undergoing salvage total laryngectomy? *J Laryngol Otol* 2014;**128**:279–83
- Virtaniemi JA, Kumpulainen EJ, Hirvikoski PP, Johansson RT, Kosma VM. The incidence and etiology of postlaryngectomy pharyngocutaneous fistulae. *Head Neck* 2001;**23**:29–33
- McCombe AW, Jones AS. Radiotherapy and complications of laryngectomy. *J Laryngol Otol* 1993;**107**:130–2
- Natvig K, Boysen M, Tausjo J. Fistulae following laryngectomy in patients treated with irradiation. *J Laryngol Otol* 1993;**107**:1136–9
- Paydarfar JA, Birkmeyer NJ. Complications in head and neck surgery: a meta-analysis of postlaryngectomy pharyngocutaneous fistula. *Arch Otolaryngol Head Neck Surg* 2006;**132**:67–72
- Gross JH, Vila PM, Simon L, Rizvi ZH, Zenga J, Jackson RS *et al.* Elective neck dissection during salvage laryngectomy: a systematic review and meta-analysis. *Laryngoscope* 2020;**130**:899–906
- NCCN Guidelines Version 3.2019, Head and Neck cancers, National Comprehensive Cancer Care Clinical Practice Guidelines in Oncology (NCCN Guidelines). In: <http://www.NCCN.org> [5 January 2020]
- National Institute for Clinical excellence (NICE). Guidance on cancer services. Improving outcomes in Head and Neck Cancers - the manual. In: https://www.nice.org.uk/guidance/csg6/resources/improving_outcomes_in_head_neck_cancer_update_pdf-773377597 [5 January 2020]
- Sharma, S, Chaukar, D, Laskar, S, Kapre, N, Deshmukh A, Pai P *et al.* Role of the pectoralis major myofascial flap in preventing pharyngocutaneous fistula following salvage laryngectomy. *J Laryngol Otol* 2016;**130**:860–4
- Sassler AM, Esclamado RM, Wolf GT. Surgery after organ preservation therapy. Analysis of wound complications. *Arch Otolaryngol Head Neck Surg* 1995;**121**:162–5
- Putten L, Bree R, Doornaert PA, Buter J, Eerenstein SE, Rietveld DH *et al.* Salvage surgery in post-chemoradiation laryngeal and hypopharyngeal carcinoma: outcome and review. *Acta Otorhinolaryngol Ital* 2015;**35**:162–72
- Sewnaik A, Keereweer S, Al-Mamgani A, Baatenburg de Jong RJ, Wieringa MH, Meeuwis CA *et al.* High complication risk of salvage surgery after chemoradiation failures. *Acta Otolaryngol* 2012;**132**:96–100
- Machtay M, Moughan J, Trotti A, Garden AS, Weber RS, Cooper JS *et al.* Factors associated with severe late toxicity after concurrent chemoradiation for locally advanced head and neck cancer: an RTOG analysis. *J Clin Oncol* 2008;**26**:3582–9
- Farrag TY, Lin FR, Cummings CW, Koch WM, Flint PW, Califano JA *et al.* Neck management in patients undergoing postradiotherapy salvage laryngeal surgery for recurrent/persistent laryngeal cancer. *Laryngoscope* 2006;**116**:1864–6
- Koss SL, Russell MD, Leem TH, Schiff BA, Smith RV. Occult nodal disease in patients with failed laryngeal preservation undergoing surgical salvage. *Laryngoscope* 2014;**124**:421–8
- Birkeland AC, Rosko AJ, Issa MR, Shuman AG, Prince ME, Wolf GT *et al.* Occult Nodal Disease Prevalence and Distribution in Recurrent Laryngeal Cancer Requiring Salvage Laryngectomy. *Otolaryngol Head Neck Surg* 2016;**154**:473–9
- Yao M, Roebuck JC, Holsinger FC, Myers JN. Elective neck dissection during salvage laryngectomy. *Am J Otolaryngol* 2005;**26**:388–92
- Ojiri H, Mendenhall WM, Stringer SP, Johnson PL, Mancuso AA. Post-RT CT results as a predictive model for the necessity of planned post-RT neck dissection in patients with cervical metastatic disease from squamous cell carcinoma. *Int J Radiat Oncol Biol Phys* 2002;**52**:420–8
- Mehanna H, Wong WL, McConkey CC, Rahman JK, Robinson M, Hartley AG *et al.* PET-NECK Trial Management Group. PET-CT surveillance versus neck dissection in advanced head and neck cancer. *N Engl J Med* 2016;**374**:1444–54
- Lin C, Puram SV, Bulbul MG, Sethi RK, Rocco JW, Old MO *et al.* Elective neck dissection for salvage laryngectomy: a systematic review and meta-analysis. *Oral Oncol* 2019;**96**:97–104
- Deganello A, Meccariello G, Bini B, Paia F, Santoro R, Mannelli G *et al.* Is elective neck dissection necessary in cases of laryngeal recurrence after previous radiotherapy for early glottic cancer? *J Laryngol Otol* 2014;**128**:1089–94
- Bohannon IA, Desmond RA, Clemons L, Magnuson JS, Carroll WR, Rosenthal EL. Management of the N0 neck in recurrent laryngeal squamous cell carcinoma. *Laryngoscope* 2010;**120**:58–61
- Hilly O, Gil Z, Goldhaber D, Amit M, Biadsee A, Popovtzer A *et al.* Elective neck dissection during salvage total laryngectomy—a beneficial prognostic effect in locally advanced recurrent tumours. *Clin Otolaryngol* 2015;**40**:9–15
- Hilly O, Stern S, Horowitz E, Leshno M, Feinmesser R. Is there a role for elective neck dissection with salvage laryngectomy? A decision-analysis model. *Laryngoscope* 2013;**123**:2706–11
- Li PJ, Hu WH, Jin T. Management of the N0 neck in recurrent laryngeal squamous cell carcinoma. *Mol Clin Oncol* 2016;**4**:70–6